

Adenosine Deaminase Inhibitors. Synthesis and Biological Evaluation of C1' and Nor-C1' Derivatives of (+)-erythro-9-(2(*S*)-Hydroxy-3(*R*)-nonyl)adenine[†]

Geraldine C. B. Harriman, Anne F. Poirot, and Elie Abushanab*

Department of Medicinal Chemistry, University of Rhode Island, Kingston, Rhode Island 02881

Rose Marie Midgett and Johanna D. Stoeckler

Division of Biology and Medicine, Brown University, Providence, Rhode Island 02912

Received February 10, 1992

The synthesis of various chiral derivatives of (+)-erythro-9-(2-hydroxy-3-nonyl)adenine, (+)-EHNA, from (2*S*,3*R*)-3-amino-1,2-*O*-isopropylidene-1,2-nonanediol by condensation with 5-amino-4,6-dichloropyrimidine is described. The compounds synthesized were C1'- and nor-C1'-(+)-EHNA derivatives. When tested with calf spleen ADA, C1'-OH- and nor-C1'-(+)-EHNA had comparable inhibitory activity that was 1 order of magnitude lower than that of (+)-EHNA. Potency was reduced further in nor-C1' derivatives.

Introduction

Adenosine deaminase (ADA, adenosine aminohydrolase, EC 3.5.4.4) is a catabolic enzyme of the purine catabolic pathway that hydrolyzes adenosine and 2'-deoxyadenosine to inosine and 2'-deoxyinosine, respectively. ADA has broad substrate specificity and deaminates a variety of adenosine analogs such as adenine arabinoside (*ara*-A), 8-azaadenosine, formycin,¹ and 2',3'-dideoxyadenosine (ddA).² Deamination may inactivate adenosine analogs and limit their utility for antiviral and cancer chemotherapy. Additionally, ADA deficiency is associated with a severe combined immunodeficiency disease that is caused by lymphotoxic accumulations of 2'-deoxyadenosine and deoxy-ATP.^{3,4} Thus, potent ADA inhibitors may be used in two ways: (1) to prevent the inactivation of chemotherapeutic adenosine analogs, thereby favoring their direct conversion to biologically active nucleotides, and (2) to treat lymphoproliferative disorders⁵⁻⁸ by mimicking ADA deficiency.

The most potent inhibitors of ADA are coformycin (CF), 2'-deoxycoformycin (dCF, Pentostatin), and erythro-9-

(2-hydroxy-3-nonyl)adenine (EHNA). The former two naturally occurring nucleosides are base-modified transition-state analogs classified as tight-binding inhibitors with K_i values of 1×10^{-11} M (CF) and 2.5×10^{-12} M (dCF).^{1,9} EHNA, on the other hand, is a synthetic semitight binding inhibitor ($K_i = 4 \times 10^{-9}$ M) that was prepared as a racemate by Schaeffer and Schwender.¹⁰

The greater potency of dCF and the report that EHNA might have an adverse effect on purine metabolism¹¹ led to the selection of dCF for clinical studies. Despite promising clinical efficacy,⁵⁻⁸ dCF causes a high incidence of serious and unpredictable toxicities involving the immune system, kidney, liver, and central nervous system.^{12,13} Inhibition of intracellular ADA by dCF is nearly irreversible, with synthesis of new enzyme required for recovery from its effects.¹⁴ It seems likely that toxicities of dCF can be ascribed to this prolonged inhibition of ADA. (\pm)-EHNA, which is rapidly metabolized^{15,16} and has a shorter duration of action, allows faster recovery of the enzyme.

Racemic EHNA was used in early biologic and metabolic studies where enhancement of the activity of *ara*-A was observed.^{17,18} The synthesis, by our group¹⁹ and others,²⁰ of chiral isomers of EHNA allowed identification of the

[†] Taken, in part, from a Ph.D. Dissertation, G. C. B. Harriman, The University of Rhode Island, 1991.

(1) Agarwal, R. P.; Cha, S.; Crabtree, G. W.; Parks, R. E., Jr. Coformycin and Deoxycoformycin: Tight-Binding Inhibitors of Adenosine Deaminase. In *Chemistry and Biology of Nucleosides and Nucleotides*; Harmon, R. E., Robins, R. K., Townsend, L. B., Eds.; Academic Press: New York, 1978; pp 159-197.

(2) Haertle, T.; Carrera, C. J.; Wasson, D. B.; Sowers, L. C.; Richman, D. D.; Carson, D. A. Metabolism and Anti-human Immunodeficiency Virus-1 Activity of 2-Halo-2',3'-dideoxyadenosine Derivatives. *J. Biol. Chem.* 1988, 263, 5870-5875.

(3) Giblett, E. R.; Anderson, J. E.; Cohen, F.; Pollara, B.; Meuwissen, J. H. Adenosine-deaminase Deficiency in Two Patients with Severely Impaired Cellular Immunity. *Lancet* 1972, 2, 1067-1069.

(4) Hirschhorn, R.; Roegner, V.; Rubinstein, A.; Papageorgiou, P. Plasma Deoxyadenosine, Adenosine, and Erythrocyte DeoxyATP are Elevated at Birth in an Adenosine Deaminase-deficient Child. *J. Clin. Invest.* 1980, 65, 768-771.

(5) Grever, M. R.; Siaw, M. F. E.; Jacob, W. F.; Niedhart, J. A.; Miser, J. S.; Coleman, M. S.; Hutton, J. J.; Balcerzak, S. P. The Biochemical and Clinical Consequences of 2'-Deoxycoformycin in Refractory Lymphoproliferative Malignancy. *Blood* 1981, 57, 406-417.

(6) Grever, M. R.; Wilson, H. E.; Kraut, E. H.; Niedhart, J. A.; Balcerzak, S. P. Deoxycoformycin in the Treatment of Refractory Chronic Lymphocytic Leukemia. *Proc. Am. Soc. Clin. Oncol.* 1981, 22, 487.

(7) Spiers, A. S. D.; Moore, D.; Cassileth, P. A.; Harrington, D. P.; Cummings, F. J.; Nieman, R. S.; Bennett, J. M.; O'Connell, M. J. Remissions in Hairy-cell Leukemia with Pentostatin (2'-Deoxycoformycin). *N. Engl. J. Med.* 1987, 316, 825-830.

(8) Johnston, J. B.; Eisenhauer, E.; Corbett, W. E. N.; Scott, J. G.; Zaentz, S. D. Efficacy of 2'-Deoxycoformycin in Hairy-cell Leukemia: A Study of the National Cancer Institute of Canada Clinical Trials. *J. Natl. Cancer Inst.* 1988, 80, 765-769.

(9) Agarwal, R. P.; Spector, T.; Parks, R. E., Jr. Tight-Binding Inhibitors-IV. Inhibition of Adenosine Deaminases by Various Inhibitors. *Biochem. Pharmacol.* 1977, 26, 359-367.

(10) Schaeffer, H. J.; Schwender, C. F. Enzyme Inhibitors: XXVI. Bridging Hydrophobic and Hydrophilic Regions on Adenosine Deaminase with Some 9-(2-Hydroxy-3-alkyl)adenines. *J. Med. Chem.* 1974, 17, 6-8.

(11) Henderson, J. F.; Brox, L.; Zombor, G.; Hunting, D.; Lomax, C. A. Specificity of Adenosine Deaminase Inhibitors. *Biochem. Pharmacol.* 1977, 26, 1967-1972.

(12) O'Dwyer, P. J.; Spiers, A. S. D.; Marsoni, S. Association of Severe Fatal Infections and Treatment with Pentostatin. *Cancer Treat. Rep.* 1986, 70, 117-1120.


(13) Gieger, J. D.; Lewis, J. L.; MacIntyre, C. J.; Nagy, J. I. Pharmacokinetics of 2'-Deoxycoformycin, an Inhibitor of Adenosine Deaminase, in the Rat. *Neuropharmacology* 1987, 26, 1383-1387.

(14) Agarwal, R. P. Recovery of 2'-Deoxycoformycin Inhibited Adenosine Deaminase of Mouse Erythrocytes and Leukemia L1210 in vivo. *Cancer Res.* 1979, 39, 1425-1427.

(15) Lambe, C. U.; Nelson, D. J. Pharmacokinetics of Inhibition of Adenosine Deaminase by erythro-9-(2-Hydroxy-3-nonyl)adenine in CBA Mice. *Biochem. Pharmacol.* 1982, 31, 535-539.

(16) McConnell, W. R.; El Dareer, S. M.; Hill, D. L. Metabolism and Disposition of erythro-9-(2-Hydroxy-3-nonyl)[¹⁴C]adenine in the Rhesus Monkey. *Drug Metab. Dispos.* 1980, 8, 5-7.


(17) Suling, W. J.; Rice, L. S.; Shannon, W. M. Effects of 2'-Deoxycoformycin and erythro-9-(2-Hydroxy-3-nonyl)adenine on Plasma Levels and Urinary Excretion of 9- β -D-Arabinofuranosyladenine in the Mouse. *Cancer Treat. Rep.* 1978, 62, 369-373.



(+)-2S,3R isomer as the most potent ADA inhibitor.²¹ (+)-EHNA proved to be the most active isomer in the enhancement of *ara*-A activity against human pancreatic and colon carcinomas.²² All four stereoisomers caused similar inhibition of purine biosynthesis in sarcoma 180 cells, but these effects were not pronounced below drug concentrations of 1×10^{-5} M.²³ The comparable anti-HSV activity of all four isomers suggests that the antiviral activity is not mediated through inhibition of ADA.²⁴ Recently, an apparent isoenzyme of ADA had been identified, ADA₂. Levels of ADA₂ are low, but are elevated in the plasma of patients with human immunodeficiency virus (HIV) infections.^{25,26} While dCF inhibits both isoenzymes, (+)-EHNA selectively inhibits the predominant ADA activity and has been used in the purification of ADA.²⁷

A shorter-acting ADA inhibitor could permit a controlled duration of chemotherapeutic efficacy while avoiding long-term enzyme inactivation. Attractive targets would be derivatives that are 1–2 orders of magnitude more potent than (+)-EHNA; such derivatives might produce enhanced, but still reversible, ADA inhibition. On the basis of his SAR studies, Schaeffer proposed that the high affinity of EHNA is derived from its ability to bridge the binding region for adenine and sites for the hexyl, hydroxyl, and 1'-methyl groups of the 9-substituent.²⁸ Subsequent attempts by others to modify the purine moiety of EHNA have met with variable success. When Baker et al. combined the nonyl chain of EHNA with 1,6-dihydro-6-

Scheme I



(hydroxymethyl)purine, they observed mutually interfering effects between these high-affinity moieties.²⁹ On the other hand, certain deaza^{30,31} as well as imidazo³² analogs have shown ADA inhibitory activity comparable to that of EHNA. A recent report on the X-ray crystal structure of ADA with a transition-state analog suggests that the exact mode of binding of ADA inhibitors may soon be revealed.³³ In the present SAR study, C1' and nor-C1' derivatives were prepared to explore the 1'-methyl binding pocket. Chirons readily available from L-ascorbic acid³⁴ facilitated the preparation of these compounds.

Chemistry

In our earlier work^{34,35} (+)-EHNA was synthesized starting from epoxide 1 through intermediates 2–4. In this approach C-4 in 1 became a methyl group (C1') and the nonyl chain was elaborated by *n*-pentylmagnesium chloride addition at C-1. Since the chirality of carbons 2 and 3 in 1 is identical, an inversion of this addition sequence would eventually lead to C1'-substituted (+)-EHNA. Thus, regiospecific opening of epoxide 1 with *n*-pentylmagnesium chloride furnished alcohol 5. Tosylation of 5 gave 6, which upon treatment with sodium azide furnished 7. This was quantitatively reduced to the key amine 8 (Scheme I).

Incorporation of amine 8 into an adenine and subsequent attempts to introduce other functionalities at C1' are

(18) Plunkett, W.; Cohen, S. S. Two Approaches That Increase the Activity of Analogs of Adenine Nucleosides in Animal Cells. *Cancer Res.* 1975, 35, 1547–1554.

(19) Bastian, G.; Bessodes, M.; Panzica, R. P.; Abushanab, E.; Chen, S.-F.; Stoeckler, J. D.; Parks, R. E., Jr. Adenosine Deaminase Inhibitors. Conversion of a Single Chiral Synthon into *erythro*- and *threo*-9-(2-Hydroxy-3-nonyl)adenines. *J. Med. Chem.* 1981, 24, 1383–1385.

(20) Baker, D. C.; Hawkins, L. D. Synthesis of Inhibitors of Adenosine Deaminase. A Total Synthesis of *erythro*-3-(Adenin-9-yl)-2-nanol and Its Isomers from Chiral Precursors. *J. Org. Chem.* 1982, 47, 2179–2184.

(21) Bessodes, M.; Bastian, G.; Abushanab, E.; Panzica, R. P.; Berman, S. F.; Marcaccio, E. J., Jr.; Chen, S.-F.; Stoeckler, J. D.; Parks, R. E., Jr. Effect of Chirality in *erythro*-9-(2-Hydroxy-3-nonyl)adenine (EHNA) on Adenosine Deaminase Inhibition. *Biochem. Pharmacol.* 1982, 31, 879–882.

(22) Chu, M. Y.; Chu, Abushanab, E.; Panzica, R. P.; Calabresi, P. C. Effects of the Chiral Isomers of *erythro*-9-(2-Hydroxy-3-nonyl)adenine on the Anti-neoplastic Activity of Adenosine Analogs Against Human Pancreatic DAN and Human Colon HCT-8. *Proc. AACR* 1988, 29, 1394.

(23) Chen, S.-F.; Stoeckler, J. D.; Choi, H.-S.; Burgess, F. W.; Marcaccio, E. J., Jr.; Steen, P. A.; Berman, S. F.; Parks, R. E., Jr.; Panzica, R. P.; Abushanab, E. Effects on the Chiral Isomers of *erythro*- and *threo*-9-(2-Hydroxy-3-nonyl)adenine on Pure Metabolism in Sarcoma 180 Cells. *Biochem. Pharmacol.* 1982, 31, 3955–3960.

(24) North, T. W.; O'Connor, L.; Abushanab, E.; Panzica, R. P. Effects of Chirality in 9-(2-Hydroxy-3-nonyl)adenine Upon DNA Synthesis in Herpes Simplex Virus-Infected Cells. *Biochem. Pharmacol.* 1983, 32, 3541–3546.

(25) Yokoyama, M. M.; Tsuboi, I. Adenosine Deaminase Isoenzymes and HIV/HTLV-1 Infections. *J. Natl. Cancer Inst.* 1988, 80, 698.

(26) Delia, S.; Mastroianni, C. M.; Massetti, A. P.; Turbessi, G.; Cirelli, A.; Cantania, S.; Vullo, V. Adenosine Deaminase and Acquired Immunodeficiency Syndrome (AIDS). *Clin. Chem.* 1987, 33, 1675.

(27) Niedzwicki, J. G.; Kouttab, N. M.; Mayer, K. H.; Carpenter, C. C. J.; Parks, R. E., Jr.; Abushanab, E.; Abernethy, D. R. Plasma Adenosine Deaminase: A Marker for Human Immunodeficiency Virus Infection. *J. Acquired Immunodef. Syndr.* 1991, 4, 178–181.

(28) Schaeffer, H. J. Design and Evaluation of Enzyme Inhibitors, Especially of Adenosine Deaminase. *Topics. Med. Chem.* 1972, 3, 1–24.

(29) Woo, P. W. K.; Baker, D. C. Inhibitors of Adenosine Deaminase. Studies in Combining High-Affinity Enzyme-Binding Structural Units. *erythro*-1,6-Dihydro-6-(hydroxymethyl)-9-(2-hydroxy-3-nonyl)purine and *erythro*-9-(2-Hydroxy-3-nonyl)purine. *J. Med. Chem.* 1982, 25, 603–605.

(30) Antonini, I.; Cristalli, G.; Franchetti, P.; Grifantini, M.; Martelli, S.; Lupidi, G.; Riva, F. Adenosine Deaminase Inhibitors: Synthesis of Deaza Analogues of *erythro*-9-(2-Hydroxy-3-nonyl)adenine. *J. Med. Chem.* 1984, 27, 274–278.

(31) Cristalli, G.; Franchetti, P.; Grafantini, M.; Vittori, S.; Lupidi, G.; Riva, F.; Bordoni, T.; Geroni, C.; Verini, M. A. Adenosine Deaminase Inhibitors. Synthesis and Biological Activity of Deaza Analogues of *erythro*-9-(2-Hydroxy-3-nonyl)adenine. *J. Med. Chem.* 1988, 31, 390–397.


(32) Cristalli, G.; Eleuteri, A.; Franchetti, P.; Grafantini, M.; Vittori, S.; Lupidi, G. Adenosine Deaminase Inhibitors. Synthesis and Structure-Activity Relationships of Imidazole Analogues of *erythro*-9-(2-Hydroxy-3-nonyl)adenine. *J. Med. Chem.* 1991, 34, 1187–1192.

(33) Wilson, D. K.; Rudolph, F. B.; Quiocho, F. A. Atomic Structure of Adenosine Deaminase Complexed with a Transition-State Analog: Understanding Catalysis and Immunodeficiency Mutations. *Science* 1991, 252, 1278–1284.

(34) Abushanab, E.; Bessodes, M.; Antonakis, K. Practical Enantioselective Synthesis of (+)-*erythro*-9-(2-Hydroxy-3-nonyl)adenine. *Tetrahedron Lett.* 1984, 25, 3841.

(35) Abushanab, E.; Vemishetti, P.; Leiby, R. W.; Singh, H. K.; Mikkilineni, A. B.; Wu, D. C.-J.; Saibaba, R.; Panzica, R. P. The Chemistry of L-Ascorbic and D-isoascorbic Acids. 1. The Preparation of Chiral Butanetriols and -tetrols. *J. Org. Chem.* 1988, 53, 2598–2602.

Scheme II



outlined in Scheme II. Condensation of 8 with 5-amino-4,6-dichloropyrimidine (ADCP) afforded 9 in good yield, which was converted to two 6-chloropurine derivatives 10 and 11. The former was formed by an acid-catalyzed exchange between the isopropylidene group and triethyl orthoformate. Amination of this mixture followed by hydrolysis of the protecting groups furnished 1'-hydroxy-(+)-EHNA (14) in near-quantitative yield.

Having obtained 14, our efforts were directed at preparing other 1'-substituted (+)-EHNA derivatives. Of special interest is the 1'-fluoro derivative whose biological activity may provide further information on binding in the methyl pocket. The introduction of a fluorine at C1' could be accomplished by fluoride-catalyzed opening of the epoxide function in 16. Thus selective tosylation of the primary hydroxyl group in 14 provided tosylate 15, which proved to be unstable on standing. It underwent intramolecular cyclization to give the tricyclic compound 17. This transformation took place in chloroform solution and was monitored by thin-layer chromatography. The ¹H NMR spectrum showed the disappearance of the tosylate methyl singlet at δ 2.41 ppm with simultaneous appearance of a methyl singlet at δ 2.31 ppm. Furthermore, the heterocyclic proton resonances at δ 7.88 and 8.15 ppm were shifted downfield to δ 8.22 and 8.44 ppm. Epoxide 16 proved to be equally unstable and could not be used for the introduction of fluorine at the C1' position.

In an attempt to reduce the nucleophilicity of the N-3 nitrogen, and thus prevent cyclization, N-6 benzoylation was undertaken. When 13 was treated with 1 equiv of benzoyl chloride in pyridine, the corresponding N-6 benzamide 18 was prepared and was hydrolyzed to diol 19. Surprisingly, conversion of diol 32 to a monotosylate could not be achieved, but mesylate 20 could be easily prepared. Unfortunately, this compound underwent cyclization to the tricyclic product 21 upon standing at room

Scheme III




Table I. Inhibition of Adenosine Deaminase

compound	K _i (μM)	compound	K _i (μM)
(+)-EHNA	2.0 × 10 ⁻³	23	1.6
14	5.6 × 10 ⁻²	25	6.0 × 10 ⁻¹
16	8.9 × 10 ⁻¹	26	3.2 × 10 ⁻¹
17	1.0	27	1.3
22	4.7 × 10 ⁻²		

temperature. Other attempts to introduce the fluorine at C1' included preparation of a fluorinated nonylamine which could be further elaborated into 1'-fluoro(+)-EHNA. Here again, intramolecular nucleophilic displacement of the fluorine atom did take place to form the corresponding cyclic intermediates.

The availability of diol 14 allowed the preparation of C1'-nor derivatives whose biological activity could provide further insight into the nature of the binding site. The modifications are also outlined in Scheme III. Sodium periodate cleavage of 14 followed by reductive workup furnished alcohol 22. This was converted to the fluoro derivative 23 upon treatment with (diethylamido)sulfur trifluoride (DAST).³⁶ On the other hand, tosylation of 14 gave 24, which upon treatment with *n*-Bu₄NBr, *N*-Bu₄NCl, and sodium azide furnished the bromo 25, chloro 26, and azido 27 derivatives, respectively.

Results and Discussion

Schaeffer and Schwender studied an extensive series of 9-alkyladenine derivatives to determine structural requirements for high affinity binding to ADA.¹⁰ Our study is an extension of this work to further probe the C2'-hydroxyl and the 1'-methyl binding sites. As shown in Table I, diminished potencies were evident in nor-(+)-EHNA derivatives. The deletion of the 1'-methyl group caused a 23-fold decrease in activity for 22. A comparable result was reported earlier for the racemate.³⁷ Derivatives of nor-(+)-EHNA, compounds 23, 25, 26, and 27 had inhibitory activities 7–34-fold lower than nor-(+)-EHNA (22) and a 160–800-fold lower than (+)-EHNA itself. The C1' hydroxy derivative 14 showed a 28-fold decrease in inhibitory activity. Thus, hydroxylation at C1' (25) and removal of the 1'-methyl group (22) had similar deleterious effects on activity. This seems to further substantiate the existence of an auxiliary methyl pocket and the importance of the hydroxyl group to binding. Interestingly, the rigid cyclic analog 17 showed higher affinity for the enzyme than the natural nucleoside substrates. The activity of 17

(36) Middleton, W. J. New Fluorinating Reagents. Dialkylaminosulfur Fluorides. *J. Org. Chem.* 1975, 40, 574–578.

(37) Schaeffer, H. J.; Schwender, C. F. Enzyme Inhibitors XXIV. Bridging Hydrophobic and Hydrophilic Regions on Adenosine Deaminase. *J. Pharm. Sci.* 1971, 60, 1204–1209.

was also comparable to that of epoxide 16, which converted slowly to 17 during testing procedures.

Experimental Section

Melting points were determined on a Büchi 535 melting point apparatus and are uncorrected. The ^1H NMR spectra were recorded on a Varian EM-390 or a Bruker AM-300 spectrometer. The chemical shifts are expressed in parts per million with respect to tetramethylsilane. Optical rotations were obtained with a Perkin-Elmer Model 141 digital readout polarimeter. All organic solutions were dried over anhydrous sodium sulfate, filtered, and evaporated to dryness under reduced pressure. Ratios of chromatography solvents are expressed in v/v. Silica gel (Davison, grade H, 230–425 mesh), suitable for flash column chromatography was purchased from Fisher Scientific. A Chromatotron (centrifugally accelerated, preparative thin-layer, radial chromatograph), Model 7924T was used to complete various separations as indicated. The 1.0- and 2.0-mm plates used were coated with silica gel PF254 containing CaSO_4 . Elemental analyses were performed by MHW Laboratories, Phoenix, AZ.

(2S,3S)-1,2-O-Isopropylidene-1,2,3-nonanetriol (5). A solution of 1 (0.11 g, 0.56 mmol) in ether (10 mL) was added to a cold (-78°C) ether solution containing *n*-pentylmagnesium chloride (0.126 mmol) and 0.1 mmol of lithium tetrachlorocuprate with stirring for 3 h. After the mixture warmed to room temperature, stirring was continued for 1 h. The reaction was quenched with a saturated solution of NH_4Cl (100 mL) and extracted with ether. Pure 5 (0.148 g, 96% yield) was obtained by silica gel flash chromatography eluting with EtOAc–hexanes (1:10): $[\alpha]_D^{25} -25.23^\circ$ (c 2.67, EtOH); ^1H NMR (CDCl_3) δ 0.87 (t, $J = 4.5$ Hz, 3 H), 1.03–1.61 (m, 16 H), 2.49–2.59 (d, $J = 4.5$ Hz, 1 H, D_2O exchangeable), 3.31–4.09 (m, 4 H). Anal. ($\text{C}_{12}\text{H}_{24}\text{O}_3$) C, H.

(2S,3S)-1,2-O-Isopropylidene-3-O-tosyl-1,2,3-nonanetriol (6). A solution of 5 (15.0 g, 0.069 mol) in CH_2Cl_2 (50 mL) was added to a stirred solution of *p*-toluenesulfonyl chloride (26.5 g, 0.131 mol), anhydrous pyridine (20 mL, 0.250 mol), and CH_2Cl_2 (100 mL), and stirring was continued at 50°C overnight. The solvents were removed, and the residue was dissolved in CH_2Cl_2 (100 mL) and washed with water (75 mL). The organic layer was dried and the solvent was removed to give a yellow viscous liquid. Purification by silica gel chromatography, eluting with EtOAc–hexanes (1:10), gave 21.0 g of pure 6 (84% yield): $[\alpha]_D^{25} -11.75^\circ$ (c 7.23, EtOH); ^1H NMR (CDCl_3) δ 0.83 (t, $J = 6$ Hz, 3 H), 1.00–1.78 (m, 16 H), 2.43 (s, 3 H), 3.43–4.72 (m, 4 H), 7.29 (d, $J = 7.5$ Hz, 2 H), 7.78 (d, $J = 7.5$ Hz, 2 H). Anal. ($\text{C}_{19}\text{H}_{30}\text{O}_5\text{S}$) C, H.

(2S,3R)-3-Azido-1,2-O-isopropylidene-1,2-nonanediol (7). Sodium azide (34.0 mg, 0.523 mmol) was added to a stirring solution of 6 (129 mg, 0.349 mmol) in anhydrous DMF (15 mL). After refluxing for 45 min, DMF was removed and pure 7 (80.0 mg, 96% yield) was obtained by silica gel flash chromatography eluting with EtOAc–hexanes (1:26): $[\alpha]_D^{25} +32.23^\circ$ (c 1.66, EtOH); ^1H NMR (CDCl_3) δ 0.88 (t, $J = 6$ Hz, 3 H), 1.00–1.69 (m, 16 H), 3.30–3.58 (m, 1 H), 3.76–4.16 (m, 3 H). Anal. ($\text{C}_{12}\text{H}_{23}\text{N}_3\text{O}_2$) C, H, N.

(2S,3R)-3-Amino-1,2-O-isopropylidene-1,2-nonanediol (8). To a stirred solution of lithium aluminum hydride (LAH, 4.00 g, 0.105 mol) in anhydrous ether (175 mL) was added dropwise a solution of 7 (5.20 g, 0.022 mol) in anhydrous ether (50 mL). The reaction mixture was then heated at reflux for 2 h and cooled to room temperature, and excess LAH was decomposed by the careful successive dropwise addition of water (4 mL), 15% NaOH (4 mL), and water (8 mL). Filtration, drying, and evaporation of the solvent gave a pure yellow liquid 8 (4.52 g, 98% yield): $[\alpha]_D^{25} -7.12^\circ$ (c 3.90, EtOH); ^1H NMR (CDCl_3) δ 0.81 (t, $J = 6$ Hz, 3 H), 1.00–1.60 (m, 18 H, 2 D_2O exchangeable), 2.72–2.95 (m, 1 H), 3.72–4.02 (m, 3 H). Anal. ($\text{C}_{12}\text{H}_{25}\text{NO}_2$) C, H, N.

5-Amino-6-chloro-4-[[1,2(S)-(isopropylidenedioxy)-3(R)-nonyl]amino]pyrimidine (9). 5-Amino-4,6-dichloropyrimidine (ADCP, 1.42 g, 8.65 mmol), *n*- Bu_3N (2.1 mL, 8.65 mmol), and 8 (1.69 g, 7.86 mmol) in anhydrous pentanol (50 mL) were heated at reflux for 30 h under N_2 atmosphere. Pentanol and *n*- Bu_3N were removed, and the residue was chromatographed over silica gel (EtOAc–hexanes 4:1) to give 9 (2.10 g, 56% yield): $[\alpha]_D^{25} +11.03^\circ$ (c 1.68, EtOH); ^1H NMR (CDCl_3) δ 0.80 (t, $J = 6$ Hz, 3 H), 1.06–1.90 (m, 16 H), 3.42–4.50 (m, 6 H, 2 D_2O exchangeable),

4.91–5.01 (d, $J = 11.4$ Hz, 1 H, D_2O exchangeable), 7.93 (s, 1 H). Anal. ($\text{C}_{18}\text{H}_{27}\text{ClN}_4\text{O}_2$) C, H, Cl, N.

6-Chloro-9-[1,2-(ethoxymethylenedioxy)-3(R)-nonyl]purine (10). 6-Chloro-9-[1,2(S)-(isopropylidenedioxy)-3(R)-nonyl]purine (11). An acidified (concentrated HCl, 0.05 mL) solution of 9 (2.59 g, 7.56 mmol) in triethyl orthoformate (75 mL) was stirred at room temperature for 24 h. The yellow oil obtained after removal of TEOF was purified by silica gel flash chromatography eluting with EtOAc–hexanes (50:3) to provide 10 (0.788 g, 28% yield) and 11 (1.54 g, 58% yield): Compound 10: ^1H NMR (CDCl_3) δ 0.74 (t, $J = 6$ Hz, 3 H), 0.93–1.60 (m, 11 H), 1.71–2.34 (m, 2 H), 3.49 and 3.51 (q, $J = 6$ Hz, 2 H), 3.73–4.21 (m, 2 H), 4.45–4.83 (m, 2 H), 5.72 and 5.78 (s, 1 H), 8.12 and 8.19 (s, 1 H), 8.68 (s, 1 H). Anal. for 22 ($\text{C}_{17}\text{H}_{25}\text{ClN}_4\text{O}_3$) C, H, Cl, N. Compound 11: $[\alpha]_D^{25} +30.46^\circ$ (c 1.65, EtOH); ^1H NMR (CDCl_3) δ 0.75 (t, $J = 6$ Hz, 3 H), 0.95–1.59 (m, 14 H), 1.84–2.29 (m, 2 H), 3.68–4.11 (m, 2 H), 4.33–4.75 (m, 2 H), 8.20 (s, 1 H), 8.65 (s, 1 H). Anal. for 11 ($\text{C}_{17}\text{H}_{25}\text{ClN}_4\text{O}_2$) C, H, Cl, N.

9-[1,2(S)-(isopropylidenedioxy)-3(R)-nonyl]adenine (13). Compound 11 (87.0 mg, 0.247 mmol) was dissolved in liquid ammonia (5 mL) and heated at 90°C in a steel bomb for 24 h. After cooling, excess ammonia was allowed to evaporate. The residue was taken up in CH_2Cl_2 (5 mL) and washed with water (2 mL). The organic layer was dried and pure 13 (76 mg, 90% yield) was obtained as a white solid from hexanes: mp 113°C ; $[\alpha]_D^{25} +7.83^\circ$ (c 1.04, EtOH); ^1H NMR (CDCl_3) δ 0.69–1.54 (m, 17 H), 1.87–2.24 (m, 2 H), 3.59–4.07 (m, 2 H), 4.34–4.68 (m, 2 H), 6.52–6.80 (bs, 2 H, D_2O exchangeable), 7.86 (s, 1 H), 8.28 (s, 1 H). Anal. ($\text{C}_{17}\text{H}_{27}\text{N}_5\text{O}_2$) C, H, N.

9-[1,2(S)-Dihydroxy-3(R)-nonyl]adenine (14). A solution of 10 and 11 (8.25 g, 10:11, 3:10) in 88% formic acid (25 mL) was stirred at room temperature overnight. The acid was removed and the pH of the solution was adjusted to 8 with 2 N NH_4OH (4 mL). The solution was stirred overnight and filtered through Celite, and the solvent was removed. Crystallization of the product with EtOAc gave pure 14 (7.72 g, 98% yield) as a white powder: mp 161.0°C ; $[\alpha]_D^{25} +36.59^\circ$ (c 1.06, EtOH); ^1H NMR ($\text{DMSO}-d_6$) δ 0.65–1.36 (m, 11 H), 1.80–2.22 (m, 2 H), 3.14–3.52 (m, 3 H, 1 H, D_2O exchangeable), 4.43–4.83 (m, 2 H, 1 H, D_2O exchangeable), 5.31 (d, $J = 6$ Hz, 1 H, D_2O exchangeable), 7.10–7.32 (bs, 2 H, D_2O exchangeable), 8.16 (s, 2 H). Anal. ($\text{C}_{14}\text{H}_{23}\text{N}_5\text{O}_2$) C, H, N.

9-[2(S)-Hydroxy-1-(tosyloxy)-3(R)-nonyl]adenine (15). To an ice-cold solution of 14 (1.10 g, 3.75 mmol) in anhydrous pyridine (25 mL) was added *p*-toluenesulfonyl chloride (786 mg, 4.12 mmol). After stirring for 2 h, the solvent was removed and intermediate 15 (500 mg, 30% yield) was isolated by silica gel flash chromatography eluting with EtOAc: ^1H NMR (CDCl_3) δ 0.78 (t, $J = 6$ Hz, 3 H), 0.92–1.34 (m, 8 H), 1.50–2.19 (m, 2 H), 2.40 (s, 3 H), 3.89–4.59 (m, 5 H, 1 D_2O exchangeable), 6.20–6.44 (bs, 2 H, D_2O exchangeable), 7.49 (AA'BB' q, $J = 9$ Hz, 4 H), 7.80 (s, 1 H), 8.14 (s, 1 H). The instability of 15 precluded obtaining correct elemental analysis.

9-[1,2(S)-Epoxy-3(R)-nonyl]adenine (16). A solution of 14 (140 mg, 0.47 mmol) and *n*- Bu_4NCl (13 mg, 0.05 mmol) in CH_2Cl_2 (4 mL) was stirred at room temperature for 15 min. A solution of saturated NaOH (2 mL) was then added and stirring was continued for an additional 10 min. While the mixture was stirred vigorously, a solution of tosyl chloride (80 mg, 0.47 mmol) in CH_2Cl_2 (2 mL) was added. The reaction was stopped after 2 h, and the residue obtained after evaporation was placed on a silica gel column eluting with MeOH–EtOAc (3:97). The epoxide 16 was obtained (28.5 mg, 22% yield) as an oil: ^1H NMR (CDCl_3) δ 0.70 (t, $J = 6$ Hz, 3 H), 1.15 (bs, 8 H), 1.8–2.15 (m, 2 H), 2.6–2.7 (m, 2 H), 3.10–3.35 (m, 1 H), 4.10–4.44 (m, 1 H), 6.30 (bs, 2 H, D_2O exchangeable), 7.75 (s, 1 H), 8.20 (s, 1 H). The instability of the compound precluded obtaining elemental analysis.

10-Amino-5,6-dihydro-4(R)-hexyl-5(S)-hydroxy-4H-pyrimido[1,2,3-*cd*]purin-5-ium Tosylate (17). A solution of 15 (118 mg, 0.264 mmol) in CHCl_3 was heated at reflux for 20 h. The solvent was removed and the residue was crystallized from MeOH to give pure 17 as a white powder (86.4 mg, 73.2% yield): mp 198°C ; ^1H NMR ($\text{MeOH}-d_4$) δ 0.85 (t, $J = 6$ Hz, 3 H), 1.04–1.87 (m, 10 H), 2.31 (s, 3 H), 4.23–4.74 (m, 4 H), 7.10 (d, $J = 9$ Hz, 2 H), 7.60 (d, $J = 9$ Hz, 2 H), 8.22 (s, 1 H), 8.44 (s, 1 H). Anal. ($\text{C}_{21}\text{H}_{29}\text{N}_5\text{O}_2\text{S}$) C, H, N, S.

N-Benzoyl-9-[1,2(*S*)-O-Isopropylidene-3(*R*)-nonyl]adenine (18). In a round-bottom flask (100-mL) protected from light, a solution of benzoyl chloride (1.65 mL, 61.8 mmol) in anhydrous pyridine (5.0 mL, 61.8 mmol) was added dropwise to a stirred solution of 13 (4.75 g, 14.2 mmol) in CH_2Cl_2 (25 mL). After 48 h the solvent was removed and pure 18 (3.91 g, 63% yield) was obtained by silica gel flash chromatography eluting with EtOAc-hexanes (4:5): $[\alpha]^{25}_{\text{D}} +30.28^\circ$ (c 1.66, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.61–1.49 (m, 17 H), 1.73–2.23 (m, 2 H), 3.61–4.01 (m, 2 H), 4.30–4.71 (m, 2 H), 7.25–7.55 (m, 3 H), 1 D_2O exchangeable), 7.80–8.10 (m, 3 H), 8.63 (s, 1 H), 9.25 (s, 1 H). Anal. ($\text{C}_{24}\text{H}_{31}\text{N}_5\text{O}_3$) C, H, N.

N-Benzoyl-9-[1,2(*S*)-dihydroxy-3(*R*)-nonyl]adenine (19). An acidified (concentrated HCl, 1 mL) solution of 18 (2.66 g, 6.07 mmol) in 95% EtOH (50 mL) was stirred at room temperature for 24 h. The solvent was removed, and pure 19 (1.87 g, 71% yield) and 14 (0.776 g, 22% yield) were obtained by silica gel flash chromatography eluting with EtOAc: mp 158.7–160.0 $^\circ\text{C}$; $[\alpha]^{25}_{\text{D}} +28.37^\circ$ (c 1.47, EtOH); $^1\text{H NMR}$ ($\text{DMSO}-d_6$) δ 0.60–1.40 (m, 11 H), 1.77–2.24 (m, 2 H), 3.13–3.45 (m, 2 H), 1 D_2O exchangeable), 3.45–3.65 (m, 1 H), 4.55–4.85 (m, 2 H), 4.95–5.45 (bs, 1 H, D_2O exchangeable), 7.32–7.66 (m, 3 H), 7.85–8.13 (m, 2 H), 8.44 (s, 1 H), 8.65 (s, 1 H). Anal. ($\text{C}_{21}\text{H}_{27}\text{N}_5\text{O}_3$) C, H, N.

N-Benzoyl-9-[2(*S*)-hydroxy-1-[(methanesulfonyl)oxy]-3(*R*)-nonyl]adenine (20). Methanesulfonyl chloride (0.100 mL, 1.36 mmol) was added dropwise to a stirred solution of 19 (541 mg, 1.36 mmol) and anhydrous pyridine (0.5 mL, 38.7 mmol) in anhydrous CH_2Cl_2 (25 mL) at 0 $^\circ\text{C}$. The reaction mixture was stirred at 0 $^\circ\text{C}$ for 3 h and at room temperature overnight. The solvent was removed and 20 (0.362 g, 56% yield) was obtained by silica gel flash chromatography eluting with EtOAc-hexanes (9:1): $^1\text{H NMR}$ (CDCl_3) δ 0.45–1.41 (m, 11 H), 1.67–2.31 (m, 2 H), 3.10 (s, 3 H), 3.74–4.33 (m, 3 H), 4.53–4.82 (m, 2 H), 1 D_2O exchangeable), 7.29–7.68 (m, 3 H), 7.81–8.12 (m, 3 H), 8.50 (s, 1 H), 8.65 (s, 1 H).

10-Benzamido-5,6-dihydro-4(*R*)-hexyl-5(*S*)-hydroxy-4H-pyrimido[1,2,3-*cd*]purin-5-ium Mesylate (21). Compound 20 (58.0 mg, 0.122 mmol) in anhydrous benzene (10 mL) was heated at reflux for 1.5 h. The solvent was removed to give pure 21 (55.0 mg, 95% yield): $[\alpha]^{25}_{\text{D}} +29.75^\circ$ (c 0.69, EtOH); $^1\text{H NMR}$ ($\text{DMSO}-d_6$) δ 0.85 (t, $J = 6$ Hz, 3 H), 1.00–1.90 (m, 10 H), 2.32 (s, 3 H), 4.40–4.90 (m, 5 H), 1 D_2O exchangeable), 7.35–7.70 (m, 3 H), 7.80–8.13 (m, 2 H), 8.79 (s, 1 H), 9.27 (s, 1 H), 11.88–12.58 (bs, 1 H, D_2O exchangeable). Anal. ($\text{C}_{22}\text{H}_{26}\text{N}_6\text{SO}_6$) C, H, N, S.

9-[1-Hydroxy-2(*R*)-octyl]adenine (22). To a stirred solution of 14 (947 mg, 3.23 mmol), $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ (70.0 mg, 0.507 mmol) in $\text{MeOH}-\text{H}_2\text{O}$ (3:1, 25 mL) was added NaIO_4 (759 mg, 3.55 mmol). The reaction was stirred at room temperature until a heavy white precipitate formed (0.5 h). The reaction mixture was filtered and the filtrate was concentrated to a volume of 15 mL. The solution was then cooled to 0 $^\circ\text{C}$ and NaBH_4 (0.500 g, 0.0132 mol) was added. The reaction was warmed to room temperature and was stirred for an additional 2 h. The solvent was then removed, and the white crystalline product was washed with small portions of water (3×2 mL) to give 22 (0.760 g, 90% yield): mp 150–154 $^\circ\text{C}$; $[\alpha]^{25}_{\text{D}} +17.75^\circ$ (c 1.88, EtOH); $^1\text{H NMR}$ ($\text{DMSO}-d_6$) δ 0.65–1.55 (m, 11 H), 1.86–2.20 (m, 2 H), 3.61–4.05 (m, 2 H), 4.38–4.77 (m, 1 H), 4.99–5.22 (bs, 1 H, D_2O exchangeable), 7.09–7.35 (bs, 2 H, D_2O exchangeable), 8.30 (s, 1 H), 8.31 (s, 1 H). Anal. ($\text{C}_{13}\text{H}_{21}\text{N}_5\text{O}$) C, H, N.

9-[1-Fluoro-2(*R*)-octyl]adenine (23). To a cold (-78 $^\circ\text{C}$) solution of (diethylamido)sulfur trifluoride (DAST, 0.15 mL, 1.14 mmol) in CH_2Cl_2 (3 mL) was slowly added a solution of 22 (113 mg, 0.430 mmol) in anhydrous CH_2Cl_2 (3 mL), and stirring was continued for 1 h. After stirring for 1 h at room temperature, excess DAST was decomposed by the careful addition of cold saturated solution of NaHCO_3 (5 mL). The crude product was extracted with CH_2Cl_2 (10 mL) and pure 23 (44.0 mg, 38% yield)

as a white powder was obtained by silica gel flash chromatography eluting with EtOAc: mp 117–118 $^\circ\text{C}$; $[\alpha]^{25}_{\text{D}} +26.77^\circ$ (c 0.79, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.85 (t, $J = 6.6$ Hz, 3 H), 1.23–1.33 (m, 8 H), 1.97–2.14 (m, 2 H), 4.58–4.96 (m, 3 H), 6.40 (bs, 2 H, D_2O exchangeable), 7.92 (s, 1 H), 8.35 (s, 1 H). Anal. ($\text{C}_{13}\text{H}_{20}\text{N}_5\text{F}$) C, H, F, N.

9-[1-(Tosyloxy)-2(*R*)-octyl]adenine (24). This compound was prepared from 19 as indicated previously for 6 in 96% yield: $[\alpha]^{25}_{\text{D}} +5.32^\circ$ (c 2.57, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.80 (t, $J = 6$ Hz, 3 H), 1.03–1.45 (m, 8 H), 1.83–2.18 (m, 2 H), 2.40 (s, 3 H), 4.22–4.84 (m, 3 H), 6.07–6.33 (bs, 2 H, D_2O exchangeable), 7.10 (d, $J = 7.5$ Hz, 2 H), 7.50 (d, $J = 7.5$ Hz, 2 H), 7.73 (s, 1 H), 8.14 (s, 1 H). Anal. ($\text{C}_{20}\text{H}_{27}\text{N}_5\text{O}_3\text{S}$) C, H, N.

9-[1-Bromo-2(*R*)-octyl]adenine (25). An anhydrous benzene solution (25 mL) of *n*-Bu₄NBr (0.325 g, 1.01 mmol) and 24 (85.0 mg, 0.204 mmol) was heated at reflux for 2.5 h. The solvent was removed and pure 25 (51.9 mg, 78% yield) as a white powder was obtained by silica gel flash chromatography eluting with EtOAc: mp 85–88 $^\circ\text{C}$; $[\alpha]^{27}_{\text{D}} +13.16^\circ$ (c 0.77, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.80 (t, $J = 6$ Hz, 3 H), 1.01–1.47 (m, 8 H), 1.89–2.32 (m, 2 H), 3.63–4.19 (m, 2 H), 4.49–4.87 (m, 1 H), 6.00–6.29 (bs, 2 H, D_2O exchangeable), 7.80 (s, 1 H), 8.29 (s, 1 H). Anal. ($\text{C}_{13}\text{H}_{20}\text{BrN}_5$) C, H, Br, N.

9-[1-Chloro-2(*R*)-octyl]adenine (26). An anhydrous benzene solution (25 mL) of *n*-Bu₄NCl (0.22 g, 0.791 mmol) and 24 (0.11 g, 0.263 mmol) was heated at reflux for 2 h. The solvent was removed and pure 26 (51.2 mg, 69% yield) as light yellow crystals was obtained by silica gel chromatography eluting with EtOAc: mp 78–79 $^\circ\text{C}$; $[\alpha]^{27}_{\text{D}} +21.26^\circ$ (c 1.93, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.79 (t, $J = 6$ Hz, 3 H), 0.96–1.48 (m, 8 H), 2.82–2.26 (m, 2 H), 3.72–4.20 (m, 2 H), 4.52–4.84 (m, 1 H), 6.40–6.66 (bs, 2 H, D_2O exchangeable), 7.81 (s, 1 H), 8.27 (s, 1 H). Anal. ($\text{C}_{13}\text{H}_{20}\text{ClN}_5$) C, H, Cl, N.

9-[1-Azido-2(*R*)-octyl]adenine (27). A solution of 24 (0.11 g, 0.262 mmol), NaN_3 (35.0 mg, 0.538 mmol), and pyridine (0.03 mL, 0.538 mmol) in DMF (13 mL) was heated at reflux for 15 h. The solvent was removed and the resulting mixture was dissolved in CH_2Cl_2 (15 mL) and washed with water (10 mL). The solvent was removed and pure 27 (44.5 mg, 59% yield) was obtained by silica gel chromatography eluting with EtOAc: $[\alpha]^{25}_{\text{D}} +21.74^\circ$ (c 1.38, EtOH); $^1\text{H NMR}$ (CDCl_3) δ 0.74 (t, $J = 6$ Hz, 3 H), 1.02–1.50 (m, 8 H), 1.80–2.30 (m, 2 H), 3.58–4.13 (m, 2 H), 4.43–4.78 (m, 1 H), 6.33–6.65 (bs, 2 H, D_2O exchangeable), 7.85 (s, 1 H), 8.33 (s, 1 H). Anal. ($\text{C}_{13}\text{H}_{20}\text{N}_6$) C, H, N.

Biological Evaluation. The activity of calf intestinal ADA (Type III, Sigma Chemical Co.) was measured at 30 $^\circ\text{C}$ by direct spectrophotometric assays at 265 nm,^{38,39} using a Gilford Model 240 spectrophotometer. The 1-mL reaction mixtures contained 0.003 unit of enzyme, 10–80 μM adenosine, and 50 mM potassium phosphate, pH 7.4. Inhibition constants below 900 nM were determined from replots of the slopes of double-reciprocal plots at multiple inhibitor concentrations. Micromolar K_i values were calculated from apparent K_i values obtained from plots of $1/\text{velocity}$ vs inhibitor concentration by the equation, $K_i^{\text{app}} = K_i(1 + [\text{S}]/K_m)$. Inhibition was assumed to be competitive and adenosine concentration was fixed at 24 μM , approximately the K_m value.

Acknowledgment. The authors thank Drs. Robert E. Parks, Jr. and Chandra Vargeese for their valuable suggestions and helpful discussions. The partial financial support of NIH Grant CA 07340 is also acknowledged.

(38) Kalokar, H. M. Differential Spectrophotometry of Purine Compounds by Means of Specific Enzymes. III Studies of the Enzymes of Purine Metabolism. *J. Biol. Chem.* 1947, 167, 461–475.

(39) Agarwal, R. P.; Parks, R. E., Jr. Adenosine Deaminase from Human Erythrocytes. *Methods Enzymol.* 1978, 51, 502–507.