

Accepted Article

Title: Perfluoroaryl Azide-Staudinger Reaction: A Fast and Bioorthogonal Reaction

Authors: Madanodaya Sundhoro, Seaho Jeon, JaeHyeung Park, Olof Ramström, and Mingdi Yan

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.201705346 Angew. Chem. 10.1002/ange.201705346

Link to VoR: http://dx.doi.org/10.1002/anie.201705346 http://dx.doi.org/10.1002/ange.201705346

WILEY-VCH

COMMUNICATION

WILEY-VCH

Perfluoroaryl Azide–Staudinger Reaction: A Fast and Bioorthogonal Reaction

Madanodaya Sundhoro,^[a] Seaho Jeon,^[a] Jaehyeung Park,^[a] Olof Ramström,^{*[a][b]} Mingdi Yan^{*[a][b]}

Abstract: We report a fast Staudinger reaction between perfluoroaryl azides (PFAAs) and aryl phosphines, occurring readily under ambient conditions. A rate constant as high as 18 M⁻¹ s⁻¹ was obtained between methyl 4-azido-2,3,5,6-tetrafluorobenzoate and methyl 2- (diphenylphosphanyl)benzoate in CD₃CN/D₂O. In addition, the iminophosphorane product was stable toward hydrolysis and aza-phosphonium ylide reactions. The PFAA-Staudinger reaction proved to be an excellent bioothorgonal reaction. PFAA-derivatized mannose and galactose were successfully transformed into cell surface glycans and efficiently labeled with phosphine-derivatized fluorescent bovine serum albumin.

The *Staudinger reaction*, the formation of an iminophosphorane from an azide and a phosphine,¹ proceeds through nucleophilic attack of the phosphine on the distal nitrogen of the azide to form the phosphazide intermediate, which decomposes to the iminophosphorane upon release of N₂.^{2,3} When exposed to water, the iminophosphorane is readily hydrolyzed to the corresponding amine and phosphine oxide. The reaction has recently gained renewed interest, owing to the search for highly efficient bioorthogonal reactions.⁴

The pioneering work applying the Staudinger reaction as a bioorthogonal reaction involves the introduction of an electrophilic ester trap on the phenyl phosphine (Fig. 1A).⁵ In the resulting Staudinger ligation reaction, the nucleophilic iminophosphorane attacks the ester intramolecularly to form a cyclic intermediate, which, upon hydrolysis, gives the conjugated amide product.⁶ The reaction has been successfully applied to metabolic engineering, where azide-derivatized N-acetylmannosamine was converted to sialic acid derivatives, allowing the expression of azido groups on the cell surface. Subsequent Staudinger ligation with biotinderivatized phosphine followed by fluorescent avidin allowed the detection of cell surface glycans by flow cytometry. Although high background noise was often observed,⁷ the Staudinger ligation is considered superior for in vivo labelling compared to the strainpromoted alkyne-azide cycloaddition (SPAAC) reaction, where, for example, the alkynes could bind strongly to murine serum albumin and were sequestered from azide-labelled tissues.8

[a] Dr. M. Sundhoro, Dr. S. Jeon, Dr. J. Park, Prof. Dr. M. Yan, Prof. Dr. O. Ramström
 Department of Chemistry, University of Massachusetts Lowell, 1

University Ave., Lowell, MA 01854, United States E-mail: mingdi_yan@uml.edu; olof_ramstrom@uml.edu Prof. Dr. M. Yan, Prof. Dr. O. Ramström

[b] Prof. Dr. M. Yan, Prof. Dr. O. Ramström Department of Chemistry, KTH – Royal Institute of Technology, Teknikringen 30, S-10044 Stockholm, Sweden E-mail: mingdi@kth.se; ramstrom@kth.se

Supporting information for this article is given via a link at the end of the document.



Figure 1. A) Staudinger ligation. B) PFAA-Staudinger reaction leading to stable iminophosphorane.

Mechanistic studies on the Staudinger reaction between phenyl azides and triphenylphosphines revealed that the reaction rate can be enhanced by introducing electron-donating groups on the phosphine or electron-withdrawing groups on the phenyl azide.^{2,6} Since electron-rich phosphines are more susceptible to oxidation in the biological environment,⁸ the alternative of using electrondeficient azides would be a more logical solution. It was reported that iminophosphoranes formed from electron-deficient azides were more stable, and had lower hydrolysis rates.^{6,9-11} The P=N bond can be stabilized by steric hindrance.^{12,13} Encouraged by these findings, we envisaged efficient Staudinger reactions using perfluoroaryl azides (PFAAs). The F atoms lower the LUMO of the aryl azides, giving PFAAs unique reactivities towards, for example, enamines, thioacids and aldehydes.14-19 Here, we report that PFAAs undergo fast Staudinger reactions with aryl phosphines under ambient conditions to give stable iminophosphoranes (Fig. **1B**). In addition, the reaction showed excellent bioothorgonality and was successfully applied for cell surface labeling.

When an equimolar amount of PFAA **1a** and phosphine **2a** (**Fig. 2A**) were mixed in acetonitrile (100 mM) at room temperature, the solution immediately turned yellow. The color remained for 4 min and was followed by evolution of nitrogen gas. The product, iminophosphorane **3aa**, was obtained in quantitative yield (\geq 99%), and the structure was confirmed by single crystal x-ray crystallography (**Fig. 2B**).

The Staudinger reaction can proceed through first- or second-order kinetics depending on whether the rate-limiting step is unimolecular decomposition or bimolecular formation of the intermediate phosphazide.²⁰ In the present case, the reaction between **1a** and **2a** followed second order as can be seen from the kinetic analysis (**Fig. 2C**). The observed rate constant, 3.68 \pm 0.03 M⁻¹s⁻¹, is six times higher than that between 1-azido-4-nitrobenzene and triphenylphosphine (0.611 M⁻¹s⁻¹),² the fastest

WILEY-VCH

COMMUNICATION

Staudinger reaction reported, and 1940 times higher than that of the classic Staudinger ligation reaction (K_{obs} = 1.9 × 10⁻³ M⁻¹s⁻¹).⁶

The effect of solvent on the reaction rate was next investigated (**Table 1**). In general, the observed rate constant increased with the polarity of the solvent (**Entries 1-4**). Consistent with this observation is the rate acceleration with the addition of D₂O (**Entries 5-7**), where an observed rate constant of 18.3 M⁻¹s⁻¹ was obtained with 50% D₂O (v/v) (**Entry 7**). These results are in agreement with a polar transition state that is more stabilized in polar solvents.²



Figure 2. A) Reaction between **1a** and **2a**. B) X-ray single crystal structure of **3aa**. C) Kinetic analysis; conditions: $[1a]_0 = 2.5 \text{ mM}$, $[2a]_0 = 2.5 \text{ mM}$, CD₃CN, 25 °C; concentrations monitored by ¹H NMR with spectra recorded every 60 s; k_{obs} (2nd order): 3.68 ± 0.03 M⁻¹ s⁻¹.

 Table 1. Effect of solvent on reaction rate between PFAA 1a and phosphine
 2a.

Entry	Solvent	<i>k</i> _{obs} (M ⁻¹ s ⁻¹) ^a	Yield (%) ^b
1	CDCl ₃	0.83 ± 0.03	79
2	CD ₃ CN	3.68 ± 0.03	91
3	(CD ₃) ₂ CO	2.37 ± 0.03	91
4	CD ₃ OD	5.7 ± 0.1	95
5	CD ₃ CN/D ₂ O (5/1)	6.5 ± 0.1	97
6	CD ₃ OD/D ₂ O (5/1)	11.3 ± 0.3	96
7	CD ₃ CN/D ₂ O (1/1)	18.3 ± 0.7	97

[a] 2nd order; reactions followed with ¹H NMR by monitoring the methyl protons in PFAA **1a** over 30-45 min. [b] NMR yield at 30 min.

The scope of the PFAA structures was studied, and high reaction rates were observed for all PFAAs tested (**Table 2**). Furthermore, the more electron-withdrawing the substituent on the PFAA core, the higher the reaction rate is consistent with the kinetics of the Staudinger reaction.² In addition, the reactions proved efficient, and high conversions (> 90%) were achieved under ambient conditions in all cases. Hammett analysis performed on the PFAAs (**Graph S1**) showed a linear correlation

($R^2 = 0.99$) with a small, positive ρ value (0.43). These results suggest a buildup of negative charge and a small influence of the PFAA's *para*-substituent on the reaction rate. The reaction was repeated with 1-azido-4-nitrobenzene (**1b**), the non-fluorinated analog of compound **1d**. This reaction was not completed after 30 min (**Entry 2**, **Table 2**), and displayed a 32 times lower rate in comparison to the reaction with its perfluorinated analog **1d** (**Entry 4**).





1	1a	3.68 ± 0.03	91	
2	1b	0.153 ± 0.001	41	
3	1c	3.92 ± 0.07	95	
4	1d	4.97 ± 0.08	95	
5	1e	2.73 ± 0.06	93	
6	1f	0.91 ± 0.01	82	

[a] 2^{nd} order; reactions were followed with ^1H NMR over the course of 30 min. [b] NMR yield at 30 min.

The scope of the phosphine was next investigated (Table 3). Introducing an electron-withdrawing methyl ester on the phenyl group increased the rate slightly (Entry 1 vs. Entry 2). Substituting the phenyl moieties with pentafluorophenyl groups (2c) strongly impeded the reaction as no product was observed after 30 min (Entry 3). This is consistent with the classic Staudinger reaction in which the rate decreases when using an electron-poor phosphine.² When electron-donating groups were introduced (2d, 2e, 2f), the phosphine was immediately converted into the phosphazide (Fig. S16A-S18A) upon mixing with PFAA 1a. The decomposition of phosphazide was in this case the ratelimiting step, following first order kinetics (Fig. S16). The rate of phosphazide decomposition decreased with the electron donating ability of the aryl substituent (Entries 4-6). For the reaction with 2d, the isolated yield was lower in comparison to the triphenyl counterpart (63%) due to the production of oxidized phosphine.

The iminophosphorane products proved stable over time. When solutions of **3aa** or **3ab** in CD₃CN were stored under ambient conditions (**Fig. S6-S7**), or when 10% D₂O was added (**Fig. S8-S9**), the ¹H NMR spectra did not change during a course of 35 days. This demonstrates the high stability of the iminophosphorane towards hydrolysis. Compounds **3aa** or **3ab** were then mixed with an excess of CS₂ or *p*-anisaldehyde in CD₃CN. After 35 days, no new peaks were detected in the ¹H

WILEY-VCH

COMMUNICATION

NMR spectra (**Fig. S10-S13**), demonstrating the inability of these iminophosphoranes to undergo aza-Wittig reactions in consequence to the low nucleophilicity of the fluorinated iminophosphoranes.

Table 3. Scope of phosphines



Entry	Phosphine	<i>k</i> _{1,obs} (M ⁻¹ s ⁻¹) ^a	<i>k</i> _{2,obs} × 10 ³ (s ⁻¹) ^a	Yield (%) ^b
1	2a	3.68 ± 0.03	-	91
2	2b	6.1 ± 0.7	4.1 ± 0.1	93
3	2c	-	-	NR°
4	2d	-	1.56 ± 0.02	96
5	2e	-	1.31 ± 0.01	89
6	2f	-	0.56 ± 0.01	62

[a] Monitored by ¹H NMR. [b] NMR yield at 30 min. [c] No reaction during the course of 30 min.

The PFAA-Staudinger reaction was next tested for its bioorthogonality (Fig. 3), employing the widely-studied Nacetylneuraminic acid (Neu5Ac) metabolic pathway that also accommodates other monosaccharides.5,21 PFAA-derivatized acetylated Man (4a), Gal (4b), and Glc (4c) (Fig. 3) were synthesized by a straightforward protocol (cf. Supporting Information). The labelling reagents, P2g-BSA-FITC or P2h-BSA-FITC, were synthesized by treating BSA with fluorescein isothiocyanate (FITC) followed by phosphines 2g or 2h, and purified by dialysis. BSA was chosen for its ability to reduce nonspecific protein adsorption. A549 cells were incubated with 4a, 4b, or 4c for 3 days,⁵ washed, and treated with P2g-BSA-FITC for 30 min. The extent of cell labelling was analysed using flow cytometry. Results showed that cells treated with 0.2 mM of 4a had ~4 times higher fluorescence intensity than the control (0 mM, Fig. 4A), indicating the successful expression of PFAA on the cell surface.

The experiment was then repeated using P_{2h}-BSA-FITC following the same protocol as above, and the labeled cells were analyzed by flow cytometry. Similar to the results of P_{2g}-BSA-FITC, cells treated with 0.2 mM of 4a displayed the highest fluorescence intensity (Fig. 4B). Furthermore, P_{2h}-BSA-FITC showed higher labeling efficiency, displayed ~11 times higher intensity than the control (0 mM) compared to P_{2g}-BSA-FITC (~4). P_{2h}-BSA-FITC was thus chosen as the labeling agent to test PFAA 4b and 4c. As can be seen from Fig. 4B, 4b was as efficient

as **4a**, whereas **4c** was much less efficient. Similar low labeling efficiency of GlcNAc derivatives was also observed by others, speculated as the restriction of GlcNAc into the sialic acid pathway, and the ability of GlcNAc analogs to produce multiple metabolic products.²¹⁻²³



Figure 3. Cell surface labelling experiment using PFAA-Staudinger reaction.



Figure 4. Fluorescence intensity of labelled cells measured by flow cytometry. (A) A549 cells were incubated with 4a at 0, 0.04, 0.09, 0.2 mM for 3 days followed by P₂₉-BSA-FITC (1 μ M) for 30 min. (B) A549 cells were incubated with 4a at 0, 0.09, 0.2 mM, 4b at 0.2 mM or 4c at 0.2 mM for 3 days followed by P_{2h}-BSA-FITC (1 μ M) for 30 min.

Fluorescence microscopy was furthermore used to image the labeled cells. For this, A549 cells were first incubated with **4a** at 0.29 mM for 3 days and then treated with **P**_{2h}-**BSA-FITC** for 1, 5, 10, 15, or 30 min. The green fluorescence that resulted from **P**_{2h}-**BSA-FITC** was clearly seen on the cells (**Fig. 5**). Successful labeling was already visible at 5 min. High fluorescence intensities were observed at 15 min, and remained unchanged after 30 min.

A549 cells were then incubated with **4a**, **4b** or **4c** for 3 days, followed by P_{2h} -**BSA-FITC** for 30 min, and the cell nuclei were stained with Hoechst 33342 which fluoresces blue. Green fluorescence was clearly seen on the cells (**Fig. 6, I**), where the signals were strong in the case of **4a** and **4b** (**Fig. 6A, 6B**), and

COMMUNICATION

much weaker in the case of **4c** (**Fig. 6C**). This is consistent with the flow cytometry results (**Fig. 4B**) showing the lower labelling efficiency for **4c**. Control samples prepared without treating the cells with **4** showed no visible green fluorescence (**Fig. 6D**).



Figure 5. Fluorescence images of labelled cells. Cells were incubated with 0.3 mM 4a for 3 days, followed by $P_{2h}\mbox{-}BSA\mbox{-}FITC$ for 1, 5, 10, 15, or 30 min.



Figure 6. Fluorescence images of A549 cells treated for 3 days with 0.3 mM of (A) 4a, (B) 4b, (C) 4c, or (D) none, followed by 1 μ M of P_{2h}-BSA-FITC at 4 °C for 30 min. The cell nuclei were stained with Hoechst 33342 dye. I: Excitation at 488 nm showing the green FITC. II: Excitation at 358 nm showing the blue stained nuclei. III: Overlay of I and II.

In summary, we have developed a fast PFAA-Staudinger reaction, with rate constants of up to 18 $M^{-1}s^{-1}$ in CD₃CN/D₂O (1/1). The iminophosphorane product was found stable towards hydrolysis thus eliminating the need of an ester trap in the Staudinger ligation. The iminophosphorane was also inert towards aza-Wittig reagents. Furthermore, PFAA-derivatized ManNAc and GalNAc were taken up by A549 cells and successfully probed by the PFAA-Staudinger reaction. The labelling reaction was fast, carried out using low concentration of

the labelling agent (1 μM) tagged with a conventional dye (FITC), affording clear images with low background noises.

Acknowledgements

We thank the National Institutes of Health (R01GM080295 and R21AI109896, to MY) for the financial support of this work. We acknowledge Dr. Andreas Fischer for his contribution in performing X-ray crystallography, and Dr. Nanjing Hao for his help in fluorescence microscopy.

Conflict of interest

The authors declare no conflict of interest.

Keywords: Staudinger reaction • perfluoroaryl azide • bioorthogonal reaction • metabolic labelling

- (1) Staudinger, H.; Meyer, J. Helv. Chim. Acta 1919, 2, 635.
- (2) Leffler, J. E.; Temple, R. D. J. Am. Chem. Soc. **1967**, *89*, 5235.
- (3) Tian, W. Q.; Wang, Y. A. *J. Chem. Theory Comput.* **2005**, *1*, 353.
- (4) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. *Angew. Chem. Int. Ed.* **2001**, *40*, 2004.
- (5) Saxon, E.; Bertozzi, C. R. Science 2000, 287, 2007.
- (6) Lin, F. L.; Hoyt, H. M.; van Halbeek, H.; Bergman, R. G.; Bertozzi, C. R. J. Am. Chem. Soc. 2005, 127, 2686.
- (7) Chang, P. V.; Prescher, J. A.; Hangauer, M. J.; Bertozzi, C. R. J. Am. Chem. Soc. 2007, 129, 8400.
- (8) Sletten, E. M.; Bertozzi, C. R. Acc. Chem. Res. 2011, 44, 666.
- (9) van Berkel, S. S.; van Eldijk, M. B.; van Hest, J. C. M. Angew. Chem. Int. Ed. **2011**, *50*, 8806.
- (10) Banks, R. E.; Sparkes, G. R. J. Chem. Soc., Perkin Trans. 1 1972, 2964.
- (11) Banks, R. E.; Prakash, A. *J. Chem. Soc., Perkin Trans.* 1 1974, 1365.
- (12) Palacios, F.; Alonso, C.; Aparicio, D.; Rubiales, G.; de los Santos, J. M. *Tetrahedron* **2007**, 63, 523.
- (13) Kennedy, R. D. Chem. Commun. 2010, 46, 4782.
- (14) Dommerholt, J.; van Rooijen, O.; Borrmann, A.; Guerra, C.
 F.; Bickelhaupt, F. M.; van Delft, F. L. Nat Commun 2014, 5.
- (15) Xie, S.; Lopez, S. A.; Ramström, O.; Yan, M.; Houk, K. N. *J. Am. Chem. Soc.* **2015**, *137*, 2958.
- (16) Xie, S.; Fukumoto, R.; Ramström, O.; Yan, M. J. Org. Chem. 2015, 80, 4392.
- (17) Xie, S.; Ramström, O.; Yan, M. Org. Lett. 2015, 17, 636.
- (18) Xie, S.; Zhang, Y.; Ramstrom, O.; Yan, M. *Chem. Sci.* **2016**, 7, 713.
- (19) Xie, S.; Manuguri, S.; Proietti, G.; Romson, J.; Fu, Y.; Inge,
 A. K.; Wu, B.; Zhang, Y.; Häll, D.; Ramström, O.; Yan, M.
 Proc. Natl. Acad. Sci. 2017.
- (20) Leffler, J. E.; Tsuno, Y. J. Org. Chem. 1963, 28, 902.
- (21) Wratil, P. R.; Horstkorte, R.; Reutter, W. Angew. Chem. Int. Ed. 2016, 55, 9482.
- (22) Du, J.; Meledeo, M. A.; Wang, Z.; Khanna, H. S.; Paruchuri, V. D. P.; Yarema, K. J. *Glycobiology* **2009**, *19*, 1382.
- (23) Saxon, E.; Luchansky, S. J.; Hang, H. C.; Yu, C.; Lee, S. C.; Bertozzi, C. R. J. Am. Chem. Soc. 2002, 124, 14893.

WILEY-VCH

COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

COMMUNICATION



Perfluoroaryl azides (PFAAs) react with aryl phosphines in a fast reaction under ambient conditions, yielding kinetically stable iminophosphorane products. The PFAA-Staudinger reaction proved to be an excellent bioorthogonal reaction and was successfully applied to cell surface labeling. Madanodaya Sundhoro,^[a] Seaho Jeon,^[a] Jaehyeung Park,^[a] Olof Ramström, ^{*[b]} and Mingdi Yan^{*[a][b]}

Page No. – Page No.

Fast, Bioorthogonal Staudinger Reaction