

Cite this: *RSC Adv.*, 2019, 9, 10091

Synthesis of 5-aminolevulinic acid with nontoxic reagents and renewable methyl levulinate†

Yuxia Zai,^a Yunchao Feng,^a Xianhai Zeng,^{ID} *^{abc} Xing Tang,^{ID} ^{abc} Yong Sun^{abc} and Lu Lin^{abc}Received 28th February 2019
Accepted 20th March 2019

DOI: 10.1039/c9ra01517e

rsc.li/rsc-advances

Synthesis of 5-aminolevulinic acid (5-ALA) was presented with novel bromination of biobased methyl levulinate (ML), followed by ammoniation and hydrolysis. Copper bromide (CuBr₂) was employed as the bromination reagent with higher selectivity and activity instead of the conventional liquid bromine (Br₂). 5-ALA was obtained in a high yield (64%) and purity (>95%) by optimum design, which is of great potential in industrialization.

5-Aminolevulinic acid (5-ALA) is generally known as an essential precursor molecule for tetrapyrrole synthesis such as porphyrin, heme, chlorophyll and vitamin B12.¹ It has been widely applied in localizing and photodynamic therapy for various cancers.^{2–4} It has also been used as a selective biodegradable insecticide, herbicide, salt tolerance agent or plant growth regulator in agricultural fields.⁵

To date, 5-ALA was mainly synthesized by microbial production methods,⁶ but the long-time and high-cost course restrict its scaled applications. On the other side, chemical routes using 2-hydroxypyridine, tetrahydrofurfurylamine and furfurylamine as starting materials involved in numerous bottleneck including toxic intermediates and rigorous reaction conditions.^{7,8} Thus, to develop a new pathway for 5-ALA production is of great significance, especially one that is a green and sustainable.

Biomass is an appealing starting material in value-added chemicals synthesis because of its advantages of renewability, sustainability and availability.⁹ Several biomass-derived platform compounds such as 5-hydroxymethylfurfural (HMF), 5-chloromethylfurfural (CMF), levulinic acid (LA) or its esters have been reported as efficient raw materials in the production of 5-ALA.^{10,11} However, the industrial manufacture of furan-type HMF and CMF cannot currently be achieved easily due to the high production and environment costs.^{12,13} Furthermore, conversion of furan-type chemicals to 5-ALA also suffers the economic problems concerning the use of expensive oxidants in the ring-opening stage. Unlike CMF and HMF, LA and its esters can be easily produced both from hemicellulose and cellulose, and its yearly tonnage is therefore available *via* the acidic

processing of biomass at a competitively low price.^{14–16} Thus, to synthesize 5-ALA from LA or its esters is exceptionally promising. Typically, 5-ALA can be effectively prepared from levulinates *via* a three-stage process including bromination, ammoniation and acidolysis.¹⁷ However, the bromination of levulinates with Br₂ in this course has low selectivity to 5-bromo derivatives. Besides, Br₂ is hazardous and environmentally unfriendly. Hence, a crucial step of the production of 5-ALA from levulinates is to explore a safe bromide agent with higher selectivity and activity.

CuBr₂, a green and low toxic brominated reagent, was usually used for the synthesis of α -bromination of cyclopentenone derivatives and its closest analogues-indanone of carbonyl compounds for its advantages of short reaction times, high selectivity of the products, high yields and easily handle procedures.^{18,19} In this content, various unsymmetrical aliphatic ketones including levulinic acid, methyl levulinate, ethyl levulinate, 5-hydroxy-2-pentanone and 2-butanone was attempted for bromizing with CuBr₂ (Table 1). Interestingly, the yields of bromination products were different, depending on the source of aliphatic ketones.

In this work, we present the synthesis of 5-ALA from biomass derived methyl levulinate (ML) under mild conditions using CuBr₂ as a greener bromine donor, and a high yield of 5-bromolevulinate (M5B) up to 85% was achieved. Furthermore, a detailed discussion of ammoniation and acidolysis was also presented, corresponding a high total 5-ALA yield over 64% (Scheme 1).


The first attempt to screen the reaction conditions for the bromination of ML with CuBr₂ are shown in Table 2. An encouraging yield of the desired product (50%) is indeed obtained using CuBr₂ as bromide agent in CH₃OH at 40 °C for 3 h (Table 2, entry 1). The investigation of the solvent indicated that CH₃OH–CHCl₃ mixed solvent was superior to ethyl acetate (EA), CHCl₃, CH₃OH, CH₃OH–EA and EA–CHCl₃ (Table 2, entry 2–6). The results may due to the fact that CH₃OH can improve the selectivity of M5B and haloalkanes are favourable to

^aCollege of Energy, Xiamen University, Xiamen 361102, China. E-mail: xianhai.zeng@xmu.edu.cn; Fax: +86-592-2880701; Tel: +86-592-2880701

^bFujian Engineering and Research Centre of Clean and High-valued Technologies for Biomass, Xiamen 361102, China

^cXiamen Key Laboratory of Clean and High-valued Utilization for Biomass, Xiamen 361102, China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra01517e



Scheme 2 The proposed mechanism of conversion of ML to M5B with CuBr_2 .

a key chemical that has been widely applied in medical and agricultural areas. CuBr_2 was applied as both catalyst and bromine atom donor and was demonstrated to be of higher selectivity and activity than the conventional hazardous Br_2 in ML bromination. Each stage proceeds in high ($\sim 85\%$) yield and affords 5-ALA in 95% purity, giving a process that could be commercially viable.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank for the financial support from the National Natural Science Foundation of China (Grant No. 21506177; 21676223), the Fundamental Research Funds for the Central Universities (Grant No. 20720160087; 20720160077; 20720170062), and the Energy Development Foundation of the College of Energy, Xiamen University (No. 2017NYFZ02).

Notes and references

- 1 K. Sasaki, M. Watanabe, T. Tanaka and T. Tanaka, *Appl. Microbiol. Biotechnol.*, 2002, **58**, 23–29.
- 2 M. C. Tetard, M. Vermandel, S. Mordon and J. P. Lejeune, *Photodiagn. Photodyn. Ther.*, 2014, **11**, 319–330.
- 3 T. Ishikawa, Y. Kajimoto, Y. Inoue, Y. Ikegami and T. Kuroiwa, *Adv. Cancer Res.*, 2015, **125**, 197.
- 4 H. Fukuda, A. Casas and A. Batlle, *Int. J. Biochem. Cell Biol.*, 2005, **37**, 272–276.
- 5 Z. J. Zhang, H. Z. Li, W. J. Zhou, Y. Takeuchi and K. Yoneyama, *Plant Growth Regul.*, 2006, **49**, 27–34.
- 6 S. L. Liu, G. M. Zhang, X. K. Li and J. Zhang, *Appl. Microbiol. Biotechnol.*, 2014, **98**, 7349–7357.

- 7 H. Kawakami, T. Ebata and H. Matsushita, *Agric. Biol. Chem.*, 1991, **55**, 1687–1688.
- 8 H. Takeya, H. Ueki, S. Miyanari, T. Shimizu and M. Kojima, *J. Photochem. Photobiol., A*, 1996, **94**, 167–171.
- 9 A. Corma, S. Iborra and A. Velty, *Chem. Rev.*, 2007, **107**, 2411–2502.
- 10 M. Mascal and S. Dutta, *Green Chem.*, 2010, **13**, 40–41.
- 11 L. Cottier, G. Descotes, L. Eymard and K. Rapp, *Cheminform.*, 1995, 303–306.
- 12 R. J. van Putten, J. C. van der Waal, E. de Jong, C. B. Rasrendra, H. J. Heeres and J. G. de Vries, *Chem. Rev.*, 2013, **113**, 1499–1597.
- 13 M. Mascal, *ChemSusChem*, 2015, **8**, 3391–3395.
- 14 S. G. Wettstein, D. M. Alonso, E. I. Gurbuz and J. A. Dumesic, *Curr. Opin. Chem. Eng.*, 2012, **1**, 218–224.
- 15 D. M. Alonso, J. Q. Bond and J. A. Dumesic, *Green Chem.*, 2010, **12**, 1493–1513.
- 16 D. J. Hayes, S. Fitzpatrick, M. H. B. Hayes and J. R. H. Ross, *The Biofine Process – Production of Levulinic Acid, Furfural, and Formic Acid from Lignocellulosic Feedstocks*, Wiley-VCH Verlag GmbH & Co., 2008, pp. 139–164.
- 17 H. J. Ha, S. K. Lee, Y. J. Ha and J. W. Park, *Synth. Commun.*, 1994, **24**, 2557–2562.
- 18 V. Z. Shirinian, D. V. Lonshakov, V. V. Kachala, I. V. Zavarzin, A. A. Shimkin, A. G. Lvov and M. K. Mikhail, Regio- and chemoselective bromination of 2,3-diarylcyclopent-2-en-1-ones, *J. Org. Chem.*, 2012, **77**, 8112–8123.
- 19 R. H. Vekariya and H. D. Patel, Synthesis of α -bromocarbonyl compounds: recent advances, *Tetrahedron*, 2014, **70**, 3949–3961.
- 20 L. Moens, Synthesis of δ -Aminolevulinic Acid, *ACS Symp. Ser.*, 2001, **784**, 37–50.
- 21 D. R. Lane, M. Mascal and P. Stroeve, *Renewable Energy*, 2016, **85**, 994–1001.
- 22 T. Liu, J. Sun, Q. Wang, L. Li and M. D. Zhou, *Eur. J. Org. Chem.*, 2017, **2017**, 1915–1921.
- 23 M. Y. Zhou, S. S. Kong, L. Q. Zhang, M. Zhao, J. A. Duan, Z. Ou-Yang and M. Wang, *Tetrahedron Lett.*, 2013, **54**, 3962–3964.
- 24 R. W. Evans, J. R. Zbieg, S. Zhu, W. Li and D. W. C. Macmillan, *J. Am. Chem. Soc.*, 2013, **135**, 16074–16077.
- 25 L. Moens, US005907058A, Lakewood and Colo, 1999.
- 26 L. Moens, US006583317B1, Lakewood and Colo, 2003.
- 27 J. J. Bozell, L. Moens, D. C. Elliott, Y. Wang, G. G. Neuenschwander, S. W. Fitzpatrick, R. J. Bilski and J. L. Jarnefeld, *Resour. Conserv. Recycl.*, 2000, **28**, 227–239.

