



Asymmetric Hetero Diels-Alder Reaction of *N*-Benzylimines Derived from *R*-Glyceraldehyde: A New Approach to Homochiral Piperidine Building Blocks and its Application to the Synthesis of (2*R*)-4-Oxopipercolic Acid

Ramón Badorrey, Carlos Cativiela*, María D. Díaz-de-Villegas* and José A. Gálvez


Instituto de Ciencia de Materiales de Aragón, Departamento de Química Orgánica, Universidad de Zaragoza-C.S.I.C.,
50009 Zaragoza, Spain. e-mail: cativiela@posta.unizar.es and loladiaz@posta.unizar.es

Abstract: The *N*-Benzyl imine derived from 2,3-di-*O*-benzyl-D-glyceraldehyde reacts with Danishefsky's diene to afford the corresponding hetero Diels-Alder adduct with a high diastereoselectivity. This compound can be transformed to enantiomerically pure (2*R*)-4-oxopipercolic acid.

© 1997 Elsevier Science Ltd.

The piperidine ring is one of the most abundant molecular fragments in both natural products and synthetic compounds with biological activity.¹ For this reason, the development of methods for the asymmetric synthesis of homochiral piperidine building blocks has recently received considerable attention.² The asymmetric hetero Diels-Alder cycloaddition using imines as dienophiles has given access to a wide variety of possibilities for the synthesis of the piperidine ring. The low reactivity of common deactivated imines is counteracted by the use of electron-rich butadiene derivatives such as Brassard's diene and Danishefsky's diene and/or the use of Lewis acids. The diastereoselectivity of this reaction has been investigated with chiral imines derived from chiral amines,³ chiral amino alcohols,⁴ chiral amino sugars,⁵ or chiral amino acid derivatives,⁶ as well as with the use of chiral Lewis acids,⁷ and more recently with the use of chiral dienes.⁸ Nevertheless, to the best of our knowledge and excluding the work of Akiba *et al.*,^{9a} that describes the use of the benzylimine derived from (*S*)-lactaldehyde as the starting material, and the work of Midland and McLoughlin,^{9b} that studies the cycloaddition reaction of a chiral imine derived from (2*R*,3*S*)-2,3-*O*-cyclohexylidene-2,3-dihydroxybutyraldehyde with Brassard's diene, there are no examples in the literature of hetero Diels-Alder cycloaddition reactions using chiral imines as dienophiles in which the chiral matrix is the carbonyl moiety.


Our recent work has been directed towards the synthesis of highly functionalised amino acids from chiral imines derived from conveniently protected D-glyceraldehyde,¹⁰ a useful starting material from the chiral pool. During the course of this research we have studied the behaviour of imine **1**, prepared from 2,3-di-*O*-benzyl-D-glyceraldehyde and benzylamine in the presence of anhydrous magnesium sulphate,^{10b} towards Danishefsky's diene.



Scheme 1

Danishefsky's diene (1.2 eq) reacts with imine **1** under standard conditions (i.e. in dichloromethane (2 ml/mmol) in the presence of 1.1 eq of a Lewis acid catalyst) to give the corresponding hetero Diels-Alder adduct in low to acceptable yields depending on the Lewis acid and reaction temperature used (Scheme 1). In most cases good stereoselectivities are obtained. The reaction in the presence of MgBr_2 or $\text{Eu}(\text{fod})_3$ works slowly, even at room temperature, and the imine decomposes in solution. However, ZnI_2 , $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and Et_2AlCl were all found to be effective catalysts to activate the imine, and good levels of induction and acceptable yields were obtained that improved on decreasing the reaction temperature from room temperature to -40°C . A rather different situation was observed when the reaction was carried out in the presence of SnCl_4 or TiCl_4 . Both catalysts promoted a rapid decomposition process, even at -78°C , and no adducts were detected in either case. From this study we conclude that the best catalyst system is ZnCl_2 in dichloromethane at -40°C . Under these reaction conditions a 90/10 diastereomeric mixture of the desired hetero Diels-Alder adducts **2**, **2'** was obtained in 65 % isolated yield. We subsequently studied the influence of solvent polarity in the hetero Diels-Alder reaction catalysed by ZnI_2 and we observed that, with aprotic solvents, an increase in the solvent polarity led to an improvement in the diastereoselectivity of the cycloaddition reaction. When the reaction was carried out in acetonitrile or nitroethane at -40°C a 95/5 diastereomeric mixture of the desired hetero Diels-Alder adducts **2**, **2'** was obtained in about 65 % isolated yield. At this point it is worth noting that the same results were obtained when a less than stoichiometric amount on ZnI_2 (0.2 eq) was used as catalyst.

The diastereomeric ratios were determined directly from the crude reaction mixtures by $^1\text{H-NMR}$ (major: δ 6.90 (dd), minor 7.02 (dd)), and the absolute configuration of the newly formed stereogenic carbon was ascertained to be (*R*) after conversion to a compound of known configuration as detailed below.




Scheme 2

Enamidone **2** can be a versatile intermediate for the construction of a wide variety of enantiomerically pure compounds containing the piperidine ring. The alkene moiety can act as a Michael acceptor in asymmetric conjugate additions, the carbonyl group can be stereoselectively transformed into several functional groups and

finally, the 1,2-diol moiety at C(2) can be conveniently elaborated to furnish a wide variety of substituents of known stereochemistry. To illustrate the synthetic utility of this compound we have performed the synthesis of conveniently protected (2*R*)-4-oxopipelic acid, an interesting compound recently used in the synthesis of selective NMDA antagonists,¹¹ in 45 % overall yield from compound 2 accordingly to scheme 2.

Conversion of compound 2 (90 % de)¹² into the corresponding saturated compound 3 was conveniently performed, in 92 % yield, with the use of L-Selectride® through a selective reduction of the double bond. Subsequent debenzoylation of the diol moiety, followed by oxidative cleavage, required prior protection of the carbonyl group. For this reason, ketone 3 was treated with ethylene glycol in the presence of *p*-toluenesulphonic acid to afford ketal 4 in 68 % yield as a single diastereoisomer as revealed by NMR spectroscopy. Compound 4 was debenzoylated by hydrogenolysis in the presence of di-*tert*-butyldicarbonate, using palladium hydroxide as a catalyst, to afford *N*-Boc derivative 5 in high yield (88 %). Oxidative cleavage of the 1,2-diol moiety of compound 5 by successive treatment with lead tetraacetate followed by Jones oxidation of the intermediate aldehyde, gave (2*R*)-4-oxopipelic acid 6 in a form conveniently protected for peptide syntheses.¹³



Scheme 3

In order to determine the absolute stereochemistry of C(2), this compound was further elaborated to give the known compound (*R*)-4-oxopipelic acid as its hydrate (scheme 3). Acid hydrolysis of 6 with a mixture of 6*N* hydrochloric acid-THF under reflux conditions for 7 days cleanly afforded the amino acid hydrochloride monohydrate 7. The specific rotation value measured for this compound ($[\alpha]_D^{25} = -4.0$ ($c = 0.55$, H_2O)) compared with the known value for (2*S*)-4-oxopipelic acid hydrochloride monohydrate¹⁴ clearly indicates an (*R*)-configuration at the stereogenic centre. Elution of the hydrochloride salt through an ion exchange chromatography column afforded the hydrate of (2*R*)-4-oxopipelic acid in enantiomerically pure form.

In conclusion, a simple method which allows the synthesis of a versatile building block for the synthesis of compounds containing the piperidine ring is described, starting from a common, commercially available precursor - D-mannitol. The synthetic utility of this chemistry has been illustrated by its application to the synthesis of enantiomerically pure (2*R*)-*N*-*tert*-butoxycarbonyl-4-(1,3-dioxolane-2-yl)-piperidine-2-carboxylic acid 6. Additional studies on the synthetic utility of this chiral building block are currently underway and will be published in due course.

Acknowledgement: This work was supported by the Dirección General de Investigación Científica y Técnica, project number PB94-0578. R. B. would like to express his gratitude to the *Ministerio de Educación y Ciencia* for a doctoral fellowship.

REFERENCES

1. (a) Baliah, V., Jeyaraman, R., Chandrasekaran, L., *Chem. Rev.*, **1983**, *83*, 379. (b) Strunz, G. M., Findlay, J. A., Pyridine and Piperidine Alkaloids. In *The Alkaloids*, Brossi, A., Ed.; Academic Press, New York, **1985**, Vol. 26, p 89. (c) Numata, A., Ibuka, I., Alkaloids from Ants and Other Insects. In *The Alkaloids*, Brossi, A., Ed.; Academic Press, New York, **1987**, Vol. 31, p 193. (d) Rubiralta, M., Giralt, E., Diez, E., In *Piperidine, Structure, Preparation, Reactivity and Synthetic Applications of Piperidine and its Derivatives*, Elsevier, Amsterdam, **1991**. (e) Pinder, A. R., *Nat. Prod. Rep.*, **1993**, *10*, 491 (f) Daly, J. W., Garraffo, H. M., Spande, T. F., In *The Alkaloids*, Cordell, G. A., Ed.; Academic Press, New York, **1993**, Vol. 43, p 185.
2. See for example: (a) Comins, D. L., Joseph, S. P., Goehring, R. R., *J. Am. Chem. Soc.*, **1994**, *116*, 4719. (b) Génisson, Y., Mehmandoust, M., Marazano, C., Das, B. C., *Heterocycles*, **1994**, *39*, 811. (c) Oppolzer, W., *Pure Appl. Chem.*, **1994**, *66*, 2127. (d) Suzuki, H. Aoyagi, S., Kibayashi, C., *J. Org. Chem.*, **1995**, *60*, 6114. (e) Munchlof, M. J., Meyers, A. I., *J. Org. Chem.*, **1995**, *60*, 7084. (f) Amat, M., Llor, N., Hidalgo, J., Hernández, A., Bosch, J., *Tetrahedron: Asymmetry*, **1996**, *7*, 977. (g) Amat, M., Llor, N., Hidalgo, J., Bosch, J., Molins, E., Miravittles, C., *Tetrahedron: Asymmetry*, **1996**, *7*, 2501. (h) De Kimpe, N., Boelens, M., Contreras, J., *Tetrahedron Lett.*, **1996**, *37*, 3171. (i) Agami, C., Couty, F., Mathieu, H., *Tetrahedron Lett.*, **1996**, *37*, 4001. (j) Herdeis, C., Kaschinski, C., Karla, R., Lotter, H., *Tetrahedron: Asymmetry*, **1996**, *7*, 867. (k) Agami, C., Hamon, L., Kadouri-Puchot, C., Le Guen, V., *J. Org. Chem.*, **1996**, *61*, 5736. (l) Froelich, O., Desos, P., Bonin, M., Quirion, J. C., Husson, H. P., *J. Org. Chem.*, **1996**, *61*, 6700.
3. Hattori, K., Yamamoto, H., *Synlett*, **1993**, 129.
4. (a) Midland, M. M., Koops, R. W., *J. Org. Chem.*, **1992**, *57*, 1158. (b) Devine, P. N., Reilly, M., Oh, T., *Tetrahedron Lett.*, **1993**, *34*, 5827.
5. (a) Pfrengle, W., Kunz, H., *J. Org. Chem.*, **1989**, *54*, 4261. (b) (a) Kunz, H., Pfrengle, W., *Angew. Chem., Int. Ed. Engl.*, **1989**, *28*, 1067.
6. (a) Waldman, H., Braun, M., *J. Org. Chem.*, **1992**, *57*, 4444. (b) Waldman, H., Braun, M., Dräger, M., *Angew. Chem., Int. Ed. Engl.*, **1990**, *29*, 1468. (c) Waldman, H., Braun, M., Dräger, M., *Tetrahedron: Asymmetry*, **1991**, *2*, 1231. (d) Lock, R., Waldman, H., *Tetrahedron Lett.*, **1996**, *37*, 2753.
7. (a) Hattori, K., Yamamoto, H., *J. Org. Chem.*, **1992**, *57*, 3264. (b) Hattori, K., Yamamoto, H., *Tetrahedron*, **1993**, *49*, 1749. (c) Ishiara, K., Miyata, M., Hattori, K., Tada, T., Yamamoto, H., *J. Am. Chem. Soc.*, **1994**, *116*, 10520.
8. Barluenga, J., Aznar, F., Ribas, C., Valdés, C., Fernández, M., Cabal, M. P., Trujillo, J., *Chem. Eur. J.*, **1996**, *2*, 805.
9. (a) Ishimaru, K., Watanabe, K., Yamamoto, Y., Akiba, K. Y., *Synlett*, **1994**, 495. (b) Midland, M. M., McLoughlin, J. I., *Tetrahedron Lett.*, **1988**, *29*, 4653.
10. (a) Cativiela, C., Díaz-de-Villegas, M. D., Gálvez, J. A., *Tetrahedron Lett.*, **1995**, *36*, 2859. (b) Cativiela, C., Díaz-de-Villegas, M. D., Gálvez, J. A., *Tetrahedron: Asymmetry*, **1996**, *7*, 529. (c) Cativiela, C., Díaz-de-Villegas, M. D., Gálvez, J. A., García, J. I., *Tetrahedron*, **1996**, *52*, 9563.
11. Machetti, F., Cordero, F. M., De Sarlo, F., Guarna, A., Brandi, A., *Tetrahedron Lett.*, **1995**, *36*, 2859.
12. All new compounds reported here were fully characterised by spectroscopic and analytical methods.
13. 6: $[\alpha]_{\text{D}}^{25} = 31.3$ (c = 1, CHCl₃), ¹H-NMR (CDCl₃, 300 Mhz, 60 °C) 1.43 (s, 9H), 1.61 (bd, 1H), 1.65-1.75 (m, 1H), 1.75-1.90 (m, 1H), 2.37 (bd (1H), 3.29 (bt, 1H), 3.85-4.10 (m, 5H), 4.95 (bs, 1H), ¹³C-NMR (CDCl₃, 75 Mhz, 60 °C) 28.3, 34.2, 34.6, 39.8, 64.3, 64.6, 80.6, 106.4, 155.4, 176.3.
14. (2S)-4-oxopipicolic acid hydrochloride monohydrate $[\alpha]_{\text{D}}^{22} = 4.1$ (c = 0.55, H₂O), Golubev, A., Sewald, N., Burger, K., *Tetrahedron Lett.*, **1996**, *37*, 4205.

(Received in UK 16 January 1997; revised 25 February 1997; accepted 28 February 1997)