

Tetrahedron: Asymmetry Vol. 6, No. 4, pp. 913-918, 1995 Elsevier Science Ltd Printed in Great Britain 0957-4166/95 \$9.50+0.00

0957-4166(95)00099-2

A Novel Chiral Cyclopentadienyl Ligand Based on Ephedrine

Adolphus A. H. van der Zeijden

Institut für Anorganische Chemie, Martin-Luther-Universität Halle-Wittenberg, Geusaer Strasse, D-06217 Merseburg, Germany

Abstract: natural (-)-(1R,2S)-ephedrine was used as a starting material for the preparation of a novel chiral cyclopentadienyl ligand in a three-step procedure. The new ligand contains a nitrogen atom suitable for chelation to a transition metal, as well as two stereogenic centres.

Introduction:

The design of novel ligands as effective chiral auxiliaries in transition metal catalysed asymmetric conversions is a rapidly growing area of interest in synthetic chemistry.¹ Apart from the traditional phosphinebased systems, chirally-substituted cyclopentadienyl (Cp) ligands become increasingly important in this sense.² This is because Cp ligands can be used as complexing agents for almost any transition metal, as well as the enormous synthetic potential to modify this type of ligand. The initial strategy in the development of chiral Cp ligands was the introduction of sterically crowded substituents, like the menthyl group, by a single C-C bond to the Cp backbone.³ Optical inductions in catalytic processes using these ligands were low, which was probably due to the dynamic nature of these ligands. Therefore, a further development was focused on the attachment of the chiral substituents to the Cp ring by more than one C-C bond, so-called annulated Cp's, in order to get a more rigid system.² Accordingly, high enantiomeric excesses were obtained during catalytic hydrogenations of olefins by titanocene derivatives.

We pursue here another strategy to obtain rigidly metal-coordinated chiral Cp systems, namely the introduction of a heteroatom in the chiral Cp, which creates a fixed chiral pocket around the complexed metal by additional intramolecular coordination. Chirality in these systems may be introduced in three ways: (1) substitution of a second group onto the Cp ring (planar chirality), (2) attachment of chiral substituents on the bridging carbons or (3) on the heteroatom.



Some scattered reports based on this approach have appeared in literature.⁴ We wish to report here our results on the synthesis of a novel chiral Cp ligand using ephedrine and pseudoephedrine as a source of chirality.

Results and discussion:

Our synthetic strategy is outlined below.



Commercially available (-)-(1R,2S)-ephedrine and (+)-(1S,2S)-pseudoephedrine were used as starting materials. Their conversion to N-methylephedrine (1**a**) and -N-methylpseudoephedrine (1**b**) was straightforward.⁵ They were reacted with SOCl₂ in CHCl₃ to obtain the alkyl chlorides (see below).⁶ With 1**a** this conversion proceeded with 15% retention and 85% inversion at C_1 (yielding 2**a** and 2**b**, respectively), whereas with 1**b** a *ca* 50:50 mixture of 2**a** and 2**b** was obtained.^{7,8} Several recrystallisations afford the 1S,2S-diastereomer 2**b** in high purity. For this reason it is better to start with 1**a**, in which case 2**b** can be obtained in an overall yield of 40-50%.



Reaction of 2b and an excess of CpNa afforded a quantitative amount of the Cp ligand (abbreviated as $Cp^{E}(phedrine)H)$.[†] It consists of a 1:1 mixture of two olefinic regioisomers. Therefore, the reaction seems to have gone stereoselectively, i. e. either with complete inversion or retention at C1. Usually, the C1-C2 ³J(H,H) coupling constant is taken to distinguish between the 1*S*,2*S* (or 1*R*,2*R*) and 1*R*,2*S* (or 1*S*,2*R*) diastereomers of (pseudo)ephedrine derivatives; its value of 10.4 Hz suggests an *S* configuration at C1. However, in Cp^EH the Cp and Ph groups at C1 are more or less isosteric and therefore it is possible that both the *R* and the *S* configurations show coupling constants in the same range. This was corroborated by the results of the reaction of a 40:60 batch of 2a and 2b (obtained from 1b and SOC12) with CpNa. The isolated mixture of Cp derivatives contained 40% of the two regioisomers of Cp^EH (with presumed 1*S*,2*S* configuration) as well as 60% of the two regioisomers of what is considered to be the 1*R*,2*S* diastereomer. The latter shows a ³J(H,H) coupling of 10.8 Hz, which indeed is close to the value of the presumed 1*S*,2*S* isomer (10.4 Hz).



Normally, this kind of alkylation proceeds by the S_N2 mechanism, and should lead to inversion at $C_{1,9}$ However, the neighbouring effect of the NMe₂ group probably causes retention at C_1 , by forming an aziridinium salt as an intermediate.⁹ This is corroborated by results of Carboni *et al.*, who showed that similar conversions of (pseudo)ephedrine chlorides with NaN₃ proceed with complete retention of configuration.⁸ We therefore believe, as suggested above, that the configuration at C_1 in Cp^EH is S.¹⁰

Similarly, we attempted to prepare the analogous indene ligand, by reaction of **2b** with an excess of an indenyl alkaline salt. The special interest for this ligand arises from the fact that it will create a third stereogenic centre upon coordination to a transition metal (principle of planar chirality).



[†]In describing the configuration around carbon, CH(Me)NMe₂, Ph, Cp, Me, and H groups have descending order of precedence.

The substitution takes place regioselective at the C_3 atom of indene, but unfortunately it is not stereoselective with respect to the ephedrine C_1 atom, producing two diastereomers.¹¹ The isomer ratio strongly depends on the alkaline metal employed (see Table I), but the outcome of the reaction can never be driven completely to one of the diastereomers. Longer reaction times do not change the isomer ratio as a possible result of base-promoted epimerization. These observations suggest a competing direct alkylating (SN2-like) pathway for this reaction. No further attempts to separate the two isomers were undertaken.

Table	I.	Rel.	an	nounts	of
diastere	om	ers o	n r	eaction	of
indenvl	Мx	vith 2	b ir	THE.1	1

М	15,25	1 <i>R</i> ,2 <i>S</i>			
Li	22%	78%			
Na	75%	25%			
K	92%	8%			

Currently, we are exploring the coordination chemistry of Cp^EH, and its application in metal mediated stereoselective transformations of organic substrates.

Acknowledgement - Financial support of the 'Deutsche Forschungsgemeinschaft' is gratefully acknowledged. Prof. Dr. Karl-Heinz Thiele is thanked for his generous support and interest.

Experimental:

General procedures - Preparations involving organometallic materials were conducted under an atmosphere of argon using carefully dried solvents. ¹H and ¹³C n.m.r. spectra were obtained using a Varian Gemini instrument operating at 300 and 75.5 MHz, respectively. Chemical shifts were referenced to those of CD₃OD (δ H 3.30; δ C 49.0 ppm), CDCl₃ (δ H 7.24; δ C 77.0 ppm) or C₆D₆ (δ H 7.15; δ C 128.0 ppm).

Starting Materials - (1R,2S)- and $(1S,2S)-\alpha$ -[1-(dimethylamino)ethyl]benzenemethanol (N-methylephedrine (1a) and -pseudoephedrine (1b)) were obtained by published methods from (-)-<math>(1R,2S)-ephedrine (Janssen) and (+)-(1S,2S)-pseudoephedrine (Aldrich), respectively.⁵

(15,25)- β -Chloro-N,N', α -trimethylbenzeneethanammonium chloride hydrate(2b) - To an ice-cold solution of 1a (62.7 g, 350 mmol) in chloroform (350 ml) was added SOCl₂ (32 ml, 440 mmol). The resulting clear, orange solution was stirred overnight at room temperature and evaporated to afford a foamy red solid. Recrystallisation from ether/ethanol (3:1 v/v) at -80 °C afforded 88 g (ca 100%) of a pale orange substance containing 85% 2b and 15% of the (1*R*,2*S*) diastereomer 2a by ¹H NMR. Several recrystallisations from acetone/ethanol (10:1 v/v) were necessary to obtain pure 2b as white needles (39 g, 44%). (Found: C, 52.8; H, 7.9; N, 5.5. C1₁H₁9Cl₂NO requires C, 52.4; H, 7.6; N, 5.5%). NMR in CD₃OD: δ H 1.07 (3H, d, 6.7 Hz, CHCH₃), 2.90/3.02 (both 3H, s, N(CH₃)₂H⁺), 4.14 (1H, dq, 10.9x6.7 Hz, CHCH₃), 5.31 (1H, d, 10.9 Hz, CHPh), 7.39-7.53 (5H, m, Ph). δ C 10.24 (CHCH₃), 36.55/43.98 (N(CH₃)₂H⁺), 62.85 (CHCH₃), 67.60 (CHPh), 129.10 (Ph,o-C), 130.28 (Ph, m-C), 130.75 (Ph, p-C), 138.37 (ipso-C).

A similar reaction with (15,25)-N-methylpseudoephedrine) and SOCl₂ in chloroform afforded a crude mixture of 50% **2a** and 50% **2b**. No attempts were made to separate the two diastereomers by recrystallisation. NMR of **2a** in CD₃OD (containing **2b**): $\delta_{\rm H}$ 1.38 (3H, d, 6.6 Hz, CHCH₃), 2.97/3.10 (both 3H, s, N(CH₃)₂H⁺), 3.92 (1H, dq, 3.7x6.6 Hz, CHCH₃), 5.82 (1H, d, 3.7 Hz, CHPh), ca 7.35-7.55 (5H, m, Ph). $\delta_{\rm C}$ 9.77 (CHCH₃), 41.66/42.57 (N(CH₃)₂H⁺), 62.48 (CHCH₃), 68.13 (CHPh), 128.57 (Ph,*o*-C), 130.09 (Ph, *m*-C), 130.25 (Ph, *p*-C), 137.59 (*ipso*-C).

(15,25)-(2-Dimethylamino-1-phenylpropyl)cyclopentadiene (' Cp^EH') - A solution of cyclopentadienylsodium was prepared by reacting sodium (18.6 g, 0.81 mol) and freshly cracked cyclopentadiene (85 ml, 1 mol) in THF (300 ml) overnight at room temperature. To this solution was added **2b** (36.1 g, 143 mmol) and stirring was continued for another night at room temperature. Water (20 ml) was added and volatiles were removed *in vacuo*. The residue was extracted with petrol (150 ml in total). Removal of solvent left a quantitative yield of a yellow-brownish viscous oil (34.6 g) containing almost pure **3**. This substance can be used for subsequent manipulations without further purification. It was identified by NMR (CDCl3) as a 1:1 mixture of 1- and 2substituted cyclopentadienes: δ_H 0.67/0.68 (3H, both d, 6.3 Hz, CHCH3), 2.20/2.22 (6H, both s, N(CH3)2), 2.91 (2H, m, CH2 of Cp ring), 3.22 (1H, m, CHCH3), 3.67/3.71 (1H, both d, 10.4 Hz, CHPh), 6.08 (s), 6.19 (d, 5.2 Hz), 6.24 (s), 6.33 (m), 6.51 (d, 5.2 Hz) (these 5 signals integrate for 3H, CH of Cp ring), 7.05-7.3 (5H, m, Ph). δ_C 9.33/9.42 (CHCH3), 40.04/40.18 (N(CH3)2), 40.87/41.67 (CH2 of Cp ring), 51.68/52.74 (C HPh), 61.58/61.87 (C HC H 3), 126.0 (Ph, *p*-C), 128.4 (br, Ph, *o*, *m*-C), 126.35/126.83/131.03/132.08/132.93/133.91 (CH of Cp ring), 143.68/143.91 (Ph, *ipso*-C), 148.96/151.18 (C of Cp ring).

A similar reaction of CpNa with a mixture of **2a** (60%) and **2b** (40%) yielded a mixture of 40% Cp^EH and 60% of the 1*R*,2*S*-analogue of Cp^EH. The latter was identified by NMR in CDCl3: $\delta_{\rm H}$ 0.86/0.88 (3H, both d, 6.4 Hz, CHCH3), 2.10/2.12 (6H, both s, N(CH3)2), 2.73 (2H, m, CH2 of Cp ring), 3.26* (1H, m, CHCH3), 3.80* (1H, d, 10.8 Hz), 6.1-6.3*, 6.46 (3H, m, CH of Cp ring), 7.1-7.3* (5H, m, Ph). $\delta_{\rm C}$ 9.58/9.71 (CHCH3), 39.91/40.1* (N(CH3)), 40.82*/41.34 (CH2 of Cp ring), 51.73/52.90 (CHPh), 60.73/61.06 (CHCH3), 127.5/128.0*/128.3* (Ph, *o*,*m*,*p*-C), 125.65/125.75/131.17/131.95*/133.22/133.83* (CH of Cp ring), 143.58/144.13 (Ph, *ipso*-C), 148.59/150.88 (C of Cp ring).

*(partly) coincides with signals of CpEH.

Reaction of 2b with indenyl salts¹² - A five-fold excess of either indenyllithium (prepared by the action of *n*-butyllithium on indene for 1h at 0 °C), -sodium (prepared by refluxing sodium with a slight excess indene for 24h) or -potassium (prepared by reaction of potassium with a slight excess of indene for 2h at room temperature) in THF was reacted with **2b** at room temperature for 24h. Work-up as above yielded a mixture that was analysed by NMR in CDCl3. Two compounds were identified: (15,25)-[(2-Dimethylamino-1-phenylpropyl)]-3-indene: $\delta_{\rm H}$ 0.97 (3H, d, 6.4 Hz, CHCH3), 2.18 (6H, s, N(CH3)2), 3.31 (2H, m, CH2 of indene), 3.45 (1H, dq, 10.4x6.4 Hz, CHCH3), 4.05 (1H, d, 10.4 Hz, CHPh), 6.38 (1H, br s, CH of indene), 7.05-7.4* (9H, m, Ph and indene). $\delta_{\rm C}$ 9.37* (CHCH3), 37.87* (CH2 of indene), 40.40 (N(CH3)2), 49.04* (CHPh), 61.51 (CHCH3), 119.45* (indene C4), 123.47 (indene C7), 124.09 (indene C6), 125.65? (indene C5), 125.9*? (Ph, *p*-C), 127.89* (Ph, *o*-C), 128.38* (Ph, *m*-C), 128.2? (indene C2), 142.18 (Ph, *ipso*-C), 144.39 (indene C8), 145.39? (indene C3), 145.95? (indene C9) (assignment of indene signals was based on those of 3-methylindene.¹³). (1*R*,25)-[(2-Dimethylamino-1-phenylpropyl)]-3-indene: $\delta_{\rm H}$ 0.72 (3H, d, 6.6 Hz, CHCH3), 2.24 (6H, s, N(CH3)2), 3.35 (3H, m, CH2 of indene and CHCH3), 4.00 (1H, d, 10.1 Hz, CHPh), 6.49 (1H, br s, CH of indene), 7.05-7.4* (9H, m, Ph and indene). $\delta_{\rm C}$ 9.37* (CHCH3), 37.87* (CH2 of indene C4), 123.64 (indene

C7), 124.45 (indene C6), 125.81 (indene C5), 125.9*? (Ph, p-C), 127.89* (Ph, o-C), 128.38* (Ph, m-C), 128.86 (indene C2), 142.76 (Ph, *ipso*-C), 144.26 (indene C8), 145.20 (indene C3), 146.37 (indene C9). *(partly) coincides with signals of the other diastereomer.

References and notes:

- 1. Ojima, I. Catalytic Asymmetric Synthesis; VCH: New York, 1993.
- 2. a) Halterman, R. L. Chem. Rev., 1992, 92, 965. b) Okudo, J. Topics Curr. Chem., 1992, 160, 97.
- 3. Cesarotti, E.; Kagan, H. B. J. Organomet. Chem., 1978, 162, 297.
- a) Huang, Q.; Qian, Y. Synthesis, 1987, 910. b) Huang, Q.; Qian, Y.; Tang, Y. J. Organomet. Chem., 1989, 368, 277. c) Huang, Q.; Qian, Y.; Tang, Y. Trans. Met. Chem., 1989, 14, 315. d) Van de Weghe, P.; Bied, C.; Collin, J.; Marçalo, J.; Santos, I. J. Organomet. Chem., 1994, 475, 121. e) Adams, H.; Bailey, N. A.; Corey, M.; Schofield, P. A.; White, C. J. Chem. Soc., Dalton Trans., 1994, 1445.
- 5. Coote, S. J.; Davies, S. G.; Goodfellow, C. L.; Middlemiss, D.; Naylor, A Tetrahedron: Asymmetry, 1990, 1, 817.
- 6. The reaction of *N*-Methylephedrine with SOCl₂ has been reported. However, no details about the diastereomeric purity of the products were given: Hukki, J.; Hirvimies, A.; Väänänen, E. Suomen Kemistilehti, **1967**, *B40*, 37.
- 7. The stereoselectivity of the reaction with SOCl₂ parallels with that of ephedrine (10% retention and 90% inversion) and pseudoephedrine (60:40): Allen, A. C.; Kiser, W. O. J. Forensic Studies, **1987**, 953.
- 8. Benalil, A.; Carboni, B.; Vaultier, M. Tetrahedron, 1991, 8177.
- 9. March, J. Advanced Organic Chemistry; John Wiley: New York, 1985.
- Meanwhile the 1S,2S configuration has been confirmed by determination of the X-ray structure of Cp^E₂Fe.2MeI: van der Zeijden, A. A. H., Sieler, J. to be published.
- 11 The two diastereomers of the indene ligands could be discriminated by ¹H NMR: for the Cp ligands the signals of the Me group in the 1*S*, 2*S* diastereomer lie 0.20 ppm upfield to those of the 1*S*, 2*R* diastereomer, whereas the signals of the NMe₂ group lie 0.10 downfield (see *Experimental*). As the indene analogues also show differences of 0.20 ppm for the Me signal and -0.10 ppm for the NMe₂ signal, and besides have the same ³J_{HH} coupling (10 Hz) as the Cp-analogue, similar three-dimensional structures are anticipated.
- 12. Weidler, A.-M. Acta Chem. Scand. 1963, 17, 2724.
- 13. Edlund, U. Chemica Scriptra 1975, 7, 85.

(Received in UK 7 February 1995)