# ARTICLE

www.rsc.org/obc

)BC

# Development of a solid-phase 'asymmetric resin-capture-release' process: application of an ephedrine chiral resin in an approach to $\gamma$ -butyrolactones<sup>†</sup>

Nessan J. Kerrigan,<sup>a</sup> Panee C. Hutchison,<sup>a</sup> Tom D. Heightman<sup>b</sup> and David J. Procter\*<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of Glasgow, The Joseph Black Building, Glasgow,

UK G12 8QQ. E-mail: davidp@chem.gla.ac.uk

<sup>b</sup> GlaxoSmithKline, New Frontiers Science Park, Third Avenue, Harlow, Essex, UK CM19 5AW

Received 11th June 2004, Accepted 8th July 2004 First published as an Advance Article on the web 4th August 2004

The potential of a solid-phase asymmetric resin-capture–release strategy for high-throughput synthesis has been evaluated. Fukuzawa's Sm(II)-mediated, asymmetric approach to  $\gamma$ -butyrolactones was selected to illustrate the feasibility of such a process.  $\alpha$ , $\beta$ -Unsaturated esters immobilised on an ephedrine chiral resin have been applied in an asymmetric approach to  $\gamma$ -butyrolactones. Lactone products are obtained in moderate isolated yields with selectivities up to 96% ee. In addition, we have shown that the ephedrine resin can be conveniently recovered and recycled although in some cases lower yields were obtained on reuse of the chiral resin. A short synthesis of a moderate DNA-binding microbial metabolite using asymmetric resin-capture–release is also described.

# Introduction

Solid phase organic synthesis in which organic substrates are manipulated whilst linked to an insoluble polymer is an important tool for the synthetic chemist who wishes to prepare collections of compounds in an efficient manner.1 The area of solid-phase synthesis continues to evolve and recently a shift towards the use of solid-supported reagents to effect the transformation of substrates in solution has been seen.2 'Resin-capture-release' is a new concept which may also play a role in the future of solid-phase synthesis.<sup>3</sup> Resin-capture-release is a hybrid technique that combines elements of traditional solid-phase synthesis and the use of supported reagents. In a conventional resin-capture-release process, a small molecule B is trapped as an activated polymer intermediate A-B, by a functionalised resin A. The addition of a second reaction partner C transforms the polymer-bound intermediate and releases a product B-C into solution whilst generating a modified polymer (Fig. 1a).<sup>3</sup>

We envisaged that the use of a functionalised *chiral* resin as an immobilised chiral auxiliary<sup>4</sup> would give rise to an 'asymmetric resin-capture–release' process (Fig. 1b). Attachment of a substrate B to a chiral resin followed by asymmetric addition of a second reaction partner C generates an activated, diastereoisomeric intermediate. This breaks down to release a non-racemic product B–C\* into solution. Crucially, the chiral resin is regenerated directly and should be recyclable.

To explore the feasibility of such a process we choose to use Fukuzawa's Sm(II)-mediated, asymmetric route to  $\gamma$ -butyrolactones<sup>5</sup> in our proof of concept studies. In Fukuzawa's studies,  $\alpha$ , $\beta$ -un-saturated esters bearing an ephedrine chiral auxiliary are reacted with aldehydes and ketones in the presence of SmI<sub>2</sub> to give  $\gamma$ -butyrolactones.<sup>5</sup> Several features of the process make it ideal for illustrating the concept of asymmetric resin-capture–release: Firstly, the ephedrine auxiliary should be amenable for linkage to a solid-phase thus giving a functionalised chiral resin. Secondly, immobilised ephedrinyl  $\alpha$ , $\beta$ -unsaturated esters should be easily accessible by acylation of a chiral ephedrine resin. Finally, after asymmetric radical addition, an unstable diastereoisomeric polymer-bound intermediate will be formed which should collapse to release an

<sup>†</sup> Electronic supplementary information (ESI) available: Additional experimental and characterisation data, <sup>1</sup>H and <sup>13</sup>C NMR for new compounds, <sup>1</sup>H NMR for literature compounds, general chiral GC conditions, chiral GC traces for lactone products and FTIR spectra of resins. See http://www.rsc.org/suppdata/ob/b4/b408836k/

enantiomerically enriched lactone product from the resin, while simultaneously regenerating the chiral ephedrine resin (Scheme 1).

Our interest in new linker systems for solid-phase synthesis<sup>6</sup> has led us to develop a pseudoephedrine resin for asymmetric solid phase synthesis.<sup>7</sup> Furthermore, we have demonstrated the potential of the chemistry for asymmetric library synthesis.<sup>7b</sup> In this contribution we describe in full our studies<sup>8</sup> on the use of an ephedrine resin in the development of an asymmetric resin-capture–release process, employing Fukuzawa's  $\gamma$ -butyrolactone synthesis<sup>5</sup> to illustrate the concept.

# **Results and discussion**

We began by preparing the chiral ephedrine resin necessary to investigate an asymmetric resin-capture-release process based on Fukuzawa's asymmetric reductive-coupling methodology.5 Inexpensive (1R, 2S)-ephedrine was immobilised on commercially available bromo Wang resin selectively through nitrogen and in one step, by adaptation of a literature procedure.9 The efficiency and chemoselectivity of the immobilisation step was clearly illustrated by the following solution phase model reaction. Benzyl bromide (1 eq.) was heated with (1R,2S)-ephedrine (2 eq.) at 85 °C in DMF. After 4 days, (1R,2S)-N-benzyl ephedrine was obtained in near quantitative yield with no trace of (1R, 2S)-O-benzyl ephedrine<sup>10</sup> observed in the crude <sup>1</sup>H NMR of the reaction. The loading of the resultant ephedrine resin 1 was found to be approximately 1 mmol g<sup>-1,7,11</sup> Straightforward esterification of the resin with acryloyl or crotonyl chloride then gave the desired resins 2 ( $v_{max}(C=O)$  1724 cm<sup>-1</sup>) and 3 ( $v_{max}(C=O)$  1734 cm<sup>-1</sup>), respectively (Scheme 2).

Using acetophenone and acrylate resin **2**, optimised conditions for the asymmetric-release step were developed. Typically, ketyl radical anion–olefin couplings of this type are carried out by adding a mixture of the carbonyl compound and the acrylate to a solution of  $SmI_2$ .<sup>12</sup> Using solution phase model studies, we found that changing the order of addition, *i.e.* adding  $SmI_2$  to a mixture of the resin and the carbonyl compound, a prerequisite for a convenient solid-phase experimental procedure, did not significantly lower the enantioselectivities obtained. For example, adding acetophenone, alcohol and acrylate to  $SmI_2$  at 0 °C, gave lactone **4** in 73% ee, while adding  $SmI_2$  to acetophenone, alcohol and acrylate, gave **4** in 68% ee. The yields obtained using both orders of addition were also similar, 84% and 79% respectively.

Temperature was found to have a dramatic effect on the efficiency of the asymmetric catch–release process. At -78 °C, only a trace of lactone **4** was detected, while at -40 °C, a disappointing

DOI: 10.1039/b408836



Fig. 1 (a) Conventional 'resin-capture-release'. (b) Asymmetric 'resin-capture-release'

Table 1



Scheme 1 'Asymmetric resin-capture–release': proof of concept studies using Fukuzawa's asymmetric approach to  $\gamma$ -butyrolactones.



**Scheme 2** *Reagents and conditions:* i, (1*R*,2*S*)-ephedrine, DMF, 85 °C; ii, acryloyl chloride or crotonyl chloride, NEt<sub>3</sub>, Et<sub>2</sub>O, rt.

23% yield of **4** was isolated. Carrying out the reaction at -15 °C using an excess of the acrylate resin **2**, however, gave a satisfactory isolated yield of lactone **4** (50%) in good enantiomeric excess (74% ee). Higher temperatures may be essential for efficient cyclative cleavage of the diastereoisomeric intermediate from the resin. Our adaptation of Fukuzawa's methodology to the solid-phase represents one of a limited number of examples of the addition of radicals to immobilised acceptors.<sup>13</sup>

Employing the optimised conditions, the asymmetric resin-capture-release process involving acrylate resin **2** and a range of ketones and aldehydes was investigated (Table 1). Phenyl alkyl ketones were found to give the corresponding lactones in moderate yield and enantiomeric excess (70–74% ee) (entries 1–3). In general, aldehydes were found to give higher enantioselectivities upon reaction with acrylate resin **2** (up to 81% ee) (entries 4–8), with more substituted aldehydes giving the highest enantioselectivities (entries 7 and 8). Significant ketyl-radical homo-coupling was not observed and few by-products were visible in the crude <sup>1</sup>H NMR spectra. In general, the *isolated* yields obtained using **2** compare favourably with the GC yields of Fukuzawa<sup>5</sup> although the use of an excess of **2** is important to give satisfactory conversion.

Treatment of the analogous crotonate resin **3** with aldehydes under optimised conditions gave the expected  $\gamma$ -butyrolactones in good yield and high enantiomeric excess. Enantioselectivities were considerably higher than those obtained using acrylate resin **2** at the same temperature (-15 °C). Again, more substituted aldehydes give the highest enantioselectivities (entries 1 and 2). In line with Fukuzawa's observations, only the *cis*-lactones were observed in the crude <sup>1</sup>H NMR spectra of the reactions with aldehydes.<sup>14</sup> In contrast, the reaction with 2-hexanone gave **17** as an inseparable 3 : 1 mixture of diastereoisomers (entry 6) (Table 2).

Entry Substrate Product Isolated yield 4 R = Me, 50%  $(74\% \text{ ee})^a$ 1 - 35 R = Et, 73%  $(70\% \text{ ee})^a$  $6 \text{ R} = \text{Pr}, 43\% (74\% \text{ ee})^a$ 4 37% (71% ee)<sup>a</sup> 5 52% (66% ee)<sup>a</sup> 6 56% (70% ee)t 7 56% (81% ee) 66% (76% ee) 8

'Asymmetric resin-capture-release' with acrylate resin 2

See Experimental Section for details. <sup>*a*</sup> Enantiomeric excess determined by chiral GC (see Electronic Supporting Information). <sup>*b*</sup> Enantiomeric excess determined by optical rotation.

Although our studies aim to illustrate the general feasibility of asymmetric resin-capture–release processes rather than to provide a solid-phase process superior to solution-phase methodology,<sup>5</sup> it is interesting to compare the two processes. Enantioselectivities obtained from reactions using acrylate resin **2** are lower than those reported in solution.<sup>5</sup> It is likely this difference originates from the different temperatures at which the two processes are carried out (-78 °C versus -15 °C). As previously stated, the yields obtained using **2** compare favourably with the GC yields of Fukuzawa<sup>5</sup> although the use of an excess of **2** is important to give satisfactory conversion. Enantioselectivities in reactions using crotonate resin **3** are comparable to those obtained in solution despite the difference in the reaction temperatures.

Published on 04 August 2004. Downloaded by University of Michigan Library on 31/10/2014 12:23:35



See Experimental Section for details. <sup>*a*</sup>Enantiomeric excess determined by chiral GC (see Electronic Supporting Information). <sup>*b*</sup>Enantiomeric excess determined by optical rotation. <sup>*c*</sup>A 3:1 mixture of *cis* (89% ee) and *trans* (76% ee) diastereoisomers was obtained. The *cis*-product was assumed to be the major diastereoisomer.

Despite the utility of Fukuzawa's asymmetric approach to  $\gamma$ butryolactones,<sup>5</sup> no explanation has been advanced for the high selectivities observed. It is clear that chelation of the ephedrine auxiliary to samarium(III) is important as it has been reported that racemic products were obtained when HMPA was employed as an additive.5 In addition, the original screen of auxiliaries, suggests that both the phenyl and the amino group play a key role.<sup>5</sup> To account for the observed enantio- and diastereoselectivity in the reaction, we therefore propose the transition structure shown in Fig. 2. Coordination of samarium(III) to both the auxiliary and to the incoming ketyl-radical anion gives rise to a well-ordered transition structure. Such chelation is only possible when the substrate adopts a s-trans conformation. In this conformation, the lower face of the substrate is blocked by the phenyl group of the ephedrine auxiliary so the radical addition occurs from the top face. The chelation not only orders the transition structure but also leads to Lewis acid activation of the substrate towards radical addition (Fig. 2).

The greater selectivities we have observed in additions to the crotonate resin 3 (average 92% ee) compared to those obtained



Fig. 2 Possible origin of selectivity in the reaction.

View Article Online

using acrylate resin 2 (average 73% ee), at -15 °C, are more difficult to explain. Such a difference in selectivity may arise from the different reactivities of the two radical acceptors. Lewis acid activation of the less reactive crotonate acceptor is likely to be important, thus addition only occurs to the chelated substrate (Fig. 2) where the diastereofacial selectivity is strictly controlled by the auxiliary. The more reactive acrylate may undergo a non-selective background reaction through a non-chelated transition structure thus eroding the enantiomeric excess of the products.

Our studies have shown that asymmetric resin-capture-release is a viable strategy for solid-phase synthesis. The efficacy of any such process clearly depends on the careful design/selection of the key reaction with regard to the asymmetry-controlling auxiliary, reagent compatibility with the polymer support, and the mechanism of product release. In our original outline of the process (Fig. 1), we highlighted that regeneration of the chiral resin after product release should allow recovery and reuse of the chiral resin. Again using Fukuzawa's methodology to illustrate this feature of the process, we have carried out preliminary studies into the feasibility of recycling the chiral ephedrine resin 1. For example, employing crotonate resin 3 with pivalaldehyde gave lactone 12 in 54% yield and 92% ee. Recovery of the resin 1 and re-esterification with crotonyl chloride gave recycled 3. Re-treatment with pivalaldehyde gave 12 in virtually identical yield and enantiomeric excess (56%, 92% ee). However, recovering and using the resin a third time gave 12 in substantially lower yield (27%) although the enantioselectivity of the reaction was still high (86% ee). (Scheme 3).

Attempts to recycle the acrylate resin 2 were less successful. For example, employing acrylate resin 2 with pivalaldehyde, gave lactone 10 in 50% yield and 79% ee. Recovery of the resin 1 and re-esterification with acryloyl chloride gave recycled 2. Treatment of the recycled resin with pivalaldehyde gave 10 in substantially lower yield and enantiomeric excess (27%, 68% ee) (Scheme 4).

Although we have shown the feasibility of recycling the chiral ephedrine resin, some difficulties remain to be addressed. The drop in yield encountered using recovered resin may be due either to a build up of inorganic salts in the polymer support or less likely, could be caused by background reduction of the acrylate and crotonate resins by SmI<sub>2</sub>.<sup>15</sup> Thus repeated re-charging and reuse of the resin would lead to the observed drop in the efficacy of the resin and the yield of the reaction. Studies are currently under way to develop a more efficient protocol for the recycling of the resin, which will allow multiple runs to be carried out, making different lactone products in each cycle.

To illustrate the utility of asymmetric resin-capture–release for the synthesis of biologically active compounds, we have undertaken a short synthesis of  $\gamma$ -butyrolactone **21**, a moderate DNA-binding metabolite isolated from Streptomyces GT61115.<sup>16</sup> Diol **18** was prepared from  $\delta$ -valerolactone according to a literature method.<sup>17</sup> Mono-protected diol **19** was prepared by bis-protection followed by cleavage of the primary TMS ether under basic conditions. Dess–Martin oxidation<sup>18</sup> of **19** then gave aldehyde **20** in excellent yield. On treatment with acrylate resin **2** and SmI<sub>2</sub> in the presence of *tert*-butanol, aldehyde **20** gave **21**, after loss of the TMS protection during work up, in 50% yield and 73% ee.<sup>19</sup> Our synthesis confirms the postulated absolute stereochemistry of **21** (Scheme 5).<sup>16</sup>

# Conclusions

In summary, we have sought to evaluate 'asymmetric resin-capture– release' as a potential process for the asymmetric, solid-phase synthesis of small molecules. We have utilised an asymmetric approach to  $\gamma$ -butyrolactones in our proof of concept studies. In the resultant process, an ephedrine chiral resin is loaded and the immobilised substrate undergoes asymmetric transformation through capture of a reactive intermediate from solution. Cyclative cleavage gives  $\gamma$ -butyrolactones in moderate yield and good enantiomeric excess. We have proposed an explanation for the origin of enantioselectivity in the process and have shown that the ephedrine chiral resin can be recovered and reused although in some cases reduced yields and enantioselectivities were observed. Finally, we have applied



Scheme 3 Reagents and conditions: i, SmI<sub>2</sub> (0.1 M in THF), THF, -15 °C, t-BuOH; ii, crotonyl chloride, NEt<sub>3</sub>, Et<sub>2</sub>O, rt.



Scheme 4 Reagents and conditions: i, SmI<sub>2</sub> (0.1 M in THF), THF, -15 °C, t-BuOH; ii, acryloyl chloride, NEt<sub>3</sub>, Et<sub>2</sub>O, rt.



Scheme 5 Reagents and conditions: i, MeLi (1.5 M in Et<sub>2</sub>O), THF, -78 °C to rt, 66%; ii, a) TMSCl, NEt<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt, b) K<sub>2</sub>CO<sub>3</sub>, MeOH, 0 °C, 96% for two steps; iii, Dess Martin periodinane, CH<sub>2</sub>Cl<sub>2</sub>, rt, 98%; iv, acrylate resin **2**, SmI<sub>2</sub> (0.1 M in THF), THF, -15 °C, *t*-BuOH, 50%.

the asymmetric resin-capture-release process in a short asymmetric synthesis of a moderate DNA-binding metabolite.

Our studies have shown that asymmetric resin-capture–release is a viable and potentially powerful strategy for solid-phase synthesis. We are currently developing other asymmetric resin-capture–release processes and investigating the application of this concept in highthroughput asymmetric synthesis.

# **Experimental**

#### **General considerations**

See the Electronic Supporting Information.<sup>†</sup>

# Preparation of (1R,2S)-N-Wang bound ephedrine 1

To a suspension of bromo Wang resin (4.23 g, 5.90 mmol, 1 equiv.) in DMF (42 mL) was added (1*R*,2*S*)-ephedrine (5.87 g, 35.5 mmol, 6 equiv.). The resulting suspension was heated at 85 °C and stirred gently for 4 days. After this time, the reaction suspension was filtered and the resin was washed with EtOH (3 × 50 mL), water (3 × 50 mL), THF–water; 1:1 (3 × 50 mL), THF (3 × 50 mL), and EtOH (3 × 50 mL). Resin **1** was then dried *in vacuo*:  $v_{max}$  (Golden Gate)/cm<sup>-1</sup> 2919m, 1606m, 1508s.

#### Preparation of (1R,2S)-N-Wang bound ephedrinyl acrylate 2

To a suspension of (1R,2S)-N-Wang bound ephedrine 1 (2.15 g, 2.69 mmol, 1 equiv.) in Et<sub>2</sub>O (21.5 mL) was added triethylamine (0.90 mL, 6.46 mmol, 2.4 equiv.) followed by acryloyl chloride (0.54 mL, 6.46 mmol, 2.4 equiv.). The resulting suspension was allowed to stir at room temperature for 24 h. After this time, the reaction suspension was filtered and the resin was washed with

THF (3 × 25 mL), THF–water; 2:1 (3 × 25 mL), THF–water; 1:1 (3 × 25 mL), THF–water; 1:2 (3 × 25 mL), (MeOH, 25 mL then CH<sub>2</sub>Cl<sub>2</sub>, 25 mL) × 3, and MeOH (3 × 25 mL). The acrylate resin **2** was then dried *in vacuo*:  $v_{max}$  (Golden Gate)/cm<sup>-1</sup> 1724s (C=O).

#### Preparation of (1R,2S)-N-Wang bound ephedrinyl crotonate 3

To a suspension of (1R,2S)-N-Wang bound ephedrine **1** (3.48 g, 4.07 mmol, 1 equiv.) in Et<sub>2</sub>O (36.0 mL) was added triethylamine (1.36 mL, 9.78 mmol, 2.4 equiv.) followed by *trans*-crotonyl chloride (1.04 mL, 9.78 mmol, 2.4 equiv.). The resulting suspension was allowed to stir at room temperature for 24 h. After this time, the reaction suspension was filtered and the resin was washed with THF (3 × 50 mL), THF–water; 2:1 (3 × 50 mL), THF–water; 1:1 (3 × 50 mL), THF–water; 1:2 (3 × 50 mL), (MeOH, 50 mL then CH<sub>2</sub>Cl<sub>2</sub>, 50 mL) × 3, and MeOH (3 × 50 mL). The crotonate resin **3** was then dried *in vacuo*:  $v_{max}$  (Golden Gate)/cm<sup>-1</sup> 1734s (C=O).

# General procedure A. Reaction of (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate/crotonate with ketones and aldehydes

A suspension of (1R,2S)-N-Wang bound ephedrinyl acrylate/ crotonate resin, 2 or 3, (3.2 equiv.) in THF (5 mL) under argon was gently stirred for 15 min prior to the addition of an aldehyde (1 equiv.) or a ketone (1 equiv.) and t-BuOH (2 equiv.). The resultant suspension was then allowed to stir for another 1 hour at room temperature before being cooled to -15 °C. A pre-cooled SmI<sub>2</sub> solution (0.1 M in THF, 5.5 equiv.) was then added and the dark blue-black solution was allowed to stir at -15 °C until TLC analysis showed that the reaction was complete. The reaction was then allowed to warm to room temperature over 5-6 hours before filtration. The filtered resin was washed repeatedly with THF. The filtrate was then concentrated to about 30 mL and washed with aqueous saturated NaCl (4 mL). The aqueous layer was separated and washed with  $Et_2O$  (3 × 15 mL). The combined organic layers were then dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in vacuo. The crude product was purified by column chromatography on silica to afford the purified product as a pale yellow or colourless oil.

#### (S)-5-Methyl-5-phenyl dihydro-furan-2-one 4<sup>5</sup>

As for general procedure A. Reaction of acetophenone (18  $\mu$ L, 0.16 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.50 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **4** (13.8 mg, 50%) as a pale yellow oil:  $[a]_{\rm D}$  –31.3 (*c* 0.63 in MeOH); lit.,<sup>5</sup> (90% ee)  $[a]_{\rm D}$  –34.1 (*c* 0.21 in

MeOH);  $\nu_{max}$  (neat)/cm<sup>-1</sup> 3028s, 1772s (C=O);  $\delta_{H}$  (400 MHz, CDCl<sub>3</sub>) 7.31–7.21 (5H, m, Ar*H* of Ph), 2.62–2.31 (4H, m, C*H*<sub>2</sub>C*H*<sub>2</sub>C=O), 1.66 (3H, s, C*H*<sub>3</sub>C);  $\delta_{C}$  (100 MHz, CDCl<sub>3</sub>) 170.6 (C=O), 144.7 (ArC), 128.9 (ArCH), 128.0 (2 × ArCH), 124.5 (2 × ArCH), 87.4 (C–O), 36.6 (CH<sub>2</sub>C=O), 29.8 (CH<sub>3</sub>C), 29.3 (CH<sub>2</sub>C); *m*/z (EI<sup>+</sup> mode) 176 (M<sup>+</sup>, 15%), 161 (100), 121 (30), 105 (23), 82 (10), 77 (18) (Found M<sup>+</sup> 176.0837. C<sub>11</sub>H<sub>12</sub>O<sub>2</sub> requires 176.0837).

74% ee, determined by chiral GC (see supporting information<sup>†</sup>).

#### (S)-5-Ethyl-5-phenyl dihydro-furan-2-one 5<sup>5</sup>

As for general procedure A. Reaction of propiophenone (21 µL, 0.16 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.50 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **5** (22.1 mg, 73%) as a pale yellow oil:  $v_{\text{max}}$  (neat)/cm<sup>-1</sup> 2963m, 1776s (C=O), 1448m, 1232m, 1196m;  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 7.39–7.21 (5H, m, Ar*H*), 2.65–2.39 (4H, m, 2 × CH<sub>2</sub>), 2.00 (2H, q, *J* 7.4, CH<sub>2</sub>CH<sub>3</sub>), 0.82 (3H, t, *J* 7.4, CH<sub>2</sub>CH<sub>3</sub>);  $\delta_{\text{C}}$  (100 MHz, CDCl<sub>3</sub>) 177.0 (C=O), 143.1 (ArC), 128.9 (2 × ArCH), 127.9 (2 × ArCH), 125.1 (ArCH), 90.2 (C–O), 35.7 (CH<sub>2</sub>CH<sub>3</sub>), 35.0 (CH<sub>2</sub>), 29.1 (CH<sub>2</sub>C=O), 8.6 (CH<sub>3</sub>C); *m/z* (EI<sup>+</sup> mode) 190 (M<sup>+</sup>, 2.5%), 161 (100), 133 (10), 105 (30) (Found M<sup>+</sup> 190.0994. C<sub>12</sub>H<sub>14</sub>O<sub>2</sub> requires 190.0994).

70% ee, determined by chiral GC (see supporting information).

#### (S)-5-Propyl-5-phenyl dihydro-furan-2-one 6

As for general procedure A. Reaction of butyrophenone (26 µL, 0.18 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.57 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **6** (15.9 mg, 43%) as a pale yellow oil: [a]<sub>D</sub> –33.0 (c 0.01 in MeOH);  $v_{max}$  (neat)/cm<sup>-1</sup>2960s, 2873m, 1780s (C=O), 1448m, 1324m, 1297m, 1228s;  $\delta_{H}$  (400 MHz, CDCl<sub>3</sub>) 7.32–7.20 (5H, m, ArH), 2.55–2.33 (4H, m, 2 × CH<sub>2</sub>), 1.91–1.81 (2H, m, CH<sub>2</sub>), 1.34–1.25 (1H, m, 1H from CH<sub>2</sub>) 1.08–0.97 (1H, m, 1H from CH<sub>2</sub>), 0.77 (3H, t, *J* 7.4, CH<sub>2</sub>CH<sub>3</sub>);  $\delta_{C}$  (100 MHz, CDCl<sub>3</sub>) 177.1 (*C*=O), 143.4 (ArC), 128.9 (2 × ArCH), 127.9 (2 × ArCH), 125.0 (ArCH), 89.9 (CO), 45.1 (CH<sub>2</sub>), 35.5 (CH<sub>2</sub>), 29.1 (CH<sub>2</sub>C=O), 17.6 (CH<sub>2</sub>CH<sub>3</sub>), 14.4 (CH<sub>3</sub>C); m/z (EI<sup>+</sup> mode) 204 (M<sup>+</sup>, 2.5%), 161 (100), 149 (2.5), 105 (25), 77 (15) (Found M<sup>+</sup> 204.1148. C<sub>13</sub>H<sub>16</sub>O<sub>2</sub> requires 204.1150).

74% ee, determined by chiral GC (see supporting information).

#### (R)-5-n-Pentyl dihydro-furan-2-one 75

As for general procedure A. Reaction of hexanal (25 µL, 0.21 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.66 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **7** (11.9 mg, 37%) as a pale yellow oil:  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.48 (1H, apparent q, *J* = 7.6, CH), 2.53 (2H, dd, *J* 6.3, 8.4, CH<sub>2</sub>C=O), 2.32 (1H, dt, *J* 6.6, 13.7, 1H from CH<sub>2</sub>CH<sub>2</sub>C=O), 1.90–1.24 (9H, m, 4 × CH<sub>2</sub>CH<sub>2</sub>, and 1H from CH<sub>2</sub>CH<sub>2</sub>C=O), 0.90 (3H, t, *J* 6.9, CH<sub>3</sub>C);  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 177.7 (*C*=O), 81.5 (CHO), 35.9 (CH<sub>2</sub>CH), 31.9 (CH<sub>2</sub>C(O)), 29.3 (CH<sub>2</sub>), 28.4 (CH<sub>2</sub>), 25.3 (CH<sub>2</sub>), 22.9 (CH<sub>2</sub>), 14.3 (CH<sub>3</sub>); *m/z* (EI<sup>+</sup> mode) 156.2 (M<sup>+</sup>, 2), 138 (3), 128 (5), 100 (5), 84 (100), 49 (10) (Found M<sup>+</sup> 156.1143. C<sub>9</sub>H<sub>16</sub>O<sub>2</sub> requires 156.1150).

71% ee, determined by chiral GC (see supporting information).

# (R)-5-n-Butyl dihydro-furan-2-one 820

As for general procedure A. Reaction of pentanal (18 µl, 0.17 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.54 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **8** (12.4 mg, 52%) as a pale yellow oil:  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.49 (1H, quintet, *J* 7.6, CHOC(O)), 2.59–2.51 (2H, m, CH<sub>2</sub>C(O)), 2.33 (1H, apparent sextet, *J* 6.8, 1H from CH<sub>2</sub>CH<sub>2</sub>C(O)), 1.91–1.83 (1H, m, 1H from CH<sub>2</sub>CH<sub>2</sub>C(O)),

1.76–1.73 (1H, m, 1H from CH<sub>2</sub>CH), 1.64–1.57 (1H, m, 1H from CH<sub>2</sub>CH), 1.48–1.32 (4H, m, 2 × CH<sub>2</sub>), 0.92 (3H, t, *J* 7.2, CH<sub>3</sub>);  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 177.3 (*C*(O)), 81.0 (CH), 35.3 (CH<sub>2</sub>CH), 28.9 (CH<sub>2</sub>C(O)), 28.0 (CH<sub>2</sub>CH<sub>2</sub>C(O)), 27.3 (CH<sub>2</sub>), 22.4 (CH<sub>2</sub>), 13.9 (CH<sub>3</sub>); *m/z* (CI<sup>+</sup> mode, isobutane) 143 ((M + H)<sup>+</sup>, 100), 81 (8) (Found (M + H)<sup>+</sup> 143.1074. C<sub>8</sub>H<sub>15</sub>O<sub>2</sub> requires 143.1072).

66% ee, determined by chiral GC (see supporting information).

# (S)-5-Isopropyl dihydro-furan-2-one 95

As for general procedure A. Reaction of isobutyraldehyde (23 µL, 0.25 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.78 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **9** (17.6 mg, 56%) as a pale yellow oil:  $[a]_{\rm D}$  +26.9 (*c* 0.02 in MeOH); lit, <sup>5</sup> (93% ee)  $[a]_{\rm D}$  +35.6 (*c* 0.2–1.0 in MeOH);  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.20 (1H, td, *J* 8.5, 7.0, CHO), 2.60–2.52 (2H, m, CH<sub>2</sub>C=O), 2.26 (1H, apparent sextet, *J* 6.5, 1H from CH<sub>2</sub>CH<sub>2</sub>C=O), 1.96–1.79 (2H, m, 1H from CH<sub>2</sub>CH<sub>2</sub>C=O and CH(CH<sub>3</sub>)<sub>2</sub>), 1.03 (3H, d, *J* 6.7, CH(CH<sub>3</sub>)<sub>2</sub>), 0.94 (3H, d, *J* 6.8, CH(CH<sub>3</sub>)<sub>2</sub>);  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 177.3 (*C*=O), 85.8 (CHO), 33.0 (CH), 29.1 (CH<sub>2</sub>C=O), 25.6 (CH<sub>2</sub>CH<sub>2</sub>C=O), 18.4 (CH<sub>3</sub>), 17.3 (CH<sub>3</sub>); *m*/z (EI<sup>+</sup> mode) 128 (M<sup>+</sup>, 25%), 95 (15), 85 (100), 69 (22), 57 (30), 56 (25) (Found M<sup>+</sup> 128.0834. C<sub>7</sub>H<sub>12</sub>O<sub>2</sub> requires 128.0837).

70% ee, determined by chiral GC (see supporting information).

#### (S)-5-tert-Butyl dihydro-furan-2-one 105

As for general procedure A. Reaction of pivalaldehyde (27 µL, 0.25 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl acrylate **2** (0.78 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **10** (19.3 mg, 56%) as a pale yellow oil:  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.19 (1H, dd, *J* 6.9, 8.8, CH), 2.55–2.50 (2H, m, CH<sub>2</sub>C=O), 2.16–2.08 (1H, m, 1H from CH<sub>2</sub>), 2.02–1.94 (1H, m, 1H from CH<sub>2</sub>), 0.95 (9H, s, 3 × (CH<sub>3</sub>)C);  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 177.8 (*C*=O), 88.6 (*C*–O), 34.2 (*C*), 29.7 (CH<sub>2</sub>C=O), 25.3 (3 × CH<sub>3</sub>), 23.3 (CH<sub>2</sub>); *m/z* (CI<sup>+</sup> mode) 143 ((M + H)<sup>+</sup>, 30%), 57 (100) (Found (M + H)<sup>+</sup>, 143.1073. C<sub>8</sub>H<sub>15</sub>O<sub>2</sub> requires 143.1072).

81% ee, determined by chiral GC (see supporting information).

#### (S)-5-Cyclohexyl dihydro-furan-2-one 115

As for general procedure A. Reaction of cyclohexanecarboxaldehyde (29 µL, 0.23 mmol, 1.0 equiv.) with (1R,2S)-N-Wang bound ephedrinyl acrylate 2 (0.75 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40-60 °C)) gave 11 (26.0 mg, 66%) as a pale yellow oil:  $[a]_D$  +29.8 (c 0.01 in MeOH); lit.<sup>5</sup> (96% ee)  $[a]_{\rm D}$  +26.2 (c 0.2–1.0 in MeOH);  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.14 (1H, dt, J 8.0, 7.4, CHO), 2.54-2.50 (2H, m, CH<sub>2</sub>C=O), 2.25 (1H, apparent sextet, J 6.7, 1H from CH<sub>2</sub>CH<sub>2</sub>C=O), 1.98–1.90 (2H, m, 1H from CH<sub>2</sub>CH<sub>2</sub>C=O and 1H from CH<sub>2</sub>), 1.80-1.76 (2H, m, CH<sub>2</sub>), 1.72-1.65 (2H, m, CH<sub>2</sub>), 1.57-1.51 (1H, m, CH), 1.29-1.16 (3H, m,  $CH_2$ ), 1.06–1.00 (2H, m,  $CH_2$ );  $\delta_C$  (100 MHz,  $CDCl_3$ ) 177.3 (C=O), 85.1 (CO), 42.7 (CH), 29.0 (CH<sub>2</sub>), 28.9 (CH<sub>2</sub>), 27.7 (CH<sub>2</sub>), 26.2 (CH<sub>2</sub>), 25.7 (CH<sub>2</sub>), 25.6 (CH<sub>2</sub>), 25.5 (CH<sub>2</sub>); m/z (EI<sup>+</sup> mode) 168 (M<sup>+</sup>, 8%), 140 (10), 111 (15), 85 (100), 67 (10), 55 (28), 41 (16) (Found M<sup>+</sup> 168.1149. C<sub>10</sub>H<sub>16</sub>O<sub>2</sub> requires 168.1150).

76% ee, determined by chiral GC (see supporting information).

#### (4R,5S)-5-tert-Butyl-4-methyl dihydro-furan-2-one 125

As for general procedure A. Reaction of pivalaldehyde (16.5 µl, 0.15 mmol, 1 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl crotonate **3** (0.49 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **12** (13.2 mg, 56%) as a yellow oil:  $v_{\text{max}}$  (neat)/cm<sup>-1</sup> 2924s, 2854m, 1736m (C=O), 1458m;  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 4.09 (1H, d, *J* 4.9, CHO), 2.76 (1H, dd, *J* 7.5, 16.6, 1H of CH<sub>2</sub>C=O), 2.69–2.64 (1H, m, CHCH<sub>3</sub>), 2.20 (1H, dd, *J* 1.6, 16.6, 1H of CH<sub>2</sub>C=O), 1.14 (3H, d, *J* 7.1, CH<sub>3</sub>CH), 1.08 (9H, s,

 $(3 \times CH_3)_3$ C);  $\delta_C$  (100 MHz, CDCl<sub>3</sub>) 177.6 (C=O), 90.6 (CH–O), 40.1 (CH<sub>2</sub>C=O), 34.6 (CHCH<sub>3</sub>), 34.2 (C), 26.9 (3 × (CH<sub>3</sub>)<sub>3</sub>C), 16.3 (CH<sub>3</sub>CH); *m/z* (EI<sup>+</sup> mode) 141 (M–CH<sub>3</sub><sup>+</sup>, 10%), 99 (100), 83 (63), 71 (44), 57 (100) and 41 (39) (Found (M – CH<sub>3</sub>)<sup>+</sup>, 141.0917. C<sub>8</sub>H<sub>13</sub>O<sub>2</sub> requires 141.0916).

93% ee, determined by chiral GC (see supporting information).

# (4R,5S)-5-Cyclohexyl-4-methyl dihydro-furan-2-one 135

As for general procedure A. Reaction of cyclohexane carboxaldehyde (18.4 µl, 0.15 mmol, 1 equiv.) with (1R,2S)-N-Wang bound ephedrinyl crotonate 3 (0.49 mmol, 3.2 equiv.) and purification by column chromatography on silica (eluting with 10% EtOAc/petroleum ether (40-60 °C)) gave 13 (18.2 mg, 66%) as a pale yellow solid:  $[a]_{D}$  +53.9 (c 0.5 in MeOH); lit.,  $[a]_{D}$  +56.2 (c 1.0 in MeOH);  $v_{max}$  (Golden Gate)/cm<sup>-1</sup> 2927s, 1747s (C=O), 1456w, 1174s;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 3.95 (1H, dd, J 4.7, 9.8, CHO), 2.65 (1H, dd, J 7.4, 16.9, 1H of CH<sub>2</sub>C=O). 2.50-2.47 (1H, m, CHCH<sub>3</sub>), 2.12 (1H, dd, J 1.1, 16.9, 1H of  $CH_2C=O$ ), 2.00–1.96 (1H, m, 1H of  $CH_2$  of cyclohexane ring), 1.71-1.62 (4H, m, 2 × CH<sub>2</sub> cyclohexane ring), 1.58-1.53 (2H, m, 1H of CH<sub>2</sub> of cyclohexane ring and CHCHO), 1.22-1.07 (2H, m, CH<sub>2</sub> of cyclohexane ring), 1.00–0.99 (1H, m, 1H of CH<sub>2</sub> of cyclohexane ring), 0.95 (3H, d, J 12.9, CH<sub>3</sub>CH), 0.92–0.83 (1H, m, 1H of  $CH_2$  of cyclohexane ring);  $\delta_C$  (100 MHz, CDCl<sub>3</sub>) 177.4 (C=O), 88.1 (CHO), 39.2 (CH<sub>2</sub>C=O), 37.8 (CHCHO), 32.4 (CHCH<sub>3</sub>), 30.7 (CH<sub>2</sub> of cyclohexane ring), 28.4 (CH<sub>2</sub> of cyclohexane ring), 26.5 (CH<sub>2</sub> of cyclohexane ring), 25.8 (CH<sub>2</sub> of cyclohexane ring), 25.7 (CH<sub>2</sub> of cyclohexane ring), 13.9 (CH<sub>2</sub> of cyclohexane ring); m/z (EI<sup>+</sup> mode) 182 (M<sup>+</sup>, 13%), 154 (28), 99 (100), 84 (41), 83 (29) and 49 (34) (Found M<sup>+</sup> 182.1306. C<sub>11</sub>H<sub>18</sub>O<sub>2</sub> requires 182.1307)

96% ee, determined by optical rotation.

#### (4R,5R)-5-Isopropyl-4-methyl dihydro-furan-2-one 14<sup>21</sup>

As for general procedure A. Reaction of isobutyraldehyde  $(20.6 \ \mu l, 0.23 \ mmol, 1 \ equiv.)$  with (1R, 2S)-N-Wang bound ephedrinyl crotonate 3 (0.73 mmol, 3.2 equiv.) and purification by column chromatography on silica (eluting with 10% EtOAc/ petroleum ether (40-60 °C)) gave 14 (17.9 mg, 55%) as a yellow oil: [a]<sub>D</sub> +11.8 (c 0.4 in CHCl<sub>3</sub>); v<sub>max</sub> (neat)/cm<sup>-1</sup> 2926s, 2875s, 2362m, 1751s (C=O), 1594m, 1457m;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 3.94 (1H, dd, J 4.8, 10.2, CHO), 2.74 (1H, dd, J 7.4, 16.9, 1H of CH<sub>2</sub>C=O), 2.58-2.53 (1H, m, CHCH<sub>2</sub>C=O), 2.21 (1H, dd, J 1.1, 16.9, 1H of CH<sub>2</sub>C=O), 1.93-1.86 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 1.09 (3H, d, J 6.5, CH<sub>3</sub>CHCH<sub>3</sub>), 1.00 (3H, d, J 7.0, CH<sub>3</sub>CHCH<sub>2</sub>), 0.91 (3H, d, J 6.6, CH<sub>3</sub>CHCH<sub>3</sub>); δ<sub>C</sub> (100 MHz, CDCl<sub>3</sub>) 177.0 (C=O), 89.1 (CHO), 39.0 (CH<sub>2</sub>C=O), 31.9 (CHCH<sub>2</sub>C=O), 28.1 (CH(CH<sub>3</sub>)<sub>2</sub>), 20.1 (CH<sub>3</sub>CHCH<sub>3</sub>), 18.4 (CH<sub>3</sub>CHCH<sub>3</sub>), 13.4 (CH<sub>3</sub>CHCH<sub>2</sub>); m/z (EI<sup>+</sup> mode) 142 (M<sup>+</sup>, 5%), 114 (16), 99 (70), 84 (100), 71 (37) and 43 (27) (Found M<sup>+</sup> 142.0996. C<sub>8</sub>H<sub>14</sub>O<sub>2</sub> requires 142.0994).

91% ee, determined by chiral GC (see supporting information).

#### (4R,5R)-5-n-Butyl-4-methyl dihydro-furan-2-one 15<sup>5</sup>

As for general procedure A. Reaction of pentanal (14 µl, 0.12 mmol, 1.0 equiv.) with (1*R*,2*S*)-*N*-Wang bound ephedrinyl crotonate **3** (0.39 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40–60 °C)) gave **15** (9.6 mg, 50%) as a pale yellow oil:  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.46–4.41 (1H, m, CHO), 2.70 (1H, dd, *J* 16.9, 7.8, 1H from CH<sub>2</sub>C=O), 2.63–2.54 (1H, m, CHCH<sub>3</sub>), 2.21 (1H, dd, *J* 16.9, 3.9, 1H from CH<sub>2</sub>C=O), 1.70–1.61 (1H, m, 1H from CH<sub>2</sub>), 1.58–1.45 (2H, m, CH<sub>2</sub>), 1.42–1.36 (3H, m, CH<sub>2</sub>), 0.95 (3H, d, *J* 7.0, CHCH<sub>3</sub>), 0.93 (3H, t, *J* 7.3, CH<sub>3</sub>CH<sub>2</sub>);  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 176.9 (C=O), 83.7 (CHO), 37.6 (CH<sub>2</sub>C=O), 33.0 (CHCH<sub>3</sub>), 29.6 (CH<sub>2</sub>), 28.1 (CH<sub>2</sub>), 22.5 (CH<sub>2</sub>), 13.9 (CH<sub>3</sub>), 13.8 (CH<sub>3</sub>); *m/z* (CI<sup>+</sup> mode) 157 ((M + H)<sup>+</sup>, 100%), 113 (6), 81 (8) (Found (M + H)<sup>+</sup> 157.1231. C<sub>9</sub>H<sub>17</sub>O<sub>2</sub> requires 157.1229).

91% ee, determined by chiral GC (see supporting information).

#### (4R,5S)-5-Cyclopentyl-4-methyl dihydro-furan-2-one 16

As for general procedure A. Reaction of cyclopentane carboxaldehyde (19.6 µl, 0.18 mmol, 1 equiv.) with (1R,2S)-N-Wang bound ephedrinyl crotonate 3 (0.57 mmol, 3.2 equiv.) and purification by column chromatography on silica (eluting with 20% EtOAc/ petroleum ether (40-60 °C)) gave 16 (13 mg, 43%) as a colourless oil: v<sub>max</sub> (neat)/cm<sup>-1</sup> 2958s, 1775s (C=O), 1454m, 1421m, 1383m, 1326m;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.15 (1H, dd, J 10.1, 5.1, CHO), 2.74 (1H, dd, J 17.0, 7.6, 1H from CH<sub>2</sub>C=O), 2.58–2.52 (1H, m, CHCH<sub>3</sub>), 2.21 (1H, dd, J 17.0, 1.9, 1H from CH<sub>2</sub>C=O), 2.16-2.12 (1H, m, CH), 1.96–1.91 (1H, m, 1H from CH<sub>2</sub>), 1.79–1.73 (1H, m, 1H from CH<sub>2</sub>), 1.68–1.56 (4H, m, CH<sub>2</sub>), 1.50–1.43 (1H, m, 1H from CH<sub>2</sub>), 1.26–1.17 (1H, m, 1H from CH<sub>2</sub>), 1.03 (3H, d, J 7.1, CH<sub>3</sub>); δ<sub>C</sub> (100 MHz, CDCl<sub>3</sub>) 177 (C=O), 88.2 (CHO), 39.5 (CH), 38.5 (CH<sub>2</sub>C=O), 32.8 (CHCH<sub>3</sub>), 30.9 (CH<sub>2</sub>), 27.8 (CH<sub>2</sub>), 25.4 (CH<sub>2</sub>), 25.3 (CH<sub>2</sub>), 14.2 (CH<sub>3</sub>); *m/z* (CI<sup>+</sup> mode, isobutane) 169 ((M + H)<sup>+</sup>, 100%) (Found  $(M + H)^+$  169.1231.  $C_{10}H_{17}O_2$  requires 169.1229).

88% ee, determined by chiral GC (see supporting information)

# (5R,4R)-5-n-Butyl-4,5-dimethyl dihydrofuran-2-one 17

As for general procedure A except, due to the lower reactivity of the ketone, 2-hexanone (3 equiv.) was reacted with the resin (1 equiv.). Reaction of 2-hexanone (78 µl, 0.62 mmol, 3.0 equiv.) with (1R,2S)-N-Wang bound ephedrinyl crotonate 3 (0.21 mmol, 1.0 equiv.) and purification by column chromatography on silica gel (eluting with 20% EtOAc/petroleum ether (40-60 °C)) gave 17 (12.7 mg, 36%) as a pale yellow oil and as an inseparable 3:1 mixture of diastereoisomers: For the major diastereoisomer:  $v_{\rm max}$  (neat)/cm<sup>-1</sup> 2959s, 2872s, 1772s (C=O), 1460m, 1382m;  $\delta_{\rm H}$ (400 MHz, CDCl<sub>3</sub>) 2.59 (1H, dd, *J* 16.6, 7.5, 1H from CH<sub>2</sub>C=O). 2.39-2.24 (2H, m, 1H from CH<sub>2</sub>C=O and CHCH<sub>3</sub>), 1.66-1.31 (6H, m, 3 × CH<sub>2</sub>), 1.39 (3H, s, CH<sub>3</sub>), 1.00 (3H, d, J 6.8, CHCH<sub>3</sub>), 0.93 (3H, t, J 7.1, CH<sub>2</sub>CH<sub>3</sub>); δ<sub>C</sub> (100 MHz, CDCl<sub>3</sub>) 176.1 (C=O), 88.7 (CO), 41.3 (CHCH<sub>3</sub>), 36.9 (CH<sub>2</sub>C=O), 34.5 (CH<sub>2</sub>), 25.6 (CH<sub>2</sub>), 24.3 (CH<sub>3</sub>), 23.2 (CH<sub>2</sub>), 14.0 (CH<sub>3</sub>), 13.9 (CH<sub>3</sub>); m/z (CI<sup>+</sup> mode, isobutene) 171 (M + H<sup>+</sup>, 100%), 113 (3), 85 (5), 71 (4) (Found (M + H)<sup>+</sup> 171.1384. C<sub>10</sub>H<sub>19</sub>O<sub>2</sub> requires 171.1385).

*Major* (89% ee) and *minor* (76% ee), the *cis*-product was assumed to be the major diastereoisomer, determined by chiral GC (see supporting information).

#### 5-Methyl-hexane-1,5-diol 1817

To a solution of  $\delta$ -valerolactone (1.46 g, 14.6 mmol, 1.0 equiv.) in THF at -78 °C was added MeLi (1.5 M in Et<sub>2</sub>O, 29.2 mL, 43.8 mmol, 3.0 equiv.) dropwise. The solution was allowed to stir at -78 °C for 30 min before being allowed to warm to room temperature overnight. The reaction was then quenched by dropwise addition of acetic acid (1.4 mL, 25.0 mmol, 1.7 equiv.) to the reaction solution. A white suspension resulted and this was stirred at room temperature for 4 h before filtration. The precipitate was washed with THF ( $3 \times 50$  mL) and the combined filtrate and washings were concentrated in vacuo to afford a crude oil. Purification by dry flash chromatography on silica (90% EtOAc/petroleum ether (40-60 °C) to 20% MeOH/EtOAc) gave 18 (1.27 g, 66%) as a colourless oil: v<sub>max</sub>(neat)/cm<sup>-1</sup> 3350s, 2968s, 2934s, 2867s, 1468m, 1379m, 1201m, 1155m;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 3.66 (2H, t, J 6.4, CH<sub>2</sub>OH), 1.74 (2H, br s, 2 × OH), 1.58 (2H, quintet, J 6.4,  $CH_2CH_2OH$ ), 1.52–1.41 (4H, m, 2 ×  $CH_2$ ), 1.22 (6H, s, 2 ×  $CH_3$ );  $\delta_{\rm C}$  (100 MHz, CDCl<sub>3</sub>) 71.0 (*C*), 62.6 (*C*H<sub>2</sub>OH), 43.3 (*C*H<sub>2</sub>), 33.0  $(CH_2CH_2OH)$ , 29.2 (2 ×  $CH_3$ ), 20.4 ( $CH_2$ ); m/z (FAB<sup>+</sup> mode) 133 ((M + H)<sup>+</sup>, 25%), 115 (100), 97 (28), 60 (8), 56 (8) (Found (M + H)<sup>+</sup> 133.1229. C7H17O2 requires 133.1230).

# 5-Methyl-5-trimethylsilyloxy-hexan-1-ol 19

To a solution of diol **18** (90 mg, 0.69 mmol, 1.0 equiv.) in  $CH_2Cl_2$  (8 mL) was added triethylamine (3.8 mL, 27.3 mmol, 40 equiv.) and trimethylsilyl chloride (1.74 mL, 13.7 mmol, 20 equiv.) and the reaction was allowed to stir at room temperature for 3 days. The reaction

was quenched with water (2 mL), the organic layer was separated and the aqueous layer was extracted with  $CH_2Cl_2$  (2 × 15 mL). The combined organics were dried (Na2SO4) and concentrated in vacuo to afford the crude product. Purification by column chromatography on silica (30% EtOAc/petroleum ether (40-60 °C)) gave a colourless oil (171.0 mg, 90%) that was shown to be composed mainly of bis-TMS protected compound by 1H NMR. To this product (144 mg, 0.52 mmol, 1 equiv.) was added potassium carbonate (144 mg, 1.04 mmol, 2.0 equiv.) and anhydrous methanol (2.0 mL). The solution was stirred at 0 °C for 2.5 h before quenching with acetic acid (0.06 mL, 1.04 mmol, 2.0 equiv.) and water (2 mL). The aqueous solution was then extracted with  $CH_2Cl_2$  (3 × 15 mL) and the combined organics were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in vacuo to afford 19 (108 mg, 92% overall) as a colourless oil, which was used without further purification:  $v_{max}(neat)/cm^{-1}$  3348s, 1468m, 1379m, 1201m, 1155m; δ<sub>H</sub> (400 MHz, CDCl<sub>3</sub>) 3.66 (2H, t, J 6.5, CH<sub>2</sub>OH), 1.57 (2H, quintet, J 6.5, CH<sub>2</sub>CH<sub>2</sub>OH), 1.54–1.37 (4H, m, 2 × CH<sub>2</sub>), 1.21 (6H, s,  $2 \times CH_3$ ), 0.11 (9H, s,  $3 \times CH_3$ Si);  $\delta_C$  (100 MHz, CDCl<sub>3</sub>) 74.0 (C), 63.0 (CH<sub>2</sub>OH), 44.5 (CH<sub>2</sub>), 33.2 (CH<sub>2</sub>CH<sub>2</sub>OH), 29.8 (2 × CH<sub>3</sub>), 20.5 (CH<sub>2</sub>), 2.6 (3 × CH<sub>3</sub>Si); m/z (FAB<sup>+</sup> mode) 189  $((M - CH_3)^+, 7\%), 131 (100), 84 (12), 73 (50), 55 (16), 43 (15)$ (Found (M - CH<sub>3</sub>)<sup>+</sup> 189.1311. C<sub>9</sub>H<sub>21</sub>O<sub>2</sub>Si requires 189.1308).

# 5-Methyl-5-trimethylsilyloxy-hexanal 20

To Dess-Martin periodinane (470 mg, 1.11 mmol, 1.1 equiv.) in CH<sub>2</sub>Cl<sub>2</sub> (4.6 mL) was added alcohol 19 (206 mg, 1.01 mmol, 1.0 equiv.) in CH2Cl2 (2 mL) via cannula. The light orange coloured solution was then stirred at room temperature for 2 h before dilution with diethyl ether (25 mL) and addition to an aqueous saturated NaHCO<sub>3</sub> (1 mL)/aqueous saturated Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (8 mL) solution. The mixture was stirred for a few minutes before the organic layer was separated. The organic layer was extracted with aqueous NaHCO<sub>3</sub> (10 mL) and water (12 mL). The organic layer was then dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in vacuo to afford 20 (199.6 mg, 98%) as a pale orange oil which was used without further purification:  $v_{max}$ (neat)/cm<sup>-1</sup> 2960s, 2715w, 1728s (C=O), 1460m, 1412m, 1383m, 1365m, 1250s;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 9.67 (1H, t, J 1.8, CH=O), 2.33 (2H, td, J 7.3, 1.8, CH<sub>2</sub>CH=O), 1.75-1.67 (2H, m, CH<sub>2</sub>CH<sub>2</sub>CH=O), 1.46-1.39 (2H, m, CH<sub>2</sub>), 1.23 (6H, s,  $2 \times CH_3$ ), 0.11 (9H, s,  $3 \times CH_3$ Si);  $\delta_C$  (100 MHz, CDCl<sub>3</sub>) 202.9 (CH=O), 73.6 (C), 44.3 (CH<sub>2</sub>CH=O), 44.2 (CH<sub>2</sub>), 29.8  $(2 \times CH_3)$ , 17.1 (CH<sub>2</sub>CH<sub>2</sub>CH=O), 2.6 (3 × CH<sub>3</sub>Si); m/z (EI<sup>+</sup> mode) 187 ((M - CH<sub>3</sub>)<sup>+</sup>, 8%), 131 (100), 129 (18), 95 (12), 73 (65), 69 (23) (Found  $(M - CH_3)^+$  187.1154. C<sub>9</sub>H<sub>19</sub>O<sub>2</sub>Si requires 187.1154).

#### 5-(4-Hydroxy-4-methyl-pentyl) dihydro-furan-2-one 21

As for general procedure A. Reaction of aldehyde 20 (48.6 mg, 0.24 mmol, 1.0 equiv.) with (1R,2S)-N-Wang bound ephedrinyl acrylate 2 (0.77 mmol, 3.2 equiv.) and purification by column chromatography on silica gel (30% EtOAc/petroleum ether (40-60 °C) to ethyl acetate) gave 21 (22.6 mg, 50%) as a pale yellow oil: [a]<sub>D</sub> +29.5 (c 0.01 in MeOH). lit.<sup>16</sup> [a]<sub>D</sub> +42.0 (c 1.0 in MeOH); v<sub>max</sub>(neat)/cm<sup>-1</sup> 3450s, 2968s, 1767s (C=O), 1462m, 1421m, 1363m, 1298w;  $\delta_{\rm H}$  (400 MHz, CDCl<sub>3</sub>) 4.55–4.48 (1H, m, CHOC(O)), 2.55 (2H, m, CH2CH2C(O)), 2.38-2.30 (1H, m, 1H from CH2C(O)), 1.92-1.85 (1H, m, 1H from CH2C(O)), 1.77-1.74 (1H, m, 1H from CH<sub>2</sub>), 1.73–1.48 (5H, m, 1H from CH<sub>2</sub>, 2 × CH<sub>2</sub>), 1.65 (6H, s,  $2 \times CH_3$ );  $\delta_C$  (100 MHz, CDCl<sub>3</sub>) 177.2 (C=O), 80.9 (CH), 70.8 (C), 43.4 (CH<sub>2</sub>), 36.1 (CH<sub>2</sub>), 29.4 (CH<sub>3</sub>), 29.2 (CH<sub>3</sub>), 28.8 (CH<sub>2</sub>C(O)), 28.0 (CH<sub>2</sub>CH<sub>2</sub>C(O)), 20.3 (CH<sub>2</sub>); m/z (CI<sup>+</sup> mode, isobutene) 187 (M<sup>+</sup>, 5%), 169 (100), 151 (10), 69 (5) (Found M<sup>+</sup> 187.1337. C<sub>10</sub>H<sub>18</sub>O<sub>3</sub> requires 187.1334).

73% ee, determined by chiral GC (see supporting information).

#### Acknowledgements

We thank the Engineering and Physical Research Council (EPSRC) (GR/R72242/01, N. J. K.), GlaxoSmithKline (CASE award,

P. C. H.) and the University of Glasgow (University Scholarship, P. C. H.) for financial support. We also thank AstraZeneca for the award of a Strategic Research Grant (D. J. P.) and Pfizer for untied funding (D. J. P.).

# References

- For recent reviews on solid phase organic synthesis see: (a) S. Booth, P. H. H. Hermkens, H. C. J. Ottenheijm and D. C. Rees, *Tetrahedron*, 1998, **54**, 15385; (b) R. C. D. Brown, *J. Chem. Soc.*, *Perkin Trans. 1*, 1998, 3293; (c) R. E. Sammelson and M. J. Kurth, *Chem. Rev.*, 2001, **101**, 137; (d) F. Guillier, D. Orain and M. Bradley, *Chem. Rev.*, 2000, **100**, 2091; (e) V. Krchnák and M. W. Holladay, *Chem. Rev.*, 2002, **102**, 61; (f) R. E. Dolle, *J. Comb. Chem.*, 2003, **5**, 693.
- 2 S. V. Ley, I. R. Baxendale, R. N. Bream, P. S. Jackson, A. G. Leach, D. A. Longbottom, M. Nesi, J. S. Scott, R. I. Storer and S. J. Taylor, *J. Chem. Soc., Perkin Trans.* 1, 2000, 3815.
- 3 For an interesting discussion of "resin-capture–release" processes, see: A. Kirschning, H. Monenschein and R. Wittenberg, *Chem. Eur. J.*, 2000, **6**, 4445.
- 4 (*a*) For a review of asymmetric solid-phase synthesis using supported chiral auxiliaries see: C. W. Y. Chung and P. H. Toy, *Tetrahedron: Asymmetry*, 2004, **15**, 387; (*b*) For a review on solid-supported catalysts and reagents which includes immobilised, chiral reagents for asymmetric synthesis see: C. A. McNamara, M. J. Dixon and M. Bradley, *Chem. Rev.*, 2002, **102**, 3275.
- 5 S. Fukuzawa, K. Seki, M. Tatsuzawa and K. Mutoh, J. Am. Chem. Soc., 1997, 119, 1482.
- 6 (a) F. McKerlie, D. J. Procter and G. Wynne, *Chem. Commun.*, 2002, 584; (b) L. A. McAllister, S. Brand, R. de Gentile and D. J. Procter, *Chem. Commun.*, 2003, 2380.
- 7 (a) P. C. Hutchison, T. D. Heightman and D. J. Procter, *Org. Lett.*, 2002,
  4, 4583; (b) P. C. Hutchison, T. D. Heightman and D. J. Procter, *J. Org. Chem.*, 2004, 69, 790.
- 8 For a preliminary account of this work, see: N. J. Kerrigan, P. C. Hutchison, T. D. Heightman and D. J. Procter, *Chem. Commun.*, 2003, 1402.
- 9 J. M. J. Fréchet, E. Bald and P. Lecavalier, J. Org. Chem., 1986, 51, 3462.
- 10 (1*R*,2*S*)-*O*-Benzyl ephedrine was prepared in 92% according to the procedure of Welch (see supporting information): J. Näslund and C. J. Welch, *Tetrahedron: Asymmetry*, 1991, **2**, 1123.
- 11 The loading of **1** was determined by esterification with thiophene carbonyl chloride followed by sulfur elemental analysis of the resin. See supporting information and ref. 7.
- 12 S. Fukuzawa, A. Nakanishi, T. Fujinami and S. Sakai, J. Chem. Soc., Perkin Trans. 1, 1988, 1669. See also ref. 5.
- 13 Few intermolecular radical additions to immobilised acceptors have been reported. For reviews of radical reactions on solid phase, see: (a) A. Ganesan, in *Radicals in Organic Synthesis*, ed. P. Renaud and M. P. Sibi, Wiley-VCH, Weinheim, 2001, vol. 2, p. 81; (b) A. Ganesan and M. P. Sibi, in *Handbook of Combinatorial Chemistry*, ed. K. C. Nicolaou, R. Hanko, W. Hartwig, Wiley-VCH, Weinheim, 2002, vol. 1, p. 227. For recent examples, see; (c) S. Caddick, D. Hamza and S. N. Wadman, *Tetrahedron Lett.*, 1999, **40**, 7285; (d) X. Zhu and A. Ganesan, *J. Comb. Chem.*, 1999, **1**, 157; (e) H. Miyabe, Y. Fujishima and T. Naito, *J. Org. Chem.*, 1999, **64**, 2174; (f) H. Miyabe, C. Konishi and T. Naito, *Org. Lett.*, 2000, **2**, 1443; (g) S. Caddick, D. Hamza, S. N. Wadman and J. D. Wilden, *Org. Lett.*, 2002, **4**, 1775; (h) D. C. Harrowven, P. J. May and M. Bradley, *Tetrahedron Lett.*, 2003, **44**, 503.
- 14 The diastereoselectivity was confirmed by comparison with Fukuzawa's  $\,^1\!\mathrm{H}$  NMR data (ref. 5) and by NOE studies.
- 15 The efficient reduction of α,β-unsaturated esters with SmI<sub>2</sub> usually requires the use of an additive such as DMA (*N*,*N*-dimethyl acetamide) or HMPA, see (*a*) J. Inanaga, S. Sakai, Y. Handa, M. Yamaguchi and Y. Yokoyama, *Chem. Lett.*, 1991, 2117; (*b*) A. Cabrera and H. Alper, *Tetrahedron Lett.*, 1992, **33**, 5007.
- 16 C. Maul, I. Sattler, M. Zerlin, C. Hinze, C. Koch, A. Maier, S. Grabley and R. Thiericke, J. Antibiot., 1999, 52, 1124.
- 17 A. S. Hernández and J. C. Hodges, J. Org. Chem., 1997, 62, 3153.
- 18 D. B. Dess and J. C. Martin, J. Org. Chem., 1983, 48, 4155.
- 19 The enantiomeric excess of lactones was determined by chiral GC (Supelco beta dex<sup>™</sup> 120 fused silica capillary column: 30 m) and comparison with authentic, racemic samples. (See the Electronic Supporting Information).
- 20 M. Utaka, H. Watabu and A. Takeda, J. Org. Chem., 1987, 52, 4363.
- 21 J. Rojo, M. García and J. C. Carretero, Tetrahedron, 1993, 49, 9787.