

## Laser-Induced Photoionization of Two-Electron-Reduced Viologen


Satoshi USUI,<sup>†</sup> Hiroshi INOUE, Hitoshi ISHIDA, and Katsutoshi OHKUBO\*

Department of Applied Chemistry, Faculty of Engineering, Kumamoto University, Kurokami, Kumamoto 860

Laser photolysis of two-electron-reduced 1,1'-bis(3-sulfonatepropyl)-4,4'-bipyridinium (PVS<sup>2-</sup>) produced by the reduction of PVS<sup>0</sup> with Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> in the cationic micelles of hexadecyltrimethylammonium chloride induced the photoionization of PVS<sup>2-</sup> to generate comparable amounts of PVS<sup>•-</sup> and hydrated electrons. Pyrene with a considerably negative reduction potential of -2.09 vs. SCE was observed to be reduced in the system.

Although viologens have been widely utilized both as an electron-transfer catalyst for reduction of various organic compounds<sup>1)</sup> and as an electron mediator in photochemical conversion of light energy to hydrogen,<sup>2)</sup> the photochemistry of two-electron-reduced viologen has hitherto been a subject of only limited investigation. The present report first describes the novel laser-induced photoionization of a two-electron-reduced viologen derivative (1,1'-bis(3-sulfonatepropyl)-4,4'-bipyridinium) in a cationic micelle in the presence or absence of electron acceptors.

1,1'-Bis(3-sulfonatepropyl)-4,4'-bipyridinium (PVS<sup>0</sup>) was synthesized according to the method described by Matsuo *et al.*<sup>3)</sup> The two-electron reduction of PVS<sup>0</sup> (50  $\mu\text{mol dm}^{-3}$ ) was carried out by the addition of more than 20 times excess Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> in a degassed hexadecyltrimethylammonium chloride (CTAC) solution (10 mmol  $\text{dm}^{-3}$ ) at pH 10.0 (carbonate buffer). The reduced viologens were photolyzed in a quartz cell (1 cm optical path length) by the 3rd harmonic light (355 nm, pulse duration 10 ns, 80 mJ / pulse) of a Nd:YAG laser (Continuum NY-60). The transient absorption



<sup>†</sup> Present address: Institute for Molecular Science, Myodaiji, Okazaki, 444.

signals were recorded by a digitizing oscilloscope (Phillips PM-3350A) and were analyzed with a microcomputer (NEC PC-9801 DA).

When  $\text{Na}_2\text{S}_2\text{O}_4$  was added to the aqueous solution of  $\text{PVS}^0$ , the solution exhibited a typical absorption spectrum of one-electron-reduced viologen<sup>4)</sup> as shown in Fig. 1a. The addition of  $\text{Na}_2\text{S}_2\text{O}_4$  to the CTAC micellar solution of  $\text{PVS}^0$ , on the other hand, gave a characteristic absorption spectrum assigned to the two-electron-reduced viologen ( $\text{PVS}^{2-}$ )<sup>4)</sup> without the absorption due to the one-electron-reduced  $\text{PVS}^0$  ( $\text{PVS}^{\bullet-}$ ) (Fig. 1b). The quantitative two-electron reduction of  $\text{PVS}^0$  in the presence of excess  $\text{Na}_2\text{S}_2\text{O}_4$  was thus possible in the hydrophobic domain of CTAC micelle; in the micellar solution, the  $\text{PVS}^{2-}$  is formed by the disproportionation reaction of  $\text{PVS}^{\bullet-}$ .<sup>5)</sup> On the other hand, such electron acceptors as pyrene (Py) in the micellar solution was not reduced by  $\text{Na}_2\text{S}_2\text{O}_4$  (Fig. 1c).

In a laser photolysis of a micellar solution containing  $\text{PVS}^{2-}$  and Py at 355 nm, a transient absorption spectrum with the absorption maxima at 493 and 603 nm was obtained immediately after the laser excitation. The transient absorptions at 493 and 603 nm are reasonably

assigned to the absorptions due to pyrene anion radical ( $\text{Py}^{\bullet-}$ )<sup>6)</sup> and  $\text{PVS}^{\bullet-}$ , respectively; the electron transfer from the photoexcited state of  $\text{PVS}^{2-}$  ( $^*\text{PVS}^{2-}$ ) to Py was found to be completed within 0.5  $\mu\text{s}$  after the laser excitation. Since the laser photolysis (at 355 nm) of micellar solution containing Py did not produce the




Fig. 1. Absorption spectra obtained by the reduction of  $\text{PVS}^0$  ( $50 \mu\text{mol dm}^{-3}$ ) with  $\text{Na}_2\text{S}_2\text{O}_4$ : (a) aqueous solution; (b)  $[\text{CTAC}] = 10 \text{ mmol dm}^{-3}$ ; (c)  $[\text{CTAC}] = 10 \text{ mmol dm}^{-3}$ ,  $[\text{Py}] = 100 \mu\text{mol dm}^{-3}$ .




Fig. 2. Transient absorption spectra obtained by the laser photolysis of  $\text{PVS}^{2-}$  ( $50 \mu\text{mol dm}^{-3}$ ) in the presence of Py.

appreciable amount of the photoexcited-state Py (see Fig. 1), the  $\text{Py}^{\bullet-}$  is not generated by the reaction of  $^*\text{Py}$  and  $\text{Na}_2\text{S}_2\text{O}_4$ . The concentration of  $\text{Py}^{\bullet-}$  obtained at 1  $\mu\text{s}$  after the laser photolysis increased with increasing the Py concentration and reached to a maximum value of  $5.5 \mu\text{mol dm}^{-3}$  in the range of  $[\text{Py}] > 100 \mu\text{mol dm}^{-3}$  (Fig. 2). The amount of  $\text{PVS}^{\bullet-}$ , on the other hand, was little affected by the Py concentration and gave constant value of  $5.3 \pm 0.6 \mu\text{mol dm}^{-3}$ . Since the concentration of  $\text{Py}^{\bullet-}$  did not exceed that of  $\text{PVS}^{\bullet-}$  even at high Py concentration, an electron transfer from  $^*\text{PVS}^{2-}$  to Py in micelle thus generated the  $\text{Py}^{\bullet-}$  species.

From the comparison with the reduction potentials of Py (-2.09 vs. SCE in DMF),<sup>7)</sup>  $\text{PVS}^{2-}$  was found to realize a laser-induced one-electron reduction of organic compound with a considerably negative reduction potential lower than -2.09 V vs. SCE.

It is of interest and significant here to investigate the photochemical behavior of  $^*\text{PVS}^{2-}$  produced by the laser photolysis of  $\text{PVS}^{2-}$  in the CTAC micelle in the absence of electron acceptors such as Py. Notably, laser flash photolysis of  $\text{PVS}^{2-}$  in the CTAC micelle without Py exhibited the absorption maxima of  $\text{PVS}^{\bullet-}$  at 603 nm and the bleaching at 400 nm due to the disappearance of  $\text{PVS}^{2-}$  (Fig. 3).

Since the photochemical system in Fig. 3 excludes the electron transfer reaction from  $^*\text{PVS}^{2-}$  to  $\text{PVS}^0$  which is quantitatively reduced to be  $\text{PVS}^{2-}$  by  $\text{Na}_2\text{S}_2\text{O}_4$  in CTAC, the possibility of photoionization is worthy to notice. In the case of the photoionization of  $\text{PVS}^{2-}$ , the photo-ejected electron should be observed as a hydrated electron ( $e_{\text{aq}}^-$ ) since there is no electron acceptor in the present system. In fact, the transient absorption spectrum of  $e_{\text{aq}}^-$  around 720 nm<sup>8)</sup> was observed at 200 ns after the laser excitation. The  $e_{\text{aq}}^-$  concentration of  $2.6 \mu\text{mol dm}^{-3}$  at the delay time of 200 ns was found to be comparable to that of  $\text{PVS}^{\bullet-}$  ( $4.3 \mu\text{mol dm}^{-3}$ ),<sup>9)</sup> the less  $e_{\text{aq}}^-$  concentration than that of  $\text{PVS}^{\bullet-}$  is probably due to the disappearance of  $e_{\text{aq}}^-$  by a reaction with water,<sup>10)</sup> which is exhibited by the rapid decay of the transient absorption at 720 nm in the time domain of 1  $\mu\text{s}$ .

In conclusion, the present laser photolysis of the two-electron-reduced viologen resulted in the novel photoionization of reduced viologen, which was confirmed by the spectroscopic observation of the hydrated




Fig. 3. Transient absorption spectra obtained by the laser photolysis of  $\text{PVS}^{2-}$  ( $50 \mu\text{mol dm}^{-3}$ ) in the CTAC micelle ( $10 \text{ mmol dm}^{-3}$ ) without Py.

electrons.

## References

- 1) H. Tomioka, K. Ueda, H. Ohi, and Y. Izawa, *Chem. Lett.*, **1986**, 1359; T. Endo, Y. Saotome, and M. Ogasawara, *J. Am. Chem. Soc.*, **106**, 1124 (1984); T. Endo, Y. Saotome, and M. Ogasawara, *Tetrahedron Lett.*, **26**, 4525 (1985); K. K. Park, C. W. Lee, and S. Y. Choi, *J. Chem. Soc., Perkin Trans. 1*, **1992**, 601.
- 2) J. H. Fendler, "Membrane Mimetic Chemistry," Wiley-Interscience, New York (1982); K. Kalyanasundaram, *Coord. Chem. Rev.*, **46**, 159 (1981); D. G. Whitten, *Acc. Chem. Res.*, **13**, 83 (1980); M. Grätzel, "Energy Resources through Photochemistry and Catalysts," Academic Press, New York (1983).
- 3) T. Nagamura, T. Kurihara, T. Matsuo, M. Sumitani, and K. Yoshihara, *J. Phys. Chem.*, **86**, 4368 (1982).
- 4) T. Watanabe and K. Honda, *J. Phys. Chem.*, **86**, 2617 (1982); M. Grätzel and J. Moser, *Proc. Natl. Acad. Sci. U.S.A.*, **80**, 3129 (1983).
- 5) R. Maidan, Z. Goren, J. Y. Becker, and I. Willner, *J. Am. Chem. Soc.*, **106**, 6217 (1984).
- 6) T. Aoyama, Y. Yamamoto, and K. Hayashi, *J. Chem. Soc., Faraday Trans. 1*, **85**, 3353 (1989).
- 7) R. Pointeau, *Ann. Chim. (Paris)*, **7**, 669 (1962).
- 8) A. Horváth, J. Szöke, and L. Wojnárovits, *Inorg. Chem. Acta*, **179**, 97 (1991).
- 9) The concentrations of  $e^-_{aq}$  and  $PVS^{•-}$  were calculated by using  $\epsilon_{603\text{ nm}} = 13000$  and  $\epsilon_{720\text{ nm}} = 20300\text{ mol}^{-1}\text{ dm}^3\text{ cm}^{-1}$  for the former<sup>8)</sup> and  $\epsilon_{603\text{ nm}} = 13800$  and  $\epsilon_{720\text{ nm}} = 2200\text{ mol}^{-1}\text{ dm}^3\text{ cm}^{-1}$  for the latter.<sup>4)</sup>
- 10) J. H. Fendler, G. W. Bogan, E. J. Fendler, G. A. Infante, and P. Jirathana, "Reaction Kinetics in Micelles," ed by E. Cordes, Plenum Press, New York-London (1973).

( Received September 14, 1993 )