in the reactants, the polar media must stabilize the reactants more efficiently than the TS. Consequently, the hydrolysis rate decreases with increasing water density.

In summary, we have demonstrated that the hydrolysis of esters takes place exclusively by ionic mechanisms, whereby hydroxide ions dissociated from SCW catalyze the reaction. This mechanism is dominant in the initial stage of the reaction, since the concentration of the generated carboxylic acid is low. As the concentration of carboxylic acid increases with the progress of the reaction, the mechanism would change to the proton-catalyzed reaction if the concentration of substrates were sufficiently high.<sup>[9]</sup> Our results clearly indicate that one must consider the involvement of hydroxide ions in other reactions in SCW, because dissociation of water produces the same amount of protons and hydroxide ions.

Received: September 6, 2001 [Z17863]

- a) O. Kajimoto, *Chem. Rev.* **1999**, *99*, 355; b) P. E. Savage, *Chem. Rev.* **1999**, *99*, 603; c) D. Bröll, C. Kaul, A. Krämer, A. Krammer, T. Richter, M. Jung, H. Vogel, P. Zehner, *Angew. Chem.* **1999**, *111*, 3180; *Angew. Chem. Int. Ed.* **1999**, *38*, 2998.
- [2] a) Green Chemistry. Frontiers in Benign Chemical Syntheses and Processes (Eds.: P. T. Anastas, T. C. Williamson), Oxford University Press, Oxford, 1998; b) C.-J. Li, T.-H. Chan, Organic Reactions in Aqueous Media, Wiley, New York, 1997.
- [3] a) M. M. Hoffmann, M. S. Conradi, J. Am. Chem. Soc. 1997, 119, 3811;
   b) Y. Ikushima, K. Hatakeda, N. Saito, M. Arai, J. Chem. Phys. 1998, 108, 5855.
- [4] M. Uematsu, E. U. Franck, J. Phys. Chem. Ref. Data 1980, 9, 1291.
- [5] a) O. Sato, Y. Ikushima, T. Yokoyama, J. Org. Chem. 1998, 63, 9100;
  b) Y. Ikushima, K. Hatakeda, O. Sato, T. Yokoyama, M. Arai, Angew. Chem. 1999, 111, 3087; Angew. Chem. Int. Ed. 1999, 38, 2910; c) Y. Ikushima, K. Hatakeda, O. Sato, T. Yokoyama, M. Arai, J. Am. Chem. Soc. 2000, 122, 1908.
- [6] a) Y. Ikushima, K. Hatakeda, O. Sato, T. Yokoyama, M. Arai, Angew. Chem. 2001, 113, 216; Angew. Chem. Int. Ed. 2001, 40, 210; b) the Canizzarro reaction in subcritical water has been also reported: Y. Tsujino, C. Wakai, N. Matsubayashi, M. Nakahara, Chem. Lett. 1999, 287.
- [7] a) M. A. Dubois, J. F. Dozol, C. Massiani, M. Ambrosio, *Ind. Eng. Chem. Res.* 1996, 35, 2743; b) S. Park, E. F. Gloyna, *Fuel* 1997, 76, 999;
  c) M. Watanabe, H. Hirakoso, S. Sawamoto, T. Adshiri, M. Arai, *J. Supercrit. Fluids* 1998, *13*, 247; d) Y. Suzuki, H. Tagaya, T. Asou, J. Kadokawa, K. Chiba, *Ind. Eng. Chem. Res.* 1999, *38*, 1391; e) C. Fromonteil, P. Bardelle, F. Cansell, *Ind. Eng. Chem. Res.* 2000, *39*, 922.
- [8] S. D. Iyer, M. T. Klein, J. Supercrit. Fluids 1997, 10, 191. It is worth mentioning that both carboxylic acid and ammonia generated by the hydrolysis are suggested as possible catalysts.
- [9] P. Krammer, H. Vogel, J. Supercrit. Fluids 2000, 16, 189.
- [10] T. Xiang, K. P. Johnston, J. Solution Chem. 1997, 26, 13.
- [11] The observed decrease in the hydrolysis rate near the critical point may also be attributed to the decrease in the ionic product of water near the critical point. Similar results have been reported for the hydrolysis of ethers: J. D. Taylor, J. I. Steinfeld, J. W. Tester, *Ind. Eng. Chem. Res.* 2001, 40, 67.
- [12] N. J. Turro, I. R. Gould, B. H. Baretz, *J. Phys. Chem.* 1983, 87, 531.
  [13] Cylindrical stainless steel (SUS316) or quartz tubes were used as reaction vessel. Similar results were obtained regardless of the vessel.
- [14] The reaction also afforded benzaldehyde and actophenone in 2-4% yield. Ethylbenzene was unstable under the reaction conditions and decomposed into various uncharacterized compounds. Stabilities of alkyl-substituted benzenes have been already studied: J. Yu, S. Eser, Ind. Eng. Chem. Res. 1998, 37, 4591.
- [15] A flow reactor equipped with an HPLC pump (JASCO PU-1580) and a pressure regulator (JASCO 880-81), connected by SUS316 tube (2.2 mm inner diameter), was used. Similar equipment has already been reported: B. M. Kabyemela, T. Adschiri, R. M. Malauan, K. Arai, *Ind. Eng. Chem. Res.* **1997**, *36*, 1552.

- [16] W. F. Marshall, E. U. Franck, J. Phys. Chem. Ref. Data 1981, 10, 295.
- [17] An alternative but less likely mechanism is the direct attack of a water molecule on the ester. Correlation of  $k_1$  with  $k'_1$  and the concentration of water suggests that about four molecules of water participate in the transition state ( $k_1 = k'_1$ [H<sub>2</sub>O]<sup>3.7</sup>).
- [18] Addition of sodium hydroxide  $(4.8 \times 10^{-7} \text{ mol L}^{-1})$  enhanced the rate of hydrolysis by a factor of about 2.5. However, the lack of data on the dissociation constant of sodium hydroxide in SCW and the difficulties in obtaining an accurate value of  $k_3$  prevented our making detailed kinetic analyses.
- [19] Because of the low solubility of 1 in water, we could not directly compare the concentration effects.
- [20] E. K. Euranto in *The Chemistry of Carboxylic Acids and Esters* (Ed.: S. Patai), Interscience Publishers, London, **1969**, pp. 505–588.
- [21] J. W. Moore, R. G. Pearson, *Kinetics and Mechanism*, 3rd ed., Wiley, New York, 1981.

### Modular Ligands for Asymmetric Synthesis: Enantioselective Catalytic Cu<sup>II</sup>-Mediated Condensation Reaction of Ethyl Pyruvate with Danishefsky's Diene\*\*

Peter I. Dalko,\* Lionel Moisan, and Janine Cossy\*

The search for new catalysts for asymmetric synthesis is an area of significant importance in modern organic synthesis, and a number of spectacular advances have been reported in recent years.<sup>[1, 2]</sup> However, there is still a need for simple and readily available catalysts, particularly for C-C bond-forming processes.<sup>[3]</sup> A modular system that allows quick access to a great diversity of ligands from simple components is valuable. Peptide-based ligands have been developed for Ti-catalyzed additions of TMSCN to meso epoxides<sup>[4]</sup> and imines (Strecker amino acid synthesis),<sup>[5]</sup> and Zr-catalyzed addition of dialkylzincs to imines.<sup>[6]</sup> Recently, peptide-based phosphane ligands have been shown to promote efficient Cu-catalyzed conjugate addition.<sup>[7]</sup> Moreover, this class of ligands allows the regioand enantioselective formation of quaternary carbon centers by using allylic reagents.<sup>[8]</sup> We report herein a simple and efficient approach that leads to the generation of a great diversity of asymmetric ligands from easily available components.[9]

The catalyst preparation is based on the condensation of 1,2-diamines<sup>[10]</sup> of type **1** with ketones or aldehydes **2** (Scheme 1), which affords imidazolidines **3** in solution in equilibrium with the open form **4**. The corresponding bisimine **5** and the starting diamine **1** are often present with compounds **3** and **4** in the reaction mixture.<sup>[11]</sup> We anticipated that chelating metals such as Cu<sup>II</sup> would shift this equilibrium toward the metallacyclic form **6** by forming a bidentate complex (Scheme 1).<sup>[12]</sup> Although no precedent for the use of

1433-7851/02/4104-0625 \$ 17.50+.50/0

Angew. Chem. Int. Ed. 2002, 41, No. 4 © WILEY-VCH Verlag GmbH, 69451 Weinheim, Germany, 2002

 <sup>[\*]</sup> Dr. P. I. Dalko, Prof. Dr. J. Cossy, L. Moisan Laboratoire de Recherches Organiques associé au CNRS, ESCPI 10 rue Vauquelin, 75231 Paris Cedex 05 Fax: (+33)1-40-79-46-60 E-mail: peter.dalko@espci.fr, janine.cossy@espci.fr

<sup>[\*\*]</sup> The Association pour la Recherche contre le Cancer (ARC) (grant no. 7579) is gratefully acknowledged for financial support.

## COMMUNICATIONS



Scheme 1. Condensation reaction between vicinal diamines and carbonyl compounds.

this type of ligand in enantioselective catalytic reactions has been reported, analogous reactions in the literature suggest that these complexes may be active in a number of reactions.<sup>[13]</sup>

To demonstrate the practical use of the catalyst system, a condensation reaction between *trans*-1-methoxy-3-(trimethyl-silyloxy)buta-1,3-diene (**7**; Danishefsky's diene) and ethyl pyruvate (**8**) was selected (Scheme 2).<sup>[14, 15]</sup> In this reaction,



Scheme 2. Reaction between Danishefsky's diene and ethyl pyruvate by using chiral Cu<sup>II</sup>-derived catalysts. Reagents and conditions: a) Cu<sup>II</sup>-L\*<sub>n</sub> (10%), 4-Å molecular sieves, THF, -72°C, 72 h; b) CF<sub>3</sub>COOH (cat.), room temperature.

dihydropyrone 9 can be formed through two distinct mechanistic pathways, that is, through a traditional hetero-Diels – Alder (HDA) cycloaddition reaction or by a Mukaiyama aldol condensation pathway followed by cyclization.<sup>[16]</sup> It is generally accepted that the Mukaiyama aldol product 10 can either be isolated, or may undergo a ring-closure reaction to give dihydropyrone 9 under the reaction conditions or by treatment with acid. Herein we show that complexes derived from chiral diamines, carbonyl compounds, and Cu<sup>II</sup> salts catalyze both reaction pathways and afford the desired adducts with high enantiomeric excesses.

The ligands were prepared by the condensation of amines **1** (Scheme 1) with ketones or aldehydes (see Table 1).<sup>[17]</sup> The chiral Lewis acid catalysts were obtained by adding a stoichiometric amount of  $Cu(OTf)_2$  to the reaction mixture, which was allowed to stand for 16 h at room temperature (premixing).<sup>[18]</sup> The premixing time significantly influenced the enantioselectivity of the condensation reaction. In gen-

626 © WILEY-VCH Verlag GmbH, 69451 Weinheim, Germany, 2002

eral, a lower *ee* value was obtained for dihydropyrone **9** when the premixing time was reduced (see below).

The condensation reaction between 7 and 8 in the presence of the chiral complexes derived from diamines 1a and 1b (10 mol%), carbonyl compounds 2a - f, and copper(II) salt at  $-72^{\circ}$ C afforded dihydropyrone 9 and the Mukaiyama aldol product 10 (Scheme 2). At this stage, the ee value of the primary cycloadduct 9 was determined<sup>[19]</sup> (Table 1), but unfortunately the ee values of 10a and 10b could not be determined. The mixture of 9 and 10 was

converted into the dihydropyrone **9** by treatment with a catalytic amount of triflic acid, and the *ee* value of the resulting dihydropyrone was measured (Table 1).<sup>[20]</sup>

Table 1. Condensation of  ${\bf 7}$  and  ${\bf 8}$  by using  ${\rm Cu^{II}}$  complexes derived from chiral amine.

Entry	Amine 1	Ketone 2	1/2	9		
-				yield [%]	ee [%] <sup>[a]</sup>	ee [%] <sup>[b]</sup>
1	(–)-1a	none		0	_	_
2	(–)-1a	cyclohexanone (2a)	1:1	72	70	(92)
3	(-)- <b>1</b> a	cyclopentanone (2b)	1:1	85	91	(92)
4	(-)- <b>1</b> a	cyclobutanone (2c)	1:1	85	94	(≥98)
5	(-)- <b>1</b> a	(-)-menthone $(2d)$	1:1	63	10	(10)
6	(+)- <b>1</b> a	(-)-menthone $(2d)$	1:1	61	12	(17)
7	(-)-1a	acetone (2e)	1:1	67	54	(70)
8	(-)- <b>1</b> a	benzaldehyde (2 f)	1:1	80	5	(18)
9	(−) <b>-1</b> a	benzaldehyde (2 f)	1:2	< 2	-	-
10	(-) <b>-1b</b>	cyclohexanone (2a)	1:1	0	-	-

<sup>[</sup>a] The *ee* values after treatment with acid. [b] The *ee* values before treatment with acid. Reaction conditions: THF, catalyst (10%), -72°C, 4 Å molecular sieves, 72 h.

As expected, the diamine – copper(II) triflate complex was inefficient in mediating the reaction (Table 1, entry 1). When ligands derived from diamine (–)-1a and cyclohexanone 2a were used (Table 1, entry 2) the reaction afforded the hetero-Diels – Alder product 9 with high *ee* (92%). However, after treatment with triflic acid, the enantiomeric excess of 9 proved to be lower (*ee* 70%).<sup>[20, 21]</sup> When the complexation time (premixing) was reduced from the usual 16 h (Table 1, entry 2) to 1 h, 9 was obtained in identical chemical yield, but the *ee* value dropped from 70% to 49%.

Upon decreasing the ring size of the ketone component from five to four carbon atoms, the *ee* value of cycloadduct **9**, before treatment with acid, increase from 92% (Table 1, entry 3) to better than 98% (Table 1, entry 4). Likewise, the *ee* value of **9** after treatment with acid increased from 91% (Table 1, entry 3) to 94% (Table 1, entry 4). Somewhat surprisingly, low enantioselectivity (10%) was observed when (-)-menthone was used in combination either with the (1*R*,2*R*)- or with the (1*S*,2*S*)-diphenylethylene diamine (Table 1, entries 5 and 6). This lack of selectivity may be a consequence of the increased steric hindrance in the proximity of the metal center in 6, which implies less selective reaction paths. Interestingly, acyclic ketones such as acetone **2e** were shown to be less selective than cyclic ketones (Table 1, entry 7). Also, imidazolidines derived from diamine and benzaldehyde **2f** resulted in low enantioselectivity (Table 1, entry 8). Although bis-imines of type **5** were shown earlier to be efficient in promoting enantioselective copper(II)-mediated HDA reactions,<sup>[2a]</sup> the bis-benzyl imine – copper complex derived from **1a** and **2f** afforded only traces of a nearly racemic product **9** under standard conditions (Table 1, entry 9).<sup>[22]</sup> Furthermore, ligands derived from **1b** and **2a** promoted neither the formation of the HDA adduct nor the Mukaiyama aldol product (Table 1, entry 10).

The Cu<sup>II</sup>/ligand stoichiometry was critical to the selectivity and reactivity. This dependence was monitored on the HDA adduct  $9^{[23]}$  Whereas a 1:1 ratio of cyclohexylidene ligand derived from (–)-**1 a**, **2 a**, and Cu(OTf)<sub>2</sub> afforded **9** in 92% *ee*, an increase in the Cu<sup>II</sup>/ligand ratio to 2:1 resulted in only 73% *ee*. On the other hand, when a twofold molar excess of ligand relative to the Cu<sup>II</sup> salt was used, the catalytic activity of the complex was inhibited.

The vicinal diamine ligands can be replaced by a chiral  $\alpha$ aminoalcohol. Although this reaction was not optimized, a reversal in enantioselectivity of the dihydropyrone **9** (67%)

yield, 35% ee) from that of the Mukaiyama aldol product **10** was observed when (*R*)-phen-ylglycinol (**11**) was used.

The foregoing three-component catalyst systems provide a simple and effective approach for generating a variety of structurally diverse

chiral complexes for use in asymmetric synthesis. Important structural features responsible for the catalytic activity are unknown at the present time. The operational simplicity, easy availability, low cost, stability, and diversity of potentially active organometallic structures all contribute to the potential usefulness of these catalysts.

#### **Experimental Section**

Diamine 1 (0.09 mmol, 0.1 equiv) and ketone 2 (0.09 mmol, 0.1 equiv) were dissolved in a small amount of dichloromethane (2 mL). After 4 h at room temperature, the solvent was evaporated, and the residue was coevaporated twice with benzene. Cu(OTf)2 (32.6 mg, 0.09 mmol, 0.1 equiv) and the ligand were introduced in a dry flask with a small amount of ground 4-Å molecular sieves (ca. 100 mg), and THF (1.5 mL) was added to the solid through a cannula. The slurry was stirred under an inert atmosphere of Ar for 16 h at room temperature. The slurry was cooled to -72 °C, ethyl pyruvate (100  $\mu L,$  0.9 mmol, 1.0 equiv) and Danishefsky's diene (215  $\mu L,$ 1.3 mmol, 1.2 equiv) were added in one portion through a syringe, and the mixture was stirred for an additional 72 h at -72°C. Trifluoroacetic acid (0.1 mL) was then introduced to the mixture, which was allowed to warm to room temperature. After 12 hours, the solution was neutralized with saturated sodium bicarbonate, the organic phase was separated, and the aqueous phase was extracted with dichloromethane  $(2 \times 10 \text{ mL})$ . The combined organic phases were dried over Na<sub>2</sub>SO<sub>4</sub>, the solvent was evaporated, and the crude reaction mixture was purified on a short column of silica gel (pentane/ethyl acetate 4:1 eluent).  $R_{\rm f} = 0.5$  (silica gel, EtOAc/ pentane 1:1); IR (film):  $\tilde{\nu} = 1740$ , 1680, 1600 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz):  $\delta = 7.36 \text{ (d, } J = 5.9 \text{ Hz}, 1 \text{ H}), 5.42 \text{ (d, } J = 5.9 \text{ Hz}, 1 \text{ H}), 4.22 \text{ (q, } J = 5.9 \text{ Hz}, 1 \text{ H})$ 7.0 Hz, 2 H), 3,00 (d, J = 16.5 Hz, 1 H), 2.68 (d, J = 16.5 Hz, 1 H), 1.66 (s, 3H), 1.27 (t, J = 7.0 Hz, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz):  $\delta = 190.1$ , 171.0, 161.9, 107.4, 82.9, 62.4, 44.8, 24.2, 14.1; MS (EI, 70 eV): *m*/*z*: 184 (27), 111 (100), 110 (24), 86 (12), 71 (89), 69 (25).

Received: June 25, 2001 Revised: November 23, 2001 [Z17345]

- For leading references, see: a) Comprehensive Asymmetric Catalysis (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, Heidelberg, **1999**; b) Asymmetric Catalysis in Organic Synthesis (Ed.: R. Noyori), Wiley, New York, **1994**; c) Asymmetric Synthesis, 2nd ed. (Ed.: I. Ojima), VCH, New York, **2000**; d) "Catalytic Asymmetric Synthesis": Acc. Chem. Res. **2000**, 33(6) (special issue); e) D. Astruc, Chimie Organométallique, EDP Sciences, Les Ulis, **2000**; f) J. Seyden Penne, Chiral Auxiliaires and Ligands in Asymmetric Synthesis, Wiley, New York, **1995**.
- [2] a) J. S. Johnson, D. A. Evans, Acc. Chem. Res. 2000, 33, 325–335;
  b) A. Pfaltz, Synlett 1999, 835–842; c) A. K. Ghosh, P. Mathivanan, J. Cappiello, Tetrahedron: Asymmetry 1998, 9, 1–45; d) E. J. Corey, C. J. Helal, Angew. Chem. 1998, 110, 2092–2118; Angew. Chem. Int. Ed. 1998, 37, 1986–2013.
- [3] A. Berkessel, R. Riedl, J. Comb. Chem. 2000, 2, 215-219.
- [4] a) B. M. Cole, K. D. Shimizu, C. A. Krueger, J. P. Harrity, M. L. Snapper, A. H. Hoveyda, *Angew. Chem.* 1996, *108*, 1776–1779; *Angew. Chem. Int. Ed. Engl.* 1996, *35*, 1668–1671; b) K. D. Shimizu, B. M. Cole, C. A. Krueger, K. W. Kuntz, M. L. Snapper, A. H. Hoveyda, *Angew. Chem.* 1997, *109*, 1782–1785; *Angew. Chem. Int. Ed. Engl.* 1997, *36*, 1704–1707.
- [5] a) C. A. Krueger, K. W. Kuntz, C. D. Dzierba, W. G. Wirschun, J. D. Gleason, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* 1999, *121*, 4248–4285; b) J. R. Porter, W. G. Wirschun, K. W. Kuntz, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* 2000, *122*, 2657–2658; c) N. S. Josephson, K. W. Kuntz, M. L. Snapper, A. H. Hoveyda, *J. Am. Chem. Soc.* 2001, *123*, 11594–11599.
- [6] J. R. Porter, J. F. Traverse, A. H. Hoveyda, M. L. Snapper, J. Am. Chem. Soc. 2001, 123, 984–985.
- [7] S. J. Dergado, H. Mizutani, A. H. Hoveyda, J. Am. Chem. Soc. 2001, 123, 755-756.
- [8] C. A. Luchaco-Cullis, H. Mizutani, K. E. Murphy, A. H. Hoveyda, Angew. Chem. 2001, 113, 1504–1508; Angew. Chem. Int. Ed. 2001, 40, 1456–1460.
- [9] For related catalyst systems, see: a) R. H. Crabtree, *Chem. Commun.* 1999, 1611–1616; b) M. T. Reetz, *Angew. Chem.* 2001, *113*, 292–320; *Angew. Chem. Int. Ed.* 2001, *40*, 284–310; c) J. Long, J. Hu, X. Shen, B. Ji, K. Ding, *J. Am. Chem. Soc.* 2002, *124*, 10–11.
- [10] a) Y. L. Bennani, S. Hanessian, Chem. Rev. 1997, 97, 3161-3195; b) D. Lucet, T. Le Gall, C. Mioskowski, Angew. Chem. 1998, 110, 2724-2772; Angew. Chem. Int. Ed. 1998, 37, 2580-2627.
- [11] C. Chapuis, A. Gauvreau, A. Klaebe, A. Lattes, J. J. Perie, Bull. Soc. Chim. Fr. 1973, 977–985.
- [12] P. Braunstein, F. Naud, Angew. Chem. 2001, 113, 702-722; Angew. Chem. Int. Ed. 2001, 40, 680-699.
- [13] Reactions that are potentially mediated by complexes derived either from bis-imines 5 or the starting diamines 1 should be kinetically disfavored transformations. A similar principle has been developed earlier in the generation of imine-based virtual combinatorial libraries; see: Y. Huc, J.-M. Lehn, *Proc. Nat. Acad. Sci. USA* 1997, 94, 2106.
- [14] For recent reviews on this topic, see: a) W. Carruthers, Cycloaddition Reactions in Organic Synthesis, Pergamon, Elmsford, NY, 1990 (Tetrahedron Organic Chemistry Series, Vol. 8); b) D. L. Boger, S. H. Weinreb, Hetero-Diels-Alder Methodology in Organic Synthesis, Academic Press, New York, 1987; c) H. Waldmann, Synthesis 1994, 535-551; d) L. F. Tietze, G. Kettschau, Top. Curr. Chem. 1997, 190, 1; d) K. A. Jørgensen, Angew. Chem. 2000, 112, 3702-3733; Angew. Chem. Int. Ed. 2000, 39, 3558-3588.
- [15] a) M. Johannsen, S. Yao, K. A. Jørgensen, *Chem. Commun.* **1997**, 2169–2170; b) S. Yao, M. Johannsen, H. Audrain, R. G. Hazell, K. A. Jørgensen, *J. Am. Chem. Soc.* **1998**, *120*, 8599–8605; c) S. Yao, M. Johannsen, H. Audrain, R. G. Hazell, K. A. Jørgensen, *J. Am. Chem. Soc.* **1998**, *120*, 8599–8605; d) M. Johannsen, S. Yao, A. Graven, K. A. Jørgensen, *Pure Appl. Chem.* **1998**, *79*, 1117; e) K. A. Jørgensen, M.

1433-7851/02/4104-0627 \$ 17.50+.50/0

Angew. Chem. Int. Ed. 2002, 41, No. 4 © WILEY-VCH Verlag GmbH, 69451 Weinheim, Germany, 2002

# COMMUNICATIONS

Johannsen, S. Yao, H. Audrain, J. Thorhauge, Acc. Chem. Res. 1999, 32,605-613.

- [16] For related aldol reactions, see: a) D. A. Evans, M. C. Kozlowski, J. A. Murry, C. S. Burgey, K. R. Campos, B. T. Connell, R. J. Staples, J. Am. Chem. Soc. 1999, 121, 669-685; b) D. A. Evans, C. S. Burgey, C. S. Kozlowski, S. W. Tregay, J. Am. Chem. Soc. 1999, 121, 686-699; c) M. Roberson, A. S. Jepsen, K. A. Jørgensen, Tetrahedron 2001, 57, 907 -913.
- [17] Best results were obtained when the diamine and the carbonyl compound were mixed in dichloromethane at room temperature without an acid catalyst. The formation of the products was monitored by 1H and 13C NMR spectroscopy.
- [18] Other Cu<sup>II</sup> salts such as Cu(SbF<sub>6</sub>)<sub>2</sub>, Cu(BF<sub>4</sub>)<sub>2</sub>, Cu(ClO<sub>4</sub>)<sub>2</sub>, Cu(OAc)<sub>2</sub>, Cu(PF<sub>4</sub>)<sub>2</sub> showed lower selectivity in this reaction.
- [19] The ee value of 9 was determined by chiral GC-MS on a Crompack Chirasil-DEX CB column.
- [20] No epimerization of 9 was observed after treatment with acid.
- [21] The absolute configuration of the newly formed stereogenic center of 9 was found to be R in each case when the (1R,2R)-diphenylethylene diamine ligand was used. The absolute configuration was determined by comparison of the optical rotation of  ${\bf 9}$  with the literature value.  $^{[15c]}$
- [22] The ligand was prepared by condensation of benzaldehyde (2 equiv) with 1a followed by addition of Cu(OTf)<sub>2</sub> (1 equiv).
- [23] The reaction was quenched at -72 °C by using trifluoroacetic acid and was allowed to stir at -10 °C for 1 h. This workup procedure allowed the isolation of the HDA and aldol products. No interconversion between the aldol and HDA products was observed under these conditions.

## A Thermotropic Mesophase Comprised of **Closed Micellar Aggregates of the Normal** Type\*\*

Petra Fuchs, Carsten Tschierske,\* Klaus Raith, Kumar Das, and Siegmar Diele

The ability of amphiphilic molecules to self-organize in aqueous systems with formation of micelles, vesicles, and lyotropic mesophases is of great importance for numerous applications of surfactants in several fields of science and technology, and it is also a prerequisite for the development of biological structures.<sup>[1, 2]</sup> In these polymolecular assemblies the hydrophilic parts (together with the solvent molecules) and the lipophilic parts of the amphiphiles are segregated into nanoscopic compartments, whereby changing the degree of curvature of the interfaces between the incompatible nano-

[\*] Prof. Dr. C. Tschierske, Dr. P. Fuchs Institute of Organic Chemistry Martin-Luther-University Halle - Wittenberg Kurt-Mothes-Straße 2, 06120 Halle (Germany) Fax: (+49) 345-55-27223 E-mail: tschierske@chemie.uni-halle.de Dr. S. Diele, Dr. K. Das Institute of Physical Chemistry Martin-Luther-University Halle-Wittenberg Mühlpforte 1, 06120 Halle (Germany) Fax: (+49) 345-55-27157 Dr. K. Raith Faculty of Pharmacy Martin-Luther-University Halle - Wittenberg (Germany)

[\*\*] This work was supported by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie.

phases gives rise to a sequence of different mesophase morphologies. Hence, in dependence on the amphiphile structure, its concentration, and the temperature, different lyotropic mesophases can result.<sup>[1-3]</sup> Arrays of alternating layers yield smectic phases (SmA), regular arrangements of cylinders give rise to hexagonal columnar phases (Col<sub>b</sub>), and closed globular or nonglobular micelles can form cubic mesophases (micellar cubic phases, Cub<sub>I</sub>). Another type of cubic mesophase, consisting of two mutually interwoven networks of branched cylinders (bicontinuous cubic mesophases,  $Cub_v$ ), occurs at the transition between smectic and columnar organization (Figure 1). For each of the nonlamellar mesophases two different types are possible. In normal phases (type 1, positive curvature of the polar/apolar interface), the stronger cohesive forces (hydrogen bonds) are located in the continuum surrounding the aggregates. In the reversed (or inverse) phases (type 2, negative interface curvature) they are



Figure 1. Dependence of the mesophase morphology of different polyhydroxy amphiphiles on the molecular structure. Abbreviations: Cub<sub>12</sub>= reversed discontinuous (micellar) cubic mesophase with Pm3n lattice; Colh2 = reversed hexagonal columnar mesophase; CubV2 = reversed bicontinuous cubic mesophase with Ia3d lattice; SmA = smectic A-phase;  $Cub_{V1} = normal-type$  bicontinuous cubic mesophase with *Ia3d* lattice;  $Col_{h1} = normal-type$  hexagonal columnar mesophase;  $M_{11} = mesophase$ comprised of discrete direct micelles with unknown lattice (the Im3m lattice is shown as one of the possible structures).

1433-7851/02/4104-0628 \$ 17.50+.50/0