

Synthesis and Insecticidal Activities of Novel Anthranilic Diamides Containing Acylthiourea and Acylurea

Ji-Feng Zhang,[†] Jun-Ying Xu,[‡] Bao-Lei Wang,[†] Yu-Xin Li,^{*,†} Li-Xia Xiong,[†] Yong-Qiang Li,[†] Yi Ma,[†] and Zheng-Ming Li^{*,†}

[†]State Key Laboratory of Elemento-Organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China

[‡]CNOOC Tianjin Chemical Research and Design Institute, Tianjin 300131, People's Republic of China

S Supporting Information

ABSTRACT: Two series of anthranilic diamides containing acylthiourea and acylurea linkers were designed and synthesized, with changed length and flexibility of the linkers to compare to known anthranilic diamide insecticides. In total, 26 novel compounds were synthesized, and all compounds were characterized by ¹H nuclear magnetic resonance and high-resolution mass spectrometry. Their insecticidal activities against oriental armyworm (*Mythimna separata*), mosquito larvae (*Culex pipiens pallens*), and diamondback moth (*Plutella xylostella*) were evaluated. The larvicidal activities against oriental armyworm indicated that the introduction of acylthiourea into some structures could retain their insecticidal activity; 8 of the 15 compounds (**13a–13e**, **14a–14e**, and **15a–15e**) exhibited 100% larvicidal activity at 10 mg/L. However, the introduction of acylurea decreased the insecticidal activity; only 3 of the 11 compounds (**17a–17k**) exhibited 100% larvicidal activity at 200 mg/L. The whole-cell patch-clamp technique indicated that compound **13b** and chlorantraniliprole exhibited similar effects on the voltage-gated calcium channel. The calcium-imaging technique was also applied to investigate the effects of compounds **13b** and **15a** on the intracellular calcium ion concentration ([Ca²⁺]_i), which indicated that they released stored calcium ions from endoplasmic reticulum. Experimental results denoted that several new compounds are potential activators of the insect ryanodine receptor (RyR).

KEYWORDS: Anthranilic diamide, insecticidal activity, acylthiourea, acylurea, calcium channel

INTRODUCTION

For years, scientists have been dedicated in searching for insecticides with new mechanisms to cope with the global food shortage. Recently, the anthranilic diamides chlorantraniliprole (Rynaxypyr; DPX-E2Y45) (compound **A** in Figure 1) and cyantraniliprole (Cyazypyr) (compound **B**) were marketed by Dupont, targeting at the insect ryanodine receptor,^{1,2} which exhibit their action by activating the uncontrolled release of the calcium store.¹ Their broad insecticidal spectra, high efficiency, and low toxicity aroused interests worldwide. Some modified structures (compound **C**) have been reported, which mainly focused on the 1-(3-chloropyridyl) pyrazole moiety (I), the aliphatic amide moiety (II), and the anthraniloyl moiety (III).^{2–11} Nevertheless, the modifications about the amide bridge part as a linker of two aryl rings were seldom reported.¹²

In crop protection and bioactive chemicals, acylthioureas and acylureas have been reported to display a variety of biological activities, such as insecticidal, fungicidal, herbicidal, antimicrobial, antitumor, etc.^{13–18} With this in mind, two series of novel anthranilic diamide derivatives (compound **D**) were designed and synthesized in this paper by introducing acylthiourea and acylurea moieties to increase the linker length and flexibility. Their synthetic routes were shown in Schemes 1–5. Their insecticidal activities against oriental armyworms, mosquito larvae, and diamondback moths were tested accordingly. The preliminary structure–activity relationship (SAR) was discussed. To provide insight to further study the biological effect

of target compounds, the whole-cell patch-clamp and calcium-imaging techniques were used to investigate the effects of compounds **13b** and **15a** on calcium channels in the central neurons of *Spodoptera exigua*.

MATERIALS AND METHODS

Instruments. ¹H nuclear magnetic resonance (NMR) spectra were recorded at 300 MHz using a Bruker AC-P300 spectrometer or 400 MHz using a Bruker AV 400 spectrometer (Bruker Co., Switzerland) in CDCl₃ or DMSO-*d*₆ solution with tetramethylsilane as the internal standard, and chemical-shift values (δ) were given in parts per million (ppm). High-resolution mass spectrometry (HRMS) data were obtained on a Varian QFT-ESI instrument. Flash chromatography was performed on CombiFlash Companion (Teledyne Isco, Inc., Lincoln, NE) with silica gel (300–400 mesh). The melting points were determined on a X-4 binocular microscope melting point apparatus (Beijing Tech Instruments Co., Beijing, China) and were uncorrected. The whole-cell patch clamp was performed using a patch-clamp amplifier (EPC-10, HEKA Elektronik, Lambrecht, Germany). Reagents were all analytically or chemically pure. All solvents and liquid reagents were dried by standard methods in advance and distilled before use. Chlorantraniliprole used in this work was synthesized according to the literature only for bioassay reference.¹⁹

General Synthetic Procedure for Compounds **6a and **6b**.** The title compounds **6a** and **6b** were prepared in our laboratory

Received: February 24, 2012

Accepted: July 19, 2012

Published: July 19, 2012





Figure 1. Chemical structures of compounds A–D.

according to the methods reported by Dong et al. (Scheme 1), with some modifications. The melting points and ^1H NMR data of all of the compounds were consistent with the literature.^{19,20}

Scheme 1. Synthesis of Title Compounds 6a and 6b^a



^aReagents and conditions: (a) $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ (80%), reflux, (b) NaOEt , EtOH , reflux, (c) POBr_3 or POCl_3 , MeCN , 80°C , (d) $\text{K}_2\text{S}_2\text{O}_8$, H_2SO_4 , MeCN , reflux, and (e) (i) aqueous NaOH , MeOH and (ii) aqueous HCl .


1,1,1-Trifluoro-4-(furan-2-yl)-4-hydroxybut-3-en-2-one (8). Sodium (1.38 g, 60 mmol) was dissolved in 30 mL of anhydrous methanol slowly. Methanol was evaporated to dryness under reduced pressure to give the white solid sodium methylate. Then, diethyl ether (50 mL), ethyl trifluoroacetate (8.52 g, 60 mmol), and 2-acetylfuran (6.60 g, 60 mmol) were successively added dropwise, stirred at room temperature for 5 h, and filtered, which was acidized with 1 M sulfuric acid to give a red liquid product. Yield = 64.9%. ^1H NMR (400 MHz, CDCl_3) δ : 13.67 (br s, 1H, OH), 7.67 (s, 1H, Ar-H), 7.33 (d, 1H, J = 3.6 Hz, Ar-H), 6.63 (d, 1H, J = 3.6 Hz, Ar-H), 6.48 (s, 1H, CH).

3-Chloro-2-(5-(furan-2-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)pyridine (9). In a three-neck 1000 mL round-bottomed flask, 1,1,1-trifluoro-4-(furan-2-yl)-4-hydroxybut-3-en-2-one (8, 27.1 g, 130 mmol) and 3-chloro-2-hydrazinylpyridine (18.9 g, 130 mmol) were added to 300 mL of acetic acid. The mixture was refluxed for 8 h, and then the excess acetic acid was removed under reduced pressure. The residue was further purified by flash column chromatography on silica gel with petroleum ether/ethyl acetate (5:1) to give intermediate 9 as a red oil. Yield = 42.1%. ^1H NMR (400 MHz, CDCl_3) δ : 8.58 (d, J = 4.8 Hz, 1H, pyridyl-H), 7.96 (d, J = 8.0 Hz, 1H, pyridyl-H), 7.51 (dd, J = 8.0, 4.8 Hz, 1H, pyridyl-H), 7.36 (s, 1H, pyrazolyl-H), 6.93 (s, 1H, furanyl-H), 6.41–6.30 (s, 1H, furanyl-H), 6.01 (d, J = 3.4 Hz, 1H, furanyl-H).

1-(3-Chloropyridin-2-yl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxylic acid (6c). To a solution of 3-chloro-2-(5-(furan-2-yl)-3-(trifluoromethyl)-1H-pyrazol-1-yl)pyridine (9, 3.88 g, 12.4 mmol) in acetone (35 mL) and water (35 mL), potassium permanganate (9.80 g, 62 mmol) was added in portion. When the addition was complete, the reaction was refluxed for 30 min. The resulting mixture was filtered. The filtrate was acidified with 2 M HCl and extracted with ethyl acetate (50 mL \times 3). The extracts were combined, washed with water (30 mL \times 2) and brine (30 mL), then dried with anhydrous sodium sulfate, and evaporated to give compound 6c as a white solid (1.85 g). Yield = 51.3%. Melting point (mp) = 176 – 179°C . ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 14.14 (s, 1H), 8.61 (dd, J = 4.8, 1.5 Hz, 1H,

pyridyl-H), 8.31 (dd, J = 8.0, 1.5 Hz, 1H, pyridyl-H), 7.74 (dd, J = 8.0, 4.8 Hz, 1H, pyridyl-H), 7.61 (s, 1H, pyrazolyl-H) (Scheme 2).

Scheme 2. Synthesis of Compound 6c^a



^aReagents and conditions: (a) NaOCH_3 , ethyl trifluoroacetate, (b) AcOH , reflux, and (c) KMnO_4 , acetone/ H_2O , reflux.

General Synthetic Procedure for Compounds 11a–11m. The compounds 11a–11m were prepared in our laboratory according to the methods reported by Dong et al. (Scheme 3 and Table 1). The melting points and ^1H NMR data of all of the compounds were consistent with the literature.^{19,20}

Scheme 3. Synthesis of Compounds 11a–11m





Table 1. Structure of Compounds 11a–11m

| compound | R_1 | R_2 | R_3 |
|----------|--------------|--------------|--------------|
| 11a | Me | Me | Cl |
| 11b | Et | Me | Cl |
| 11c | i-Pr | Me | Cl |
| 11d | i-Pr | H | Cl |
| 11e | n-Pr | H | Cl |
| 11f | n-Bu | Me | Cl |
| 11g | t-Bu | Me | Cl |
| 11h | cyclopropyl | Me | Cl |
| 11i | cyclohexyl | Me | Cl |
| 11j | i-Pr | Me | Br |
| 11k | benzyl | Me | Br |
| 11l | i-Pr | Me | H |
| 11m | i-amyl | Me | H |

General Synthetic Procedure for Compounds 13a–13e, 14a–14e, and 15a–15e. To a suspension of 1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxylic acid derivative (6a–6c) (1 mmol) in 25 mL of dichloromethane, oxalyl chloride (2.0 mmol) and a drop of N,N -dimethylformamide (DMF) were added. After stirring at room temperature for 3 h, the solution was evaporated. The resulting acyl chloride was dissolved in 20 mL of acetonitrile and added to a solution of potassium thiocyanate in 15 mL of acetonitrile with two drops of polyethylene glycol-400 (PEG-400). After stirring at room temperature for 30 min, the mixture was filtered to give the acyl isothiocyanate derivatives 12a–12c, which were used without further

Scheme 4. Synthesis of Compounds 13a–13e, 14a–14e, and 15a–15e^a

^aReagents and conditions: (a) (i) oxalyl chloride, CH₂Cl₂ and (ii) KSCN, PEG-400, CH₃CN and (b) 11a–11m, CH₃CN.

purification, to which the 2-amino-3-methylbenzamide derivatives (11a–11m) (1.0 mmol) were added and stirred overnight, then filtered, and further purified by recrystallization using methanol or by a silica-gel column eluted with petroleum ether/ethyl acetate (4:1) to obtain the title compounds 13a–13e, 14a–14e, and 15a–15e.

1-(3-Chloropyridin-2-yl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide (16). To a suspension of 3-trifluoromethyl-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxylic acid (6c) (2.40 g, 8.23 mmol) in 50 mL of dichloromethane, oxalyl chloride (17.0 mmol) and a drop of DMF were added. After stirring at room temperature for 3 h, the solution was evaporated. The resulting acyl chloride was dissolved in 50 mL of tetrahydrofuran (THF) and added to aqueous ammonia (25%) (5.76 g, 41.15 mmol) at 0 °C. After stirring overnight, a large amount of solid formed and was filtered, which was further purified by a silica-gel column eluted with petroleum ether/ethyl acetate (2:1) to give the title compound as a white solid (1.85 g). Yield = 77.3%. mp = 183–185 °C. ¹H NMR (400 MHz, CDCl₃) δ: 8.56 (d, *J* = 4.8 Hz, 1H, pyridyl-H), 8.00 (d, *J* = 8.0 Hz, 1H, pyridyl-H), 7.53 (dd, *J* = 8.0, 4.8 Hz, 1H, pyridyl-H), 7.38 (s, 1H, pyrazolyl-H), 6.78 (s, 1H, NH), 5.77 (s, 1H, NH).

General Synthetic Procedure for Compounds 17a–17k. A suspension of compound 16 (0.29 g, 1 mmol) in 20 mL of 1,2-dichloroethane and oxalyl chloride (3.0 mmol) was heated to reflux and kept for 5 h. Then, the mixture was evaporated, and the residue was dissolved in acetonitrile, to which the 2-amino-3-methylbenzamide derivatives (11a–11m) (1 mmol) were added and stirred overnight. Then, the produced precipitate was filtered and washed with acetonitrile (5 mL × 3) to give the title compounds 17a–17k.

Larvicidal Activity against Oriental Armyworm (*M. separata*). The larvicidal activities of compounds 13a–13e, 14a–14e, 15a–15e, 17a–17k, and chlorantraniliprole were evaluated using the reported procedure.¹⁹ The insecticidal activity against oriental armyworms was tested by foliar application; individual corn (*Tangyu 10*, *Zea mays* L.) leaves were placed on moistened pieces of filter paper in Petri dishes. The leaves were then sprayed with the test solution and allowed to dry. The dishes were infested with 10 fourth-instar oriental armyworm larvae. Percentage mortalities were evaluated 2 days after treatment. Each treatment was replicated 3 times.

Larvicidal Activity against Mosquito Larvae (*C. pipiens pallens*). The larvicidal activities of compounds 13a–13e, 14a–14e, and chlorantraniliprole against mosquito larvae were evaluated by the reported procedure.²² The compounds 13a–13e, 14a–14e, and chlorantraniliprole were prepared to different concentrations by dissolving compounds in acetone and adding distilled water. Then, 20 fourth-instar mosquito larvae were placed in 10 mL of test solution and raised for 3 days. Each treatment was performed 3 times, and the average value of the three tested values was calculated. The results were expressed by death percentage.

Larvicidal Activity against Diamondback Moth (*P. xyloste-la*). The larvicidal activity of compounds 15a–15e and chlorantraniliprole was tested by the leaf-dip method. At first, a solution of each test sample in DMF (AR, purchased from Alfa Aesar) at a concentration of 200 mg/L was prepared and then diluted to the required concentration with water (distilled). Leaf disks (6 × 2 cm) were cut from fresh cabbage leaves and then sprayed with the test solution for 3 s and allowed to dry. The resulting leaf disks were placed individually into glass tubes. Each disk was infested with 30 s-instar

diamondback moth larvae. Percentage mortalities were evaluated 2 days after treatment. Each treatment was performed 3 times.

Electrophysiological Recording and Calcium Imaging. The currents of calcium (*I*_{Ca}) were recorded using the reported procedure.²³ Calibration of the fluorescence signal was achieved using the method by Iakahashi et al., with some modifications.²⁴ The attached neurons were rinsed in standard physiological saline [150 mM NaCl, 4 mM KCl, 2 mM MgCl₂, 2 mM CaCl₂, and 10 mM *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES), buffered to pH 6.9] and then incubated in the dark at 28 °C in standard external saline containing the dye fluo-3-AM (Sigma, 10 μM) for 45 min or incubated in the dark at 28 °C in external saline [150 mM NaCl, 4 mM KCl, 2 mM MgCl₂, 2 mM ethylene glycol bis(2-aminoethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA), and 10 mM HEPES, buffered to pH 6.9] containing the dye fluo-5N (Invitrogen, 5 μM) for 5 h. After dye loading, cells were again rinsed in physiological saline twice.


For full depletion of thapsigargin-sensitive stores, cells were incubated with 1 μM thapsigargin for 10 min. Calcium ratio imaging studies were conducted using the imaging system coupled to an inverted fluorescence microscope with a Fluor 40× oil immersion objective (Olympus IX71). Cells which excited at 488 and 530 nm under fluorescence emission were acquired using a charge-coupled device (CCD) camera (Image Pri-6.0).

Each experiment was repeated at least 6 times. The data were analyzed using SPSS, Inc., version 17.0, and Microcal Origin, version 8.0 (Origin Lab Corp., Northampton, MA). Results were expressed as the mean ± standard deviation (SD) (*n* = number of cells). Statistical significance was determined by Student's paired or unpaired *t* tests. Fluorescence values were expressed as *F*/*F*₀, with *F*₀ being the resting (or baseline) fluorescence and *F* being the change in fluorescence from baseline after the drug application.

RESULTS AND DISCUSSION

Synthesis. The carboxylic acids 6a and 6b were prepared using the procedure reported by Dong et al. (Scheme 1),^{19,20} with some modifications, for the synthesis of intermediates. With the reported procedure, intermediate 2 was prepared in 69% yield by refluxing 2,3-dichloropyridine (1) and hydrazine hydrate (50%) in ethanol for 36 h. In our experiments, hydrazine hydrate (80%) was used to directly reflux with compound 1 for 5–6 h without solvent ethanol. By the improvement, compound 2 could be obtained in a 93% high yield.

The title compounds 13a–13e, 14a–14e, and 15a–15e were synthesized, as shown in Scheme 4. Carboxylic acid (6a–6c) was treated with oxalyl chloride and then added to potassium thiocyanate in acetonitrile with two drops of PEG-400. The corresponding acyl isothiocyanate derivative was filtered after stirring at room temperature for 30 min and reacted with 1 equiv of amines 11a–11m to afford the title compounds 13a–13e, 14a–14e, and 15a–15e, without further purification. Unlike the reported procedure that a reflux condition was usually needed,¹⁴ this reaction was worked out

Scheme 5. Synthesis of Compounds 17a–17k^a

^aReagents and conditions: (a) (i) oxalyl chloride, CH₂Cl₂ and (ii) NH₃·H₂O and (b) (i) oxalyl chloride, ClCH₂CH₂Cl, reflux and (ii) 11a–11m, CH₃CN.

Table 2. Insecticidal Activities of Compounds 13a–13e, 14a–14e, 15a–15e, 17a–17k, and Chlorantraniliprole against Oriental Armyworms

| compound | R ₁ | R ₂ | R ₃ | R ₄ | larvicidal activity (%) at a concentration of mg/L | | | | | |
|---------------------|----------------|----------------|----------------|-----------------|--|-----|-----|-----|-----|-----|
| | | | | | 200 | 100 | 50 | 20 | 10 | 5 |
| 13a | Me | Me | Cl | Cl | 100 | 100 | 100 | 100 | 40 | |
| 13b | i-Pr | Me | Cl | Cl | 100 | 100 | 100 | 100 | 100 | 40 |
| 13c | i-Pr | Me | Br | Cl | 100 | 80 | 40 | | | |
| 13d | i-Pr | Me | H | Cl | 100 | 100 | 100 | 100 | 100 | 0 |
| 13e | i-amyl | Me | H | Cl | 60 | | | | | |
| 14a | Me | Me | Cl | Br | 100 | 100 | 100 | 100 | 100 | 50 |
| 14b | i-Pr | Me | Br | Br | 100 | 100 | 100 | 100 | 100 | 0 |
| 14c | t-Bu | Me | Cl | Br | 100 | 100 | 100 | 100 | 100 | 40 |
| 14d | cyclohexyl | Me | Cl | Br | 0 | | | | | |
| 14e | i-amyl | Me | H | Br | 100 | 30 | | | | |
| 15a | Me | Me | Cl | CF ₃ | 100 | 100 | 100 | 100 | 100 | 50 |
| 15b | Et | Me | Cl | CF ₃ | 100 | 100 | 100 | 100 | 100 | 0 |
| 15c | cyclohexyl | Me | Cl | CF ₃ | 40 | | | | | |
| 15d | cyclopropyl | Me | Cl | CF ₃ | 100 | 100 | 100 | 100 | 100 | 30 |
| 15e | benzyl | Me | Br | CF ₃ | 100 | 100 | 100 | 100 | 0 | |
| 17a | Me | Me | Cl | CF ₃ | 100 | 20 | | | | |
| 17b | cyclopropyl | Me | Cl | CF ₃ | 20 | | | | | |
| 17c | n-Bu | Me | Cl | CF ₃ | 40 | | | | | |
| 17d | t-Bu | Me | Cl | CF ₃ | 60 | | | | | |
| 17e | cyclohexyl | Me | Cl | CF ₃ | 0 | | | | | |
| 17f | i-Pr | Me | Cl | CF ₃ | 70 | | | | | |
| 17g | cyclopropyl | Me | Br | CF ₃ | 100 | 0 | | | | |
| 17h | n-Pr | H | Cl | CF ₃ | 0 | | | | | |
| 17i | cyclopropyl | Me | H | CF ₃ | 100 | 80 | | | | |
| 17j | cyclopropyl | H | Cl | CF ₃ | 30 | | | | | |
| 17k | n-Pr | Me | H | CF ₃ | 60 | | | | | |
| chlorantraniliprole | | | | | 100 | 100 | 100 | 100 | 100 | 100 |

at room temperature under the catalysis of a small amount of PEG-400 with high yield.

The title compounds 17a–17k were synthesized, as shown in Scheme 5. Carboxylic acid 6c was converted to amide 16 by reacting with oxalyl chloride and then aqueous ammonia. Compound 16 was then refluxed with oxalyl chloride in 1,2-dichloroethane to give its corresponding acyl isocyanate derivative, which coupled with amines 11a–11m to afford the title compounds 17a–17k.

The ¹H NMR data of acylthiourea and acylurea derivatives were characteristic in all of the target compounds. In compounds 13a–13e and 14a–14e, the two active proton signals of NHCS and NHCO on carbonyl thiourea moiety were observed in DMSO-*d*₆ at 12.15–12.04 and 11.48–11.37 ppm, respectively. In compounds 15a–15e, the introduction of a trifluoromethyl group increased the hydrophobicity of the whole molecule; therefore, the ¹H NMR determination was carried out in CDCl₃. The chemical shifts of these two active protons were at 11.34–11.21 and 10.10–9.77 ppm. In

compounds 17a–17k, when R₂ = Me, the two active proton signals of CONHCO and NHCO on the carbonyl acylurea moiety appeared at 11.55–11.46 and 9.93–9.72 ppm, respectively. Whereas their values shifted to 11.54–11.52 and 11.16–11.11 ppm, respectively, when R₂ = H (17h and 17j) in DMSO-*d*₆.

SAR. Larvicidal Activities against Oriental Armyworms (*M. separata*). The larvicidal activities of compounds 13a–13e, 14a–14e, 15a–15e, 17a–17k, and commercial chlorantraniliprole against oriental armyworms were summarized in Table 2. In general, most of the compounds exist in moderate to good pesticidal activities. Particularly, eight compounds (13a–13e, 14a–14e, and 15a–15e) exhibited 100% larvicidal activities at 10 mg/L against oriental armyworm. From the activities against oriental armyworms, we could obviously find that the compounds containing the acylthiourea moiety (13a–13e, 14a–14e, and 15a–15e) gave higher insecticidal activities than the corresponding acylurea derivatives (17a–17k). When compounds 15a and 15d are compared to compounds 17a

and **17b**, we could find that one atom difference between acylthiourea and acylurea can cause a great change in its insecticidal activity. When different substitutions in the aliphatic amide moiety of title compounds **13a–13e**, **14a–14e** and **15a–15e** are compared, it was found that large substituents, such as *i*-amyl and cyclohexyl (**13e**, **14d**, **14e**, and **15c**) decreased the insecticidal activity against oriental armyworms. Furthermore, compounds with the *i*-propyl and *t*-butyl substituents showed the best larvicidal activity against oriental armyworms in our research, which was consistent with the results reported in previous SARs.^{2,25}

Biological Assay. Insecticidal activities against oriental armyworms (*Mythimna separata*), mosquito larvae (*Culex pipiens pallens*), and diamondback moths (*Plutella xylostella*) were performed on test organisms reared in a greenhouse. The bioassay was replicated at 25 ± 1 °C according to statistical requirements. Assessments were made on a dead/alive basis, and mortality rates were corrected applying Abbott's formula.²¹ Evaluation was based on a percentage scale of 0–100, where 0 equals no activity and 100 equals total kill. Error of the experiments was 5%. For comparative purposes, chlorantraniliprole was tested as a reference. The insecticidal activity was summarized in Tables 2–4.

Larvicidal Activities against Mosquito Larvae (*C. pipiens pallens*). The larvicidal activities of compounds **13a–13e** and **14a–14e** against mosquito larvae were shown in Table 3. Most

Table 3. Insecticidal Activities of Compounds 13a–13e, 14a–14e, and Chlorantraniliprole against Mosquito Larvae

| compound | larvicidal activity (%) at a concentration of mg/L | | |
|---------------------|--|-----|-----|
| | 2 | 1 | 0.5 |
| 13a | 70 | 30 | |
| 13b | 40 | | |
| 13c | 90 | 60 | 20 |
| 13d | 20 | | |
| 13e | 10 | | |
| 14a | 60 | 20 | |
| 14b | 90 | 70 | 30 |
| 14c | 100 | 90 | 30 |
| 14d | 40 | | |
| 14e | 20 | | |
| chlorantraniliprole | 100 | 100 | 100 |

of the compounds tested showed good larvicidal activities, while compounds with big substituents *i*-amyl and cyclohexyl in the aliphatic amide moiety revealed somewhat less activity.

Larvicidal Activity against Diamondback Moths (*P. xylostella*). The larvicidal activities against diamondback moths of compounds **15a–15e** and chlorantraniliprole were shown in Table 4. From it, we can see that most of the compounds tested showed excellent larvicidal activities. Compounds **15d** and **15e** showed the best activities (both 100% at 0.01 mg/L). Similarly, compound **15e** with a large substituent (R_2 = benzyl) also showed good larvicidal activity against diamondback moths.

The toxicity profile (LC_{50} values) of compounds **15d** and **15e**, which were found to be the most active insecticides of these two series of compounds against diamondback moths (100% at 0.01 mg/L), was shown in Table 5. The LC_{50} values of compounds **15d** and **15e** were 0.000 25 and 0.000 65 mg/L,

Table 4. Insecticidal Activities of Compounds 15a–15e and Chlorantraniliprole against Diamondback Moths

| compound | larvicidal activity (%) at a concentration of mg/L | | | | |
|---------------------|--|-----|-----|------|-------|
| | 20 | 1 | 0.1 | 0.01 | 0.001 |
| 15a | 100 | 100 | 100 | 71 | 43 |
| 15b | 100 | 100 | 100 | 86 | 29 |
| 15c | 30 | 0 | | | |
| 15d | 100 | 100 | 100 | 100 | 71 |
| 15e | 100 | 100 | 100 | 100 | 57 |
| chlorantraniliprole | 100 | 100 | 100 | 100 | 100 |

respectively, but still higher than that of chlorantraniliprole (0.000 012 3 mg/mL; Table 5).

Effects of Compounds 13b and 15a on Calcium Channels of Neurons from *S. exigua*. Figure 2 shows the change rate of peak current amplitude versus recording time in 0.1 and 0.001 mmol/L compound **13b**- and 0.1 mmol/L chlorantraniliprole-treated and control neurons. The peak currents were reduced to $54.71 \pm 4.95\%$ ($n = 6$), $74.63 \pm 5.37\%$ ($n = 5$), $39.43 \pm 5.02\%$ ($n = 8$), and $85.44 \pm 1.14\%$ ($n = 6$) of the initial value by the end of the 10 min recording, when the neurons were treated with 0.1 and 0.001 mmol/L compound **13b**, 0.1 mmol/L chlorantraniliprole, and the control, respectively. The peak current was reduced to $54.71 \pm 4.95\%$ ($n = 6$) when the neurons were treated with 0.1 mmol/L compound **13b**. In comparison to the control ($85.44 \pm 1.14\%$), compound **13b** at 0.001 mmol/L has a weak effect on $Ca^{2+}_{(L)}$ currents. The recorded peak currents of calcium channels treated with compound **13b** were in a concentration-dependent manner (Figure 2 and Table 6).

There was a significant change in maximal value of the peak current when the neurons were held at -70 mV with the treatment of 0.1 mmol/L compound **13b** (Figure 3). The calcium current decreased when compound **13b** was injected in the surrounding of the neuron cells. The maximal value of I_{Ca} (-2.1303 ± 0.0237 nA) shifted to the negative direction by approximately 10 mV after the neurons were treated with compound **13b** for 1 min. By the end of the 5 and 10 min recordings, I_{Ca} decreased to -1.6356 ± 0.0641 and -1.2738 ± 0.882 nA, respectively, and the $I-V$ curves were also shifted to the negative direction by approximately 10 mV during the recording. Unlike the results for I_{Ca} described above, the maximal value of I_{Ca} did not shift to a negative direction when the neurons were treated with 0.001 mM compound **13b**, from which we concluded that, when the concentration was ≤ 0.001 mM, there was no effect on calcium channels in the central neurons of *S. exigua* third larvae.

These results indicated that $Ca^{2+}_{(L)}$ channels of *S. exigua* neurons were modulated by compound **13b** and partly closed after the action. Although there was no notable change on activation voltage, the negatively shifted $I-V$ relationship curves indicate that voltage dependence was influenced by compound **13b**. All of these results suggest that the $Ca^{2+}_{(L)}$ channels of *S. exigua* neurons were the possible target of compound **13b**.

Figure 4 illustrated the change of $[Ca^{2+}]_i$ versus the recording time when the neurons were treated with compounds **13b**, **15a**, and chlorantraniliprole. The peak of $[Ca^{2+}]_i$ was $116.863 \pm 2.33\%$ ($n = 10$), $112.408 \pm 1.26\%$ ($n = 8$), and $123.292 \pm 2.17\%$ ($n = 21$) of the initial value by the end of the 10 min recording when the cells were treated with 1000 mg/L compound **13b**, 1000 mg/L compound **15a**, and 1000 mg/L

Table 5. LC₅₀ Values of Compounds 15d, 15e, and Chlorantraniliprole against Diamondback Moths

| compound | $y = a + bx$ | LC ₅₀ (mg/L) | R | 95% confidence interval |
|---------------------|---------------------|-------------------------|--------|---|
| 15d | $y = 11.04 + 1.68x$ | 0.00025 | 0.9974 | 0.00023–0.00028 |
| 15e | $y = 12.90 + 2.48x$ | 0.00065 | 0.9723 | 0.00053–0.00081 |
| chlorantraniliprole | $y = 11.50 + 1.32x$ | 0.0000123 | 0.9995 | 1.16×10^{-5} – 1.31×10^{-5} |




Figure 2. Variation of the peak current for the whole-cell calcium channels in neurons (treated with different concentrations of compound 13b and chlorantraniliprole) at different times during patch-clamp recording in comparison to the peak value at 0 min.

chlorantraniliprole, respectively. In comparison to the control ($99.91 \pm 2.56\%$), compounds 13b and 15a induced a $[Ca^{2+}]_i$ increase without extracellular Ca^{2+} . It indicated that compounds 13b and 15a could activate the calcium release channel in the endoplasmic reticulum (ER) membrane. Figure 3 also indicated that the recorded $[Ca^{2+}]_i$ (F/F_0) had a good positive correlation with bioactivities.

There were two kinds of calcium release channels in the ER membrane, namely, RyR and IP₃R Ca^{2+} channels.²⁶ To test which pathway was involved in the elevation of $[Ca^{2+}]_i$, the primary cultured neurons were dyed, loading with fluo-5N, then treated with heparin (10 mg/mL, a competitive antagonist of IP₃) for 20 min, and incubated with 1 μ M thapsigargin for 10 min. When external Ca^{2+} was free, IP₃ receptors were blocked using heparin and the intracellular calcium store was depleted with thapsigargin; the decrease of $[Ca^{2+}]_i$ was only attributed to compound 13b (1000 mg/L). These data indicated that RyRs would be the possible action target of this series of novel compounds.




Figure 3. Current–voltage relationship curves of whole-cell calcium channels recorded in neurons of *S. exigua* (treated with 0.1 mmol/L compound 13b and chlorantraniliprole) at different intervals (in minutes).




Figure 4. Change of $[Ca^{2+}]_i$ versus recording time when the neurons were treated with 1000 mg/L compounds 13b, 15a, and chlorantraniliprole.

In summary, two novel series of anthranilic diamides containing acylthiourea and acylurea moieties were synthesized,

Table 6. Analysis of the Peak Current Change Rate with Time in the Neurons of *S. exigua* (Treated by Compound 13b and Chlorantraniliprole)^a

| time (min) | 0.1 mM compound 13b | 0.001 mM compound 13b | 0.1 mM chlorantraniliprole | control |
|------------|---------------------|-----------------------|----------------------------|---------------------|
| 0 | 100 | 100 | 100 | 100 |
| 1 | 91.492 ± 3.47^b | 104.964 ± 1.34^c | 89.62 ± 2.16^b | 104.47 ± 2.33^b |
| 2 | 84.128 ± 2.16^b | 97.215 ± 7.58^b | 74.01 ± 8.01^b | 102.75 ± 2.45^c |
| 3 | 80.727 ± 4.77^b | 95.803 ± 7.26^b | 69.57 ± 9.43^b | 101.76 ± 2.7^c |
| 4 | 74.437 ± 5.31^b | 90.134 ± 6.41^b | 63.36 ± 7.71^b | 99.66 ± 1.64^c |
| 5 | 70.252 ± 8.48^b | 84.442 ± 5.48^b | 55.79 ± 4.73^b | 98.06 ± 1.01^c |
| 8 | 61.013 ± 3.28^b | 81.209 ± 3.69^b | 44.31 ± 2.62^b | 90.85 ± 0.65^b |
| 10 | 54.715 ± 4.95^b | 74.632 ± 6.12^b | 39.685 ± 5.02^b | 85.44 ± 1.14^b |

^aValues are the mean \pm SD. ^bSignificant difference at $p < 0.01$. ^cSignificant difference at $p < 0.05$.

and their larvicidal activities against oriental armyworm, mosquito larvae, and diamondback moth were evaluated. The results indicated that the introduction of acylthiourea moiety into some structures could retain their insecticidal activity; 8 of the 15 compounds (13a–13e, 14a–14e, and 15a–15e) exhibited 100% larvicidal activity at 10 mg/L against oriental armyworm. However, the introduction of acylurea moiety decreased the insecticidal activity; only 3 of the 11 compounds (17a–17k) exhibited 100% larvicidal activity at 200 mg/L against oriental armyworm. We speculated that it might be the solubility factor involved because compounds containing acylthiourea moiety exhibited better solubility in organic solvents during our experiments. The effects on calcium channels of neurons from *S. exigua* indicated that the title compounds influenced the same target RyRs as chlorantraniliprole. Also, the experiment of intracellular calcium of neurons provided us a rapid detection for the activity of the target compound.

■ ASSOCIATED CONTENT

● Supporting Information

¹H NMR, HRMS, and melting point data for compounds 13a–13e, 14a–14e, 15a–15e, and 17a–17k. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: liyx128@nankai.edu.cn (Y.-X.L.); nkzml@vip.163.com (Z.-M.L.).

Funding

This work has been supported by the National Basic Research Program of China (973 Project 2010CB126106) and the National Natural Science Foundation of China (NNSFC) (31000861). We are also grateful for the financial support from the National Key Technologies Research and Development Program (2011BAE06B05), the Fundamental Research Funds for the Central Universities, and the Natural Science Foundation Project of Tianjin Ministry of Science and Technology of China (11JCYBJC04500).

Notes

The authors declare no competing financial interest.

■ REFERENCES

- (1) Lahm, G. P.; Selby, T. P.; Frendenberger, J. H.; Stevenson, T. M.; Myers, B. J.; Seburyamo, G.; Smith, B. K.; Flexner, L.; Clark, C. E.; Cordova, D. Insecticidal anthranilicdiamides: A new class of potent ryanodine receptor activators. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4898–4906.
- (2) Feng, Q.; Liu, Z. L.; Xiong, L. X.; Wang, M. Z.; Li, Y. Q.; Li, Z. M. Synthesis and insecticidal activities of novel anthranilicdiamides containing modified *n*-pyridylpyrazoles. *J. Agric. Food Chem.* **2010**, *58*, 12327–12336.
- (3) Clark, D. A.; Lahm, G. P.; Smith, B. K.; Barry, J. D.; Clagg, D. G. Synthesis of insecticidal fluorinated anthranilicdiamides. *Bioorg. Med. Chem.* **2008**, *16*, 3163–3170.
- (4) Lahm, G. P.; Cordova, D.; Barry, J. D. New and selective ryanodine receptor activators for insect control. *Bioorg. Med. Chem.* **2009**, *17*, 4127–4133.
- (5) Lahm, G. P.; Stevenson, T. M.; Selby, T. P.; Freudenberger, J. H.; Cordova, D.; Flexner, L.; Bellin, C. A.; Dubas, C. M.; Smith, B. K.; Hughes, K. A.; Hollingshaus, J. G.; Clark, C. E.; Benner, E. A. Rynaxypyr: A new insecticidal anthranilicdiamide that acts as a potent and selective ryanodine receptor activator. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6274–6279.
- (6) Li, B.; Yang, H. B.; Wang, J. F.; Yu, H. B.; Zhang, H.; Li, Z. N.; Song, Y. Q. 1-Substituted pyridyl-pyrazolyl amide compounds and uses thereof. U.S. Patent 201146186 A1, 2011.
- (7) Alig, B.; Fischer, R.; Funke, C.; Gesing, E. F.; Hense, A.; Malsam, O.; Drewes, M. W.; Gorgens, U.; Murata, T.; Wada, K.; Arnold, C.; Sanwald, E. Anthranilic acid diamide derivative with hetero-aromatic and hetero-cyclic substituents. WO Patent 2007144100 A1, 2007.
- (8) Loiseleur, O.; Hall, R. G.; Stoller, A. D.; Graig, G. W.; Jeanguenat, A.; Edmunds, A. Novel insecticides. WO Patent 2009024341 A2, 2009.
- (9) Loiseleur, O.; Durieux, P.; Trah, S.; Edmunds, A.; Jeanguenat, A.; Stoller, A.; Hughes, D. J. Pesticides containing a bicyclic bisamide structure. WO Patent 2007093402 A1, 2007.
- (10) Kruger, B.; Hense, A.; Alig, B.; Fischer, R.; Funke, C.; Gesing, E. R.; Malsam, A.; Drewes, M. W.; Arnold, C.; Lummen, P.; Sanwald, E. Dioxazine- and oxadiazine substituted arylamides. WO Patent 2007031213 A1, 2007.
- (11) Dumas, D. J. Process for preparing 2-amido-5-cyanobenzoic acid derivatives. WO Patent 2009111553 A1, 2009.
- (12) Hughes, D. J.; Peace, J. E.; Riley, S.; Russell, S.; Swanborough, J. J.; Jeanguenat, A.; Renold, P.; Hall, R. G.; Loiseleur, O.; Trah, S.; Wenger, J. Synergistic pesticidal mixtures with nitrogen-containing component. WO Patent 2007009661 A2, 2007.
- (13) Sun, R. F.; Zhang, Y. L.; Chen, L.; Li, Y. Q.; Li, Q. S.; Song, H. B.; Huang, R. Q.; Bi, F. C.; Wang, Q. M. Design, synthesis, bioactivity, and structure–activity relationship (SAR) studies of novel benzoylphenylureas containing oxime ether group. *J. Agric. Food Chem.* **2008**, *56*, 11376–11391.
- (14) Saeed, A.; Batool, M. Synthesis and bioactivity of some new 1-tolyl-3-aryl-4-methylimidazole-2-thiones. *Med. Chem. Res.* **2007**, *16*, 143–154.
- (15) Cesarini, S.; Spallarossa, A.; Ranise, A.; Schenone, S.; Rosano, C.; La Colla, P.; Sanna, G.; Busonera, B.; Loddo, R. N-Acylated and N,N'-diacylated imidazolidine-2-thione derivatives and N,N'-diacylated tetrahydropyrimidine-2(1H)-thione analogues: Synthesis and antiproliferative activity. *Eur. J. Med. Chem.* **2009**, *44*, 1106–1118.
- (16) Rao, X. P.; Wu, Y.; Song, Z. Q.; Shang, S. B.; Wang, Z. D. Synthesis and antitumor activities of unsymmetrically disubstitutedacylthioureas fused with hydrophenanthrene structure. *Med. Chem. Res.* **2011**, *20*, 333–338.
- (17) Sun, C. W.; Huang, H.; Feng, M. Q.; Shi, X. L.; Zhang, X. D.; Zhou, P. A novel class of potent influenza virus inhibitors: Polysubstitutedacylthiourea and its fused heterocycle derivatives. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 162–166.
- (18) Hallur, G.; Jimeno, A.; Dalrymple, S.; Zhu, T.; Jung, M. K.; Hidalgo, M.; Isaacs, J. T.; Sukumar, S.; Hamel, E.; Khan, S. R. Benzoylphenylurea sulfur analogues with potent antitumor activity. *J. Med. Chem.* **2006**, *49*, 2357–2360.
- (19) Dong, W. L.; Xu, J. Y.; Xiong, L. X.; Liu, X. H.; Li, Z. M. Synthesis, structure and biological activities of some novel anthranilic acid esters containing *N*-pyridylpyrazole. *Chin. J. Chem.* **2009**, *27*, 579–586.
- (20) Dong, W. L.; Liu, X. H.; Xu, J. Y.; Li, Z. M. Design and synthesis of novel anthranilicdiamides containing 5,7-dimethyl[1,2,4]triazolo-[1,5-*a*]pyrimidine. *J. Chem. Res.* **2008**, *9*, 530–533.
- (21) Abbott, W. S. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* **1925**, *18*, 265–267.
- (22) Chen, L.; Huang, Z. Q.; Wang, Q. M.; Shang, J.; Huang, R. Q.; Bi, F. C. Insecticidal benzoylphenylurea-S-carbamate: A new propesticide with two effects of both benzoylphenylureas and carbamates. *J. Agric. Food Chem.* **2007**, *55*, 2659–2663.
- (23) Li, Y. X.; Mao, M. Z.; Li, Y. M.; Xiong, L. X.; Li, Z. M.; Xu, J. Y. Modulations of high-voltage activated Ca²⁺ channels in the central neurones of *Spodoptera exigua* by chlorantraniliprole. *Physiol. Entomol.* **2011**, *36*, 230–234.
- (24) Takahashi, A.; Camacho, P.; Lechleiter, J. D.; Herman, B. Measurement of intracellular calcium. *Physiol. Rev.* **1999**, *79*, 1089–1125.
- (25) Liu, A. P.; Hu, Z. B.; Wang, Y. J.; Wang, X. G.; Huang, M. Z.; Ou, X. M.; Mao, C. H.; Pang, H. L.; Huang, L. Preparation of pyrazole

containing 2-amino-*N*-oxybenzamide compounds as pesticides. CN Patent 101337959 A, 2009.

(26) Striggow, F.; Ehrlich, B. E. Ligand-gated calcium channels inside and out. *Curr. Opin. Cell. Biol.* **1996**, *8*, 490–495.