

Tetrahedron Letters 40 (1999) 8255-8259

TETRAHEDRON LETTERS

## Unusual biomimetic oxidations of indoles: synthesis of suaveoline and an alkaloid G analogue

Patrick D. Bailey,\* Keith M. Morgan, Georgina M. Rosair and Rhodri Ll. Thomas Department of Chemistry, Heriot-Watt University, Riccarton, Edinburgh, EH14 4AS UK

Received 26 July 1999; accepted 31 August 1999

## Abstract

The one-pot conversion of ajmaline 1 into the nitrile 4 (X-ray structure) and suaveoline 2a is reported; both reactions utilise hydroxylamine hydrochloride in a refluxing alcohol, and involve unusual multi-step mechanisms in which the product ratio can be controlled by the choice of the alcohol. In an additional unusual oxidation, the synthetic intermediate 14 used in a synthesis of suaveoline can be oxidatively cyclized to 16. © 1999 Elsevier Science Ltd. All rights reserved.

Keywords: alkaloids; indoles; oxidation.

The interconversion of natural products remains an important aspect of synthetic work on biologically active compounds. It can not only provide a rapid route to advanced intermediates for the semi-synthesis of natural products or analogues thereof, but can also sometimes shed light on biosynthetic pathways. Since its discovery in 1972,<sup>1</sup> the biosynthetic origin of suaveoline **2a** has been investigated by several groups, and it has been shown that ajmaline **1** (which is also found in the same plants) is converted into suaveoline **2a**, *N*-methylsuaveoline **2b** and raumacline **3** in vivo (Scheme 1).<sup>2</sup>



Scheme 1. Known biotransformations from ajmaline 1

We have been interested in the indole alkaloids of this family for some time, not only because of their biological activity and potential, but also because of the synthetic challenge posed by their common 1,3-disubstituted tetrahydro- $\beta$ -carboline ring system. Initial mechanistic studies<sup>3</sup> led us to develop a simple

<sup>\*</sup> Corresponding author.

<sup>0040-4039/99/\$ -</sup> see front matter © 1999 Elsevier Science Ltd. All rights reserved. *P11:* S0040-4039(99)01684-6

stereoselective method of gaining access to such tricyclic systems,<sup>4</sup> and we have subsequently applied this methodology to a number of total syntheses.<sup>5</sup>

In ongoing synthetic work, we wished to obtain advanced intermediates for work on ajmaline, suaveoline, and related alkaloids. Some of the published degradation/inter-conversion procedures on ajmaline are summarized in Scheme 2. We were anxious to obtain the cyano-alcohol 4, because it can be re-converted into ajmaline,<sup>6,7</sup> and is very similar to an advanced synthetic intermediate 7 that we already had in hand;<sup>5a</sup> we also wanted to prepare a sample of suaveoline (which is not readily available) for comparison with our synthetic material.<sup>5a</sup>



Scheme 2. Reagents and conditions: (i) BzCl, PhH, then  $Pb(OAc)_4$ ; (ii) reflux in MeI, then  $NH_3$  (aq);<sup>1</sup> (iii) NaBH<sub>4</sub>, H<sub>2</sub>O-MeOH, rt, then ZCl, 1 M NaOH-DCM, 0°C; (iv) Pb(OAc)<sub>4</sub>, then H<sub>2</sub>, Pd-C, MeOH;<sup>2</sup> (v) NH<sub>2</sub>OH·HCl, reflux in H<sub>2</sub>O, then AcOH, Ac<sub>2</sub>O, HCl, then KOH, MeOH; (vi) LiAlH<sub>4</sub>, Et<sub>2</sub>O, rt<sup>6</sup>

To our surprise, treatment of ajmaline 1 with hydroxylamine hydrochloride in refluxing ethanol yielded the cyano-alcohol 4 directly in 89% yield, whose structure was confirmed by single crystal X-ray structure determination (Fig. 1).<sup>8</sup> This reaction presumably proceeded via the oxime 8, and the unexpectedly facile dehydration might have involved intramolecular proton abstraction, as suggested in Scheme 3.

We were further surprised to discover that this reaction was acutely solvent dependent, as indicated in Table 1. The most useful and unexpected result was the direct conversion of ajmaline 1 into suaveoline **2a** (30% yield) on refluxing the former in *n*-butanol with hydroxylamine hydrochloride. As this is an oxidation reaction, we speculated that oxygen was the oxidant, and this was supported by the suppression of suaveoline formation when the reaction was conducted under argon. Further insight into the mechanism was gained by the observation that ammonium chloride/air did not effect the conversion of  $1 \rightarrow 2a$  in refluxing butanol, strongly implicating an oxidized form of hydroxylamine as the actual



Figure 1. X-Ray crystal structure of 4



Scheme 3. Reagents and conditions: see Table 1

 Table 1

 Conditions for the conversion of ajmaline 1 into 2 or 4

Entry	Conditions	Product(s)
1	1, NH2-OH.HCl,	
	reflux in EtOH, air	4 (89%)
2	1, NH <sub>2</sub> -OH.HCl,	
	reflux in n-PrOH, air	2a (<5%) + 4 (31%)
3	1, NH <sub>2</sub> -OH.HCl,	
	reflux in n-BuOH, air	r <b>2a</b> (30%)
4	1, NH <sub>2</sub> -OH.HCl,	. ,
	reflux in PhMe, air	essentially no reaction
5	1, NH <sub>2</sub> -OH.HCl,	-
	reflux in n-BuOH, As	r <b>2a</b> (<5%) + mixture
6	1, NH4Cl,	
	reflux in <i>n</i> -BuOH, air	r 1 (65%) (no 2a or 4)

oxidizing agent. In further support of this, the diimine of the protected di-aldehyde **11b** does cyclize to a pyridine derivative under similar conditions;<sup>5 a</sup> we are therefore confident that, had NH<sub>4</sub>Cl/air/butanol reflux generated any of the dialdehyde **11a**, then we would have observed the formation of suaveoline **2a**. On the basis of these results, it would appear that the hydroxylamine is involved in the oxidative mechanism, perhaps forming [NO]<sup>+</sup> which is trapped as the nitrite, and further oxidized to the nitrate, as indicated in Scheme 4.



Scheme 4. Proposed mechanism for conversion of 1 into 2a

With a sample of suaveoline in hand, we attempted to complete a synthesis of this alkaloid from the advanced intermediate 13 that we had prepared earlier from L-tryptophan in nine steps.<sup>5b</sup> To achieve this (see Scheme 5), we reduced the nitrile 13 to the amine with LiAlH<sub>4</sub>, and protected the amine with the Boc group to give 14, prior to oxidation of the secondary alcohol. Surprisingly, both TPAP and Swern oxidation conditions led to virtually no reaction, but oxidation with PCC generated a major and minor product, both possessing a ketone functionality. The major product (31%) was identified from extensive NMR analysis as the bridged polycycle 16,<sup>9</sup> whilst the desired ketone 15a was formed in only 16% yield.



Scheme 5. Reagents and conditions: (i) LiAlH<sub>4</sub>, THF, rt, O/N; (ii) Boc<sub>2</sub>O, NaHCO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 30 min (64% from **13**); (iii) PCC, Florisil<sup>®</sup> (1:1), CH<sub>2</sub>Cl<sub>2</sub>, O/N (47%); (iv) TFA (100%); (v) *n*-PrCHO, O<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, MS, reflux; (vi) H<sub>2</sub>, Pd(OH)<sub>2</sub>-C (96%);<sup>5</sup>a (vii) POCl<sub>3</sub>/py then DIBAL-H (-78°C) (86%);<sup>5</sup> (viii) see Fu and Cook<sup>10</sup>

The polycycle 16 is of interest because of the existence of indole alkaloids cyclized at this position such as Alkaloid G 17.<sup>11</sup> As far as we are aware, this is the first example of transition metal oxidation leading to derivatization at this position, with DDQ being the usual reagent of choice;<sup>12</sup> presumably, one-electron oxidation processes operate in both cases, offering two alternative reagents for accessing compounds of this type.

Deprotection of the Boc-derivative 15a gave an unstable amino-ketone 15b, which was refluxed with butanal in the presence of molecular oxygen; from the product mixture, *N*-benzylsuaveoline 2c could be detected in small amounts by comparison (TLC and mass spectrometry) with authentic material. This constitutes a new total synthesis of suaveoline 2a, although we have already published a more efficient route from 13 to 2a.<sup>5b</sup>

In summary, we have used ajmaline 1 as an advanced yet cheap indolic building block for the construction of the nitrile 4, for use in the semi-synthesis of other targets in this family, and we have effected a one step conversion of ajmaline 1 into suaveoline 2a. We have also identified an unexpected and useful cyclization procedure at the 3'-indole position using PCC.

## Acknowledgements

We thank Dr. A. S. F. Boyd and Dr. R. Fergusson for NMR and mass spectra, and EPSRC for financial support.

## References

- 1. Majumdar, S. P.; Potier, P.; Poisson, J. Tetrahedron Lett. 1972, 13, 1563-1566.
- 2. Polz, L.; Stöckigt, J. Tetrahedron Lett. 1990, 31, 6693-6696.
- 3. (a) Bailey, P. D. Journal of Chemical Research 1987, 202–203; (b) Bailey, P. D.; Hollinshead, S. P.; McLay, N. R. Tetrahedron Lett. 1987, 28, 5177–5180.
- (a) Bailey, P. D.; Hollinshead, S. P.; McLay, N. R.; Morgan, K. M.; Palmer, S. J.; Prince, S. N.; Reynolds, C. D.; Wood, S. D. J. Chem. Soc., Perkin Trans. 1 1993, 431–439; (b) Bailey, P. D.; Moore, M. H.; Morgan, K. M.; Smith, D. I.; Vernon, J. M. Tetrahedron Lett. 1994, 35, 3587–3588.
- (a) Bailey, P. D.; Morgan, K. M. Chem. Commun. 1996, 1479–1480; (b) Bailey, P. D.; Collier, I. D.; Hollinshead, S. P.; Moore, M. H.; Morgan, K. M.; Smith, D. I.; Vernon, J. M. J. Chem. Soc., Perkin Trans. 1 1997, 1209–1214; (c) Bailey, P. D.; McLay, N. R. J. Chem. Soc., Perkin Trans. 1 1993, 441–449; (d) Bailey, P. D.; Hollinshead, S. P.; McLay, N. R.; Everitt, J. H.; Reynolds, C. D.; Wood, S. D.; Giordano, F. J. Chem. Soc., Perkin Trans. 1 1993, 451–458.
- 6. Anet, F. A. L.; Chakravarti, D.; Robinson, R.; Schlittler, E. J. Chem. Soc. 1954, 1242-1260.
- 7. Masamune, S.; Ang, S. K.; Egli, C.; Nakatsuka, N.; Sarkar, S. K.; Yasunari, Y. J. Am. Chem. Soc. 1967, 89, 2506–2507.
- 8. Crystal data for 4: yellow crystal (0.6×0.4×0.25 mm) from CHCl<sub>3</sub>/methanol mounted in epoxy resin glue in a thin-walled glass capillary, C<sub>20</sub>H<sub>25</sub>N<sub>3</sub>O, *M*=323.43, orthorhombic, space group *P*2<sub>1</sub>2<sub>1</sub>2<sub>1</sub>, *a*=6.8037(4) *b*=11.4296(8), *c*=22.018(3) Å, *V*=1712.2(2) Å<sup>3</sup>, *Z*=4, *D<sub>c</sub>*=1.255 g cm<sup>-3</sup>, *m*=0.079 mm<sup>-1</sup>, 2183 reflections were measured on a Siemens P4 diffractometer with Mo Ka radiation, l=0.71073 Å, w mode, q range 1.85 to 24.99° at 293 K. The structure was refined to *R*<sub>1</sub>=0.0457, *wR*<sub>2</sub>=0.1217 and goodness of fit 1.053 for 1855 unique observed [*I*>2σ(*I*)] data and 219 parameters with maximum and minimum residual electron density peaks of 0.32 and -0.32 eÅ<sup>3</sup>, respectively. The Flack x parameter was refined to 1(3) i.e. absolute structure cannot be determined from X-ray data. Data collection and reduction were performed using the program XSCANS [Siemens Analytical Instruments, Inc., Madison, Wisconsin, USA, 1994]. Direct methods solution and refinements (full-matrix least squares on *F*<sup>2</sup>) were performed using SHELXTL/PC version 5.03 [G. M. Sheldrick, 1994, SHELXTL/PC. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA]. Atomic coordinates, displacement parameters and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre (CCDC).
- 9. Selected key data for **16**:  $\delta_C$  (50 MHz, CDCl<sub>3</sub>) 28.5(q), 28.7(q), 43.3(t), 44.8(t), 48.7(d), 49.5(d), 50.1(d), 57.7(t), 60.5(d), 79.5(s), 107.2(s), 108.6(d), 119.9(d), 121.5(d), 121.9(d), 126.2(s), 127.6(d), 128.6(d), 134.5(s), 137.9(2×s), 155.0(s), 206.7(s); M<sup>+</sup> 457.2359, calcd for C<sub>28</sub>H<sub>31</sub>N<sub>3</sub>O<sub>3</sub>: 457.2365.
- 10. Fu, X.; Cook, J. M. J. Org. Chem. 1993, 58, 661-672.
- 11. Endress, S.; Takayama, H.; Suda, S.; Kitajima, M.; Aimi, N.; Sakai, S.; Stöckigt, J. Phytochemistry 1993, 32, 725-730.
- 12. Li, J.; Cook, J. M. J. Org. Chem. 1998, 63, 4166-4167.