

- vasiliou, *Science*, **196**, 549 (1977).
 (13) L. C. Tang and G. C. Cotzias, *Proc. Natl. Acad. Sci. U.S.A.*, **74**, 2126 (1977).
 (14) G. C. Cotzias and L. C. Tang, *Science*, **197**, 1094 (1977).

- (15) L. C. Tang and G. C. Cotzias, *Proc. Natl. Acad. Sci. U.S.A.*, **74**, 1242 (1977).
 (16) J. G. Cannon, J. P. O'Donnell, T. Lee, and C. R. Hoppin, *J. Med. Chem.*, **18**, 1212 (1975).

Study of the Structural Requirements for Dopa Potentiation and Oxotremorine Antagonism by L-Prolyl-L-leucylglycinamide

Rodney L. Johnson,* Edward E. Smissman,¹

Department of Medicinal Chemistry, School of Pharmacy, University of Kansas, Lawrence, Kansas 66045

and Nicholas P. Plotnikoff

Division of Medicinal Chemistry and Pharmacology, Abbott Laboratories, North Chicago, Illinois 60064.

Received June 13, 1977

A number of analogues of the tripeptide L-prolyl-L-leucylglycinamide (1) were synthesized and evaluated in the Dopa potentiation and oxotremorine antagonism tests. The replacement of the glycinamide residue with either the glycine methylamide, glycine, aminoacetonitrile, amino-2-propanone, semicarbazide, or β -alaninamide residues resulted in a loss of activity in both tests. A 1:1 mixture of L-prolyl-L-leucyl-(-)-thiazolidine-2-carboxamide (8) and L-prolyl-L-leucyl-(+)-thiazolidine-2-carboxamide (9) showed marked activity in the Dopa potentiation test but was unable to antagonize the tremors induced by oxotremorine. L-Prolyl-L-leucyl-L-prolinamide (11), on the other hand, was active in the oxotremorine antagonism test but inactive in the Dopa potentiation test. The replacement of the pyrrolidine ring of 1 with either a thiazolidine or cyclopentane ring system caused a loss of activity. The cyclopentanecarboxylic acid analogue 13, however, was found to have moderate activity in the serotonin potentiation test.

Several of the hypothalamic releasing and release-inhibiting hormones have recently been shown to have effects on the central nervous system which are independent of their endocrine effects.² Among these polypeptides is L-prolyl-L-leucylglycinamide (1). This tripeptide has been


effects of deserpidine,¹¹ potentiates the behavioral effects of apomorphine,¹² attenuates puromycin-induced amnesia,¹³ and facilitates the development of morphine dependence.¹⁴ Moreover, a number of preliminary clinical studies have indicated that 1 may be of potential value in the treatment of Parkinson's disease.^{15,16}

In an effort to determine the structural features of 1 that enable this tripeptide to potentiate the behavioral effects of L-Dopa and antagonize the tremors induced by oxotremorine, we have synthesized several analogues of 1 in which the prolyl and glycinamide residues have been modified. In order to determine the importance of the primary carboxamide moiety, the glycinamide residue has been replaced with the glycine methylamide, glycine, aminoacetonitrile, and amino-2-propanone residues, analogues 2–5, respectively. In the case of analogue 6 an imino group has been substituted for the methylene group while in compound 7 an additional methylene group has been inserted into the peptide chain. In addition to these modifications, several analogues (8–11) have been synthesized where either a thiazolidine or pyrrolidine ring system has been incorporated into the glycinamide portion of the molecule.

The prolyl residue has been modified in an attempt to determine the importance of the pyrrolidine ring system. Castensson et al.¹⁰ have reported that the replacement of the prolyl residue with a pyroglutamyl residue provides a compound (14) which is an even better antagonist of oxotremorine-induced tremors than 1. We have thus synthesized compounds 12 and 13 whereby the pyrrolidine ring has been replaced by a thiazolidine and cyclopentane ring system, respectively.


Results and Discussion

Chemistry. The protected dipeptide Z-Pro-Leu-OH (15), which was prepared in a manner similar to that previously described by Cash,¹⁷ served as the key intermediate in the synthesis of 2–11. Aminoacetonitrile, semicarbazide, β -alaninamide, L-prolinamide, and methyl



postulated to be the hypothalamic hormone that inhibits the release of melanocyte-stimulating hormone from the anterior pituitary gland.³⁻⁶ The extra-endocrine effects of 1 were first demonstrated by Plotnikoff et al.⁷ with Everett's Dopa potentiation test.⁸ In this test 1 was found to potentiate the behavioral effects of L-3,4-dihydroxyphenylalanine (L-Dopa) in both normal and hypophysectomized mice. Subsequent studies with 1 have also shown that this tripeptide antagonizes the central and peripheral effects of oxotremorine,^{9,10} reverses the sedative

* Address correspondence to this author at the University of Kansas Medical Center, Department of Pharmacology, Kansas City, Kansas 66103.



glycinate were coupled to **15** using the mixed anhydride method.^{18,19} The benzyloxycarbonyl-protecting group was removed from Z-Pro-Leu-NHCH₂CN (**19**), Z-Pro-Leu-NHNHCONH₂ (**22**), Z-Pro-Leu-β-Ala-NH₂ (**23**), and Z-Pro-Leu-Pro-NH₂ (**27**) by hydrogenolysis to give directly analogues **4**, **6**, **7**, and **11**. The protected ester Z-Pro-Leu-Gly-OCH₃ (**16**), on the other hand, was treated with either methylamine or aqueous sodium hydroxide to give Z-Pro-Leu-Gly-NHCH₃ (**17**) and Z-Pro-Leu-Gly-OH (**18**), respectively. Removal of the benzyloxycarbonyl group by hydrogenolysis provided **2** and **3**.

Amino-2-propanone semicarbazone was obtained from glycine by way of the Dakin-West reaction²⁰ and coupled to **15** via the mixed anhydride method to give Z-Pro-Leu-NHCH₂C(CH₃)=NNHCONH₂ (**20**). Treatment of **20** with aqueous hydrochloric acid afforded Z-Pro-Leu-NHCH₂COCH₃ (**21**) which was subsequently converted to **5** with hydrobromic acid in acetic acid.

The synthesis of diastereoisomers **8** and **9** required the prior synthesis of the thiazolidine amino acid residue. Thus, 2-aminoethanethiol was condensed with glyoxylic acid to give racemic thiazolidine-2-carboxylic acid (**24**). Treatment of **24** with thionyl chloride and ethanol yielded racemic ethyl thiazolidine-2-carboxylate (**25**) which was easily resolved using the *d*- and *l*-tartaric acids. Previous attempts by us to resolve the primary amide of **24** with the tartaric and di-*p*-toluoyltartaric acids proved unsuccessful. Both (–)- and (+)-**25** were coupled to **15** using *N,N*-dicyclohexylcarbodiimide. The resulting tripeptide esters were converted to primary amides and then decarboxylated with hydrobromic acid to give **8** and **9**, respectively. The isomeric thiazolidine analogue **10** was prepared from L-thiazolidine-4-carboxamide (**26**) using similar coupling and deprotection procedures employed for the synthesis of **8** and **9**.

In the synthesis of **12**, methyl L-leucylglycinate was coupled to *N*-*tert*-butyloxycarbonyl-L-thiazolidine-4-carboxylic acid by the mixed anhydride method. After the resulting ester was converted to a primary amide with ammonia, the *tert*-butyloxycarbonyl protecting group was removed with hydrobromic acid in acetic acid to give **12**. Compound **13**, on the other hand, was prepared by reacting the *N*-hydroxysuccinimide ester of cyclopentanecarboxylic acid with L-leucylglycinamide.

Pharmacology. Compounds **2**–**13** were evaluated in the Dopa potentiation⁸ and oxotremorine antagonism tests.^{9,21} The results of these tests are depicted in Tables I and II, respectively. They indicate that there are strict structural requirements for the ability of **1** to potentiate the behavioral effects of L-Dopa and antagonize the central and peripheral effects of oxotremorine. In particular, the primary carboxamide moiety appears to be necessary for the activity of **1** since the replacement of this group with the –CONHCH₃, –COOH, –CN, and –COCH₃ moieties, analogues **2**–**5**, respectively, results in a loss of activity in both the Dopa potentiation test and the oxotremorine

Table I. Activity of L-Prolyl-L-leucylglycinamide (**1**) and Its Analogues in the Dopa Potentiation Test^a

Compd	Dose, mg/kg ip	Rating	Compd	Dose, mg/kg ip	Rating
1	0.1	3	8	2.5	1
	1	3		10	2
	10	3	9	10	1
2	10	1	8 + 9^b	0.625	1
3	10	1		2.5	2
4	2.5	1		10	3
	10	2	10	10	1
5	2.5	2	11	10	1
	10	2	12	10	1
6	10	1	13	2.5	1
7	10	1		10	2

^a See Experimental Section for a description of the test and ratings. ^b Ratio of **8** and **9** is 1:1.

Table II. Activity of L-Prolyl-L-leucylglycinamide (**1**) and Its Analogues in the Oxotremorine Antagonism Test^a

Compd	Dose, mg/kg ip	Rating	Compd	Dose, mg/kg ip	Rating
1	1	22	8	20	19
	4	12	9	20	22
	20	7	8 + 9^b	20	22
2	20	22	10	20	22
3	20	20	11	10	22
4	20	22		20	14
5	20	22	12	10	22
6	20	22		20	17
7	20	19	13	20	22

^a See Experimental Section for a description of the test and ratings. ^b Ratio of **8** and **9** is 1:1.

antagonism test. These results suggest that the carboxamide group is important for the interaction of **1** with its receptor. The results also show that the substitution of a semicarbazide or β-alaninamide residue for the glycine residue, as was done in compounds **6** and **7**, also results in a loss of pharmacological activity.

In the case of the thiazolidine-2-carboxamide analogues, diastereoisomer **8** was found to be moderately active in the Dopa potentiation test at a dose of 10 mg/kg while **9** was completely inactive. Interestingly, a 1:1 mixture of **8** and **9** markedly potentiated the behavioral effects of L-Dopa. The reason for this surprising result is not known although a possible explanation may be that the inactive diastereoisomer inhibits the metabolism of the active one. Neither of the diastereoisomers nor the mixture antagonized the effects of oxotremorine, however. In contrast, the isomeric thiazolidine analogue **10** was inactive in both tests while the prolinamide analogue, compound **11**, was found to be capable of antagonizing the effects of oxotremorine but incapable of potentiating the behavioral effects of L-Dopa. The observed differences in activity between the thiazolidine-2-carboxamide and prolinamide analogues suggest that the mechanism of action of these analogues is different.

Although Castensson et al.^{10,22} have shown that the replacement of the prolyl residue with the pyroglutamyl residue yields a compound with enhanced activity in the oxotremorine antagonism test, our results indicate that the pyrrolidine ring cannot be replaced by the larger thiazolidine ring system or by the cyclopentane ring system without losing biological activity. Surprisingly, the cyclopentanecarboxylic acid analogue **13** at a dose of 10 mg/kg was found by us to have moderate activity in the serotonin potentiation test.²³ Although thyrotropin-releasing hormone (<Glu-His-Pro-NH₂) has previously been shown to be active in the serotonin potentiation test, **1** has

been found to be inactive.²⁴ None of the other compounds prepared in this study were found to have serotonin potentiating activity.

Experimental Section

Melting points were determined on a Thomas-Hoover Unimelt apparatus and on a Mettler FP2-FP51 system. Specific rotations were measured with a Perkin-Elmer 141 polarimeter. Microanalyses were performed either on a Hewlett-Packard 185B C, H, N analyzer, University of Kansas, by Midwest MicroLab Inc., Indianapolis, Ind., or by the Australian Microanalytical Service, University of Melbourne, Melbourne, Australia. Where analyses are indicated by symbols of the elements, analytical results obtained were within $\pm 0.4\%$ of the theoretical values. IR spectra were recorded on either a Beckman IR-33 or a Perkin-Elmer 257 spectrophotometer. NMR spectra were recorded on either a Perkin-Elmer R-12 spectrophotometer or Varian Associates T-60 and EM-360 spectrophotometers with either Me₄Si or sodium trimethylsilylpropanesulfonate as the internal standards. TLC was carried out on Merck 0.25-mm silica gel 60 F-254 plates.

N-Benzoyloxycarbonyl-L-prolyl-L-leucine (15). The method of Cash¹⁷ was modified in the following manner. After the THF was removed in vacuo from the acidified reaction mixture, the aqueous mixture remaining was extracted with chloroform. The chloroform extract was washed with water and dried over Na₂SO₄. Removal of the chloroform in vacuo gave a viscous oil which when dissolved in ether and allowed to stand at room temperature afforded crystalline product: mp 136.5–137 °C; $[\alpha]_D^{24}$ –65.2° (c 0.8, MeOH) [lit.¹⁷ mp 138.5–139.5 °C; $[\alpha]_D^{22}$ –63.0° (c 5, MeOH)].

N-Benzoyloxycarbonyl-L-prolyl-L-leucylglycine Methylamide (17). Methyl *N*-benzyloxycarbonyl-L-prolyl-L-leucylglycinate²⁵ (16, 3.5 g, 8 mmol) was dissolved in 30 mL of ethanol saturated with CH₃NH₂. The solution was stirred at room temperature for 24 h and then stripped of solvent and excess CH₃NH₂ under reduced pressure to give a colorless oil. Trituration in Et₂O afforded 3.4 g (98%) of the methylamide: mp 148–150 °C. An analytical sample was obtained by recrystallizing a portion of this material from EtOAc–acetone: mp 149–151 °C; $[\alpha]_D^{22}$ –60.5° (c 0.5, MeOH). Anal. (C₂₂H₃₂N₄O₅) C, H, N.

L-Prolyl-L-leucylglycine Methylamide (2). A solution of 17 (2.0 g, 4.6 mmol) in 65 mL of MeOH was added to a flask containing 400 mg of 5% Pd/C and hydrogen was bubbled into the reaction mixture for 30 min. The mixture was filtered and the filtrate stripped of solvent in vacuo to yield 1.3 g of crude product as a colorless oil. This material was chromatographed on a silica gel dry column (Woelm activity III, 75 × 2.5 cm) with MeOH as the developing solvent. The portion of the column between the *R_f* values of 0.4 and 0.2 was isolated and extracted with CHCl₃. Evaporation of the CHCl₃ yielded 700 mg (51%) of 2 as a hygroscopic solid: mp 63–66 °C; $[\alpha]_D^{27}$ –37.6° (c 0.9, MeOH). Anal. (C₁₄H₂₆N₄O₃·H₂O) C, H, N.

The oxalate salt of 2 provided a suitable nonhygroscopic derivative and was prepared by treating a solution of oxalic acid (40 mg, 0.44 mmol) in THF with a solution of 2 (100 mg, 0.34 mmol) in THF. Recrystallization of the resulting precipitate from isopropyl alcohol yielded 100 mg (76%) of the oxalate salt: mp 168–169.5 °C; $[\alpha]_D^{22}$ –40.2° (c 0.54, H₂O). Anal. (C₁₆H₂₈N₄O₇) C, H, N.

L-Prolyl-L-leucylglycine (3). *N*-Benzyloxycarbonyl-L-prolyl-L-leucylglycine²⁶ (18, 700 mg, 1.67 mmol) was dissolved in MeOH (40 mL). This solution was added to a suspension of 500 mg of 5% Pd/C in 5% AcOH (10 mL). Hydrogen was bubbled into the reaction mixture until TLC analysis revealed no starting material remaining (5 h). The reaction mixture was filtered and the filtrate evaporated to dryness under reduced pressure to give a white solid. Recrystallization of this material from a mixture of isopropyl alcohol and H₂O afforded 360 mg (76%) of 3: mp 222–222.5 °C; $[\alpha]_D^{20}$ –73° (c 0.2, H₂O). Anal. (C₁₃H₂₃N₃O₄) C, H, N.

N-Benzoyloxycarbonyl-L-prolyl-L-leucylaminoacetonitrile (19). The mixed anhydride prepared from 15 (15 g, 41 mmol), NEt₃ (4.15 g, 41 mmol), and isobutyl chloroformate (5.65 g, 41 mmol) was treated with a solution of aminoacetonitrile hydrochloride (3.8 g, 41 mmol) and NEt₃ (4.15 g, 41 mmol) in H₂O (30 mL). The reaction mixture was stirred at room temperature for 2 h, acidified with concentrated HCl, stripped of THF in vacuo,

and extracted with CHCl₃. The CHCl₃ extract was washed with 5% NaHCO₃ and dried (MgSO₄). The CHCl₃ was evaporated to give a colorless oil. An ethereal solution of this material on standing at room temperature provided 12.5 g (76%) of crystalline product: mp 120–122 °C. An analytical sample was obtained by recrystallizing a portion of this material from CHCl₃–Et₂O: mp 121–122 °C; $[\alpha]_D^{24}$ –86.0° (c 0.5, CHCl₃). Anal. (C₂₁H₂₈N₄O₄) C, H, N.

L-Prolyl-L-leucylaminoacetonitrile (4). A solution of 19 (8 g, 20 mmol) and AcOH (1.2 g, 20 mmol) in MeOH (75 mL) was hydrogenated for 2 h over 1.5 g of 5% Pd/C. The mixture was filtered and the filtrate evaporated to dryness. The residue was partitioned between CHCl₃ and 10% HCl. The aqueous layer was made basic with 2 N NaOH and extracted with CHCl₃ (4 × 20 mL). The combined CHCl₃ extracts were dried (Na₂SO₄) and the solvent was evaporated to give the desired product as a yellow oil. Trituration in Et₂O yielded 3.2 g (60%) of crystalline nitrile: mp 95–100 °C. Recrystallization of this material from CHCl₃–Et₂O provided 2.4 g of pure product: mp 101–103 °C; $[\alpha]_D^{20}$ –125.8° (c 1.24, CHCl₃). Anal. (C₁₃H₂₂N₄O₂) C, H, N.

N-Benzoyloxycarbonyl-L-prolyl-L-leucylamino-2-propanone Semicarbazone (20). A solution of amino-2-propanone semicarbazone hydrochloride²⁰ (1.7 g, 11 mmol) and NEt₃ (1.1 g, 11 mmol) in H₂O (25 mL) was added to the mixed anhydride prepared in THF from 15 (3.6 g, 10 mmol), *N*-methylmorpholine (1.0 g, 10 mmol), and isobutyl chloroformate (1.4 g, 10 mmol). After the mixture was stirred at room temperature for 2 h, it was stripped of THF in vacuo and the residue partitioned between CHCl₃ and 10% Na₂CO₃. The organic phase was washed with 10% HCl and H₂O and dried (Na₂SO₄). Removal of the CHCl₃ in vacuo yielded 4.43 g (93%) of 20: mp 102–104 °C; $[\alpha]_D^{21}$ –54.4° (c 0.3, MeOH). Anal. (C₂₃H₃₄N₆O₅) C, H, N.

N-Benzoyloxycarbonyl-L-prolyl-L-leucylamino-2-propanone (21). A solution of 20 (1 g, 2 mmol) in MeOH (10 mL) was treated with 10% HCl (15 mL). The mixture was stirred at room temperature for several hours and then extracted with CHCl₃ (4 × 20 mL). The combined CHCl₃ extracts were dried (MgSO₄) and the CHCl₃ was evaporated to give 790 mg (90%) of the ketone: mp 142.5–143 °C; $[\alpha]_D^{20}$ –89.3° (c 1.0, MeOH). Anal. (C₂₂H₃₁N₃O₅) C, H, N.

L-Prolyl-L-leucylamino-2-propanone Oxalate (5). A 30% solution of HBr in AcOH (3 mL) was added to 21 (700 mg, 1.7 mmol) and the mixture stirred for 30 min. The addition of Et₂O gave a hygroscopic precipitate which was collected, washed with Et₂O, and dissolved in 10 mL of H₂O. The aqueous solution was washed with EtOAc (10 mL), made alkaline with 2 N NaOH, saturated with NaCl, and extracted with CHCl₃ (3 × 8 mL). The combined CHCl₃ extracts were dried (Na₂SO₄) and the CHCl₃ was removed in vacuo to give 450 mg (95%) of 5 as a hygroscopic glassy solid. This material was converted to the oxalate salt by treating a solution of the free base with a solution of oxalic acid (200 mg) in THF. The precipitate was washed with Et₂O and recrystallized from isopropyl alcohol to give 450 mg of the oxalate salt: mp 166–168 °C; $[\alpha]_D^{25}$ –58.2° (c 1.0, H₂O). Anal. (C₁₆H₂₇N₃O₇) C, H, N.

N-Benzoyloxycarbonyl-L-prolyl-L-leucylsemicarbazide (22). The mixed anhydride prepared from 15 (3.6 g, 10 mmol), NEt₃ (1.0 g, 10 mmol), and isobutyl chloroformate (1.4 g, 10 mmol) was treated with a solution of semicarbazide hydrochloride (1.5 g, 13.5 mmol) and NEt₃ (1.4 g, 13.5 mmol) in H₂O (20 mL). The product was isolated in the same manner as described for 19. Recrystallization from MeOH–Et₂O afforded 4.15 g (99%) of semicarbazide: mp 193–195 °C; $[\alpha]_D^{25}$ –80.0° (c 0.36, MeOH). Anal. (C₂₀H₂₉N₅O₅) C, H, N.

L-Prolyl-L-leucylsemicarbazide Acetate (6). The benzyloxycarbonyl-protecting group was removed from 22 (2.4 g, 8 mmol) in a manner analogous to that used for 3 to yield 2.6 g of crude material. Recrystallization from MeOH–Et₂O gave 1.8 g (64%) of the acetate salt: mp 160.5–161.5 °C; $[\alpha]_D^{23}$ –54.3° (c 0.76, MeOH). Anal. (C₁₄N₂₇N₅O₅) C, H, N.

N-Benzoyloxycarbonyl-L-prolyl-L-leucyl-β-alaninamide (23). The mixed anhydride prepared in the THF from 15 (3.6 g, 10 mmol), NEt₃ (1.0 g, 10 mmol), and isobutyl chloroformate (1.4 g, 10 mmol) was mixed with an aqueous solution of β-alaninamide hydrochloride (1.3 g, 10 mmol) and NEt₃ (1.0 g, 10 mmol). The product was isolated in the same manner as described

for 19. After evaporation of the solvent, the residue was recrystallized from isopropyl alcohol to give 2.3 g (53%) of product: mp 177–179 °C; $[\alpha]_D^{25}$ -79.1° (c 1.0, EtOH). Anal. (C₂₂H₃₂N₄O₅) C, H, N.

L-Prolyl-L-leucyl-β-alaninamide Hydrochloride (7). A solution of the protected amide 23 (1.3 g, 3 mmol) in 30 mL of MeOH containing concentrated HCl (0.5 mL) was hydrogenated for 1 h over 70 mg of 5% Pd/C. After the mixture was filtered, the filtrate was stripped of solvent in vacuo and the residue recrystallized from isopropyl alcohol to afford 830 mg (83%) of the hydrochloride salt: mp 198–200 °C; $[\alpha]_D^{25}$ -62.0° (c 1.0, MeOH). Anal. (C₁₄H₂₇N₃O₆Cl) C, H, N.

(±)-Thiazolidine-2-carboxylic Acid (24). The method used was analogous to that reported by Yurev et al.²⁷ for the synthesis of substituted thiazolidine-2-carboxylic acids. A solution of 2-aminoethanethiol hydrochloride (19.2 g, 0.17 mmol) in MeOH was treated with NaOCH₃ (9.1 g, 0.17 mmol). After the solvent was removed in vacuo, the free base was treated with a solution of glyoxylic acid (12.5 g, 0.17 mmol) in absolute EtOH (100 mL). This mixture was shaken vigorously for several minutes and then cooled in an ice bath. The yellow precipitate was collected and recrystallized from 70% EtOH to afford 11.2 g (50%) of the amino acid: mp 190–191 °C dec (lit.²⁸ 181 °C); NMR (CDCl₃) δ 5.22 (s, 1 H, NCHS), 3.58–3.92 (m, 2 H, SCH₂), 3.15–3.40 (m, 2 H, NCH₂). Anal. (C₄H₇NO₂S) C, H, N.

(±)-Ethyl Thiazolidine-2-carboxylate Hydrochloride (25). To an ice-cold suspension of 24 (15.8 g, 0.12 mmol) in 50 mL of absolute EtOH was added in a dropwise manner SOCl₂ (14.5 g, 0.12 mmol). The reaction mixture was heated with constant stirring at 60 °C for 14 h after which time the solvent was removed under reduced pressure to yield an oil which solidified on standing. Recrystallization of this solid from a mixture of EtOH and Et₂O yielded 18.95 g (81%) of 25: mp 103–104.5 °C; NMR (CDCl₃) δ 5.56 (s, 1 H, NCHS). Anal. (C₆H₁₂NO₂SCl) C, H, N.

Resolution of (±)-Ethyl Thiazolidine-2-carboxylate. An aqueous solution of 25 (7.5 g, 38 mmol) was made basic with 10% Na₂CO₃ and extracted with Et₂O (4 × 15 mL). The combined ethereal extracts were dried (Na₂SO₄) and stripped of solvent in vacuo to give 5.91 g (36.7 mmol) of the free base. A solution of this material in anhydrous Et₂O (10 mL) was added to a hot solution of (+)-tartaric acid (5.5 g, 36.7 mmol) in absolute EtOH (30 mL). On cooling a white crystalline product was obtained. Recrystallization of this material to optical purity from EtOH–Et₂O (3:1 v/v) gave 5.39 g (94% of theory) of the (+)-tartrate salt of (–)-ethyl thiazolidine-2-carboxylate: mp 120–122 °C; $[\alpha]_D^{18}$ -65.5° (c 0.2, MeOH). Anal. (C₁₀H₁₇NO₈S) C, H, N, S.

An aqueous solution of the (+)-tartrate salt of (–)-ethyl thiazolidine-2-carboxylate (1.75 g, 5.6 mmol) was made alkaline with 10% Na₂CO₃ and extracted with Et₂O (4 × 15 mL). After the combined ethereal extracts were dried (Na₂SO₄), the Et₂O was removed in vacuo to afford 0.87 g (97%) of (–)-ethyl thiazolidine-2-carboxylate: mp 47–49 °C; $[\alpha]_D^{25}$ -107.0° (c 1.7, EtOAc). Anal. (C₆H₁₁NO₂S) C, H, N.

The mother liquors from the above recrystallization were combined and stripped of solvent in vacuo. The residue was dissolved in 10% Na₂CO₃ and extracted with Et₂O (2 × 20 mL). The combined ethereal extracts were dried (MgSO₄) and the Et₂O was removed in vacuo to give 1.85 g (11.5 mmol) of crude (+)-ethyl thiazolidine-2-carboxylate. An ethereal solution of this material was added to a hot solution of (–)-tartaric acid (1.73 g, 11.5 mmol) in absolute EtOH. The crude (–)-tartrate salt which precipitated on cooling was treated in the same manner as the (+)-tartrate salt to give 2.07 g (36% of theory) of the (–)-tartrate salt of (+)-ethyl thiazolidine-2-carboxylate: mp 121–123 °C; $[\alpha]_D^{18}$ +65.0° (c 0.2, MeOH). Anal. (C₁₀H₁₇NO₈S) C, H, N, S.

(+)-Ethyl thiazolidine-2-carboxylate was isolated from its (–)-tartrate salt (0.75 g, 2.4 mmol) in the same manner as was the (–) isomer to yield 330 mg (85%) of (+) ester: mp 51–53 °C; $[\alpha]_D^{25}$ +109.7° (c 2.0, EtOAc).

L-Prolyl-L-leucyl-(–)-thiazolidine-2-carboxamide (8). To an ice-cold solution of 15 (1.82 g, 5 mmol) and (–)-ethyl thiazolidine-2-carboxylate (0.81 g, 5 mmol) in EtOAc (50 mL) was added *N,N*-dicyclohexylcarbodiimide (1.03 g, 5 mmol). The mixture was stirred at 0 °C for 6 h and then overnight at room temperature. The mixture was filtered and the filtrate washed

with 10% Na₂CO₃, 10% HCl, and saturated NaCl solution. After the organic layer was dried (Na₂SO₄), the solvent was evaporated to give 2.16 g (85%) of ester as an oil. This material was dissolved in MeOH (50 mL) saturated with NH₃. When TLC [CHCl₃–CH₃OH (9:1)] showed no starting material to be present, the solvent was removed in vacuo to give 2.0 g of amide which was used without further purification.

The amide (1.1 g, 2.3 mmol) was treated with 4 mL of a 30% solution of HBr in acetic acid. After 30 min at room temperature Et₂O was added. The salt was collected by filtration, washed with Et₂O, and dissolved in H₂O (10 mL). The aqueous solution was washed with EtOAc (10 mL), made basic with 2 N NaOH, saturated with NaCl, and extracted with CHCl₃ (4 × 10 mL). The combined CHCl₃ extracts were dried (Na₂SO₄) and the solvent was removed in vacuo to give 610 mg (77%) of 8: mp 75–78 °C; $[\alpha]_D^{25}$ -90.9° (c 0.55, CHCl₃); NMR (CDCl₃) δ 5.57 (s, 1 H, NCHS). Anal. (C₁₅H₂₆N₄O₃S) C, H, N; calcd, 16.36; found, 15.74.

L-Prolyl-L-leucyl-(+)-thiazolidine-2-carboxamide (9). Following the same procedures that were used for 8, (+)-ethyl thiazolidine-2-carboxylate (330 mg, 2 mmol) was coupled to 15 (725 mg, 2 mmol) with *N,N*-dicyclohexylcarbodiimide. The same reaction conditions and work-up as above gave 300 mg (44%) of 9 after recrystallization from EtOAc: mp 158–158.5 °C; $[\alpha]_D^{25}$ -23.9° (c 1.5, CHCl₃); NMR (CDCl₃) δ 5.50 (s, 1 H, NCHS). Anal. (C₁₅H₂₆N₄O₃S) C, H, N, S.

L-Thiazolidine-4-carboxamide (26). Methyl L-thiazolidine-4-carboxylate (12.5 g, 85 mmol) was treated with a methanolic solution of NH₃ for 1 day. The solvent was removed in vacuo and the residue recrystallized from EtOH to give 8 g (71%) of amide: mp 97–98.5 °C; $[\alpha]_D^{21}$ -125.4° (c 1.0, MeOH). Anal. (C₄H₈N₂OS) C, H, N, S.

L-Prolyl-L-leucyl-L-thiazolidine-4-carboxamide (10). Compound 26 (390 mg, 3 mmol) in EtOAc (40 mL) was coupled to 15 (1.09 g, 3 mmol) with *N,N*-dicyclohexylcarbodiimide (620 mg, 3 mmol). After the reaction was stirred for 5 h at 0 °C, it was filtered. The filtrate was washed with 10% HCl, 5% NaHCO₃, and saturated NaCl solution. The organic layer was dried (Na₂SO₄) and the solvent removed in vacuo to yield 940 mg (66%) of protected tripeptide: mp 77–80 °C. Without further purification, the benzyloxycarbonyl-protecting group was removed with a 30% solution of HBr in acetic acid. The reaction conditions and work-up were the same as those described for 8. Recrystallization of the crude product from EtOAc gave 348 mg (50%) of 10: mp 168–170 °C; $[\alpha]_D^{29}$ -181.2° (c 0.5, MeOH). Anal. (C₁₅H₂₆N₄O₃S) C, H, N, S.

L-Prolyl-L-leucyl-L-prolinamide (11). The mixed anhydride prepared in THF from 15 (1.9 g, 5.3 mmol), NEt₃ (0.53 g, 5.3 mmol), and isobutyl chloroformate (0.72 g, 5.3 mmol) was treated with a solution of L-prolinamide (0.6 g, 5.3 mmol) in THF (10 mL). The mixture was stirred at room temperature for 2 h, filtered, and stripped of THF under reduced pressure. The residue was dissolved in EtOAc (50 mL) and washed successively with 10% Na₂CO₃, 10% HCl, and H₂O. After the organic layer was dried (Na₂SO₄), the solvent was removed in vacuo to give 2.0 g (87%) of Z-Pro-Leu-Pro-NH₂ (27) as a clear colorless oil. This material was hydrogenated for 1.5 h over 750 mg of 5% Pd/C in MeOH (75 mL) containing concentrated HCl (0.8 mL). The mixture was filtered and the salt which was obtained after evaporation of the solvent was dissolved in CHCl₃. Ammonia was bubbled into the solution and the precipitate of NH₄Cl was removed by filtration. The residue which remained after the CHCl₃ was evaporated was recrystallized from EtOAc to give 530 mg (38%) of pure 11: mp 192–193.5 °C; $[\alpha]_D^{21}$ -120° (c 0.1, MeOH). Anal. (C₁₆H₂₈N₄O₃) C, H, N.

Methyl *N*-tert-Butyloxycarbonyl-L-thiazolidine-4-carboxyl-L-leucylglycinate (28). The mixed anhydride prepared in THF from 1.17 g (5 mmol) of *tert*-butyloxycarbonyl-L-thiazolidine-4-carboxylic acid [prepared by the general procedure of Hofmann et al.²⁹ mp 130–130.5 °C; $[\alpha]_D^{19}$ -114.1° (c 1.0, MeOH)], *N*-methylmorphine (0.5 g, 5 mmol), and isobutyl chloroformate (0.7 g, 5 mmol) was treated with a solution of Leu-Gly-OCH₃-HBr (1.7 g, 6 mmol) and NEt₃ (0.6 g, 6 mmol) in DMF (7 mL). The mixture was stirred at 25 °C for 30 min and then evaporated to dryness. A solution of the residue in CHCl₃ (30 mL) was washed with 5% NaHCO₃, 10% HCl, and saturated NaCl solution. The organic layer was dried (Na₂SO₄) and stripped

of solvent in vacuo to give 1.0 g (50%) of product: mp 133–135 °C. Recrystallization from MeOH–hexane provided an analytically pure sample: mp 135–137 °C; $[\alpha]_D^{27}$ –131.8° (c 1.2, MeOH). Anal. (C₁₈N₃N₃O₈S) C, H, N.

N-tert-Butyloxycarbonyl-L-thiazolidine-4-carbonyl-L-leucylglycinamide (29). Compound 28 (596 mg, 1.4 mmol) was dissolved in MeOH (30 mL) saturated with NH₃ and placed in a refrigerator for 1 week. The solvent was removed in vacuo and the residue recrystallized from EtOAc–Et₂O to give 445 mg (77%) of 29: mp 149–151 °C; $[\alpha]_D^{25}$ –122.6° (c 0.73, MeOH). Anal. (C₁₇H₂₀N₄O₅S) C, H, N.

L-Thiazolidine-4-carbonyl-L-leucylglycinamide (12). Compound 29 (550 mg, 1.4 mmol) was deprotected with 2 mL of 30% HBr in acetic acid. The reaction conditions and work-up were similar to those described for 8. Recrystallization of the crude material from MeOH–EtOAc gave 140 mg (33%) of pure 12: mp 154–155 °C; $[\alpha]_D^{23}$ –51.7° (c 0.6, MeOH). Anal. (C₁₂H₂₇N₄O₃S) C, H, N.

N-Cyclopentanecarbonyl-L-leucylglycinamide (13). *N*,*N*-Dicyclohexylcarbodiimide (4.1 g, 20 mmol) was added to a solution of cyclopentanecarboxylic acid (2.3 g, 20 mmol) and *N*-hydroxysuccinimide (2.3 g, 20 mmol) in dioxane (50 mL). The mixture was stirred at 25 °C for 6 h. The solid was removed by filtration and the filtrate evaporated to dryness under reduced pressure. A solution of the residue in EtOAc was washed with 5% NaHCO₃ and dried (Na₂SO₄). Evaporation of the solvent gave a semisolid which was recrystallized from isopropyl alcohol to give 2.25 g (53%) of the activated ester: mp 68–70 °C.

A solution of 640 mg (3 mmol) of the *N*-hydroxysuccinimide ester of cyclopentanecarboxylic acid, NEt₃ (300 mg, 3 mmol), and L-leucylglycinamide acetate (750 mg, 3 mmol) in DMF (10 mL) was stirred overnight at 25 °C. The DMF was removed in vacuo and the residue partitioned between CHCl₃ and 10% HCl. The CHCl₃ layer was washed with 10% Na₂CO₃ and saturated NaCl solution. The organic layer was dried (Na₂SO₄) and stripped of solvent in vacuo to give after recrystallization from MeOH–EtOAc 370 mg (44%) of 13: mp 170–172 °C; $[\alpha]_D^{24}$ –20.6° (c 1.0, MeOH). Anal. (C₁₄H₂₅N₃O₃) C, H, N.

Dopa Potentiation Test. Male ICR mice (A. R. Schmidt, Sprague–Dawley), weighing 16–20 g, were pretreated orally 4 h before the test with a low dose of the monoamine oxidase inhibitor, pargyline hydrochloride (40 mg/kg). The test compound was then administered intraperitoneally. Four mice were tested at each dose. After 1 h a challenge dose of *dl*-Dopa (200 mg/kg) was given intraperitoneally and the mice were observed for 1 h. The changes in activity and aggressive behavior were recorded and rated according to the following system: (1) slight—piloerection, slight salivation, slight increased motor activity; (2) moderate—piloerection, salivation, marked increased activity, and irritability; and (3) marked—piloerection, profuse salivation, marked increased irritability and reactivity, jumping, squeaking, and aggressive fighting. Mice which were given only pargyline hydrochloride (40 mg/kg) and *dl*-Dopa (200 mg/kg) served as the controls. These mice had a behavioral rating of 1.

Oxotremorine Antagonism Test. Normal (ICR) male mice (16–20 g) were pretreated intraperitoneally with varying doses of the test drug 1 h before the administration of oxotremorine (0.5 mg/kg ip). Four mice were tested at each dose of the test drug. The responses of the mice were recorded by observation techniques and were compared with those observed in mice which received only oxotremorine as a control. The responses were graded as 0 (no effect), 1 (slight), 2 (moderate), and 3 (marked) for the effect of oxotremorine on each group of mice. The graded responses for the central effects of oxotremorine (including decreased motor activity, tremors, head twitch, limb abduction, and ataxia) were added together with the graded responses of the

peripheral effects (lachrymation, salivation, and diarrhea) to produce a total score. Oxotremorine itself results in an average total score of 22 out of a possible maximum total score of 24. A compound is considered active if the total score is 16/22 or less.

References and Notes

- (1) Deceased July 14, 1974.
- (2) (a) A. J. Kastin, N. P. Plotnikoff, A. V. Schally, and C. A. Sandman, *Rev. Neurosci.*, **2**, 111 (1976); (b) A. J. Prange, Jr., I. C. Wilson, G. R. Breese, and M. A. Lipton, *Prog. Brain Res.*, **42**, 1 (1975).
- (3) R. M. G. Nair, A. J. Kastin, and A. V. Schally, *Biochem. Biophys. Res. Commun.*, **43**, 1376 (1971).
- (4) A. J. Kastin, A. V. Schally, and S. Viosca, *Proc. Soc. Exp. Biol. Med.*, **137**, 1437 (1971).
- (5) M. E. Celis, S. Taleisnik, and R. Walter, *Proc. Natl. Acad. Sci. U.S.A.*, **68**, 1428 (1971).
- (6) M. E. Celis, R. Macagno, and S. Taleisnik, *Endocrinology*, **93**, 1229 (1973).
- (7) N. P. Plotnikoff, A. J. Kastin, M. S. Anderson, and A. V. Schally, *Life Sci.*, **10**, 1279 (1971).
- (8) G. M. Everett, *Antidepressant Drugs*, *Proc. Int. Symp.*, **1st**, 1966, **122**, 164 (1966).
- (9) N. P. Plotnikoff, A. J. Kastin, M. S. Anderson, and A. V. Schally, *Proc. Soc. Exp. Biol. Med.*, **140**, 811 (1972).
- (10) S. Castensson, H. Sievertsson, B. Lindeke, and C. Y. Sum, *FEBS Lett.*, **44**, 101 (1974).
- (11) N. P. Plotnikoff, A. J. Kastin, M. S. Anderson, and A. V. Schally, *Neuroendocrinology*, **11**, 67 (1973).
- (12) N. P. Plotnikoff and A. J. Kastin, *Arch. Int. Pharmacodyn. Ther.*, **211**, 211 (1974).
- (13) R. Walter, P. L. Hoffman, J. B. Flexner, and L. B. Flexner, *Proc. Natl. Acad. Sci. U.S.A.*, **72**, 4180 (1975).
- (14) J. M. van Ree and D. de Wied, *Life Sci.*, **19**, 1331 (1976).
- (15) A. J. Kastin and A. Barbeau, *Can. Med. Assoc. J.*, **107**, 1079 (1972).
- (16) P. A. Fischer, E. Schnieder, P. Jacobi, and H. Maxion, *Eur. Neurol.*, **12**, 360 (1974).
- (17) W. D. Cash, *J. Org. Chem.*, **26**, 2136 (1961).
- (18) G. W. Anderson, J. E. Zimmerman, and F. M. Callahan, *J. Am. Chem. Soc.*, **88**, 1338 (1966).
- (19) G. W. Anderson, J. E. Zimmerman, and F. M. Callahan, *J. Am. Chem. Soc.*, **89**, 5012 (1967).
- (20) J. D. Hepworth, "Organic Syntheses", Collect. Vol. V, Wiley, New York, N.Y., 1973, p 27.
- (21) N. P. Plotnikoff and A. J. Kastin, *Pharmacol. Biochem. Behav.*, **2**, 417 (1974).
- (22) S. Bjorkman, S. Castensson, B. Lindeke, and H. Sievertsson, *Acta Pharm. Suec.*, **13**, 289 (1976).
- (23) A. Carlsson, J. Jonason, and M. Lindquist, *J. Pharm. Pharmacol.*, **21**, 769 (1969).
- (24) N. P. Plotnikoff and A. J. Kastin in "Peptides: Chemistry, Structure and Biology", R. Walter and J. Meienhofer, Ed., Ann Arbor Science Publishers, Ann Arbor, Mich., 1975.
- (25) H. C. Beyerman, H. Hindriks, and E. W. B. De Leer, *Chem. Commun.*, 1668 (1968).
- (26) C. Ressler and V. du Vigneaud, *J. Am. Chem. Soc.*, **76**, 3107 (1954).
- (27) Ya. K. Yur'ev, E. V. Dyatlovitzkaya, and A. E. Sokol, *J. Gen. Chem. USSR*, **29**, 3844 (1958).
- (28) J. P. Fourneau, O. Efimovski, J. C. Gangnalt, R. Jacquier, and C. Ridant, *C. R. Hebd. Seances Acad. Sci., Ser. C*, **272**, 1515 (1971).
- (29) K. Hofmann, W. Haas, M. J. Smithers, R. D. Wells, Y. Wolman, N. Yanaihara, and G. Zanetti, *J. Am. Chem. Soc.*, **87**, 620 (1965).