


Scheme I



with triethyl orthoformate, following a recently reported procedure for the esterification of phosphonic acids.⁷

Hydrogenolysis of 5 produced the free amino ester 6. Although removal of the carbobenzoxy group could also be achieved with HBr-HOAc, this method was found to give a low yield of impure product. Coupling of the amino ester 6 with carbobenzoxy-D-alanine (7) was accomplished in good yield with *N,N'*-dicyclohexylcarbodiimide (DCC) affording the carbobenzoxy dipeptide diester 8.

Removal of the carbobenzoxy group from 8 with HBr-HOAc was accompanied by cleavage of the phosphonate ester groups. The resultant hydrobromide 9 was converted to the free base with moist Ag_2O and residual silver ion removed by an ion exchange resin giving the free dipeptide 1 as a mixture of diastereomers. Tlc of 1 on cellulose plates with a variety of solvent systems indicated the presence of the two isomers but a clean separation was not achieved. Further work aimed at the resolution of these isomers is planned.

Compound 1 was tested for antimicrobial activity against a variety of organisms on both complete and minimal media using D-cycloserine and 1-aminoethylphosphonic acid (3) as standards for comparison. The only activity observed was against *Pseudomonas aeruginosa* on minimal media and this activity was of the same order of magnitude as that exhibited by 3. These results could mean that 1 is being incorporated into cell wall synthesis and that 3 instead of D-alanine is released during the cross-linking of the linear peptide strands. Another possibility is that 1 is a weak antimetabolite of D-alanyl-D-alanine. In order to gain additional information on this we are preparing the dipeptide similar to 1 but having 1-aminoethylphosphonic acid at the N-terminal part of the molecule.

Experimental Section

Melting points were taken on a Mel-Temp melting point apparatus and are corrected or uncorrected as indicated. Infrared spectra were taken on a Perkin-Elmer Model 257 spectrophotometer, nmr spectra on a Jeolco C-60HL spectrometer (Me_4Si , DSS), mass spectra on a Du Pont Model 21-492 spectrometer, and optical rotations on a Perkin-Elmer Model 141 polarimeter. Elemental analyses were performed by either Galbraith Laboratories, Inc., Knoxville, Tenn., or Chemalytics, Inc., Tempe, Ariz.

1-Aminoethylphosphonic Acid (3). 1-Aminoethylphosphonic acid (3) was prepared according to the procedure of Chambers and Isbell⁶ using triethyl 2-phosphonopropionate⁸ (119 g, 0.5 mol) to give 45.7 g (73%) of 3: mp (uncor) 274–278° dec (lit. mp 283–284°,⁶ mp 278°).

N-Carbobenzoxy-1-aminoethylphosphonic Acid (4). A well-stirred solution of 5 g (0.04 mol) of 3 in 30 ml of H_2O was adjusted to pH 9.5 with 4 N NaOH. To this was added 15 ml of Et_2O and 10.8 g (0.06 mol at 95%) of benzyl chloroformate and the mixture was cooled in an ice bath. Stirring was continued 11.5 hr and the pH maintained at 9–9.5 by intermittent addition of 4 N NaOH. The final mixture was extracted with Et_2O (discarded). The aqueous layer was poured slowly onto a mixture of 30 ml of H_2O , 30 ml of concentrated HCl, and 100 g of ice. This was extracted with Et_2O . The combined, dried (MgSO_4) Et_2O extracts were concentrated *in vacuo* and the residue was recrystallized from EtOAc -*n*-hexane giving 6.1 g (58.6%) of crystals of 4: mp (cor) 111–113° dec; ir (Nujol) 1260 ($\text{P}=\text{O}$), 1690 ($\text{C}=\text{O}$), and 3295 cm^{-1} (NH); nmr (D_2O -NaOD) δ 7.34 (s, 5, C_6H_5), 5.1 (s, 2, $\text{C}_6\text{H}_5\text{CH}_2\text{O}$), 4.03–3.36 (m, 1, CH_3CHP), and 1.4 ppm (dd, 3, $\text{J}_{\text{PH}} = \text{J}_{\text{HH}} = 13$ Hz, CH_3CHP). Anal. ($\text{C}_{10}\text{H}_{14}\text{NO}_5\text{P}$) C, H, N, P.

Diethyl N-Carbobenzoxy-1-aminoethylphosphonate (5). A stirred suspension of 5.2 g (0.02 mol) of 4 and 25 g (0.17 mol) of $(\text{EtO})_3\text{CH}$ was slowly heated. The ethyl formate and EtOH by-products were continuously removed by distillation thereby allowing the temperature to rise to 135° where it was held for 1 hr. The cooled mixture was filtered. The excess $(\text{EtO})_3\text{CH}$ was removed from the filtrate under a high vacuum leaving 6 g (95.3%) of 5 as a viscous gum: ir (neat) 1170 (POEt), 1250 ($\text{P}=\text{O}$), 1725 ($\text{C}=\text{O}$), and 3250 cm^{-1} (NH); nmr (CCl_4) δ 7.1 (s, 5, C_6H_5), 4.94 (s, 2, $\text{C}_6\text{H}_5\text{CH}_2\text{O}$), 2.47–3.6 (m, 5, POCH₂ and CH_3CHP), and 1.48–0.99 ppm (m, 9, POCH₂CH₃ and CH_3CHP). The ester was used without further purification.

Diethyl 1-Aminoethylphosphonate (6). A mixture of 6 g (0.019 mol) of crude 5 and 1 g of 10% Pd/C in 75 ml of absolute MeOH was hydrogenated at 3.37 kg/cm² for 24 hr. The catalyst was filtered and the filtrate was concentrated *in vacuo* (room temperature) to give 2.4 g (70.6%) of the amino ester 6:¹⁰ ir (neat) 1120 (POEt), 1240 ($\text{P}=\text{O}$), and 3300, 3400 cm^{-1} (NH₂); nmr (CCl_4) δ 4.05 (m, 4, $\text{J} = 7$ Hz, POCH₂), 3.24–2.67 (m, unresolved, 1, CH_3CHP), 1.84 (s, 2, NH₂), and 1.43–1.03 ppm (m, 9, CH_3CHP and POCH₂CH₃); mass spectrum (70 eV) *m/e* (rel intensity) 181 (30), 138 (57), 44 (100).

Diethyl N-Carbobenzoxy-D-alanyl-1-aminoethylphosphonate (8). To a stirred solution of 4.89 g (0.027) of 6, 5.3 g (0.027 mol) of DCC, and 50 ml of CH_2Cl_2 was added 6.02 g (0.027 mol) of Cbz-D-Ala (7),¹¹ prepared according to the procedure of Bergmann and Zervas,¹² in 30 ml of CH_2Cl_2 . Stirring was continued for 24 hr, the suspension filtered, and the filtrate concentrated *in vacuo*. The residue was redissolved in 30 ml of CCl_4 and filtered, and the filtrate was concentrated to give 10.3 g (99%) of the carbobenzoxy peptide diethyl ester 8 as a glassy gum: ir (neat) 1170 (POEt), 1240 ($\text{P}=\text{O}$), 1675 (amide $\text{C}=\text{O}$), 1730 (urethane $\text{C}=\text{O}$), and 3300 cm^{-1} (NH); nmr (CDCl_3) δ 7.84–7.5 (broad, 1, CONH), 7.19 (s, 5, C_6H_5), 6.24–5.9 (broad, 1, NHCO_2), 5 (s, 2, $\text{C}_6\text{H}_5\text{CH}_2\text{O}$), 4.67–3.73 (m, 6, POCH₂CH₃, CHCON, and CHP), and 1.52–1.11 ppm (m, 12, POCH₂CH₃, CH_3CHCO , and CH_3CHP); mass spectrum (70 eV) *m/e* (rel intensity) 386 (100), 341 (9), 251 (27), 249 (89), 208 (95), 206 (85), 180 (77), 178 (50), 135 (36).

N-D-Alanyl-1-aminoethylphosphonic Acid (1). To a stirred solution of 8.3 g (0.022 mol) of 8 in 50 ml of glacial HOAc was added 50 ml of 45% HBr-HOAc and stirring continued 2 hr. To this solution was added 750 ml of absolute Et_2O . After standing 4 hr, the solvents were decanted and last traces removed *in vacuo* leaving the HBr salt 9 as a gummy mass. This was taken up in 200 ml of H_2O and treated with 5 g of Ag_2O for 0.5 hr in a flask protected from light. The suspension was filtered through a pad of Celite, the filtrate passed through an Amberlite IRC-50 (2 × 45 cm), and the eluate concentrated *in vacuo* to give 4.2 g (99%) of a white solid. Trituration with several portions of hot MeOH afforded the dipeptide 1: mp (uncor) 220–230° dec; ir (Nujol) 1260 ($\text{P}=\text{O}$) and 1670 cm^{-1} ($\text{C}=\text{O}$); nmr (D_2O) δ 4.34–3.66 (m, 2, CH_3CHCO and NCHP) and 1.6–1.13 ppm (m, 6, CH_3CHCO and CH_3CHP); $[\alpha]_D^{25} -14.5^\circ$ (c 2, H_2O). Anal. ($\text{C}_5\text{H}_{13}\text{N}_2\text{O}_4\text{P}$) C, H, N; P: calcd, 15.79; found, 15.31.

Primary Antimicrobial Screening of 1. Antimicrobial screening of 1 was carried out by an agar well diffusion method utilizing both complete and minimal media. D-Cycloserine and 1-aminoethylphosphonic acid (3) were used as standards for comparison. Cultures employed for screening were obtained from the American Type Culture Collection (ATCC), Rockville, Md., and included the following organisms: *Staphylococcus aureus* (ATCC 6538), *Bacillus subtilis* (ATCC 6633), *Escherichia coli* (ATCC 11775), *Proteus vulgaris* (ATCC 9489), *Pseudomonas aeruginosa* (ATCC 15442), *Mycobacterium smegmatis* (ATCC 607), *Candida albicans* (ATCC 10231), *Saccharomyces cerevisiae* (ATCC 9763),

Aspergillus niger (ATCC 16888), and *Trichophyton mentagrophytes* (ATCC 9129).

The primary screening plates were prepared using Eugon agar (BBL) for bacteria and Mycophil agar (BBL) for fungi as complete media and Davis minimal agar (Difco) for bacteria and Czapek agar (Difco) for fungi as the minimal media.

The agar culture media were autoclaved, cooled to 45°, inoculated with bacterial or fungal suspensions, mixed thoroughly, and poured into Petri plates (100 × 15 mm) so that each plate contained 15 ml of agar culture media seeded with the test microorganism. Agar wells 10 mm in diameter were prepared in the plates and the samples (0.1 ml) were pipetted into the well. The solutions used were: D-cycloserine, 0.2 mg/ml; 3, 20 mg/ml; 1, 20 mg/ml.

The plates were incubated at 37° for bacteria or 25° for fungi and read for activity at 24-hr intervals. The results were expressed as the distance from the edge of the well to the outer edge of the clear zone of inhibition.

Acknowledgment. This work was supported by the Research Institute of Pharmaceutical Sciences, School of Pharmacy, University of Mississippi. We wish to acknowledge Dr. Stephen Billets for aid in mass spectral interpretations and Mrs. Sheida White for assistance in biological evaluations.

References

- (1) T. J. Franklin and G. A. Snow, "Biochemistry of Antimicrobial Action," Academic Press, New York, N.Y., 1971, pp 35-45.
- (2) J. L. Strominger, E. Ito, and R. H. Threnn, *J. Amer. Chem. Soc.*, **82**, 998 (1960).
- (3) K. Yamauchi, M. Kinoshita, and M. Imoto, *Bull. Chem. Soc. Jap.*, **45**, 2528 (1972).
- (4) K. Yamauchi, M. Kinoshita, and M. Imoto, *Bull. Chem. Soc. Jap.*, **45**, 2531 (1972).
- (5) W. F. Gilmore and H. A. McBride, *J. Pharm. Sci.*, **63**, 1087 (1974).
- (6) J. R. Chambers and A. F. Isbell, *J. Org. Chem.*, **29**, 832 (1964).
- (7) D. A. Nicholson, W. A. Cilley, and O. T. Quimby, *J. Org. Chem.*, **35**, 3149 (1970).
- (8) G. M. Kosolapoff and J. S. Powell, *J. Amer. Chem. Soc.*, **72**, 4198 (1950).
- (9) K. D. Berlin, N. K. Roy, R. T. Claunch, and D. Bude, *J. Amer. Chem. Soc.*, **90**, 4494 (1968).
- (10) M. E. Chalmers and G. M. Kosolapoff, *J. Amer. Chem. Soc.*, **75**, 5278 (1953).
- (11) M. Hunt and V. duVigneaud, *J. Biol. Chem.*, **124**, 669 (1935).
- (12) M. Bergmann and L. Zervas, *Chem. Ber.*, **65**, 1192 (1932).

Centrally Acting Emetics. 8. Conformational Aspects of Certain Dihydrophenanthrene Congeners of Apomorphine^{1,†}


Joseph G. Cannon,* Robert V. Smith, Mohd. A. Aleem,

Division of Medicinal Chemistry and Natural Products, College of Pharmacy

and John Paul Long

Department of Pharmacology, College of Medicine, The University of Iowa, Iowa City, Iowa 52242. Received August 1, 1974

A prior communication¹ described the synthesis and biological effects of a ring-cleavage derivative (1) of apomorphine (3). The low order of emetic activity of 1, which is




also closely related structurally to the potent emetic 5,6-dihydroxy-2-dimethylaminotetralin (2), was rationalized on conformational grounds. Rekker, *et al.*,² presented arguments that an anti disposition (4) of the catechol ring and the amino group of dopamine is the biologically active one. The dopamine portions of 2 and apomorphine exhibit this anti disposition.¹ It was proposed¹ that serious steric interaction between the dimethylamino group in a pseudo-equatorial conformation and the ethyl group at position 8 in compound 1 induces a conformational "flip"

in the dihydrophenanthrene ring, to place the amino group in the less sterically crowded pseudo-axial arrangement. This conformation presents the dopamine portion of 1 in a gauche arrangement (5), and the molecule cannot interact optimally with the emetic receptor.

Inspection of molecular models led to the speculation that a phenanthrene system (6) lacking the ethyl group at position 8 would accommodate the dimethylamino group in the presumably more stable pseudo-equatorial disposition, in which the catechol ring and the amino group are in the biologically favorable anti conformation. Accordingly, it was predicted that 6 would exhibit potent emetic and peripheral apomorphine-like effects. Preparation of 6 was accomplished by a multistep sequence beginning with 3,4-dimethoxyphenanthrene-9-carboxylic acid. Spectral (ir, nmr) data were consistent with proposed structures of all compounds prepared.

Pharmacology. Compound 6 was evaluated in five dogs anesthetized with barbitol sodium (200 mg/kg). The arterial pressure was measured using the right femoral artery and the compound was administered into the left femoral vein. In doses up to 2 mg/kg, 6 did not alter the blood



[†]This investigation was supported by Grant NS04349, National Institute of Neurological Diseases and Stroke.