SPECIAL FEATURE: ARTICLE





Chiral aminophosphines derived from hydroxyproline and their application in allene–imine [4 + 2] annulation

San N. Khong¹ · Changmin Xie¹ · Xinyi Wang^{1,2} · Hao Tan^{1,2} · Ohyun Kwon¹

Received: 20 February 2019 / Revised: 20 March 2019 / Accepted: 22 March 2019 $\ensuremath{\mathbb{S}}$ The Author(s), under exclusive licence to the Japan Antibiotics Research Association 2019

Abstract

A robust synthetic route from L-hydroxyproline (L-Hyp) to phosphines has established an expandable library of six chiral aminophosphines, which were then applied to the phosphine-catalyzed [4 + 2] allene–imine annulation. The enantioinduction in the annulations—induced by a purely steric effect—were moderate (up to 57% ee). A switch of the reaction site from the γ - to the β '-carbon atom of the allenoate was observed during the annulations performed using sterically demanding chiral phosphines.

Introduction

Piperidines are among the nitrogen heterocycles most frequently used in pharmaceuticals [1]. Apart from their appearance in pharmaceutically interesting substances [2], tetrahydropyridines, and piperidines are present as common structural motifs in many bioactive natural products [3–5]. Because of the interesting biological effects exerted by tetrahydropyridine- and piperidine-containing compounds, many methods have been developed to access these valuable structures [2, 6]. In 2003, we disclosed that functionalized tetrahydropyridines could be synthesized through a PBu₃-catalyzed [4 + 2] reaction of allenoates and imines [7, 8]. Subsequently, Fu [9], Zhao [10], Guo [11], and Sasai [12] realized asymmetric versions of the [4 + 2] reaction using various chiral phosphines. In the last 50 years, major

These authors contributed equally: San N. Khong, Changmin Xie

Dedicated to Professor Samuel J. Danishefsky and his great scientific contributions to total syntheses of highly complex and biologically important natural products.

Supplementary information The online version of this article (https://doi.org/10.1038/s41429-019-0181-0) contains supplementary material, which is available to authorized users.

Ohyun Kwon ohyun@chem.ucla.edu efforts have been exerted in the development of chiral phosphine ligands [13–15]. The chirality of a phosphine may reside on the phosphorus center, in the carbon backbone, or in the phosphine's molecular framework [16]. As part of our efforts in phosphine organocatalysis, we have developed a class of structurally unique *P*-chiral [2.2.1] bicyclic phosphines [17, 18]. Recently, we applied one such bridged bicyclic *P*-chiral phosphine, exo-(*p*-anisyl)-Hyp-Phos, in the asymmetric allene–imine [4 + 2] annulation to afford valuable chiral piperidines in good yields and with excellent enantioselectivities [19]. Herein, we report a class of new chiral phosphines derived from L-hydroxyproline (L-Hyp) and their application in the allene–imine [4 + 2] annulation.

The abundance and ready availability of chiral α -amino acids have made them desirable sources of chirality. Indeed, many bifunctional chiral aminophosphines have been derived from α -amino acids [14]. In previous studies, the presence of a free NH group, responsible for hydrogen bonding, has been the major focus when designing chiral aminophosphines for asymmetric phosphine organocatalysis (Scheme 1a) [14]. Progress in the field of enamine and iminium catalysis has revealed that steric interaction is another useful mode of asymmetric induction [20-23]. Nevertheless, steric interaction has not previously been considered a promising platform for the design of α -amino acid-derived aminophosphines. L-Hyp, an abundant nonproteinogenic amino acid having a proline skeleton and an additional OH group for further steric tuning, has been chosen as the platform for our catalyst design (Scheme 1a). In addition to introducing sterically tunable protecting

¹ Department of Chemistry and Biochemistry, University of California, Los Angeles, CA, USA

² College of Chemistry, Nankai University, Tianjin, China



Scheme 1 Synthetic design of chiral aminophosphines 2 from L-Hyp

groups at the pyrroline nitrogen atom and the 4-OH group, a hydrogen-bonding site could also be installed—for example, as a urea or thiourea motif—at the pyrroline nitrogen atom. We envisioned that the aminophosphines **2a–f** could be obtained from L-Hyp by installing a diphenylphosphino group through the displacement of the mesylate in **1** and subsequent functional group manipulations (Scheme 1b). The synthesis of the mesylate **1** also appeared to be straightforward. Starting from commercially available L-Hyp, various alcohol and amine protecting groups could be installed prior to substitution of the mesylate by a diphenylphosphino group.

Results and discussion

The synthesis of compound **2a** is illustrated in Scheme 2. First, the Boc-protection of L-Hyp proceeded to provide *N*-Boc-L-Hyp **3**. Benzylation of the free OH group followed by borane-mediated reduction of the carboxyl group resulted in the primary alcohol **5**. Only one column chromatographic purification through SiO₂ was necessary to prepare 16.3 g of **5** in 70% yield in the three-step process. Mesylation of **5** led to the key intermediate **1** for phosphine synthesis. The mesylate **1** was converted into the phosphine **2a** in high yield after slow addition of an excess of potassium diphenylphosphide at low temperature.

Having prepared the aminophosphine 2a, the acid-liable N-Boc protecting group was readily replaced with several

acyl groups to afford the chiral aminophosphines **2b–f** of varying steric bias (Scheme 3). After Boc-deprotection, an excess of triethylamine was added to neutralize TFA and promote the acylation with the acyl chloride. Using this one-pot procedure, the four aminophosphines **2b–e** were prepared in high yields. When phenyl isocyanate was used instead of an acyl chloride, the phosphine **2f**, presenting a hydrogen-bonding site, was obtained in 77% yield.

The six chiral phosphines with varied N-substituents were then applied to our [4+2] annulation [7]. We first examined the [4+2] annulation of the activated allenoate 8 and the imine 7 mediated by the phosphine 2a in CH₂Cl₂ (Table 1, entry 1). An inseparable mixture of the tetrahydropyridine 9 and the tetrahydropyridine 10 was obtained in 68% yield, with the ee of 9 being 30%. When the solvent was switched to benzene, the ee of 9 increased to 42%, but 10 was now the major product (entry 2). We then tested the other phosphines for the annulation in CH₂Cl₂ (entries 3–6). When using the catalyst 2b for the annulation in CH₂Cl₂, the ee of 9 increased to 42% compared with that of 2a in the same solvent (cf. entries 3 and 1). Under otherwise identical conditions, the ratio of 10 had also increased. The phosphine 2d, featuring a bulkier acyl substituent on the nitrogen atom, produced the tetrahydropyridine 9 with a higher ee of 53% (entry 5). Interestingly, in the reaction catalyzed by the phosphine 2e presenting an adamantylcarbonyl group, the selectivity of the product was completely reversed-now favoring the tetrahydropyridine 10, which was isolated in 80% yield and contaminated with only a



Scheme 3 Preparation of the phosphines 2b-f

trace of 9 (entry 6). The NH group in the phosphine 2f seemed to have little effect on the enantioselectivity, but the formation of 10 was largely inhibited (entry 7). From these preliminary results, it appeared that the enantioselectivity increased upon increasing the bulk of the N-acyl substituent, presumably through a mechanism involving steric control.

It was also interesting that more sterically demanding phosphines could facilitate the formation of **10** over **9**. The reversed regioselectivity can be explained by two possible pathways of the imine–allene [4 + 2] reaction (Scheme 4) [7, 19, 24]. At the onset of the [4 + 2] reaction, addition of the phosphine **2** to the allene **8** generates intermediate **A**, which is in resonance with structure **B**, which, in turn, is in

equilibrium with intermediate **B**'. In path I, γ -addition of intermediate **B** to the imine **7** affords intermediate **C**, followed by the formation of **9**. In path II, β' -addition of intermediate **B**' to the imine **7** generates intermediate **C**', followed by the formation of **10**. When a bulkier phosphine was used in the reaction, the γ -position of intermediate **B** was more encumbered, relative to its β' -position. In this case, path II, which afforded the product **10**, became the favored pathway.

The same catalysts were then employed in the allene–imine [4+2] annulation of the less activated α -methylallenoate **11** (Table 2). Because the reaction was slower with this allene **11**, the catalyst loading was increased to 30 mol% and 4-Å MS were added to minimize

CO₂Et catalyst (20 mol%) ß' CO₂Et Ts CO₂Et rt, 2 days O₂Ft O₂Et CO₂Et 9 10 ORTEP 10 Entry Catalyst Solvent Conv Dr^c Ratio^c Yield ee (%)^d (9 (%)^b (cis-9/ (9/10)(%)^e trans-+10)(9) 9) CH_2Cl_2 5.9:1 1 2a >99 3.0:1 68 30 2 97 9.1:1 1:3.1 35 42 2a benzene >99 42 3 2b CH_2Cl_2 5.5:1 1:2.3 44 >99 6.2:1 45 4 20 CH_2Cl_2 1:4.4 66 5 2d CH₂Cl₂ >99 7.7:1 1:12.5 70 53 6 2e CH₂Cl₂ >99 1:63.0 80 57^f _ 7 2f CH₂Cl₂ 49 6.7:1 15:1 39 27

Table 1 Phosphines 2-catalyzed [4+2] annulation of the allenoate 8^a

^aReaction conditions: 7 (1 equiv, 0.1 mmol), 8 (1.2 equiv), the phosphine 2 (20 mol%), and the solvent (2 mL)

^bConversion determined using mesitylene as an internal standard

^cDetermined from the ¹H NMR spectrum of the crude mixture

^dIsolated yield

 $^{\circ}$ The ee value of **9** was determined through chiral HPLC analysis; its absolute configuration was determined through comparison with the data of the known compound in ref. [9]

^fThe ee of compound **10**; the absolute configuration was not determined

Scheme 4 Possible reaction pathways for the allene–imine [4+2] annulation



hydrolysis of the imine 7. Although the conversion of the imine 7 and the yield of 12 were not good, we observed a similar trend in the induction of enantioselectivity as that in Table 1. Increasing the bulk of the N-acyl substituent in the phosphine 2 afforded greater enantioselectivity for the formation of the tetrahydropyridine 12. When using the catalyst 2d, an ee of 41% was induced (entry 4). The bulky phosphines again facilitated the formation of the β' -addition product, in this case 13 (cf. entries 4 vs. 1 and 2). These

preliminary findings, obtained from nonoptimized reaction conditions, were consistent with the model of steric control.

Conclusion

Taking advantage of the ready availability of naturally occurring chiral α -amino acids, we have chosen L-Hyp, a nonproteinogenic—but abundantly available—amino acid,

	Ph → N. Ts + Υ 7	$\overset{\beta'}{\longleftarrow} CO_2Et \frac{\text{catalyst}}{4 \text{ Å MS}}_{3}$	(30 mol%) , CH ₂ Cl ₂ , rt Ph Y C days	O_2Et Ts N β' CO_2Et CO_2	₂Et
Entry	Catalyst	Conv (%) ^b	Ratio ^{c,d} (12:13)	Yield (%) ^e (12)	ee (%) ^f (12)
1	2a	56	10:1	23	20
2	2b	74	>50:1	9	29
3	2c	77	7.1:1	10	28
4	2d	70	1.9:1	11	41
5	2e	80	14.1:1	26	23
6	2f	47	>50:1	6	14

Table 2 Phosphines 2–catalyzed [4+2] annulation of the α -methylallenoate 11^a

^aReaction conditions: 7 (1 equiv, 0.1 mmol), 11 (2 equiv), the phosphine 2 (30 mol%), 4-Å MS (10 mg), and CH₂Cl₂ (2 mL)

^bConversion was determined using mesitylene as an internal standard

^cDetermined from the ¹H NMR spectrum of the crude mixture

^d12 and 13 were obtained as a mixture; no pure 13 was isolated, but its structure was inferred according to our mechanistic proposal in Scheme 4 ^eNMR yield (internal standard: mesitylene)

^fDetermined through chiral HPLC analysis; the absolution configuration was determined through comparison with the data of the known compound in ref. [19].

for the design of several chiral phosphines. With the goal of exploiting steric interactions for enantioinduction, we have developed a synthetic route toward a number of L-Hyp-derived chiral phosphines. The preliminary application of these phosphines in the allene–imine [4 + 2] annulation has proven moderately successful in terms of enantioselectivities. When using bulky phosphines, an interesting switch of the reaction site occurred: from the γ -to the β' -carbon atom of the allenoate. The clear trend in the enantioinduction hints at the potential of developing chiral phosphines that operate with steric tuning as the sole source of enantioselectivity control in the [4 + 2] annulation.

Experimental

4-Benzyloxy-2-methanesulfonyloxymethylpyrrolidine-1-carboxylic acid tert-butyl ester (1)

Boc₂O (18.3 g, 83.9 mmol) was added slowly to a solution of L-Hyp (10.0 g, 76.3 mmol) in 1,4-dioxane (80 mL) and 1 N aqueous NaOH (80 mL) at 0 °C. The reaction mixture was stirred for 8 h at room temperature, then concentrated under reduced pressure, acidified to pH 1 using 2 N aqueous HCl, and extracted with EtOAc. The extract was washed with brine, dried (Na₂SO₄), and concentrated to provide the Boc-protected L-Hyp **3** [25] as a clear viscous oil. This

crude product was used in the next step without further purification.

The Boc-protected L-Hyp 3 was dissolved in THF (100 mL) and then cannulated into a slurry of NaH (60% in mineral oil, 11.2 g, 155 mmol) in THF (200 mL) at 0 °C. Benzyl bromide (15.0 g, 87.7 mmol) was added dropwise to the reaction mixture. The flask was warmed to room temperature and then the mixture was heated under reflux for 6 h. The reaction mixture was cooled to room temperature and poured over ice. The organic solvent was evaporated under reduced pressure and the aqueous solution was washed with EtOAc. The aqueous phase was acidified with 2N HCl until the pH was 2, and then it was extracted with EtOAc. The organic phase was concentrated under reduced pressure to vield 4 as a brownish vellow oil. This brownish vellow oil was dissolved in dry THF (300 mL) and cooled to 0 °C. BH₃·DMS (5.7 mL, 58 mmol) was added dropwise to the reaction mixture, which was then kept stirring at 0 °C for an additional 1 h. The flask was removed from the ice bath, and the mixture stirred overnight at room temperature. The reaction mixture was poured over ice and sequentially extracted with EtOAc, washed with saturated aqueous NaHCO₃, washed with brine, and dried (Na₂SO₄). The solution was concentrated under reduced pressure and purified through flash column chromatography (FCC; gradient EtOAc/hexanes, from 30 to 50%) to yield 5 [26] as a slightly yellow oil (16.3 g, 70% over three steps).

The slightly yellow oil from the previous step (16.1 g, 52.3 mmol) was dissolved in dry CH_2Cl_2 (160 mL) and cooled to 0 °C. Et₃N (7.8 mL, 56.0 mmol) was added and then MsCl (6.3 g, 54.9 mmol) was added slowly. After the reaction had reached completion (checked using TLC), the solvent was evaporated under reduced pressure. The residue was dissolved in EtOAc and washed sequentially with water, saturated aqueous NaHCO₃, and brine, then dried (Na₂SO₄). The product was isolated through FCC (EtOAc/ hexanes, 30%) to provide **1** [26] (20.0 g, 99%) as a slightly yellow viscous oil.

(2S,4R)-tert-butyl 4-(benzyloxy)-2-((diphenylphosphino)methyl)pyrrolidine-1carboxylate (2a)

Compound 1 (771 mg, 2 mmol) was placed in a flame-dried round-bottom flask containing a stirrer bar. Dry THF (20 mL) was added via syringe under argon. The flask was cooled to $-45\ensuremath{\,^\circ C}$ (MeCN/dry ice bath) and then $0.5\,N$ Ph₂PK solution in THF (6 mL, 3 mmol) was added over 2 h via syringe pump. When the addition was complete, the mixture was stirred at the same temperature for an additional 7 h. The flask was warmed to room temperature and then H₂O (1-1.5 mL) was added until the reaction color disappeared. CH₂Cl₂ (20 mL) was added and then the organic phase was collected. The aqueous phase was extracted twice with CH₂Cl₂. The combined organic phases were dried (Na₂SO₄) and concentrated under reduced pressure. The residue was purified through FCC (EtOAc/ hexanes, from 1:10 to 1:5) to afford 2a (846 mg, 89%) as a clear viscous oil. $\left[\alpha\right]_{D}^{26}$ -47.4 (c 0.35, acetone); ¹H NMR (500 MHz, CDCl₃) δ 7.61 (br s, 1H), 7.54 (br s, 1H), 7.46 (app t, J = 6.6 Hz, 2H), 7.38–7.30 (m, 11H), 4.49 (br s, 2H), 4.27-4.10 (m, 2H), 3.90-3.43 (m, 2H), 3.04-2.88 (m, 1H), 2.33-2.12 (m, 3H), 1.49 (s, 9H); ¹³C NMR (125 MHz, $CDCl_3$) δ 154.4, 137.8, 132.7 (d, J = 19.2Hz), 132.3 (d, J = 19.5 Hz), 128.6, 128.3 (d, J = 7.1 Hz), 128.2, 127.4, 127.3, 79.4, 70.6, 54.1 (d, J = 18.9 Hz), 53.2, 38.6, 35.0 (d, J = 10.8 Hz), 33.5 (br s), 28.3; ³¹P NMR (202 MHz, CDCl₃) δ -22.9; IR (film) ν_{max} 2975, 2930, 2362, 2338, 1690, 1393, 1171, 739, 697 cm⁻¹; HRMS-DART: calcd for $C_{29}H_{35}NO_{3}P$ ([M+H]⁺) *m/z* 476.2349; found 476.2345.

General procedure for preparing 2b-e

The phosphine **2a** (0.20 g, 0.42 mmol) was dissolved in CH_2Cl_2 (1.5 mL) in a flame-dried round-bottom flask containing a stirrer bar. TFA (0.5 mL) was added dropwise to the solution at 0 °C and then the mixture was stirred at room temperature for 1 h. CH_2Cl_2 (3 mL) was added and then the mixture was cooled to 0 °C. NEt₃ (1.2 mL) was added dropwise, and then an acyl chloride (1.2 eq) was added

slowly. The mixture was stirred at room temperature overnight and then washed with H_2O . The organic phase was dried (Na₂SO₄) and concentrated under reduced pressure. The residue was purified through FCC.

((2S,4R)-4-(Benzyloxy)-2-((diphenylphosphino) methyl)pyrrolidin-1-yl)(3,5-bis(trifluoromethyl) phenyl)methanone (2b)

Obtained as a white solid (70%). $[\alpha]_{D}^{24}$ -58.77 (c 0.65, acetone); ¹H NMR (500 MHz, CDCl₃) δ 7.87 (s, 1H), 7.60 (s, 2H), 7.57–7.53 (m, 4H), 7.38–7.16 (m, 11H), 4.78–4.70 (m, 1H), 4.46 (d, J = 11.9Hz, 1H), 4.30 (d, J = 11.9Hz, 1H), 4.14 (br s, 1H), 3.53 (dd, J = 11.3, 3.3 Hz, 1H), 3.32 (d, J = 11.3 Hz, 1H), 2.84-2.76 (m, 2H), 2.48 (dd, J = 13.2,7.9 Hz, 1H), 2.34 (br s, 1H); 13 C NMR (125 MHz, CDCl₃) δ 167.1, 138.4, 137.6, 137.5, 137.49, 132.8 (d, *J* = 21.3 Hz), 132.4 (d, J = 18.6 Hz), 131.5 (g, J = 33.7 Hz), 129.1, 128.63, 128.56 (d, J = 7.5 Hz), 128.54 (d, J = 6.8 Hz), 128.4, 127.7, 127.6 (br s), 127.4, 123.6 (sp, J = 3.6 Hz), 122.8 (q, J = 273.7 Hz), 77.1, 70.8, 55.6, 55.0 (d, J = 16.9 Hz), 36.9 (d, J = 10.6 Hz), 32.3 (d, J = 12.9 Hz); ³¹P NMR (202 MHz, CDCl₃) δ –26.4; IR (film) ν_{max} 3057, 2926, 2867, 1642, 1363, 1280, 1137, 906, 740, 697 cm⁻¹; HRMS-DART: calcd for $C_{33}H_{29}F_6NO_2P$ ([M+H]⁺) *m*/*z* 616.1830; found 616.1830.

1-((2S,4R)-4-(benzyloxy)-2-((diphenylphosphino) methyl)pyrrolidin-1-yl)-2,2-dimethylpropan-1-one (2c)

Obtained as a white solid (78%). $[\alpha]_D^{26}$ –4.0 (*c* 0.75, acetone); ¹H NMR (500 MHz, CDCl₃) δ 7.60 (t, *J* = 7.5 Hz, 2H), 7.44–7.41 (m, 2H), 7.37 (app t, *J* = 7.0 Hz, 2H), 7.34–7.26 (m, 9H), 4.52–4.43 (m, 3H) (apparent overlapping peaks), 4.17 (app s, 1H), 3.90 (d, *J* = 11.4 Hz, 1H), 3.56 (dd, *J* = 11.4, 4.4 Hz, 1 H), 2.91 (d, *J* = 13.9 Hz, 1H), 2.24–2.19 (m, 2 H), 2.16–2.10 (m, 1H), 1.17 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ 176.7, 137.8, 132.8 (d, *J* = 18.8 Hz), 132.7 (d, *J* = 19.4 Hz), 128.6, 128.5 (d, *J* = 7.0 Hz), 128.35 (d, *J* = 7.0 Hz), 128.32, 127.6, 127.3, 77.4, 70.6, 55.7 (d, *J* = 17.5 Hz), 53.1, 38.8, 35.5 (d, *J* = 9.3 Hz), 32.8 (d, *J* = 13.7 Hz), 27.5; ³¹P NMR (202 MHz, CDCl₃) δ –23.5; IR (film) ν_{max} 2969, 2930, 1620, 1407, 1093, 1074, 738, 697 cm⁻¹; HRMS-DART: calcd for C₂₉H₃₅NO₂P ([M+H]⁺) *m/z* 460.2400; found 460.2397.

1-((2S,4R)-4-(benzyloxy)-2-((diphenylphosphino) methyl)pyrrolidin-1-yl)-2,2-diphenylethanone (2d)

Obtained as a pale-yellow solid (73%). $[\alpha]_D^{27}$ –50.14 (*c* 0.17, acetone); ¹H NMR (500 MHz, CDCl₃) δ (in aliphatic region, major rotamer) 5.00 (s, 1H), 4.52–4.50 (m,

1H) (overlapping peaks), 4.25 (d, J = 12.2 Hz, 1H), 4.17 (d, J = 11.9 Hz, 1H), 4.13–4.10 (m, 1H), 3.66 (d, J = 10.2 Hz, 1H), 3.49 (dd, J = 11.0, 4.7 Hz, 1H), 3.26 (dt, J = 13.5, 4.5 Hz, 1H), 2.24–2.14 (m, 2H), 2.05–2.00 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) δ (major rotamer) 170.3, 139.4, 138.9, 137.7, 133.0 (d, J = 19.3 Hz), 132.7 (d, J = 19.3 Hz), 129.1, 128.9, 128.8, 128.6, 128.59, 128.56, 128.43, 128.41, 128.35, 128.32, 127.6, 127.5, 126.9 (d, J = 8.0 Hz), 76.5 (d, J = 1.8 Hz), 70.7, 57.0, 54.9 (d, J = 19.2 Hz), 51.9, 36.9 (d, J = 9.8 Hz), 32.7 (d, J = 12.5 Hz); ³¹P NMR (202 MHz, CDCl₃) δ –22.5 (major rotamer), -23.2 (minor rotamer); IR (film) ν_{max} 3060, 3028, 1645, 1411, 1192, 738, 697 cm⁻¹; HRMS-DART: calcd for C₃₈H₃₇NO₂P ([M+H]⁺) *m/z* 570.2556; found 570.2554.

(1S,3R)-adamantan-1-yl((2S,4R)-4-(benzyloxy)-2-((diphenylphosphino)methyl)pyrrolidin-1-yl) methanone (2e)

Obtained as a colorless sticky compound (98%). $[\alpha]_D^{27}+2.7$ (c 1.20, acetone); ¹H NMR (500 MHz, CDCl₃) δ 7.60 (t, J = 7.3 Hz, 2H), 7.44-7.41 (m, 2H), 7.39-7.28 (m, 11H),4.54–4.50 (m, 2H) (overlapping peaks), 4.44 (d, J = 12 Hz, 1H), 4.19 (br s, 1H), 3.99 (d, J = 11.2 Hz, 1H), 3.62 (dd, J = 11.3, 4.4 Hz, 1H), 2.88 (d, J = 13.3 Hz, 1H), 2.19(dd, J = 13.4, 9.5 Hz, 2H), 2.12 (br s, 1H), 1.99 (s, 3H), 1.87 (s, 6H), 1.71–1.65 (app q, J = 12.4 Hz, 6H); ¹³C NMR $(125 \text{ MHz}, \text{ CDCl}_3) \delta 176.1, 137.9, 132.8 \text{ (d, } J = 18.8 \text{ Hz}\text{)},$ 132.7 (d, J = 19.6 Hz), 128.53, 128.48, 128.34, 128.30, 127.6, 127.5, 77.6, 70.8, 55.9 (d, *J* = 16.3 Hz), 53.1, 41.6, 38.0, 36.5, 35.2 (d, *J* = 8.2 Hz), 32.9 (d, *J* = 12.0 Hz), 28.3; ³¹P NMR (202 MHz, CDCl₃) δ –21.8; IR (film) ν_{max} 3053, 2905, 2850, 2235, 1610, 1391, 1073, 909, 734, 697 cm⁻¹; HRMS-DART: calcd for $C_{35}H_{41}F_6NO_2P$ ([M + H]⁺) m/z538.2869; found 538.2872.

(2S,4R)-4-(benzyloxy)-2-((diphenylphosphanyl) methyl)-N-phenylpyrrolidine-1-carboxamide (2f)

The phosphine **2a** (76.0 mg, 0.158 mmol) was dissolved in CH_2Cl_2 (0.5 mL) in a flame-dried round-bottom flask containing with a stirred bar. TFA (0.17 mL) was added dropwise to the solution at 0 °C and then the mixture was stirred at room temperature for 1 h. CH_2Cl_2 (1 mL) was added and then the mixture was cooled to 0 °C. NEt₃ (0.4 mL) was added dropwise and then the reaction was quenched through the addition of H_2O (2 mL). The aqueous phase was exacted with CH_2Cl_2 . The combined organic phases were washed with brine, dried (Na₂SO₄), and concentrated. The residue was dissolved in CH_2Cl_2 (1 mL). Phenyl isocyanate (21.0 mg, 0.176 mmol) was added and then the mixture was stirred overnight. The solution was concentrated and the residue purified through FCC (hexanes/EtOAc, 5:1) to give 2f as a white semisolid. $[\alpha]_{D}^{28}$ -39.8 (c 1.00, acetone); ¹H NMR (500 MHz, d₆-acetone) δ 8.12 (s, 1H), 7.66–6.87 (m, 20H), 4.47 (s, 2H), 4.37–4.21 (m, 2H), 3.68 (d, J = 4.1 Hz, 2H), 3.05 (dt, J =13.5, 3.4 Hz, 1H), 2.28–2.20 (m, 1H), 2.18–2.06 (m, 2H). ¹³C NMR (100 MHz, d_6 -acetone) δ 154.4, 140.5, 139.6 (d, J = 12.9 Hz, 138.7, 138.2 (d, J = 13.0 Hz), 132.8 (d, J =19.6 Hz), 132.6 (d, J = 20.0 Hz), 128.7, 128.6 (d, J = 2.5Hz), 128.5, 128.4 (d, J = 7.0 Hz), 128.3, 128.25 (d, J = 8.1 Hz), 127.4, 121.9 (d, J = 16.7 Hz), 119.5, 118.5, 77.0 (d, J = 1.7 Hz), 70.4, 54.70 (d, J = 20.4 Hz), 51.9, 37.3 (d, J =8.4 Hz), 33.7 (d, J = 13.9 Hz). ³¹P NMR (121 MHz, d_{6} acetone) δ –23.12. IR (film) ν_{max} 3324, 3054, 2926, 2426, 1700, 1633, 1443, 1385, 1246, 750, 695 cm⁻¹; HRMS-DART: calcd for $C_{31}H_{32}N_2O_2P$ ([M+H]⁺) m/z 495.2196; found 485.2219.

General experimental procedure for the allene-imine [4 + 2] annulation (with the allene 8)

The allenoate (0.12 mmol) was added in one portion to a solution of *N*-tosylimine (0.1 mmol) and the aminophosphine catalyst (0.02 mmol) in the solvent (2 mL). The flask was capped with a Teflon cap under a N₂ flow. The mixture was stirred at room temperature for 2 days. The resulting solution was concentrated and the residue purified through FCC on silica gel (eluent: 20–30% EtOAc in hexane) to give the product.

Diethyl (25,35)-2-phenyl-1-tosyl-1,2,3,6tetrahydropyridine-3,4-dicarboxylate (10)

Obtained as a white solid (80%). 57% ee—determined using an SFC instrument, a Daicel ChiralPak IC-3 column, and 20% MeCN; t_R (minor) = 3.74 min; t_R (major) = 4.81 min; $[\alpha]_D^{25}$ -38.0 (c 1.10, acetone); ¹H NMR (400 MHz, CDCl₃) δ 7.64 (d, J = 8.3 Hz, 2H), 7.28–7.16 (m, 7H), 6.94 (dd, J = 3.7, 2.7 Hz, 1H), 5.93 (s, 1H), 4.24–4.12 (m, 4H), 4.11–3.97 (m, 2H), 3.69 (dt, J = 19.9, 2.4 Hz, 1H), 2.38 (s, 3H), 1.26 (t, J = 7.4 Hz, 3H), 1.22 (t, J = 7.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.5, 165.2, 143.5, 137.8, 136.9, 135.2, 129.4, 128.6, 127.8, 127.4, 126.8, 126.4, 61.6, 61.1, 56.1, 44.9, 41.7, 21.4, 14.08, 13.97; IR (film) ν_{max} 2981, 1717, 1344, 1260, 1160, 1096 cm⁻¹. HRMS-DART: calcd for C₂₄H₂₈NSO₆ ([M + H]⁺) m/z 458.1632; found 458.1628.

General experimental procedure for the allene–imine [4+2] annulation (with the allene 11)

The allenoate (0.2 mmol) was added in one portion to a mixture of *N*-tosylimine (0.1 mmol), the aminophosphine catalyst (0.03 mmol), and 4-Å MS (10 mg) in CH_2Cl_2

(2 mL). The flask was capped with a Teflon cap under a N_2 flow. The mixture was stirred at room temperature for 3 days. The resulting solution was concentrated and mesitylene was added as an internal standard to determine the yield and conversion. This mixture was dissolved in THF and then 1 N HCl was added to hydrolyze the imine. The aqueous phase was exacted with CH₂Cl₂. The combine extracts were washed with brine, dried (Na₂SO₄), and concentrated. The residue was purified through PTLC (hexane/EtOAc, 5:1) to give the product for ee determination.

Acknowledgements Financial support for this study was provided by the NIH (R01GM071779). We thank Dr. Saeed Khan (UCLA) for the crystallographic analyses. C.X. thanks Prof. Neil K. Garg and Sarah Anthony (UCLA) for sharing their SFC instrument.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Vitaku E, Smith DT, Njardarson JT. Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among U.S. FDA approved pharmaceuticals. J Med Chem. 2014;57:10257–74.
- Rubiralta M, Giralt E, Diez A. Piperidine. Structure, preparation, reactivity and synthetic applications of piperidine and its derivatives. Elsevier: Amsterdam; 1991.
- Michael JP. Indolizidine and quinolizidine alkaloids. Nat Prod Rep. 2004;21:625–49.
- Michael JP. Simple indolizidine alkaloids. In: Cordell GA, editor. The Alkaloids. Vol. 55. San Diego: Academic Press; 2001.
- Dewick PM. Medicinal natural products. Chichester: John Wiley and Sons; 1997. Chapter 6.
- 6. Buffat MGP. Synthesis of piperidines. Tetrahedron. 2004;60:1701–29.
- Zhu X-F, Lan J, Kwon O. An expedient phosphine-catalyzed [4+2] annulation: synthesis of highly functionalized tetrahydropyridines. J Am Chem Soc. 2003;125:4716–7.

- Lu K, Kwon O. Phosphine-catalyzed [4+2] annulation: synthesis of ethyl 6-phenyl-tosyl-1,2,5,6-tetrahydropyridine-3-carboxylate. Org Synth. 2009;86:212–24.
- Wurz RP, Fu GC. Catalytic asymmetric synthesis of piperidine derivatives through the [4+2] annulation of imines with allenes. J Am Chem Soc. 2005;127:12234–5.
- Xiao H, et al. Bifunctional N-acyl-aminophosphine-catalyzed asymmetric [4+2] cycloadditions of allenoates and imines. Chem Eur J. 2011;17:10562–5.
- Yu H, et al. Phosphine-catalyzed [4+2] cycloaddition of sulfamatederived cyclic imines with allenoates: synthesis of sulfamate-fused tetrahydropyridines. Tetrahedron. 2014;70:340–8.
- 12. Takizawa S, Arteaga FA, Yoshida Y, Suzuki M, Sasai H. Enantioselective organocatalyzed formal [4+2] cycloaddition of ketimines with allenoates: easy access to a tetrahydropyridine framework with a chiral tetrasubstituted stereogenic carbon center. Asian J. Org. Chem. 2014;3:412–5.
- Guo H, Fan YC, Sun Z, Wu Y, Kwon O. Phosphine organocatalysis. Chem Rev. 2018;118:10049–293.
- Ni H, Chan W-L, Lu Y. Phosphine-catalyzed asymmetric organic reactions. Chem Rev. 2018;118:9344–411.
- Fu W, Tang W. Chiral monophosphorus ligands for asymmetric catalytic reactions. ACS Catal. 2016;6:4814–58.
- Tang W, Zhang X. New chiral phosphorus ligands for enantioselective hydrogenation. Chem Rev. 2003;103:3029–70.
- Henry CE, et al. Hydroxyproline-derived pseudoenantiomeric [2.2.1] bicyclic phosphines: asymmetric synthesis of (+)- and (-)-pyrrolines. J Am Chem Soc. 2014;136:11890–3.
- Smaligo AJ, Vardhineedi S, Kwon O. Carvone-derived P-stereogenic phosphines: design, synthesis, and use in allene–imine [3 +2] annulation. ACS Catal. 2018;8:5188–92.
- 19. Xu Q, et al. Catalytic enantioselective synthesis of guvacine derivatives through [4 + 2] annulations of imines with α -methylallenoates. Org Lett. 2018;20:6089–93.
- Bertelsen S, Jørgensen KA. Organocatalysis: after the gold rush. Chem Soc Rev. 2009;38:2178–89.
- Jensen KL, Dickmeiss G, Jiang H, Albrecht Ł, Jørgensen KA. The diarylprolinol silyl ether system: a general organocatalyst. Acc Chem Res. 2012;45:248–64.
- Jiang H, Albrecht Ł, Jørgensen KA. Aminocatalytic remote functionalization strategies. Chem Sci. 2013;4:2287–2300.
- Burés J, Armstrong A, Blackmond DG. Curtin–Hammett paradigm for stereocontrol in organocatalysis by diarylprolinol ether catalysts. J Am Chem Soc. 2012;134:6741–50.
- 24. Tran YS, Kwon O. Phosphine-catalyzed [4+2] annulation: synthesis of cyclohexenes. J Am Chem Soc. 2007;129:12632–3.
- Moriarty KJ, et al. N-heterocyclic inhibitors of TNF-ALPHA expression, US2002/65270 (A1); 2002.
- 26. Lee K, et al. Melanocortin receptor agonists, US2010120783 (A1); 2010.