

GLC Determination of Degradation of Two Related Amine Uptake Inhibitors

REX W. SOUTER* and ALAN DINNER

Abstract □ A convenient GLC method was developed to assay the chemically related CNS agents *dl*-3-(*p*-trifluoromethylphenoxy)-*N*-methyl-3-phenylpropylamine (Ia) and *dl*-3-(*o*-methoxyphenoxy)-*N*-methyl-3-phenylpropylamine (Ib). Subsequently, this GLC method was used to monitor the stability of Ia and Ib in acidic and basic media. These CNS agents were stable in basic solution but labile under acid conditions, and the degradations of both compounds approximated pseudo-first-order kinetics. Products formed during the acid hydrolysis of Ia and Ib were identified and were shown not to interfere in the GLC assays of the parent ethers.

Keyphrases □ GLC—stability determination, 3-phenylpropylamines, CNS agents □ CNS agents—3-phenylpropylamines, GLC stability determination

After the synthesis of a potential new drug, a suitable analytical assay procedure is needed for stability studies. Ideally, this method should be facile, able to detect the drug at low levels, and valid in the presence of chemical degradation products of the drug.


This paper reports a method for the assays of the selective serotonin uptake inhibitor (1-3) *dl*-3-(*p*-trifluoromethylphenoxy)-*N*-methyl-3-phenylpropylamine (Ia) and the norepinephrine uptake inhibitor (2) *dl*-3-(*o*-methoxyphenoxy)-*N*-methyl-3-phenylpropylamine (Ib). Both compounds are currently under study as inhibitors of uptake into monoaminergic neurons in brain.

EXPERIMENTAL

Reagents—The solvents used were all of analytical reagent quality. The 99% *n*-pentacosane¹ used as internal standard, the OV-225 liquid phase², and the acid-washed, dimethyldichlorosilane-treated Chromosorb G³ were all purchased materials.

Apparatus—A high-precision gas chromatograph was used⁴. Screw-capped vials⁵ with septums⁶ were used for all sample extractions.

Chromatographic Conditions—The column was a 0.9-m × 0.83-cm (3-ft × 0.25-in.) o.d. acid-washed, dimethyldichlorosilane-treated glass U-tube packed with 4% OV-225 on 80-100-mesh



Chromosorb G. The oven was operated at 200° for studies of Ia and at 215° for studies of Ib. Helium was used as the carrier gas (after passing through a drying trap packed with 5A molecular sieves) at an approximate flow rate of 60 ml/min.

Procedure—Peak areas were calculated using an on-line calculation program from the expanded RTE 2100 computer system⁴. Area ratios of sample-internal standard usually agreed within 1% for replicate injections.

For the degradation studies, a known amount (150-300 mg) of Ia or Ib as the hydrochloride salt was added to 100 ml of refluxing 0.1 N HCl. At various time intervals, aliquots of solution were removed and cooled in ice water. A 2-ml sample of each cooled aliquot was pipetted into the screw-capped septum vial. To this vial was added 7.4 N NH₄OH (to bring the pH to 7.5-8), and the solutions were mixed. Finally, 2.0 ml of a chloroform solution of the internal standard *n*-pentacosane (0.5-0.6 mg/ml) was added, the vial was capped, and the mixture was shaken for 2 min.

The two layers were allowed to separate, and then samples of the chloroform portion were injected into the GLC column. Plots of log [ether] versus time were constructed using a least-squares program available through a computer⁷, and the slopes of the resulting straight lines were used to calculate the pseudo-first-order rate constants. Reactions were followed for at least two half-lives (C and D in Fig. 1). NMR spectra were obtained⁸ using tetramethylsilane as the internal reference. Mass spectra were obtained on a gas chromatograph-mass spectrometer⁹ unit at 70 ev.




Figure 1—Kinetic plots for acid degradation of Ia and Ib. Key: A, Ia, initial measured concentration = 2.03 mg/ml, slope = -0.078; B, Ia, initial measured concentration = 1.50 mg/ml, slope = -0.097; C, Ib, initial measured concentration = 2.63 mg/ml, slope = -0.33; and D, Ib, initial measured concentration = 1.97 mg/ml, slope = -0.46.

¹ Lachat Chemicals, Mequon, Wis.

² Ohio Valley Specialty Chemicals, Marietta, Ohio.

³ Johns-Manville, Denver, Colo.

⁴ Model F&M 402, Hewlett-Packard, Avondale, Pa.


⁵ No. 13028, Pierce Chemical Co., Rockford, Ill.

⁶ Lined with Teflon, Canton Biomedical Products, Boulder, Colo.

⁷ DEC-10, Digital Equipment Corp., Marlborough, Me.

⁸ Varian T-60A, Varian Instrument Division, Palo Alto, Calif.

⁹ LKB 9000S, LKB Instruments, Rockville, Md.



Decomposition of *dl*-3-(*p*-Trifluoromethylphenoxy)-*N*-methyl-3-phenylpropylamine (Ia)—A solution of 300 mg (1 mmole) of Ia in 100 ml of 0.1 *N* aqueous hydrochloric acid was allowed to reflux for 24 hr. The solution was cooled, adjusted to pH 8 with 7.4 *N* NH₄OH, and then extracted with three 50-ml portions of chloroform. The chloroform layers were combined, dried over anhydrous magnesium sulfate, and finally concentrated on a rotary evaporator at 25°.

A portion of the concentrated chloroform solution was streaked onto a 500- μ m TLC plate¹⁰, and the plate was developed using methanol-acetic acid (90:10) as the eluant. Three zones were observed under 254-nm UV light (*R*_f 0.83, 0.65, and 0.55). The zones were scraped from the plate and eluted with chloroform, and the resulting solutions were evaporated. The fastest-moving zone was identified as *p*-trifluoromethylphenol (IIa); NMR (carbon tetrachloride): δ 7.65 (d, 2H, *J* = 8 Hz, aromatics), 6.9 (d, 2H, *J* = 8 Hz, aromatics), and 6.3 (s, 1H, OH); mass spectrum molecular ion at *m/e* 162, calculated for C₇H₅F₃O, 162; prominent fragmentation peak at *m/e* 143 (*p*⁺ - F). TLC comparison with an authentic sample gave the same *R*_f values in two different solvent systems.

The middle zone was identified *via* IR comparison as unreacted starting material. The slowest-moving zone was identified as *N*-methyl-3-phenyl-3-hydroxypropylamine (III); NMR (deuterated methanol): δ 7.4 (s, 5H, aromatics), 4.9 (t, 1H, *J* = 6 Hz, benzylic H), 3.1 (t, 2H, *J* = 6 Hz, NCH₂—), 2.6 (s, 3H, NCH₃), and 2.1 (q, 2H, *J* = 6 Hz, CHCH₂—); mass spectrum molecular ion at *m/e* 165, calculated for C₁₀H₁₅NO, 165; major fragment at *m/e* 147 (*M*⁺ - H₂O).

Decomposition of *dl*-3-(*o*-Methoxyphenoxy)-*N*-methyl-3-





Figure 2—Typical gas chromatogram of partially degraded Ia. Key: a, *p*-trifluoromethylphenol; b, V; c, VI; d, Ia; and e, *n*-pentacosane.

phenylpropylamine (Ib)—This material was degraded and worked up in a manner similar to that of Ia. Development of the thin-layer chromatogram revealed the presence of three zones (*R*_f 0.69, 0.65, and 0.55). The fastest-moving zone was shown to be guaiacol (*o*-methoxyphenol, IIb) *via* IR comparison with authentic material. The middle zone was unreacted starting material (IR comparison), and the slowest-moving zone was identified (as for Ia) as III.

RESULTS AND DISCUSSION

Calibration curves for Ia and Ib were obtained, and they re-

¹⁰ Silica gel F-254, EM Labs., Elmsford, N.Y.




Figure 3—Typical gas chromatogram of partially degraded *Ib*. Key: a, *IIb*; b, *IV*; c, *n*-pentacosane; and d, *Ib*.

vealed a linear relationship between the peak area of compound—peak area of internal standard and the concentration of sample injected between 1.0×10^{-2} and $2.4 \mu\text{g}/\mu\text{l}$. Quantities as low as 10 ng were detectable for either compound.

The kinetics of the acidic degradation of solutions of ethers *Ia*

and *Ib* at two different starting concentrations are shown in Fig. 1. The degradations of both compounds approximate pseudo-first-order kinetics, with a k value of 0.20 hr^{-1} for *Ia* and of 0.90 hr^{-1} for *Ib*. Although these conditions are generally too mild for the cleavage of either aryl alkyl (4, 5) or dialkyl ethers (6, 7), the hydrolysis of aryl benzyl ethers is quite facile (8, 9). Benzyl phenyl ether itself has been cleaved in various solvents with hydrobromic acid at room temperature (10).

Literature reports of the solvolyses of other benzyl phenyl ethers (9, 10) suggest that the cleavage rates of the two compounds examined were not unusual.

The products of the acid-catalyzed cleavage of *Ia* were *p*-trifluoromethylphenol¹¹ (*IIa*) and *N*-methyl-3-phenyl-3-hydroxypropylamine (*III*), while *Ib* yielded *o*-methoxyphenol (*IIb*) and *III*. As expected, no benzyl halide was found since it would be hydrolyzed immediately under the reaction conditions (11) to the observed benzyl alcohol; no *O*-demethylation of *Ib* was noted.

The NMR, IR, and mass spectral data were consistent with the proposed structures. A combined GLC-mass spectral analysis of the product mixture from either *Ia* or *Ib* also revealed a small amount of material (1%) at m/e 147. The major fragments at m/e 146 and 117 ($\text{C}_6\text{H}_5\text{CH}=\text{CHCH}_2^+$) strongly suggest *IV* as the likely structure. Injection of purified *III* into the gas chromatograph revealed no decomposition of *IV*. Hence, *IV* is generated from the acid-catalyzed dehydration of *III* in the reaction mixture.

Figures 2 and 3 represent typical chromatograms of partially degraded samples of *Ia* and *Ib*. Clearly, the degradation products do not interfere with the starting material or with the internal standard. In Fig. 2, peaks b and c represent impurities present in the *Ia* used. The identities of b and c were determined by use of GLC-mass spectrometry and by comparisons of chromatographic retention times with authentic samples. Peak b is 3-(*p*-trifluoromethylphenoxy)-*N,N*-dimethyl-3-phenylpropylamine (*V*), and peak c is 3-(*m*-trifluoromethylphenoxy)-*N*-methyl-3-phenylpropylamine (*VI*).

Attempted basic degradation of either *Ia* or *Ib* with 0.1 *N* aqueous sodium hydroxide at 100° resulted in complete recovery of starting material after 3 days. In general, alkali metals are required for the basic cleavage of benzyl aryl ethers (12).

REFERENCES

- (1) D. Wong, J. Horn, F. Bymaster, K. Hauser, and B. Molloy, *Life Sci.*, **15**, 471(1974).
- (2) D. Wong *et al.*, *Fed. Proc.*, **33**, 255(1974).
- (3) R. Fuller, K. Perry, and B. Molloy, *Life Sci.*, **15**, 1161(1974).
- (4) J. Bunnett, E. Bunzel, and K. Nahakedian, *J. Amer. Chem. Soc.*, **84**, 4136(1962).
- (5) J. Bunnett and E. Bunzel, *ibid.*, **83**, 117(1961).
- (6) R. Burwell, Jr., and M. Fuller, *ibid.*, **79**, 2332(1957).
- (7) D. Jaques and J. Leiston, *J. Chem. Soc.*, **1961**, 4963.
- (8) K. Kratzl and G. Mischke, *Monatsch. Chem.*, **94**, 753(1963).
- (9) H. Hart and R. Elias, *J. Amer. Chem. Soc.*, **83**, 895(1961).
- (10) A. Drummond and A. Eastham, *ibid.*, **79**, 3689(1957).
- (11) R. Burwell, Jr., *Chem. Rev.*, **54**, 615(1954).
- (12) P. Schorigin, *Chem. Ber.*, **57**, 1627(1924).

ACKNOWLEDGMENTS AND ADDRESSES

Received February 11, 1975, from Lilly Research Laboratories, Indianapolis, IN 46206

Accepted for publication June 4, 1975.

The authors thank Mr. Joseph Gregory and Mr. John Brunson for technical assistance and Mr. Kenneth Hauser for providing authentic samples of *V* and *VI*.

* To whom inquiries should be directed.

¹¹ This material was shown to be a metabolite of *Ia* in the rat; C. J. Parli, unpublished results.