

Acta Cryst. (1999). **C55**, 573–574

Bis(dipropylthiocarbamoyl) disulfide†

FANGFANG JIAN,^a LIJIAN JIANG,^a HOONG-KUN FUN,^b
KANDASAMY CHINNAKALI,^{b,‡} IBRAHIM ABDUL RAZAK^b AND
XIAOZENG YOU^a

^aCoordination Chemistry Institute & State Key Laboratory of Coordination Chemistry, Nanjing University, Nanjing 210093, People's Republic of China, and ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia. E-mail: hkfun@usm.my


(Received 23 October 1998; accepted 24 November 1998)

Abstract

In the title compound, $C_{14}H_{28}N_2S_4$, the two dithiocarbamate groups, which are perpendicular to each other, are bridged by an S—S bond. The C—N bonds in the dithiocarbamate groups show partial double-bond character.

Comment

The bis(dialkylthiocarbamoyl)disulfide compounds are highly effective in breaking the dormancy of plant seeds, bulbs and tubers. These compounds can be used as herbicides or in the cultivation of crop plants during the off season (Hideo *et al.*, 1974). Gasoline basestocks are blended with 0.001–0.5 wt% of these compounds to obtain a stable compound suitable for long-term storage without sludge deposition (Kenichiro & Michiro, 1992). They are also used as an additive for electrolytes of secondary lithium batteries. The use of these electrolytes prevents growth of Li dendrites and renders the batteries a longer life (Masayuki, 1996).



The title molecule, (I), consists of two dipropylthiocarbamate units bridged by an S—S bond. The two planar dithiocarbamate units, containing a C—S and a C=S bond, are oriented perpendicular to each other with a dihedral angle of 89.98 (4)°. The shorter N1—C1 and N2—C8 bond distances [1.326 (3) and 1.337 (3) Å] in the dithiocarbamate units are indicative of considerable double-bond character. The S—C,

S=C and C—N bond distances are comparable with those observed in the related structures (Sharma *et al.*, 1991; Gimeno *et al.*, 1996; Jian *et al.*, 1999). Short C2...S1 [2.893 (3) Å], C5...S2 [3.039 (3) Å], C9...S3 [2.910 (3) Å] and C12...S4 [3.061 (3) Å] intramolecular contacts are observed in this structure. The crystal structure is stabilized by van der Waals interactions.




Fig. 1. The structure of the title compound showing 50% probability displacement ellipsoids and the atom-numbering scheme.

Experimental

To a heated aqueous solution of *N,N'*-dipropylthiocarbamate sodium salt was added a solution of lanthanum chloride with stirring. The white precipitate which formed was collected by filtration. The deposit was recrystallized from an acetonitrile solution.

Crystal data

$C_{14}H_{28}N_2S_4$
 $M_r = 352.62$
Orthorhombic
Pbca
 $a = 11.0232$ (2) Å
 $b = 15.1645$ (2) Å
 $c = 24.2826$ (1) Å
 $V = 4059.11$ (9) Å³
 $Z = 8$
 $D_x = 1.154$ Mg m⁻³
 D_m not measured

Mo $K\alpha$ radiation
 $\lambda = 0.71073$ Å
Cell parameters from 8192 reflections
 $\theta = 2.69$ – 33.18°
 $\mu = 0.462$ mm⁻¹
 $T = 293$ (2) K
Block
 $0.52 \times 0.40 \times 0.34$ mm
Colourless

Data collection

Siemens SMART CCD area-detector diffractometer
 ω scans
Absorption correction: empirical (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.79$, $T_{\max} = 0.86$
24592 measured reflections

4662 independent reflections
3205 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$
 $\theta_{\max} = 27.50^\circ$
 $h = 0 \rightarrow 14$
 $k = 0 \rightarrow 19$
 $l = 0 \rightarrow 31$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.143$

$(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.36$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³

† Alternative name: dithiobis(*N,N*-dipropylthioformamide).

‡ On leave from: Department of Physics, Anna University, Chennai 600 025, India.

$S = 1.06$
 4662 reflections
 185 parameters
 H atoms constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0558P)^2 + 1.6893P]$
 where $P = (F_o^2 + 2F_c^2)/3$

Extinction correction: none
 Scattering factors from
*International Tables for
 Crystallography* (Vol. C)

Sheldrick, G. M. (1997). *SHELXTL. Structure Determination Programs*. Version 5.10. Bruker Analytical X-ray Systems, Madison, Wisconsin, USA.
 Siemens (1996). *SMART and SAINT. Area-Detector Control and Integration Software*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Spek, A. L. (1990). *Acta Cryst.* **A46**, C-34.

Table 1. Selected geometric parameters (Å, °)

S1—C1	1.814 (3)	N1—C2	1.475 (3)
S1—S3	1.997 (1)	N1—C5	1.478 (3)
S2—C1	1.653 (2)	N2—C8	1.337 (3)
S3—C8	1.820 (3)	N2—C9	1.472 (3)
S4—C8	1.646 (3)	N2—C12	1.472 (3)
N1—C1	1.326 (3)		
C1—N1—C2—C3	92.0 (3)	C8—N2—C9—C10	−96.9 (3)
C1—N1—C5—C6	91.4 (3)	C8—N2—C12—C13	−90.0 (3)

The data collection covered a hemisphere of reciprocal space by a combination of three sets of exposures; each set had a different φ angle (0, 88 and 180°) for the crystal and each exposure of 30 s covered 0.3° in ω . The crystal-to-detector distance was 4 cm and the detector swing angle was −35°. Crystal decay was monitored by repeating 30 initial frames at the end of data collection and analysing the intensity of duplicate reflections, and was found to be negligible.

All H atoms were geometrically fixed and allowed to ride on the attached atoms; rotating group refinement was used for the methyl groups. Atoms C4 and C11 of the propyl groups show high thermal vibrations but no indications of disorder were observed.

Data collection: *SMART* (Siemens, 1996). Cell refinement: *SAINT* (Siemens, 1996). Data reduction: *SAINT*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1997). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 1990).

This work was supported by the NSFC (project 29501001) and NSF of Jiangsu Province (project BK95024401). The authors would like to thank the Malaysian Government for research grant R&D No. 190/9609/2801. KC thanks the Universiti Sains Malaysia for a Visiting Post-Doctoral Fellowship.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1256). Services for accessing these data are described at the back of the journal.

References

- Gimeno, M. C., Jambrina, E., Laguna, A., Laguna, M., Murray, H. H. & Terroba, R. (1996). *Inorg. Chim. Acta*, **249**, 69–77.
 Hideo, O., Kaoru, I. & Tetsuya, O. (1974). Japanese Patent 73 18, 450.
 Jian, F. F., Zhu, D. R., Fun, H.-K., Chinnakali, K., Razak, I. A. & You, X. Z. (1999). *Acta Cryst.* **C55**. In the press.
 Kenichiro, S. & Michiro, M. (1992). Japanese Patent 04 359, 091.
 Masayuki, M. (1996). Japanese Patent 06 36, 797.
 Sharma, S., Bohra, R. & Mehrotra, R. C. (1991). *J. Crystallogr. Spectrosc. Res.* **21**, 61–64.
 Sheldrick, G. M. (1996). *SADABS. Program for Empirical Absorption Correction of Area-Detector Data*. University of Göttingen, Germany.

Acta Cryst. (1999). **C55**, 574–576

p-Hydroxybenzaldehyde nicotinoyl-hydrazone monohydrate

HOONG-KUN FUN,^a KANDASAMY CHINNAKALI,^{a†} IBRAHIM ABDUL RAZAK,^a ZHONG-LIN LU^b AND BEI-SHENG KANG^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bThe Institute of Physical Chemistry, School of Chemistry and Chemical Engineering, Zhongshan University, Guangzhou 510 275, People's Republic of China. E-mail: hkfun@usm.my


(Received 30 June 1998; accepted 25 November 1998)

Abstract

The title compound, C₁₃H₁₁N₃O₂·H₂O, is in the hydrazone tautomeric form and the configuration of the azomethine C=N double bond [1.275 (3) Å] is *E*. In the solid state, the molecules are stacked parallel to (110) planes and form a hydrogen-bonded column-like supramolecular structure involving two independent water molecules on twofold axes.

Comment

Arylhydrazones have been investigated extensively in recent years because of their biological activity, especially as potent inhibitors for many enzymes (Ma *et al.*, 1994; Lu *et al.*, 1996; Fun, Lu, Duan, Tian, You, Gong & Guo, 1997; Fun, Lu, Duan, Tian, You, Guo & Gong, 1997). As part of our work on the synthesis and characterization of arylhydrazone compounds, we report here the structure of *p*-hydroxybenzaldehyde nicotinoyl-hydrazone monohydrate, (I).



The geometry of the molecule agrees well with those found in related structures (Fun, Lu, Duan, Tian, You,

† On leave from: Department of Physics, Anna University, Chennai 600 025, India.