



A Journal of



Accepted Article

Title: 2-Bromo-2-chloro-3-arylpropanenitriles as C-3 Synthons for the Synthesis of Functionalized 3-Aminothiophenes

Authors: Sviatoslav Batsyts, Maksym Shehedyn, Evgeny Goreshnik, Mykola D. Obushak, Andreas Schmidt, and Yurii Ostapiuk

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Org. Chem.* 10.1002/ejoc.201901512

Link to VoR: <http://dx.doi.org/10.1002/ejoc.201901512>

Supported by



WILEY-VCH

FULL PAPER Accepted Article

2-Bromo-2-chloro-3-arylpropanenitriles as C-3 Synthons for the Synthesis of Functionalized 3-Aminothiophenes


Sviatoslav Batsyts,^[a,b] Maksym Shehedyn,^[a] Evgeny Goreshnik,^[c] Mykola D. Obushak,^[a] Andreas Schmidt,^{*[b]} and Yurii Ostapiuk^{*[a]}

Abstract: 2-Bromo-2-chloro-3-arylpropanenitriles can be prepared by Meerwein reaction from 2-chloroacrylonitrile and various aryl diazonium salts under copper(II) bromide catalysis. They proved to be stable compounds which form 2-chlorocinnamonnitriles upon treatment with bases. Reaction of the title compounds with substituted thioglycolates gave substituted 3-aminothiophenes which have not yet been accessible by other routes. Three-component reactions with the title compound, sodium sulfide and bromonitromethane, chloroacetonitrile, or ethyl 4-chloroacetoacetate gave 2-nitro- and 2-cyano-substituted 3-aminothiophenes, and thienopyridinediones in high yields and in one single step, respectively.

Introduction


Undoubtedly thiophenes^[1] belong to the most important heterocyclic ring systems, as thiophene derivatives have found numerous applications^[2] as biologically active compounds^[3] or as structure increments of new materials. Thus, thiophenes proved to be active as antifungal,^[4] channel blocking,^[5] anti-tumor^[6] (Raltitrexed, Tomudex®), and anti-parasitic compounds.^[7] Among the latter mentioned, antileishmanial activities have been reported.^[8] They were also developed as inhibitors of the NorA multidrug transporter of bacteria^[9] and 17β-hydroxysteroid dehydrogenase type 1,^[10] and as agonists of the sphingosine-1-phosphate 1 receptor.^[11] Concerning materials chemistry, liquid crystals,^[12] non-linear optical materials,^[13] conducting polymers,^[14] photoluminescent compounds,^[15] semiconductors,^[16] and light-emitting materials^[17] on the basis of thiophenes have been described. Thiophenes can be prepared by the formation of one, two or three bonds from suitable starting materials by ring closure reactions, concerted cycloadditions, rearrangements, or functionalization of an existing thiophene ring.^[1] Among these

methods, the Gewald reaction^[18] is very convenient to prepare 2-aminothiophenes with a high degree of functionalization (Scheme 1), and nowadays, the chemistry of 2-aminothiophenes is still one of the most extensive and dynamic fields of thiophene research.^[19] Gewald's thiophenes possess the stabilizing β-enaminocarbonyl chromophore which also allows for numerous subsequent functionalizations.^[20]



Scheme 1. Gewald synthesis of 2-aminothiophenes.

In comparison to the syntheses of 2-aminothiophenes, the preparations of 3-aminothiophenes, which are also underrepresented in nature,^[21] are less developed. Nevertheless, 3-aminothiophenes constitute partial structures of numerous bioactive molecules such as the herbicide and fungicide dimethenamid^[22] or the local anesthetic articaine,^[23] among many others (Scheme 2). Fused systems such as thieno[3,2-*b*]pyridine-5(4*H*)-ones were developed very recently as tunable fluorescent dyes.^[24]



Scheme 2. 3-Aminothiophenes as biologically active compounds and new materials

[a] Prof. Dr. Yurii Ostapiuk, Prof. Dr. Mykola D. Obushak, M. Sc. Maksym Shehedyn, Ivan Franko National University of Lviv Department of Organic Chemistry, Kyryla i Mefodiy Str. 6, 79005 Lviv, Ukraine, E-mail: y.ostapiuk@gmail.com http://chem.lnu.edu.ua

[b] Prof. Dr. Andreas Schmidt, Dr. Sviatoslav Batsyts, Clausthal University of Technology, Institute of Organic Chemistry, Leibnizstrasse 6, D-38678 Clausthal-Zellerfeld, Germany E-mail: schmidt@ioe.tu-clausthal.de

[c] Dr. Evgeny Goreshnik Jozef Stefan Institute, Department of Inorganic Chemistry and Technology, Jamova 39, SI-1000 Ljubljana, Slovenia

Supporting information for this article is given via a link at the end of the document.

First prepared in 1926^[25] 3-aminothiophenes can be obtained by Hoffmann or Curtius rearrangements^[21] or by Pd-catalyzed amination starting from 3-halothiophenes.^[26] Very recently, a 6π-electrocyclization involving a vinyl sulfide linked to an *in-situ* generated keteniminium salt for the synthesis of 3-aminothiophenes was developed by which one bond is formed to close the ring.^[21] Concerning two- or three-component reactions, 3-aminothiophenes can be obtained from unsaturated functionalized nitriles I - III by Thorpe-Ziegler-type cyclizations, where the 3-aminothiophenes are often 5-aryl-substituted (Figure 1). By far the mostly applied starting materials are substituted nitriles I,^[27] whereas only a few applications of α-substituted

FULL PAPER Accepted Article

nitriles **II**^[28] or 1-cyanoalkynes **III**^[29] have been described. The syntheses of the precursors **I** and **II**, however, are limited to suitably substituted acetophenones^[27] and benzaldehydes^[28] as starting materials, respectively. In most cases, the syntheses of the nitriles **I** and **II** are restricted to donor-substituents attached to the aromatic ring, whereas acceptor-substituents result in strongly decreased yields of the nitriles formed. A few examples using β -oxo-substituted nitriles as alternative to **I** have been described.^[30] The starting materials are available by Meerwein reaction, *i.e.* the coupling of arenediazonium salts with substituted alkenes using copper(II) salts as catalyst,^[31] or by its variations.^[32] Aryl-alkenes as well as aryl-alkanes can thus be prepared from arenediazonium salts which combine several advantages as starting materials in organic synthesis. They can easily be prepared from a broad variety of – often inexpensive – aniline derivatives at ambient reaction conditions. It is not astonishing that the Meerwein arylation has developed to a broadly applicable and versatile synthetic approach for the functionalization of alkenes^[33] for the preparation of interesting target molecules such as natural products and drugs.^[34]





Figure 1. Synthetic approaches to functionalized thiophenes.

In continuation of our interest in heterocycle synthesis^[35] including thiophene functionalization,^[27c,27d,36] and Meerwein reactions^[37] we report here on a useful synthesis of 3-aminothiophene derivatives starting from 2-bromo-2-chloro-3-phenylpropanenitriles. To the best of our knowledge, this substance class has not yet been described. It allows for the introduction of a large number of functional groups in thiophene syntheses which are otherwise difficult or even impossible to prepare.

Results and Discussion

To investigate optimal reaction conditions of the Meerwein reaction for the synthesis of 2-bromo-2-chloro-3-phenylpropanenitriles, aniline **1a** and 2-chloroacrylonitrile were used as model compounds (Scheme 3). 2-Chloroacrylonitrile was obtained in two steps from acrylonitrile by chlorination followed by dehydrochlorination of the formed dichloro compound. First, aniline **1a** was added to an aqueous solution of hydrogen bromide (2.2 equiv) and then cooled to -5 °C. A saturated solution of sodium nitrite (1.1 equiv) in water was added to the resulting anilinium bromide within 5 min in the temperature range from -5 to +5 °C to form the phenyl diazonium bromide **2** *in situ* which was directly used for the arylation of 2-chloroacrylonitrile without prior isolation.



	catalyst (mol%)	solvent ^[a]	by-product ^[b]	yield ^[c] (%)
1	CuBr (15)	acetone	50	15
2	CuBr (10)	acetone	45	17
3	CuBr (5)	acetone	40	26
4	CuBr (1)	acetone	37	30
5	CuBr ₂ (15)	acetone	16	40
6	CuBr ₂ (5)	acetone	17	39
7	CuBr ₂ (1)	acetone	15	40
8	CuBr ₂ (0.45)	acetone	14	41
9	CuBr ₂ (0.45)	EtOH	PhOH	0
10	CuBr ₂ (0.45)	MeOH	PhOH	0
11	CuBr ₂ (0.45)	AcOH	-	25
12	CuBr ₂ (0.45)	DMF	-	20
13	CuBr ₂ (0.45)	1,4-dioxane	-	5


^[a] solvent/water = 1:1. ^[b] Sandmeyer product (bromobenzene).
^[c] isolated yield.

Scheme 3. Optimization of reaction conditions.

During optimization of the reaction conditions we found that propanenitrile **3a** was formed in 41% yield, when the reaction was carried out in acetone/water (1:1) in the presence of CuBr₂ (0.45 mol%). The product **3a** was finally separated by distillation under reduced pressure. Using other polar organic solvents such as dioxane, acetic acid, DMF, or DMSO, respectively, diminished the yield of nitrile **3a**. In comparison to copper(II) bromide, the reaction with copper(I) bromide also decreased the yield of the product due to the formation of bromobenzene by Sandmeyer reaction. Applying the optimized conditions, the arylation of 2-chloroacrylonitrile **2** with other electron-poor and electron-rich arenediazonium salts **1b-g** was performed (Scheme 4). The resulting propanenitriles **3a-g** were extracted with dichloromethane from the reaction mixtures and distilled under reduced pressure (1 - 3 torr). Distillations under slightly higher pressures (~10 torr) have led to a partial loss of HBr and formation of cyanonitriles **4a-g** which are also available on treatment of **3a-g** with NEt₃ in toluene in almost quantitative yields. The previously undescribed nitriles **3a-g** have a pungent smell and


FULL PAPER Accepted Article

proved to be stable on storage at 4 °C over a period of 5 years without any traces of decomposition. The hydrogen atoms of the methylene groups are diastereotopic due to the asymmetric environment of the adjacent chiral carbon atom of **3a-g** and give two overlapped doublets with coupling constants 2J of approximately 14.2 Hz in the chemical shift range from 3.77 to 4.17 ppm, depending on the substitution pattern. A comparison of NMR data of the 2-chlorocinnamonnitriles **4** with literature values^[38] shows that **4a,b,e,g** are formed exclusively as *Z*-isomers, whereas the other derivatives gave mixtures of *Z/E*-isomers. Compound **4a** has been prepared before starting from benzaldehyde and chloroacetonitrile^[39] or trichloroacetonitrile,^[40] by the reaction of a dioxazaborocane derivative with zinc(II) cyanide in the presence of copper(II) nitrate,^[41] and by treatment of cinnamonnitrile with HCl/DMF/oxone in low yield.^[42] The compounds **4a** and **4c** are also available starting from the corresponding 2-cyano-3-phenylacrylates^[43] or the benzaldehydes and 4-azido-3-chloro-5-methoxy-2(*H*)-furanone.^[44] They have not been synthesized before as described here. To the best of our knowledge, the cinnamonnitriles **4b**, **4f**, and **4g** are new compounds.




Scheme 4. Syntheses of propanenitriles 3 and 2-chlorocinnamonnitriles 4.

The 2-bromo-2-chloro-3-phenylpropanenitriles **3a-g** underwent smooth reactions with a broad variety of thioglycolates **5A-F** in the presence of sodium methanolate in methanol within 1 h. Thus, the 3-aminothiophenes **6Aa-Fg** were prepared in good to excellent yields which differ in their substitution pattern in positions C-2 and C-5 (Scheme 5). The thiophene **6Aa** has *i.a.* been prepared before according to route I (Figure 1) in 54% yield,^[45] and a similar yield was given in a patent.^[46] An alternative approach starts from acetophenone which is subjected to a Vilsmeier-Haak reaction, followed by condensation with hydroxylamine and subsequent treatment with methyl mercaptoacetate.^[47] According to a literature research, the entire series from **6Ba**, **6Ca**, **6Da**, **6Ea** to **6Fa** are new compounds. Their syntheses seem to be very difficult or even impossible to accomplish by established methods.



Scheme 5. Formation of 3-amino-5-aryltiophenes with various substituents at C-2 and C-5 position (6Aa-Fg).

In continuation of this project, we performed three-component reactions for the preparation 3-aminothiophenes **6Ga-g** and **6Ha-g** starting from 2-bromo-2-chloro-3-phenylpropanenitriles **3a-g** employing sodium sulfide and bromonitromethane **5G** or chloroacetonitrile **5H** (Scheme 6). Best results were achieved when methanol as solvent and freshly prepared sodium hydrosulfide were used. Exchanging the solvent to DMF and applying harsher reaction conditions such as reflux temperature caused decreased yields. In comparison to literature-known procedures to synthesize 2-nitro-3-amino-5-phenylthiophene **6Ga** according to route I (Figure 1) which gave very low yields,^[48] the method described here is very useful. Alternative methods to prepare 3-aminothiophene-2-carbonitriles such as **6Ha**, however, gave better yields. They use S-(isocyanomethyl) ethanethioate and bromoethene^[49] or 3-phenylpropiolonitrile^[50] as starting materials, respectively.




Scheme 6. Three component reaction for the synthesis of 3-aminothiophenes.

Replacing the methanol and sodium methanolate by ethanol and sodium ethoxide gave the opportunity to perform the reaction with ethyl 4-bromocrotonate **5I** or ethyl 4-chloroacetoacetate **5J**, because transesterification reactions during the three-component

FULL PAPER Accepted Article

reaction are not disturbing. Interestingly, the reaction with **5I** stopped on the step of the Thorpe-Ziegler-type cyclization, whereas **5Ja-g** induced a cascade reaction which resulted in the formation of thienopyridinediones **6Ja-g** (Scheme 7). The latter proved to be poorly soluble in all NMR solvents tested. According to the ¹H NMR spectra, the compounds **6J** exist in the enol form and the corresponding proton gives singlets in the range from 5.58 to 5.76 ppm in the ¹H NMR spectra.



Scheme 7. Formation of thiophenes **6** by three-component reactions.

The structure of compound **6Ja** was also proven by a single crystal X-ray analysis the molecular drawing of which is shown in Figure 2. Single crystals of 2-phenylthieno[3,2-*b*]pyridine-5,7(4H,6H)-dione **6Ja** were obtained by diffusion of methanol into a saturated solution of the compound in DMF.




Figure 2. View of the molecular structure derived from the single-crystal X-ray analysis (displacement parameters are drawn at 50% probability level).

The compound **6Ja** crystallized in the monoclinic space group *P*2₁/*n* with one pyridinedione molecule and one DMF molecule in the asymmetric unit. The molecule of **6Ja** is almost planar, the

dihedral angle between the phenyl ring and the thiophene core is 10.8(2)^o. The distance between two adjacent parallel molecules of **6Ja** is 359.7 pm, which corresponds to the distance of $\pi\cdots\pi$ stacking interactions. The hydroxyl hydrogen atom is involved in O-H \cdots O hydrogen bonding with the carbonyl oxygen atom of the neighboring molecule, connecting them into an infinite zigzag H-bonded chain (Fig S2). In contrast, the N-bonded hydrogen atom of the pyridinedione core forms an N-H \cdots O hydrogen bond with the DMF molecule. The distance C5–C6 of 138.1(2) pm indicates a considerable double bond character and proves the enol form in solid-state.

Conclusions

The reported method of the bromoarylation of 2-chloroacrylonitrile by Meerwein reaction for the synthesis of the hitherto undescribed title compounds, 2-bromo-2-chloro-3-arylpropanenitriles, is reliable and affords the desired products in good yields under mild conditions. These propanenitriles are very stable compounds and proved to be useful in two- or three-component syntheses of functionalized thiophenes which are otherwise difficult or even impossible to prepare.

Experimental Section

Melting points are uncorrected and were determined in an apparatus according to Dr. Tottoli (Büchi). The ATR-IR spectra were obtained on a Bruker Alpha in the range of 400 to 4000 cm⁻¹. ¹H and ¹³C NMR spectra were recorded on a Bruker 170 Avance 500 spectrometer (at 500 MHz for ¹H NMR, 126 MHz for ¹³C NMR), a Bruker Avance III 600 MHz spectrometer (at 600 MHz for ¹H NMR, 151 MHz for ¹³C NMR), and a Bruker Avance 400 MHz spectrometer (at 400 MHz for ¹H NMR, 101 MHz for ¹³C NMR). Multiplicities are described by using the following abbreviations: s = singlet, d = doublet, t = triplet, q = quartet, and m = multiplet. Signal orientations in DEPT experiments were described as follows: o = no signal; + = up (CH, CH₃); - = down (CH₂). The mass spectra were measured with a Varian 320 MS Triple Quad GC/MS/MS (EIMS) or with an Agilent LCMSD series HP 1100 with APIES at fragmentor voltages as indicated. Elemental analysis was performed on a Carlo Erba 1106 analyzer.

Crystal structure determination of **6Ja**

Single-crystal data for **6Ja** were collected on an Agilent Gemini+ four-circle diffractometer equipped with an Atlas CCD detector, using graphite monochromatized MoK α radiation. Data were treated using the CrysAlis PRO program.^[51] The structure was solved using SHELXS and refined by least squares method on F^2 by SHELXL with the following graphical user interfaces of OLEX².^[52] N-bonded and O-bonded H atoms in **6Ja** were derived from the Fourier difference syntheses and refined in an isotropic mode. The other hydrogen atoms were placed on geometrically calculated positions and refined as riding atoms with relative isotropic displacement parameters.

6Ja: yellow needles, C₁₃H₉NO₂S·C₃H₇NO, M_r = 316.37, crystal size 0.15 × 0.41 × 0.44 mm, monoclinic, space group *P*2₁/*n* (no. 14), temperature 150(2) K, wavelength 0.71073 Å, *a* = 7.3297(3) Å, *b* = 8.8005(3) Å, *c* = 23.6124(11) Å, β = 95.155(4)^o, *V* = 1516.96(11) Å³, *Z* = 4, ρ = 1.385 g/cm³,

FULL PAPER Accepted Article

absorption coeff. = 0.227 mm⁻¹, $F(000)$ = 664, θ range for data collection 2.47–28.80°, limiting indices $-9 \leq h \leq 9$, $-11 \leq k \leq 11$, $-26 \leq l \leq 31$, refinement method: full-matrix least-squares on F^2 , measured reflections = 13943, used in refinement = 3400, free parameters = 209, goodness-of-fit on F^2 = 1.060, $R(F^2 > 2\sigma(F^2))$ = 0.0405, $wR(F^2)$ = 0.0961, largest diff. peak / hole = 0.28 / -0.34 e Å⁻³.

CCDC 1957206 (**6Ja**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

General Procedure of the Meerwein Arylation: The corresponding aniline **1**, (0.10 mol) was added to 40% aqueous HBr (34.3 mL, 0.22 mol). The mixture was cooled to –5°C, and a saturated solution of 7.6 g (0.11 mol) of sodium nitrite was added dropwise at such a rate that the temperature did not exceed 5°C. A cold solution of freshly-prepared diazonium salt **2** was added dropwise under stirring to a mixture of CuBr₂ (0.100 g, 0.448 mmol), 2-chloroacrylonitrile (0.8 mL, 0.100 mol), and acetone (50 mL). The rate of the addition was selected so that nitrogen evolved at a rate of 2–3 bubbles per second (addition time 0.5–1 h). The mixture was then stirred until nitrogen no longer evolved, 200 mL of water was added, the organic layer was separated, and the aqueous layer was extracted with DCM (3×15 mL). The combined extract were dried over MgSO₄, evaporated, and the residue was distilled under reduced pressure.

2-Bromo-2-chloro-3-phenylpropanenitrile (3a): Yield 9.780 g, 40%, light yellow oil, b.p. 116–120 °C/2 torr. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.50–7.46 (m, 2H), 7.46–7.41 (m, 3H), 4.06 (d, J = 14.2 Hz, 1H, CH₂), 4.00 (d, J = 14.2 Hz, 1H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 133.2, 131.1, 128.7, 128.5, 116.3, 51.7, 51.2 ppm. IR (ATR): 3033, 2930, 1757, 1741, 1515, 1427, 1258, 1202, 1081, 1023, 1011, 958, 882, 812, 771, 726, 699, 627, 602, 577, 513, 496, 479, 425 cm⁻¹. MS (ESI): m/z = 244.0 [M+H]⁺. Elemental analysis calcd for C₉H₇BrClN: C 44.21, H 2.89, N 5.73, found: C 44.03, H 2.81, N 5.60.

2-Bromo-2-chloro-3-(3-methylphenyl)propanenitrile (3b): Yield 14.993 g, 58%, a light yellow oil, b.p. 119–120 °C/2 torr, m.p. 16 °C. ¹H NMR (500 MHz, CDCl₃): δ = 7.35–7.30 (m, 1H, C₆H₄), 7.27–7.23 (m, 3H, C₆H₄), 3.82 (d, J = 14.3 Hz, 1H, CH₂), 3.77 (d, J = 14.3 Hz, 1H, CH₂), 2.42 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 138.4, 132.2, 131.6, 129.7, 128.6, 128.0, 115.8, 54.0, 51.4, 21.4 ppm. IR (ATR): 3028, 2920, 2219, 1608, 1429, 1331, 1169, 1131, 803, 759, 718, 697, 638, 441 cm⁻¹. MS (ESI): m/z = 258.0 [M+H]⁺. Elemental analysis calcd for C₁₀H₉BrClN: C 46.46, H 3.51, N 5.42, found: C 46.32; H 3.48; N 5.50.

2-Bromo-2-chloro-3-(4-methylphenyl)propanenitrile (3c): Yield 11.891 g, 46%, white crystals, b.p. 123–125 °C/2 torr, m.p. 49 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.34 (d, J = 7.8, 2H, C₆H₄), 7.21 (d, J = 7.8, 2H, C₆H₄), 3.97 (d, J = 14.3, 1H, CH₂), 3.92 (d, J = 14.3, 1H, CH₂), 2.31 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 138.0, 131.0, 130.2, 129.1, 116.4, 51.5, 51.3, 20.9 ppm. IR (ATR): 3029, 2932, 2923, 1915, 1708, 1612, 1515, 1427, 1318, 1223, 1113, 1070, 1023, 958, 884, 812, 770, 726, 706, 627, 602, 513, 496, 478, 425 cm⁻¹. MS (ESI): m/z = 258.0 [M+H]⁺. Elemental analysis calcd for C₁₀H₉BrClN: C 46.46, H 3.51, N 5.42, found: C 46.35, H 3.44, N 5.39.

2-Bromo-2-chloro-3-(4-fluorophenyl)propanenitrile (3d): Yield 18.637 g, 71%, white crystals, b.p. 105–107 °C/3 torr, m.p. 29 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.51 (dd, J_{HH} = 8.3, J_{HF} = 5.7 Hz, 2H C₆H₄), 7.24 (t, J = 8.8 Hz, 2H, C₆H₄), 4.04 (d, J = 14.3 Hz, 1H, CH₂), 3.99 (d, J = 14.3 Hz, 1H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 162.4 (d, J = 245.4 Hz), 133.3 (d, J = 8.4 Hz), 129.5 (d, J = 3.0 Hz), 116.2, 115.4 (d, J = 21.5 Hz), 51.2 (d, J = 2.1 Hz), 50.6 ppm. IR (ATR): 2219, 1600, 1508, 1430, 1419, 1225, 1161, 1101, 1014, 967, 885, 833, 786, 706, 624, 602, 583, 538, 519, 498, 486, 433 cm⁻¹. MS (ESI): m/z = 261.9 [M+H]⁺. Elemental analysis calcd for C₉H₇BrClFN: C 41.18, H 2.30, N 5.34, found: C, 41.09, H 2.22, N 5.29.

2-Bromo-2-chloro-3-(3-(trifluoromethyl)phenyl)propanenitrile (3e):

Yield 24.375 g, 78%, light yellow oil, b.p. 106–107 °C/2 torr. ¹H NMR (500 MHz, CDCl₃): δ = 7.76–7.68 (m, 2H, C₆H₄), 7.66 (d, J = 7.5 Hz, 1H, C₆H₄), 7.60–7.54 (m, 1H, C₆H₄), 3.91 (d, J = 14.4 Hz, 1H, CH₂), 3.87 (d, J = 14.4 Hz, 1H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 135.2, 134.4, 129.4, 129.1 (d, J = 31.8 Hz), 127.7 (q, J = 3.8 Hz), 125.3 (q, J = 3.7 Hz), 124.0 (q, J = 272.3 Hz), 116.1, 50.4 ppm. IR (ATR): 2219, 1725, 1600, 1510, 1452, 1432, 1333, 1312, 1201, 1165, 1122, 1101, 1075, 1039, 970, 912, 810, 768, 701, 660, 585, 421 cm⁻¹. MS (ESI): m/z = 311.9 [M+H]⁺. Elemental analysis calcd for C₁₀H₆BrClF₃N: C 38.43, H 1.94, N 4.48, found: C 38.40, H 1.87, N 4.39.

3-(3-Acetylphenyl)-2-bromo-2-chloropropanenitrile (3f): Yield 19.768 g, 69%, white crystals, b.p. 170–172 °C/2 torr, m.p. 48 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.08 (s, 1H, C₆H₄), 8.01 (d, 1H, J = 7.5 Hz, C₆H₄), 7.73 (d, J = 7.4 Hz, 1H, C₆H₄), 7.61–7.56 (m, 1H, C₆H₄), 4.16 (d, J = 14.3 Hz, 1H, CH₂), 4.10 (d, J = 14.3 Hz, 1H, CH₂), 2.59 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.6, 136.9, 135.8, 133.8, 130.7, 128.9, 128.7, 116.3, 50.9, 50.8, 26.8 ppm. IR (ATR): 3343, 3041, 2973, 2935, 1680, 1600, 1429, 1358, 1268, 1190, 1098, 1070, 1021, 954, 881, 806, 755, 695, 653, 587, 495, 465, 420 cm⁻¹. MS (ESI): m/z = 286.0 [M+H]⁺. Elemental analysis calcd for C₁₁H₉BrCINO: C 46.11, H 3.17, N 4.89, found: C 45.98, H 3.08, N 4.75.

Methyl 3-(2-bromo-2-chloro-2-cyanoethyl)benzoate (3g): Yield 17.847 g, 69%, white crystals, b.p. 167–169 °C/2 torr, m.p. 76 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.10 (s, 1H, C₆H₄), 8.00 (d, J = 7.8 Hz, 1H, C₆H₄), 7.75 (d, J = 7.6 Hz, 1H, C₆H₄), 7.58 (dd, J = 4.6, 10.8 Hz, 1H, C₆H₄), 4.17 (d, J = 14.1 Hz, 1H, CH₂), 4.12 (d, J = 14.1 Hz, 1H, CH₂), 3.87 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.9*, 136.1, 136.0*, 135.9, 134.9, 133.9*, 133.0, 132.0, 131.9*, 131.7, 129.89, 129.83*, 129.77, 129.5, 129.4*, 129.3, 129.01, 128.96*, 128.90, 116.9, 116.2*, 115.4, 52.34*, 51.25, 50.77*, 50.74*, 50.1 ppm. IR (ATR): 3029, 2219, 1722, 1608, 1437, 1425, 1203, 1114, 800, 625, 495, 220 cm⁻¹. MS (ESI): m/z = 302.0 [M+H]⁺. Elemental analysis calcd for C₁₁H₉BrCINO₂: C 43.67, H 3.00, N 4.63, found: C 43.56, H 2.97, N 4.50. * - most stable rotamer. Signals of three rotamers are present in ¹³C NMR spectra of this compound.

General Procedure for the Preparation of the 2-chloro-3-arylprop-2-enenitriles 4: To a solution of propanenitrile **3** (50 mmol, 1 eq.) in toluene (30 mL) NEt₃ (7.63 mL, 55 mmol, 1.1 eq.) was added dropwise under stirring. The resulting mixture was stirred for 1 h under reflux. The mixture was then cooled to r.t. and water (30 mL) was added, the organic layer was separated, dried over MgSO₄, and evaporated. The residue was distilled under reduced pressure.

2-Chloro-3-phenylprop-2-enenitrile (4a): Yield 7.525 g, 92%, light yellow oil, b.p. 106–110 °C/2 torr. ¹H NMR (500 MHz, CDCl₃): δ = 7.78–7.71 (m, 2H), 7.47–7.45 (m, 3H), 7.34 (s, 1H) ppm. ¹³C NMR (126 MHz, CDCl₃): δ = 142.4, 131.9, 131.4, 130.5, 129.0, 116.5, 101.6 ppm. IR (ATR): 2933, 2219, 1452, 1333, 1201, 1165, 1124, 1075, 810, 768, 701, 660 cm⁻¹. Spectroscopic data are in agreement with those reported in the literature.^[41]

2-Chloro-3-(3-methylphenyl)prop-2-enenitrile (4b): Yield 8.347 g, 94%, a light yellow oil, b.p. 96–99 °C/2 torr. ¹H NMR (500 MHz, CDCl₃): δ = 7.64–7.47 (m, 2H), 7.39–7.25 (m, 3H), 2.42 (s, 3H, CH₃) ppm. ¹³C NMR

FULL PAPER Accepted Article

(126 MHz, CDCl₃): δ = 145.6, 142.6, 139.1, 132.2, 129.4, 129.1, 125.9, 112.4, 100.0, 21.4 ppm. IR (ATR): 3029, 2922, 2861, 2219, 1608, 1484, 1451, 1429, 1380, 1331, 1285, 1245, 1168, 1130, 1074, 967, 921, 888, 785, 761, 690, 601, 548, 441 cm⁻¹. MS (ESI): *m/z* = 178.0 [M+H]⁺. Elemental analysis calcd for C₁₀H₈CIN: C 67.62, H 4.54, N 7.89, found: C 67.49, H 4.45, N 7.78.

2-Chloro-3-(4-methylphenyl)prop-2-enenitrile (4c): Yield 8.347 g, 94%, a light yellow oil, b.p. 96–99 °C/2 torr. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.93 (s, 0.3H, CH), 7.87 (s, 0.7H, CH), 7.74 (d, *J* = 8.1 Hz, 0.6H, C₆H₄), 7.64 (d, *J* = 8.1 Hz, 1.4H, C₆H₄), 7.32 (d, *J* = 7.9 Hz, 2H, C₆H₄), 2.35 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 146.4, 144.3, 143.3, 142.0, 141.8, 130.6, 129.9, 129.6, 128.7, 116.8, 115.5, 98.3, 97.4, 21.3, 21.2 ppm. Spectroscopic data are in agreement with those reported in the literature.^[53]

2-Chloro-3-(4-fluorophenyl)prop-2-enenitrile (4d): Yield 8.621 g, 95%, white crystals, b.p. 84–87 °C/2 torr, m.p. 28 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.00 (s, 0.4H, CH), 7.94 (s, 0.6H, CH), 7.90 (dd, *J*_{HF} = 5.6 Hz, *J*_{HF} = 8.6 Hz, 0.8H, C₆H₄), 7.81 (dd, *J*_{HF} = 5.5 Hz, *J*_{HF} = 8.5 Hz, 1.2H, C₆H₄), 7.39–7.34 (m, 2H, C₆H₄) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.5 (d, *J* = 251.0 Hz), 163.4 (d, *J* = 251.8 Hz), 145.2, 142.2, 133.1 (d, *J* = 9.1 Hz), 131.2 (d, *J* = 9.0 Hz), 128.3 (d, *J* = 3.3 Hz), 128.0 (d, *J* = 3.3 Hz), 116.4 (d, *J* = 22.1 Hz), 116.2 (d, *J* = 22.0 Hz), 115.4, 115.2, 99.0 (d, *J* = 2.3 Hz), 98.4 (d, *J* = 2.4 Hz) ppm. IR (ATR): 3077, 3031, 2220, 1599, 1507, 1415, 1287, 1237, 1162, 1111, 1012, 968, 830, 788, 686, 623, 583, 566, 538, 515, 414 cm⁻¹. MS (ESI): *m/z* = 182.0 [M+H]⁺. Elemental analysis calcd for C₉H₅CIFN: C 59.53, H 2.78, N 7.71, found: C 59.42, H 2.71, N 7.63. To the best of our knowledge, no spectroscopic data have been reported.

2-Chloro-3-(3-(trifluoromethyl)phenyl)prop-2-enenitrile (4e): Yield 8.621 g, 95%, a light yellow oil, b.p. 106–108 °C/5 torr. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.17 – 8.02 (m, 3H, CH, C₆H₄), 7.89 (d, *J* = 7.7 Hz, 1H, C₆H₄), 7.79–7.75 (m, 1H, C₆H₄) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 146.5, 143.5, 137.1, 131.5, 131.0 (q, *J* = 32.6 Hz), 126.0 (q, *J* = 3.6 Hz), 125.0 (q, *J* = 272.8 Hz), 122.6 (q, *J* = 4.2 Hz), 115.3, 98.6 ppm. IR (ATR): 3032, 2924, 2219, 1609, 1430, 1331, 1202, 1168, 1128, 1074, 909, 802, 761, 692, 659, 601, 442 cm⁻¹. MS (ESI): *m/z* = 232.0 [M+H]⁺. Elemental analysis calcd for C₁₀H₅CIF₃N: C 51.86, H 2.18, N 6.05, found: C 51.64, H 2.11, N 5.97. To the best of our knowledge, no spectroscopic data have been reported.

3-(3-Acetylphenyl)-2-chloroprop-2-enenitrile (4f): Yield 8.621 g, 95%, white crystals, b.p. 141–146 °C/2 torr, m.p. 43–44 °C. ¹H NMR (400 MHz, [D₆]DMSO), (two isomers 3:7): δ = 8.37 (s, 0.3H, CH), 8.33 (s, 0.7H, CH), 8.13 (s, 0.3H, C₆H₄), 8.11–8.05 (m, 2H, C₆H₄), 7.99 (d, *J* = 7.7 Hz, 0.7H, C₆H₄), 7.72–7.67 (m, 1H, C₆H₄), 2.62 (s, 1.2H, CH₃), 2.61 (s, 1.8H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.3, 197.2, 149.1, 145.7, 142.8, 137.3, 137.1, 134.3, 132.5, 131.8, 130.8, 130.7, 129.8, 129.7, 129.4, 128.3, 116.2, 115.0, 100.5, 99.8, 26.7 ppm. IR (ATR): 3034, 2217, 1677, 1594, 1577, 1493, 1425, 1357, 1274, 1199, 1069, 1013, 958, 939, 917, 891, 794, 712, 677, 589, 551, 521, 489 cm⁻¹. MS (ESI): *m/z* = 206.0 [M+H]⁺. Elemental analysis calcd for C₁₁H₈CINO: C 64.25, H 3.92, N 6.81, found: C 64.08, H 3.84, N 6.77.

Methyl 3-(2-chloro-2-cyanoethyl)benzoate (4g): Yield 8.621 g, 95%, white crystals, b.p. 146–150 °C/2 torr, m.p. 74 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.18 (s, 1H), 8.02 (d, *J* = 7.5 Hz, 1H), 7.97 (d, *J* = 7.1 Hz, 1H), 7.68 (t, *J* = 7.5 Hz, 1H), 3.88 (s, 3H) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.7, 147.9, 142.6, 136.8, 133.4, 130.5, 129.8, 129.0, 116.3, 101.0, 26.7 ppm. IR (ATR): 3075, 3061, 3043, 2960, 2214, 1722, 1607, 1579, 1437, 1424, 1305, 1280, 1205, 1115, 1087, 1067, 1042, 1012, 799, 747, 676, 604, 589, 550, 495 cm⁻¹. MS (ESI): *m/z* = 222.0 [M+H]⁺.

Elemental analysis calcd for C₁₁H₈CINO₂: C 59.61, H 3.64, N 6.32, found: C 59.44, H 3.54, N 6.20.

General method for preparation of methyl 3-amino-5-arylthiophene-2-carboxylates (6Aa–Ag), 3-amino-N-R-5-arylthiophene-2-carboxamides (6Ba–Eg), and 1-(3-amino-5-arylthiophen-2-yl)ethanones (6Fa–Fg). A solution of propanenitrile 3 (3 mmol) in methanol (5 mL) was added dropwise under stirring to a mixture of appropriate thiol (5A–F) (3.02 mmol) with a solution of MeONa (9 mL, 1M) in methanol at ambient temperature. The resulting mixture was stirred under reflux over a 1 h. The solvent was evaporated under reduced pressure. Acetic acid (20 mL, ω = 2%) was added to residue. The product was filtered off and recrystallized from (methanol or methanol-DMF). In the case of 6B–F the reaction was performed at rt.

Methyl 3-amino-5-phenylthiophene-2-carboxylate (6Aa): Yield 0.524 g, 75%, white crystals, m.p. 99–100 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.62–7.65 (m, 2H, C₆H₅), 7.38–7.48 (m, 3H, C₆H₅), 7.00 (s, 1H, thiophene), 4.29 (br.s, 2H, NH₂), 3.74 (s, 3H, CH₃) ppm. Spectroscopic data are in agreement with those reported in the literature.^[54]

Methyl 3-amino-5-(3-methylphenyl)thiophene-2-carboxylate (6Ab): Yield 0.592 g, 80%, white crystals, m.p. 83–84 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.48–7.37 (m, 2H, C₆H₄), 7.31 (t, *J* = 7.5 Hz, 1H, C₆H₄), 7.20 (d, *J* = 7.2 Hz, 1H, C₆H₄), 6.97 (s, 1H, thiophene), 6.60 (s, 2H, NH₂), 3.72 (s, 3H, OCH₃), 2.33 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.0, 155.4, 147.8, 138.6, 132.7, 129.9, 129.2, 126.1, 122.8, 116.2, 96.8, 51.0, 21.0 ppm. Spectroscopic data are in agreement with those reported in the literature.^[55]

Methyl 3-amino-5-(4-methylphenyl)thiophene-2-carboxylate (6Ac): Yield 0.526 g, 71%, white crystals, m.p. 141–142 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.51 (d, *J* = 8.2 Hz, 2H, C₆H₄), 7.24 (d, *J* = 7.9 Hz, 2H, C₆H₄), 6.93 (s, 1H, thiophene), 6.58 (s, 2H, NH₂), 3.73 (s, 3H, OCH₃), 2.32 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 163.9, 155.6, 147.7, 138.8, 130.0, 129.7, 125.4, 115.6, 96.1, 50.9, 20.8 ppm. IR (ATR): 3491, 3372, 2951, 1660, 1589, 1549, 1514, 1461, 1291, 1276, 1188, 1115, 1083, 1012, 961, 841, 804, 782, 764, 714, 655, 587, 471, 437 cm⁻¹. Spectroscopic data are in agreement with those reported in the literature.^[56]

Methyl 3-amino-5-(4-fluorophenyl)thiophene-2-carboxylate (6Ad): Yield 0.633 g, 84%, white crystals, m.p. 159–160 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.67 (dd, *J*_{HH} = 6.5 Hz, *J*_{HF} = 5.5 Hz, 2H, C₆H₄), 7.27 (dd, *J*_{HH} = 8.4 Hz, *J*_{HF} = 7.4 Hz, 2H, C₆H₄), 6.94 (s, 1H, thiophene), 6.61 (s, 2H, NH₂), 3.73 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.0, 162.6 (d, *J* = 247.0 Hz), 155.6, 146.5, 129.5 (d, *J* = 3.1 Hz), 127.9 (d, *J* = 8.4 Hz), 116.4, 116.2 (d, *J* = 22.0 Hz), 96.7, 51.0 ppm. Spectroscopic data are in agreement with those reported in the literature.^[27]

Methyl 3-amino-5-(3-(trifluoromethyl)phenyl)thiophene-2-carboxylate (6Ae): Yield 0.740 g, 82%, white crystals, m.p. 95–96 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.92 (d, *J* = 7.7 Hz, 1H, C₆H₄), 7.89 (s, 1H, C₆H₄), 7.74 (d, *J* = 7.7 Hz, 1H, C₆H₄), 7.67 (t, *J* = 7.7 Hz, 1H, C₆H₄), 7.13 (s, 1H, thiophene), 6.62 (s, 2H, NH₂), 3.74 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.9, 155.4, 145.4, 133.9, 130.5, 130.0 (q, *J* = 31.8 Hz), 129.7, 125.6 (q, *J* = 3.3 Hz), 123.9 (q, *J* = 272.6 Hz), 121.9 (q, *J* = 3.7 Hz), 117.7, 97.6, 51.1 ppm. Spectroscopic data are in agreement with those reported in the literature.^[56]

Methyl 5-(3-acetylphenyl)-3-aminothiophene-2-carboxylate (6Af): Yield 0.709 g, 86%, white crystals, m.p. 96–97 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.08 (s, 1H, C₆H₄), 7.94 (d, *J* = 7.4 Hz, 1H, C₆H₄), 7.85 (d,

FULL PAPER Accepted Article

$J = 7.0$ Hz, 1H, C₆H₄), 7.60–7.54 (m, 1H, C₆H₄), 7.11 (s, 1H, thiophene), 6.63 (s, 2H, NH₂), 3.72 (s, 3H, OCH₃), 2.61 (s, 3H, C(O)CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 197.6, 163.9, 155.2, 146.4, 137.5, 133.2, 130.0, 129.8, 128.8, 124.8, 117.2, 97.5, 51.1, 26.9$ ppm. IR (ATR): 3418, 3344, 1710, 1642, 1582, 1518, 1449, 1324, 1301, 1274, 1184, 1104, 1080, 1034, 834, 814, 753, 731, 681, 644, 510 cm⁻¹. MS (ESI): $m/z = 276.1$ [M+H]⁺. Elemental analysis calcd for C₁₄H₁₃NO₃S: C 61.07, H 4.76, N 5.09, found: C 60.99, H 4.65, N 4.94.

Methyl 3-amino-5-(3-(methoxycarbonyl)phenyl)thiophene-2-carboxylate (6Ag): Yield 0.715 g, 82%, off-white crystals, m.p. 124–126 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 8.11$ (d, $J = 1.3$ Hz, 1H, 2H-C₆H₄), 7.95 (dd, $J = 7.7, 1.1$ Hz, 1H, 6H-C₆H₄), 7.93–7.87 (m, 1H, C₆H₄), 7.61 (t, $J = 7.8$ Hz, 1H, C₆H₄), 7.09 (s, 1H, thiophene), 5.82–5.34 (br.s, 2H, NH₂), 3.88 (s, 3H, CH₃), 3.74 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 165.7, 163.9, 155.8, 146.0, 133.3, 130.6, 130.2, 130.0, 129.6, 125.8, 117.1, 98.5, 52.5, 51.1$ ppm. IR (ATR): 3433, 1670, 1634, 1590, 1550, 1515, 1463, 1428, 1365, 1313, 1268, 1235, 1075, 948, 900, 832, 782, 748, 690, 677, 590, 492, 434 cm⁻¹. MS (ESI): $m/z = 292.1$ [M+H]⁺. Elemental analysis calcd for C₁₄H₁₃NO₄S: C 57.72, H 4.50, N 4.81, found: C 57.59, H 4.43, N 4.74.

3-Amino-N,5-diphenylthiophene-2-carboxamide (6Ba): Yield 0.714 g, 81%, white crystals, m.p. 203–204 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.33$ (s, 1H, NH), 7.68 (d, $J = 8.4$ Hz, 2H, Ph), 7.65 (d, $J = 8.1$ Hz, 2H, Ph), 7.47 (t, $J = 7.2$ Hz, 2H, Ph), 7.42–7.37 (m, 1H, Ph), 7.29 (t, $J = 7.3$ Hz, 2H, Ph), 7.06–7.01 (m, 2H, Ph, thiophene), 6.82–6.54 (br.s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.2, 155.1, 144.5, 139.4, 133.0, 129.4, 129.0, 128.4, 125.4, 123.0, 120.7, 116.9, 100.2$ ppm. IR (ATR): 3473, 3364, 3250, 3054, 1616, 1585, 1514, 1489, 1464, 1432, 1320, 1307, 1274, 1233, 1097, 1072, 907, 824, 747, 684, 636, 510 cm⁻¹. MS (ESI): $m/z = 295.1$ [M+H]⁺. Elemental analysis calcd for C₁₇H₁₄N₂OS: C 69.36, H 4.79, N 9.52, found: C 69.28, H 4.70, N 9.40.

3-Amino-5-(3-methylphenyl)-N-phenylthiophene-2-carboxamide (6Bb): Yield 0.803 g, 87%, light yellow crystals, m.p. 171–172 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.30$ (s, 1H, NH), 7.68 (d, $J = 7.7$ Hz, 2H, Ar), 7.48–7.42 (m, 2H, Ar), 7.35 (t, $J = 7.5$ Hz, 1H, Ar), 7.29 (t, $J = 7.8$ Hz, 2H, Ar), 7.21 (d, $J = 7.6$ Hz, 1H, Ar), 7.03 (t, $J = 7.3$ Hz, 1H, Ar), 7.00 (s, 1H, thiophene), 6.68 (s, 2H, NH₂), 2.36 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.2, 155.1, 144.7, 139.4, 138.6, 133.0, 129.6, 129.2, 128.4, 126.0, 123.0, 122.6, 120.7, 116.8, 100.0, 21.0$ ppm. IR (ATR): 3471, 3362, 3264, 1622, 1581, 1533, 1499, 1456, 1434, 1316, 1267, 1234, 1176, 1093, 1072, 903, 872, 826, 781, 748, 722, 686, 636, 616, 543, 513, 489, 432 cm⁻¹. MS (ESI): $m/z = 309.1$ [M+H]⁺. Elemental analysis calcd for C₁₈H₁₆N₂OS: C 70.10, H 5.23, N 9.08, found: C 70.00, H 5.14, N 8.97.

3-Amino-5-(4-methylphenyl)-N-phenylthiophene-2-carboxamide (6Bc): Yield 0.720 g, 78%, light yellow crystals, m.p. 205–206 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.29$ (s, 1H, NH), 7.68 (d, $J = 7.6$ Hz, 2H, Ar), 7.54 (d, $J = 8.1$ Hz, 2H, Ar), 7.32–7.27 (m, 4H, Ar), 7.03 (t, $J = 7.4$ Hz, 1H, Ar), 6.97 (s, 1H, thiophene), 6.68 (s, 2H, NH₂), 2.33 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.2, 155.2, 144.7, 139.4, 138.6, 130.3, 129.9, 128.4, 125.4, 123.0, 120.7, 116.3, 99.7, 20.9$ ppm. IR (ATR): 3470, 3358, 1616, 1582, 1520, 1499, 1460, 1433, 1319, 1270, 1232, 1177, 1100, 1074, 1018, 811, 750, 692, 567, 541, 511 cm⁻¹. MS (ESI): $m/z = 309.1$ [M+H]⁺. Elemental analysis calcd for C₁₈H₁₆N₂OS: C 70.10, H 5.23, N 9.08, found: C 69.96, H 5.15, N 8.94.

3-Amino-5-(4-fluorophenyl)-N-phenylthiophene-2-carboxamide (6Bd): Yield 0.814 g, 87%, white crystals, m.p. 226–227 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.32$ (s, 1H, NH), 7.75–7.63 (m, 4H, Ar), 7.35–7.26 (m, 4H, Ar), 7.03 (t, $J = 7.3$ Hz, 1H, Ar), 6.98 (s, 1H, thiophene), 6.69 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.2, 162.4$ (d, $J =$

246.7 Hz), 155.1, 143.4, 139.3, 129.7 (d, $J = 2.9$ Hz), 128.4, 127.6 (d, $J = 8.2$ Hz), 123.0, 120.7, 117.1, 116.3 (d, $J = 21.9$ Hz), 100.2 ppm. IR (ATR): 3473, 3363, 3252, 3202, 1615, 1581, 1524, 1462, 1435, 1320, 1270, 1229, 1158, 1096, 1074, 1012, 911, 837, 821, 754, 694, 611, 576, 563, 541, 511, 495, 440 cm⁻¹. MS (ESI): $m/z = 313.1$ [M+H]⁺. Elemental analysis calcd for C₁₇H₁₃FN₂OS: C, 65.37, H 4.19, N, 8.97, found: C 65.40, H 4.07, N 8.90.

3-Amino-N-phenyl-5-(3-(trifluoromethyl)phenyl)thiophene-2-carboxamide (6Be):

Yield 0.944 g, 87%, cream-colored crystals, m.p. 167–168 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.38$ (s, 1H, NH), 7.96 (d, $J = 7.6$ Hz, 1H, Ar), 7.91 (s, 1H, Ar), 7.76 (d, $J = 7.6$ Hz, 1H, Ar), 7.72 (d, $J = 7.7$ Hz, 1H, Ar), 7.71–7.66 (m, 2H, Ar), 7.30 (t, $J = 7.9$ Hz, 2H, Ar), 7.18 (s, 1H, thiophene), 7.04 (t, $J = 7.3$ Hz, 1H, Ar), 6.71 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.1, 155.1, 142.3, 139.3, 134.1, 130.7, 130.1$ (q, $J = 32.5$ Hz), 129.4, 128.5, 125.3 (q, $J = 3.4$ Hz), 123.1, 121.8 (q, $J = 27.5$ Hz), 121.7 (q, $J = 3.6$ Hz), 120.7, 118.3, 101.0 ppm. IR (ATR): 3473, 1616, 1575, 1464, 1436, 1319, 1124, 803, 749, 725, 693, 676, 510, 428, 415 cm⁻¹. MS (ESI): $m/z = 363.1$ [M+H]⁺. Elemental analysis calcd for C₁₈H₁₃F₃N₂OS: C 59.66, H 3.62, N 7.73, found: C 59.49, H 3.54, N 7.68.

5-(3-Acetylphenyl)-3-amino-N-phenylthiophene-2-carboxamide (6Bf):

Yield 0.786 g, 78%, light yellow crystals, m.p. 187–188 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.37$ (s, 1H, NH), 8.16 (s, 1H, 2H-C₆H₄), 7.98 (d, $J = 7.7$ Hz, 1H, Ar), 7.92 (d, $J = 8.4$ Hz, 1H, Ar), 7.69 (d, $J = 7.7$ Hz, 2H, Ar), 7.63 (t, $J = 7.7$ Hz, 1H, Ar), 7.30 (t, $J = 7.9$ Hz, 2H, Ar), 7.14 (s, 1H, thiophene), 7.04 (t, $J = 7.3$ Hz, 1H, Ar), 6.72 (s, 2H, NH₂), 2.64 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 197.6, 163.1, 155.2, 143.3, 139.3, 137.6, 133.4, 129.9, 129.7, 128.7, 128.5, 124.6, 123.1, 120.7, 117.6, 100.5, 26.9$ ppm. IR (ATR): 3433, 3400, 3311, 1671, 1634, 1590, 1551, 1515, 1463, 1428, 1365, 1313, 1269, 1235, 1071, 948, 900, 830, 786, 748, 717, 690, 677, 592, 492, 437 cm⁻¹. MS (ESI): $m/z = 337.1$ [M+H]⁺. Elemental analysis calcd for C₁₉H₁₆N₂O₂S: C 67.84, H 4.79, N 8.33, found: C 67.73, H 4.70, N 8.25.

Methyl 3-(4-amino-5-(phenylcarbamoyl)thiophen-2-yl)benzoate (6Bg): Yield 0.792 g, 75%, light yellow crystals, m.p. 152–153 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.37$ (s, 1H, NH), 8.17 (s, 1H, 2H-C₆H₄), 8.00–7.91 (m, 2H, Ar), 7.69 (d, $J = 7.2$ Hz, 2H, Ar), 7.65–7.56 (m, 1H, Ar), 7.37–7.24 (m, 2H, Ar), 7.13 (s, 1H, thiophene), 7.07–6.99 (m, 1H, Ar), 6.72 (s, 2H, NH₂), 3.90 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 165.8, 163.1, 155.2, 142.9, 139.3, 133.5, 130.6, 130.0, 129.9, 129.3, 128.5, 125.7, 123.1, 120.7, 117.6, 100.5, 52.5$ ppm. IR (ATR): 3413, 3351, 3314, 1718, 1634, 1576, 1520, 1436, 1321, 1301, 1277, 1229, 1108, 1082, 1003, 959, 904, 819, 752, 688, 642, 548, 508, 469, 409 cm⁻¹. MS (ESI): $m/z = 353.1$ [M+H]⁺. Elemental analysis calcd for C₁₉H₁₆N₂O₃S: C 64.76, H 4.58, N 7.95, found: C 64.61, H 4.49, N 7.90.

3-Amino-N-(2-fluorophenyl)-5-phenylthiophene-2-carboxamide (6Ca):

Yield 0.692 g, 74%, light yellow crystals, m.p. 148–149 °C. ¹H NMR (500 MHz, [D₆]DMSO): $\delta = 9.17$ (s, 1H, NH), 7.65 (d, $J = 7.3$ Hz, 2H, Ar), 7.54 (t, $J = 7.5$ Hz, 1H, Ar), 7.47 (t, $J = 7.5$ Hz, 2H, Ar), 7.40 (t, $J = 7.2$ Hz, 1H, Ar), 7.28–7.13 (m, 3H, Ar), 7.02 (s, 1H, thiophene), 6.64 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): $\delta = 163.0, 156.0$ (d, $J = 246.6$ Hz), 154.9, 144.9, 133.0, 129.4, 129.0, 127.1, 126.4 (d, $J = 7.8$ Hz), 126.1 (d, $J = 12.0$ Hz), 125.5, 124.2 (d, $J = 3.5$ Hz), 117.0, 115.7 (d, $J = 20.2$ Hz), 100.2 ppm. IR (ATR): 3424, 3313, 1646, 1617, 1595, 1520, 1459, 1446, 1327, 1255, 1186, 1117, 1074, 1032, 936, 826, 766, 745, 726, 691, 534, 499, 457 cm⁻¹. MS (ESI): $m/z = 313.1$ [M+H]⁺. Elemental analysis calcd for C₁₇H₁₃FN₂OS: C 65.37, H 4.19, N 8.97, found: C 65.39, H 4.11, N, 8.88,

3-Amino-N-(2-fluorophenyl)-5-(3-methylphenyl)thiophene-2-carboxamide (6Cb):

Yield 0.596 g, 61%, yellow crystals, m.p. 131–132 °C. ¹H NMR (400 MHz, [D₆]DMSO): $\delta = 9.14$ (s, 1H, NH), 7.60–7.53 (m, 1H, Ar), 7.49–7.42 (m, 2H, Ar), 7.35 (t, $J = 7.6$ Hz, 1H, Ar), 7.29–7.15 (m, 4H,

FULL PAPER Accepted Article

Ar), 7.02 (s, 1H, thiophene), 6.64 (s, 2H, NH₂), 2.37 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 162.9 (o), 155.8 (d, J = 246.2 Hz, o), 154.9 (o), 145.0 (o), 138.5 (o), 132.9 (o), 129.6 (+), 129.1 (+), 126.9 (d, J = 1.6 Hz, +), 126.2 (d, J = 7.8 Hz, +), 126.0 (d, J = 14.3 Hz, o), 125.9 (+), 124.1 (d, J = 3.4 Hz, +), 122.6 (+), 116.7 (+), 115.6 (d, J = 20.0 Hz, +), 100.0 (o), 20.9 (+) ppm. IR (ATR): 3383, 3030, 1646, 1521, 1496, 1454, 1426, 1388, 1364, 1333, 1116, 996, 762, 725, 692, 623, 553, 506, 486, 449 cm⁻¹. MS (ESI): m/z = 327.1 [M+H]⁺. Elemental analysis calcd for C₁₈H₁₅FN₂OS: C 66.24, H 4.63, N 8.58, found: C 66.17, H 4.57, N 8.44.

3-Amino-N-(2-fluorophenyl)-5-(4-methylphenyl)thiophene-2-carboxamide (6Cc): Yield 0.655 g, 67%, light yellow crystals, m.p. 191–192 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 9.11 (s, 1H, NH), 7.53 (d, J = 8.1 Hz, 3H, Ar), 7.27 (d, J = 7.9 Hz, 2H, Ar), 7.25–7.12 (m, 3H, Ar), 6.97 (s, 1H, thiophene), 6.62 (s, 2H, NH₂), 2.33 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.0, 155.9 (d, J = 246.3 Hz), 155.0, 145.1, 138.7, 130.3, 129.9, 127.0, 126.3 (d, J = 7.5 Hz), 126.2 (d, J = 12.2 Hz), 125.4, 124.2 (d, J = 3.1 Hz), 116.4, 115.7 (d, J = 19.9 Hz), 99.7, 20.9 ppm. IR (ATR): 3419, 3312, 3024, 1646, 1616, 1592, 1515, 1445, 1321, 1267, 1185, 1108, 1032, 932, 840, 809, 775, 745, 718, 637, 531, 501, 449 cm⁻¹. MS (ESI): m/z = 327.1 [M+H]⁺. Elemental analysis calcd for C₁₈H₁₅FN₂OS: C 66.24, H 4.63, N 8.58, found: C 66.11, H 4.56, N 8.48.

3-Amino-N-(2-fluorophenyl)-5-(4-fluorophenyl)thiophene-2-carboxamide (6Cd): Yield 0.821 g, 83%, light yellow crystals, m.p. 174–175 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 9.16 (s, 1H, NH), 7.69 (dd, J_{HH}= 8.6, J_{HF}= 5.3 Hz, 2H, 4-FC₆H₄), 7.59–7.49 (m, 1H, 2-FC₆H₄), 7.31 (t, J = 8.8 Hz, 2H, 4-FC₆H₄), 7.26–7.14 (m, 3H, 2-FC₆H₄), 6.98 (s, 1H, thiophene), 6.64 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.0, 162.4 (d, J = 246.7 Hz), 156.0 (d, J = 245.1 Hz), 155.0, 143.7, 129.7 (d, J = 3.1 Hz), 127.7 (d, J = 8.4 Hz), 127.1, 126.4 (d, J = 7.8 Hz), 126.1 (d, J = 11.8 Hz), 117.1, 116.3 (d, J = 21.9 Hz), 115.7 (d, J = 20.0 Hz), 100.1 ppm. IR (ATR): 3487, 3370, 3236, 3212, 1626, 1583, 1511, 1450, 1315, 1274, 1257, 1193, 1156, 1097, 1032, 1011, 934, 836, 818, 749, 714, 630, 563, 533, 488, 461, 449, 434 cm⁻¹. MS (ESI): m/z = 331.1 [M+H]⁺. Elemental analysis calcd for C₁₇H₁₂F₂N₂OS: C 61.81, H 3.66, N 8.48, found: C 61.70, H 3.60, N 8.39.

3-Amino-N-(2-fluorophenyl)-5-(3-(trifluoromethyl)phenyl)thiophene-2-carboxamide (6Ce): Yield 0.912 g, 80%, yellow crystals, m.p. 167–168 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 9.25 (s, 1H, NH), 7.95 (d, J = 7.6 Hz, 1H, Ar), 7.92 (s, 1H, Ar), 7.80–7.69 (m, 2H, Ar), 7.57–7.51 (m, 1H, Ar), 7.29–7.15 (m, 4H, Ar, thiophene), 6.66 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 162.9, 156.0 (d, J = 246.2 Hz), 154.9, 142.6, 134.1, 130.7, 130.1 (q, J = 31.7 Hz), 129.5, 127.2, 126.5 (d, J = 8.0 Hz), 126.0 (d, J = 12.1 Hz), 125.4 (q, J = 3.5 Hz), 124.2 (d, J = 3.5 Hz), 124.0 (q, J = 272.7 Hz), 121.7 (q, J = 3.5 Hz), 118.3, 115.7 (d, J = 20.0 Hz), 100.9 ppm. IR (ATR): 3477, 3421, 3403, 3351, 1647, 1580, 1513, 1449, 1316, 1268, 1250, 1163, 1118, 1070, 1000, 971, 934, 886, 822, 803, 773, 751, 723, 692, 676, 640, 533, 490, 438 cm⁻¹. MS (ESI): m/z = 381.1 [M+H]⁺. Elemental analysis calcd for C₁₈H₁₂F₄N₂OS: C 56.84, H 3.18, N 7.36, found: C 56.77, H 3.10, N 7.39.

5-(3-Acetylphenyl)-3-amino-N-(2-fluorophenyl)thiophene-2-carboxamide (6Cf): Yield 0.902 g, 85%, yellow crystals, m.p. 174–175 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 9.23 (s, 1H, NH), 8.15 (s, 1H, Ar), 7.98 (d, J = 7.7 Hz, 1H, Ar), 7.91 (d, J = 7.6 Hz, 1H, Ar), 7.63 (t, J = 7.7 Hz, 1H, Ar), 7.58–7.50 (m, 1H, Ar), 7.31–7.16 (m, 3H, Ar), 7.14 (s, 1H, thiophene), 6.66 (s, 2H, NH₂), 2.64 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.6, 162.9, 156.0 (d, J = 246.6 Hz), 155.0, 143.7, 137.6, 133.4, 129.9 (d, J = 7.5 Hz), 128.7, 127.2, 126.4 (d, J = 7.1 Hz), 126.0 (d, J = 12.1 Hz), 124.7, 124.2, 124.2, 117.7, 115.7 (d, J = 20.0 Hz), 100.5, 26.9 ppm. IR (ATR): 3441, 3418, 3407, 3306, 1674, 1640, 1593, 1547, 1512, 1447, 1320, 1268, 1255, 1217, 1189, 1105, 1077, 1023, 904, 828, 786, 745, 716, 679,

645, 591, 488, 448 cm⁻¹. MS (ESI): m/z = 355.1 [M+H]⁺. Elemental analysis calcd for C₁₉H₁₅FN₂O₂S: C 64.39, H 4.27, N 7.90, found: C 64.41, H 4.19, N 7.82.

Methyl 3-(4-amino-5-((2-fluorophenyl)carbamoyl)thiophen-2-yl)benzoate (6Cg): Yield 0.832 g, 75%, light yellow crystals, m.p. 153–154 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 9.08 (s, 1H, NH), 8.22–8.09 (m, 1H, Ar), 8.01–7.87 (m, 2H, Ar), 7.69–7.54 (m, 2H, Ar), 7.29–7.15 (m, 3H, Ar), 7.13 (s, 1H, thiophene), 6.58 (s, 2H, NH₂), 3.91 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.8, 162.9, 156.0 (d, J = 246.7 Hz), 155.0, 143.3, 133.4, 130.6, 130.03, 129.97, 129.3, 127.2, 126.5 (d, J = 7.8 Hz), 126.0 (d, J = 11.3 Hz), 125.7, 124.2 (d, J = 3.1 Hz), 117.7, 115.7 (d, J = 20.2 Hz), 100.5, 52.5 ppm. IR (ATR): 3456, 3419, 3344, 2948, 1709, 1641, 1584, 1518, 1449, 1324, 1301, 1273, 1184, 1104, 1081, 1034, 834, 812, 753, 731, 681, 644, 512, 448 cm⁻¹. MS (ESI): m/z = 371.1 [M+H]⁺. Elemental analysis calcd for C₁₉H₁₅FN₂O₃S: C 61.61, H 4.08, N 7.56, found: C 61.52, H 4.01, N 7.44.

3-Amino-N-benzyl-5-phenylthiophene-2-carboxamide (6Da): Yield 0.600 g, 65%, white crystals, m.p. 135–136 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.11 (s, 1H, NH), 7.59 (d, J = 6.9 Hz, 2H, Ar), 7.44 (t, J = 7.0 Hz, 2H, Ar), 7.40–7.17 (m, 6H, Ar), 6.97 (s, 1H, thiophene), 6.50 (s, 2H, NH₂), 4.37 (d, J = 5.2 Hz, 2H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.4, 153.9, 143.5, 140.4, 133.2, 129.3, 128.7, 128.2, 127.2, 126.6, 125.4, 117.0, 100.5, 42.1 ppm. IR (ATR): 3405, 3282, 2923, 1619, 1585, 1528, 1436, 1364, 1318, 1296, 1242, 1171, 1092, 970, 837, 822, 787, 756, 743, 701, 685, 613, 560, 515, 473, 446 cm⁻¹. MS (ESI): m/z = 309.1 [M+H]⁺. Elemental analysis calcd for C₁₈H₁₆N₂OS: C 70.10, H 5.23, N 9.08, found: C 69.7, H 5.15, N 8.98.

3-Amino-N-benzyl-5-(3-methylphenyl)thiophene-2-carboxamide (6Db): Yield 0.685 g, 71%, light yellow crystals, m.p. 150–151 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.11 (s, 1H, NH), 7.55 (t, J = 6.9 Hz, 1H, Ar), 7.49–7.40 (m, 2H, Ar), 7.34 (t, J = 7.6 Hz, 1H, Ar), 7.26–7.16 (m, 5H, Ar), 7.00 (s, 1H, thiophene), 6.62 (s, 2H, NH₂), 4.37 (d, J = 5.6 Hz, 2H, CH₂), 2.36 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.0, 154.9, 145.5, 138.6, 133.0, 129.7, 129.2, 127.0, 126.3, 126.1, 124.2, 122.7, 116.8, 115.7, 100.1, 42.1, 21.0 ppm. IR (ATR): 3533, 3419, 3309, 1639, 1615, 1592, 1516, 1445, 1323, 1251, 1183, 1110, 1080, 1031, 930, 826, 786, 745, 708, 689, 648, 602, 525, 478, 448, 427 cm⁻¹. MS (ESI): m/z = 323.1 [M+H]⁺. Elemental analysis calcd for C₁₉H₁₈N₂OS: C 70.78, H 5.63, N 8.69, found: C 70.80, H 5.55, N 8.73.

3-Amino-N-benzyl-5-(4-methylphenyl)thiophene-2-carboxamide (6Dc): Yield 0.830 g, 86%, light yellow crystals, m.p. 154–155 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.06[†] (t, J = 5.2 Hz, 1H, NH), 7.48 (d, J = 7.5 Hz, 2H, Ar), 7.34–7.28 (br, s, 4H, Ar), 7.27–7.16 (m, 3H, Ar), 6.91 (s, 1H, thiophene), 6.48 (s, 2H, NH₂), 4.37 (d, J = 5.6 Hz, 2H, CH₂), 2.32 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.4, 153.9, 143.7, 140.4, 138.3, 130.5, 129.8, 128.2, 127.2, 126.6, 125.3, 116.5, 100.1, 42.1, 20.9 ppm. IR (ATR): 3473, 3295, 3028, 2913, 1606, 1589, 1520, 1450, 1310, 1240, 1155, 1097, 1028, 974, 872, 809, 715, 693, 512, 445 cm⁻¹. MS (ESI): m/z = 323.1 [M+H]⁺. Elemental analysis calcd for C₁₉H₁₈N₂OS: C 70.78, H 5.63, N 8.69, found: C 70.66, H 5.56, N 8.58.

3-Amino-N-benzyl-5-(4-fluorophenyl)thiophene-2-carboxamide (6Dd): Yield 0.743 g, 76%, cream-colored crystals, m.p. 171–172 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.10[†] (t, J = 5.7 Hz, 1H, NH), 7.63 (dd, J = 8.2, 5.2 Hz, 2H, C₆H₄), 7.37–7.08 (m, 7H, Ar), 6.92 (s, 1H, thiophene), 6.50 (s, 2H, NH₂), 4.37 (d, J = 5.7 Hz, 2H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.3, 162.2 (d, J = 243.5 Hz), 153.9, 142.3, 140.4, 129.9 (d, J = 3.6 Hz), 128.2, 127.5 (d, J = 8.4 Hz), 127.2, 126.6, 117.2, 116.2 (d, J = 22.1 Hz), 95.5, 42.1 ppm. IR (ATR): 3454, 3340, 3281, 1611, 1588, 1525, 1450, 1363, 1317, 1230, 1160, 1095, 1029, 1013, 977, 872, 841,

FULL PAPER Accepted Article

825, 716, 695, 563, 495, 448 cm⁻¹. MS (ESI): *m/z* = 327.1 [M+H]⁺. Elemental analysis calcd for C₁₈H₁₅FN₂OS: C 66.24, H 4.63, N 8.58, found: C 66.17, H 4.55, N 8.43.

3-Amino-N-benzyl-5-(3-(trifluoromethyl)phenyl)thiophene-2-carboxamide (6De): Yield 1.003 g, 89%, light yellow crystals, m.p. 148–149 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.19[†] (t, *J* = 5.9 Hz, 1H, NH), 7.90 (d, *J* = 7.7 Hz, 1H, Ar), 7.85 (s, 1H, Ar), 7.75–7.65 (m, 2H, Ar), 7.33–7.29 (br. s, 4H, Ar), 7.26–7.19 (m, 1H, Ar), 7.13 (s, 1H, thiophene), 6.52 (s, 2H, NH₂), 4.38 (d, *J* = 5.9 Hz, 2H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 164.2, 153.8, 141.3, 140.3, 134.2, 130.6, 130.0 (q, *J* = 31.2 Hz), 129.3, 128.2, 127.3, 126.6, 125.1 (q, *J* = 3.8 Hz), 121.9 (q, *J* = 276.7 Hz), 121.6 (q, *J* = 3.8 Hz), 118.4, 101.4, 42.2 ppm. IR (ATR): 3410, 3298, 2981, 1634, 1583, 1549, 1524, 1443, 1422, 1327, 1315, 1277, 1232, 1163, 1117, 1072, 1027, 980, 963, 894, 829, 800, 738, 725, 692, 675, 607, 518, 447 cm⁻¹. MS (ESI): *m/z* = 377.1 [M+H]⁺. Elemental analysis calcd for C₁₉H₁₅F₃N₂OS: C 60.63, H 4.02, N 7.44, found: C 60.50, H 3.94, N 7.32.

5-(3-Acetylphenyl)-3-amino-N-benzylthiophene-2-carboxamide (6Df): Yield 0.840 g, 80%, light yellow crystals, m.p. 186–187 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.18[†] (t, *J* = 5.9 Hz, 1H, NH), 8.10 (s, 1H, C₆H₄), 7.95 (d, *J* = 7.7 Hz, 1H, Ar), 7.86 (d, *J* = 7.8 Hz, 1H, Ar), 7.60 (t, *J* = 7.8 Hz, 1H, Ar), 7.36–7.26 (m, 4H, Ar), 7.25–7.19 (m, 1H, Ar), 7.09 (s, 1H, thiophene), 6.52 (s, 2H, NH₂), 4.38 (d, *J* = 5.9 Hz, 2H, CH₂), 2.63 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.6, 164.2, 153.9, 142.3, 140.3, 137.6, 133.6, 129.8, 129.7, 128.4, 128.2, 127.3, 126.6, 124.5, 117.8, 101.0, 42.17, 26.9 ppm. IR (ATR): 3450, 3329, 3266, 1681, 1593, 1531, 1428, 1359, 1315, 1271, 1220, 1175, 1105, 1081, 1029, 981, 948, 912, 872, 838, 792, 762, 725, 695, 684, 639, 589, 514, 461, 432 cm⁻¹. MS (ESI): *m/z* = 351.1 [M+H]⁺. Elemental analysis calcd for C₂₀H₁₈N₂O₂S: C 68.55, H 5.18, N 7.99, found: C 68.48, H 5.10, N 7.82.

Methyl 3-(4-amino-5-(benzylcarbamoyl)thiophen-2-yl)benzoate (6Dg): Yield 0.878 g, 80%, light yellow crystals, m.p. 141–142 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.20[†] (t, *J* = 6.0 Hz, 1H, NH), 8.12 (t, *J* = 1.6 Hz, 1H, Ar), 7.98–7.87 (m, 2H, Ar), 7.60 (t, *J* = 7.8 Hz, 1H, Ar), 7.35–7.28 (m, 4H, Ar), 7.27–7.18 (m, 1H, Ar), 7.08 (s, 1H, thiophene), 6.53 (s, 2H, NH₂), 4.38 (d, *J* = 6.0 Hz, 2H, CH₂), 3.88 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 166.1, 164.5, 154.2, 142.2, 140.6, 133.9, 130.9, 130.3, 130.1, 129.4, 128.5, 127.6, 126.9, 125.9, 118.1, 101.3, 52.8, 42.5 ppm. IR (ATR): 3452, 3340, 3262, 3025, 1719, 1596, 1580, 1527, 1427, 1312, 1291, 1277, 1229, 1174, 1114, 1106, 1021, 1010, 983, 911, 873, 834, 752, 722, 695, 682, 638, 611, 516, 461, 436 cm⁻¹. MS (ESI): *m/z* = 367.1 [M+H]⁺. Elemental analysis calcd for C₂₀H₁₈N₂O₃S: C 65.56, H 4.95, N 7.64, found: C 65.42, H 4.88, N 7.50.

[†] Mutual splitting of signals of NH and CH₂ groups is present in the ¹H NMR spectra of **6Dc–6Dg**.

2-(Morpholin-4-ylcarbonyl)-5-phenylthiophen-3-amine (6Ea): Yield 0.414 g, 48%, cream-colored crystals, m.p. 106–107 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.59 (d, *J* = 7.1 Hz, 2H, C₆H₅), 7.42 (t, *J* = 7.2 Hz, 2H, C₆H₅), 7.38–7.27 (m, 1H, C₆H₅), 6.95 (s, 1H, thiophene), 6.12 (s, 2H, NH₂), 3.65–3.61 (m, 4H, CH₂), 3.61–3.56 (m, 4H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.3, 152.4, 144.2, 133.0, 129.2, 128.7, 125.4, 116.9, 101.2, 66.3, 45.6 ppm. IR (ATR): 3419, 3308, 2959, 2892, 2843, 1575, 1533, 1456, 1444, 1392, 1303, 1255, 1214, 1149, 1112, 1073, 1014, 934, 911, 879, 849, 824, 756, 724, 687, 584, 515, 451 cm⁻¹. MS (ESI): *m/z* = 289.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₆N₂O₂S: C 62.48, H 5.59, N 9.71, found: C 62.37, H 5.47, N 9.66.

5-(3-Methylphenyl)-2-(morpholin-4-ylcarbonyl)thiophen-3-amine (6Eb): Yield 0.724 g, 80%, light yellow crystals, m.p. 180–181 °C. ¹H NMR

(500 MHz, [D₆]DMSO): δ = 7.45–7.34 (m, 2H, C₆H₄), 7.30 (t, *J* = 7.4 Hz, 1H, C₆H₄), 7.17 (d, *J* = 7.0 Hz, 1H, C₆H₄), 6.93 (s, 1H, thiophene), 6.14 (s, 2H, NH₂), 3.62 (s, 4H, CH₂), 3.59 (s, 4H, CH₂), 2.33 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.4, 152.7, 144.4, 138.5, 132.9, 129.4, 129.1, 125.9, 122.6, 116.8, 100.9, 66.3, 45.6, 21.0 ppm. IR (ATR): 3441, 3299, 2966, 2913, 2891, 2853, 1579, 1537, 1493, 1466, 1456, 1426, 1378, 1299, 1258, 1157, 1112, 1016, 938, 885, 839, 780, 764, 719, 692, 579, 531, 483, 455, 428 cm⁻¹. MS (ESI): *m/z* = 303.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₈N₂O₂S: C 63.55, H 6.00, N 9.26, found: C 63.43, H 5.90, N 9.18.

5-(4-Methylphenyl)-2-(morpholin-4-ylcarbonyl)thiophen-3-amine (6Ec):

Yield 0.724 g, 80%, white crystals, m.p. 174–175 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.47 (d, *J* = 7.4 Hz, 2H, C₆H₄), 7.22 (d, *J* = 7.2 Hz, 2H, C₆H₄), 6.90 (s, 1H, thiophene), 6.13 (s, 2H, NH₂), 3.62 (s, 4H, CH₂), 3.59 (s, 4H, CH₂), 2.31 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.4, 152.7, 144.4, 138.3, 130.2, 129.8, 125.3, 116.3, 100.5, 66.3, 45.6, 20.8 ppm. IR (ATR): 3476, 3366, 2980, 2913, 2886, 2849, 1583, 1539, 1509, 1462, 1449, 1428, 1386, 1360, 1304, 1269, 1253, 1215, 1147, 1108, 1007, 932, 877, 836, 805, 757, 717, 661, 578, 514, 473, 447 cm⁻¹. MS (ESI): *m/z* = 303.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₈N₂O₂S: C 63.55, H 6.00, N 9.26, found: C 63.46, H 5.92, N 9.16.

5-(4-Fluorophenyl)-2-(morpholin-4-ylcarbonyl)thiophen-3-amine

(6Ed): Yield 0.771 g, 84%, white crystals, m.p. 159–160 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.63 (s, 2H, C₆H₄), 7.25 (t, *J* = 7.2 Hz, 2H, C₆H₄), 6.90 (s, 1H, thiophene), 6.12 (s, 2H, NH₂), 3.62 (s, 4H, CH₂), 3.58 (s, 4H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.2, 162.2 (d, *J* = 246.2 Hz), 152.4, 143.1, 129.6, 127.6 (d, *J* = 8.5 Hz), 117.0, 116.2 (d, *J* = 21.8 Hz), 101.2, 66.3, 45.6 ppm. IR (ATR): 3439, 3377, 3324, 2985, 2851, 1575, 1533, 1505, 1451, 1392, 1305, 1256, 1232, 1162, 1110, 1071, 1009, 933, 876, 828, 805, 754, 719, 579, 513, 488, 478, 454 cm⁻¹. MS (ESI): *m/z* = 307.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₅FN₂O₂S: C 58.81, H 4.94, N 9.14, found: C 58.74, H 4.87, N 9.06.

2-(Morpholin-4-ylcarbonyl)-5-(3-(trifluoromethyl)phenyl)thiophen-3-amine (6Ee):

Yield 0.822 g, 77%, light yellow crystals, m.p. 115–116 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.89 (d, *J* = 7.2 Hz, 1H, C₆H₄), 7.85 (s, 1H, C₆H₄), 7.75–7.60 (m, 2H, C₆H₄), 7.10 (s, 1H, thiophene), 6.11 (s, 2H, NH₂), 3.63 (s, 4H, CH₂), 3.59 (s, 4H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.0, 152.1, 142.0, 134.0, 130.5, 130.0 (q, *J* = 33.6 Hz), 129.4, 125.0 (q, *J* = 3.4 Hz), 124.0 (q, *J* = 272.2 Hz), 121.5 (q, *J* = 4.2 Hz), 118.3, 102.3, 66.3, 45.6 ppm. IR (ATR): 3422, 3327, 2961, 2844, 1578, 1537, 1457, 1436, 1418, 1399, 1330, 1305, 1257, 1229, 1163, 1114, 1073, 1015, 964, 880, 828, 801, 749, 725, 692, 665, 586, 516, 467, 436 cm⁻¹. MS (ESI): *m/z* = 357.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₅F₃N₂O₂S: C 53.93, H 4.24, N 7.86, found: C 53.85, H 4.18, N 7.75.

1-(3-(4-Amino-5-(morpholin-4-ylcarbonyl)thiophen-2-yl)phenyl)ethanone (6Ef):

Yield 0.792 g, 80%, light yellow crystals, m.p. 132–133 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.06 (s, 1H, 2H-C₆H₄), 7.94 (d, *J* = 7.3 Hz, 1H, C₆H₄), 7.84 (d, *J* = 7.0 Hz, 1H, C₆H₄), 7.60–7.55 (m, 1H, 5H-C₆H₄), 7.06 (s, 1H, thiophene), 6.13 (s, 2H, NH₂), 3.63 (s, 4H, CH₂), 3.60 (s, 4H, CH₂), 2.62 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.7, 165.1, 152.3, 143.0, 137.6, 133.4, 129.8, 128.4, 124.5, 117.7, 101.8, 66.3, 45.6, 26.9 ppm. IR (ATR): 3449, 3320, 2993, 2970, 2946, 2856, 1674, 1590, 1542, 1491, 1464, 1428, 1394, 1348, 1302, 1275, 1259, 1220, 1180, 1163, 1105, 1062, 1006, 946, 899, 879, 828, 788, 748, 717, 677, 606, 593, 519, 455 cm⁻¹. MS (ESI): *m/z* = 331.1 [M+H]⁺. Elemental analysis calcd for C₁₇H₁₈N₂O₃S: C 61.80, H 5.49, N 8.48, found: C 61.69, H 5.40, N 8.45.

Methyl 3-(4-amino-5-(morpholin-4-ylcarbonyl)thiophen-2-yl)benzoate (6Eg):

Yield 0.830 g, 80%, white crystals, m.p. 140–141 °C. ¹H NMR (500

FULL PAPER Accepted Article

MHz, [D₆]DMSO): δ = 8.08 (s, 1H, 2H-C₆H₄), 7.92 (d, J = 7.5 Hz, 1H, C₆H₄), 7.87 (d, J = 7.1 Hz, 1H, C₆H₄), 7.61–7.56 (m, 1H, C₆H₄), 7.05 (s, 1H, thiophene), 6.14 (s, 2H, NH₂), 3.88 (s, 3H, CH₃), 3.63 (s, 4H, CH₂), 3.60 (s, 4H, CH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 165.8, 165.1, 152.3, 142.6, 133.4, 130.6, 129.9 (overlapped), 129.1, 125.5, 117.7, 101.8, 66.3, 52.5, 45.6 ppm. IR (ATR): 3423, 3334, 2956, 2910, 2891, 2852, 1716, 1598, 1547, 1489, 1454, 1429, 1398, 1356, 1295, 1266, 1254, 1220, 1157, 1110, 1080, 1010, 985, 948, 906, 878, 838, 812, 753, 741, 688, 583, 474 cm⁻¹. MS (ESI): m/z = 347.1 [M+H]⁺. Elemental analysis calcd for C₁₇H₁₈N₂O₄S: C 58.94, H 5.24, N 8.09, found: C 58.80, H 5.16, N 7.96.

1-(3-Amino-5-phenylthiophen-2-yl)ethanone (6Fa): Yield 0.514 g, 79%, yellow crystals, m.p. 133–134 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.66 (d, J = 6.8 Hz, 2H, C₆H₅), 7.49–7.39 (m, 3H, C₆H₅), 7.22 (s, 2H, NH₂), 7.01 (s, 1H, thiophene), 2.28 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 189.2, 155.5, 147.7, 132.6, 129.4, 129.2, 125.6, 116.2, 108.5, 28.2 ppm. IR (ATR): 3413, 3305, 1589, 1525, 1489, 1462, 1362, 1306, 1222, 1181, 1164, 1128, 1016, 958, 927, 824, 759, 724, 683, 638, 617, 588, 549, 466, 449, 436 cm⁻¹. MS (ESI): m/z = 218.1 [M+H]⁺. Elemental analysis calcd for C₁₂H₁₁NOS: C 66.33, H 5.10, N 6.45, found: C 63.22, H 5.01, N 6.39.

1-(3-Amino-5-(3-methylphenyl)thiophen-2-yl)ethanone (6Fb): Yield 0.388 g, 56%, yellow crystals, m.p. 125–126 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.47–7.42 (m, 2H, C₆H₄), 7.32 (t, J = 7.5 Hz, 1H, C₆H₄), 7.27–7.11 (m, 3H, C₆H₄, NH₂), 7.00 (s, 1H, thiophene), 2.34 (s, 3H, CH₃), 2.27 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 189.6, 156.0, 148.4, 139.0, 133.1, 130.5, 129.6, 126.6, 123.3, 116.6, 108.9, 28.6, 21.4 ppm. IR (ATR): 3420, 3276, 3161, 2917, 1594, 1521, 1464, 1361, 1310, 1286, 1217, 1168, 1119, 1090, 1017, 971, 943, 891, 866, 836, 774, 706, 685, 673, 642, 572, 549, 440, 425 cm⁻¹. MS (ESI): m/z = 232.1 [M+H]⁺. Elemental analysis calcd for C₁₃H₁₃NOS: C 67.50, H 5.66, N 6.06, found: C 67.37, H 5.55, N 5.99.

1-(3-Amino-5-(4-methylphenyl)thiophen-2-yl)ethanone (6Fc): Yield 0.589 g, 85%, dark yellow crystals, m.p. 129–130 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.53 (d, J = 7.9 Hz, 2H, C₆H₄), 7.26–7.20 (m, 4H, C₆H₄, NH₂), 6.96 (s, 1H, thiophene), 2.31 (s, 3H, CH₃), 2.26 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 189.5, 156.1, 148.5, 139.6, 130.4, 130.2, 126.0, 116.1, 108.6, 28.6, 21.3 ppm. IR (ATR): 3405, 3278, 3172, 2919, 1592, 1538, 1501, 1463, 1359, 1309, 1240, 1215, 1184, 1106, 1019, 956, 927, 830, 806, 716, 692, 638, 587, 562, 507, 487, 430 cm⁻¹. MS (ESI): m/z = 232.1 [M+H]⁺. Elemental analysis calcd for C₁₃H₁₃NOS: C 67.50, H 5.66, N 6.06, found: C 67.36, H 5.59, N 5.97.

1-(3-Amino-5-(4-fluorophenyl)thiophen-2-yl)ethanone (6Fd): Yield 0.571 g, 81%, yellow crystals, m.p. 122–123 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.68 (dd, J_{HH} = 7.8, J_{HF} = 5.5 Hz, 2H, C₆H₄), 7.29–7.23 (m, 4H, C₆H₄, NH₂), 6.96 (s, 1H, thiophene), 2.26 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 189.7, 163.1 (d, J = 247.3 Hz), 156.0, 147.1, 129.8 (d, J = 3.0 Hz), 128.4 (d, J = 8.5 Hz), 116.8, 116.7 (d, J = 22.0 Hz), 109.0, 28.6 ppm. IR (ATR): 3408, 3272, 3169, 1591, 1534, 1501, 1462, 1361, 1318, 1229, 1159, 1096, 1012, 957, 924, 833, 818, 799, 717, 690, 667, 632, 585, 559, 543, 515, 480, 435 cm⁻¹. MS (ESI): m/z = 236.1 [M+H]⁺. Elemental analysis calcd for C₁₂H₁₀FNOS: C 61.26, H 4.28, N 5.95, found: C 61.18, H 4.19, N 5.84.

1-(3-Amino-5-(3-(trifluoromethyl)phenyl)phenyl)thiophen-2-yl)ethanone (6Fe): Yield 0.769 g, 90%, yellow crystals, m.p. 134–135 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.04–7.85 (m, 2H, C₆H₄), 7.75 (d, J = 7.6 Hz, 1H, C₆H₄), 7.68 (t, J = 7.6 Hz, 1H, C₆H₄), 7.22 (s, 2H, NH₂), 7.15 (s, 1H, thiophene), 2.29 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 189.9, 155.8, 145.9, 134.2, 130.9, 130.5 (q, J = 31.6 Hz), 130.1, 126.1 (q, J = 3.7 Hz), 124.3 (q, J = 272.4 Hz), 122.4 (q, J = 3.7 Hz), 118.1, 109.6,

28.7 ppm. IR (ATR): 3402, 3270, 3170, 1595, 1524, 1494, 1454, 1406, 1318, 1221, 1110, 1073, 1021, 972, 943, 894, 823, 794, 724, 686, 657, 634, 580, 550, 510, 492, 481, 432 cm⁻¹. MS (ESI): m/z = 286.0 [M+H]⁺. Elemental analysis calcd for C₁₃H₁₀F₃NOS: C 54.73, H 3.53, N 4.91, found: C 54.69, H 3.44, N 4.79.

1-(3-(5-Acetyl-4-aminothiophen-2-yl)phenyl)ethanone (6Ff): Yield 0.699 g, 90%, yellow crystals, m.p. 154–155 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.14 (t, J = 1.6 Hz, 1H, C₆H₄), 8.02–7.97 (m, 1H, C₆H₄), 7.94–7.90 (m, 1H, C₆H₄), 7.62 (t, J = 7.8 Hz, 1H, C₆H₄), 7.31–7.17 (brs, 2H, NH₂), 7.12 (s, 1H, thiophene), 2.64 (s, 3H, CH₃), 2.30 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 197.5 (o), 189.4 (o), 155.4 (o), 146.4 (o), 137.5 (o), 133.0 (o), 130.0 (+), 129.8 (+), 129.0 (+), 124.8 (+), 117.0 (+), 108.8 (o), 28.2 (+), 26.8 (+) ppm. IR (ATR): 3169, 1676, 1622, 1595, 1450, 1398, 1361, 1311, 1280, 1222, 1170, 1116, 1082, 944, 908, 833, 749, 707, 678, 644, 594, 560, 454, 420 cm⁻¹. MS (ESI): m/z = 260.1 [M+H]⁺. Elemental analysis calcd for C₁₄H₁₃NO₂S: C 64.84, H 5.05, N 5.40, found: C 64.70, H 5.00, N 5.41.

Methyl 3-(5-acetyl-4-aminothiophen-2-yl)benzoate (6Fg): Yield 0.544 g, 66%, dark yellow crystals, m.p. 129–130 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 8.10 (s, 1H, C₆H₄), 7.94 (d, J = 7.5 Hz, 1H, C₆H₄), 7.90 (d, J = 7.6 Hz, 1H, C₆H₄), 7.58 (t, J = 7.7 Hz, 1H, C₆H₄), 7.21 (s, 2H, NH₂), 7.07 (s, 1H, thiophene), 3.87 (s, 3H, OCH₃), 2.28 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 189.8, 166.0, 155.8, 146.5, 133.6, 131.0, 130.6, 130.3, 130.1, 126.2, 117.5, 109.3, 52.8, 28.7 ppm. IR (ATR): 3427, 3320, 2950, 1716, 1593, 1524, 1485, 1461, 1436, 1398, 1361, 1282, 1214, 1111, 1002, 960, 935, 901, 873, 823, 751, 665, 640, 577, 548, 475, 436 cm⁻¹. MS (ESI): m/z = 276.1 [M+H]⁺. Elemental analysis calcd for C₁₄H₁₃NO₃S: C 61.07, H 4.76, N 5.09, found: C 60.94, H 4.68, N 4.99.

General method for preparation of 5-aryl-2-nitrothiophen-3-amines (6Ga-Gg), 3-amino-5-phenylthiophene-2-carbonitriles (6Ha-Hg), ethyl 3-(3-amino-5-aryltiophen-2-yl)prop-2-enoates (6la-Ig), and 2-aryltiopheno(3,2-b)pyridine-5,7(4H,6H)-diones (6Ja-Jg). Reactions were carried out under an atmosphere of nitrogen in flame or oven-dried glassware. The corresponding reagent (**5G-J**) (3.3 mmol) was added dropwise under stirring to a solution of sodium hydrosulfide (0.185 g, 3.3 mmol) in methanol (15 mL) at -5 °C. The mixture was then stirred at -5 °C within 1 h. Then a solution of MeONa (9 mL, 1M) in methanol was added. After that solution of propanenitrile **3** (3 mmol) in methanol (5 mL) was added dropwise under stirring at -5 °C. The resulting mixture was stirred for 2 h at -5 °C and additional 2 h at ambient temperature. The solvent was evaporated under reduced pressure. Acetic acid (20 mL, ω = 2%) was added to the residue. The product was filtered off and recrystallized from methanol. For the case of syntheses of **6la-Ig** and **6Ja-Jg** ethanol and EtONa were used instead of methanol and MeONa.

5-Phenyl-2-nitrothiophen-3-amine (6Ga): Yield 0.475 g, 72%, yellow crystals, m.p. 222–223 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.40–8.03 (br.s, 2H, NH₂), 7.73–7.67 (m, 2H, C₆H₅), 7.54–7.44 (m, 3H, C₆H₅), 7.02 (s, 1H, thiophene) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 152.0 (o), 149.6 (o), 131.4 (o), 130.6 (+), 129.4 (+), 126.0 (+), 120.2 (o), 115.0 (+) ppm. IR (ATR): 3427, 3287, 3156, 1600, 1552, 1447, 1367, 1318, 1219, 1163, 1115, 1071, 1010, 998, 988, 908, 825, 762, 707, 684, 646, 514, 444 cm⁻¹. MS (ESI): m/z = 221.0 [M+H]⁺. Elemental analysis calcd for C₁₀H₈N₂O₂S: C 54.53, H 3.66, N 12.72, found: C 54.40, H 3.55, N 12.66.

5-(3-Methylphenyl)-2-nitrothiophen-3-amine (6Gb): Yield 0.393 g, 56%, white crystals, m.p. 210–211 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.27–8.07 (br.s, 2H, NH₂), 7.53–7.47 (m, 2H, C₆H₄), 7.38 (t, J = 7.9 Hz, 1H, C₆H₄), 7.31 (d, J = 7.5 Hz, 1H, C₆H₄), 7.00 (s, 1H, thiophene), 2.36 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 152.0 (o), 149.8 (o), 138.9 (o), 131.4 (+, two overlapped signals), 129.3 (+), 126.4 (+), 123.2 (+),

FULL PAPER Accepted Article

120.1 (o), 114.9 (+), 20.8 (+) ppm. IR (ATR): 3416, 3302, 1610, 1553, 1501, 1445, 1387, 1305, 1261, 1223, 1119, 1033, 998, 965, 894, 831, 777, 745, 697, 681, 653, 557, 515, 458, 421 cm⁻¹. MS (ESI): *m/z* = 235.1 [M+H]⁺. Elemental analysis calcd for C₁₁H₁₀N₂O₂S: C 56.39, H 4.30, N 11.96, found: C 56.28, H 4.18, N 11.81.

5-(4-Methylphenyl)-2-nitrothiophen-3-amine (6Gc): Yield 0.569 g, 81%, yellow crystals, m.p. 225–226 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.35–8.00 (br.s, 2H, NH₂), 7.60 (d, *J* = 8.0 Hz, 2H, C₆H₄), 7.31 (d, *J* = 7.9 Hz, 2H, C₆H₄), 6.97 (s, 1H, thiophene), 2.35 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 152.1 (o), 149.9 (o), 140.8 (o), 132.5 (o), 130.0 (+), 128.7 (o), 125.9 (+), 114.3 (+), 20.9 (+) ppm. IR (ATR): 3437, 3308, 1605, 1554, 1525, 1445, 1389, 1313, 1150, 1116, 1016, 933, 900, 802, 787, 744, 702, 634, 592, 571, 485, 466, 410 cm⁻¹. Spectroscopic data are in agreement with those reported in the literature.^[57]

5-(4-Fluorophenyl)-2-nitrothiophen-3-amine (6Gd): Yield 0.543 g, 76%, yellow crystals, m.p. 290–291 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.36–8.06 (m, 2H, NH₂), 7.77 (dd, *J*_{HH} = 7.7, *J*_{HF} = 5.7 Hz, 2H, C₆H₄), 7.34 (t, *J* = 8.4 Hz, 2H, C₆H₄), 6.98 (s, 1H, thiophene) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 163.3 (d, *J* = 249.0 Hz), 151.9, 148.4, 134.0, 128.5, 126.4 (d, *J* = 7.6 Hz), 120.2, 116.4 (d, *J* = 22.1 Hz) ppm. IR (ATR): 3442, 3299, 3081, 1646, 1626, 1595, 1526, 1514, 1487, 1449, 1391, 1317, 1234, 1163, 1119, 1027, 1013, 970, 935, 886, 827, 803, 751, 692, 677, 492, 447 cm⁻¹. Spectroscopic data are in agreement with those reported in the literature.^[57]

2-Nitro-5-(3-(trifluoromethyl)phenyl)thiophen-3-amine (6Ge): Yield 0.527 g, 61%, yellow crystals, m.p. 197–198 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.3–8.12 (br. s, 2H, NH₂), 8.02–7.98 (m, 2H, C₆H₄), 7.86 (d, *J* = 7.6 Hz, 1H, C₆H₄), 7.78–7.70 (m, 1H, C₆H₄), 7.17 (s, 1H, thiophene) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 151.6 (o), 147.2 (o), 132.5 (o), 130.7 (+), 130.1 (+), 130.1 (q, *J* = 32.5 Hz, o), 126.9 (q, *J* = 3.6 Hz, +), 123.7 (q, *J* = 272.6 Hz, o), 122.4 (q, *J* = 3.8 Hz, +), 120.7 (o), 116.8 (+) ppm. IR (ATR): 3418, 3307, 1625, 1563, 1510, 1446, 1395, 1314, 1229, 1166, 1071, 960, 887, 832, 799, 746, 717, 689, 664, 510, 420 cm⁻¹. MS (ESI): *m/z* = 289.0 [M+H]⁺. Elemental analysis calcd for C₁₁H₇F₃N₂O₂S: C 45.84, H 2.45, N 9.72, found: C 45.75, H 2.37, N 9.66.

5-((3-Acetylphenyl)phenyl)-2-nitrothiophen-3-amine (6Gf): Yield 0.511 g, 65%, off-yellow crystals, m.p. 180–181 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.35–8.12 (br.s, 2H, NH₂), 7.77 (s, 1H, C₆H₄), 7.68 (d, *J* = 7.9 Hz, 1H, C₆H₄), 7.55 (d, *J* = 8.5 Hz, 1H, C₆H₄), 7.31 (t, *J* = 7.8 Hz, 1H, C₆H₄), 7.00 (s, 1H, thiophene), 2.62 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 197.5, 148.9, 140.5, 137.6, 132.5, 130.0, 129.9, 126.1, 123.0, 119.9, 116.7, 26.9 ppm. IR (ATR): 3285, 3169, 1676, 1622, 1597, 1569, 1450, 1398, 1361, 1312, 1277, 1222, 1170, 1118, 1082, 1028, 944, 908, 832, 792, 749, 707, 678, 645, 594, 561, 454, 412 cm⁻¹. MS (ESI): *m/z* = 263.0 [M+H]⁺. Elemental analysis calcd for C₁₂H₁₀N₂O₃S: C 54.95, H 3.84, N 10.68, found: C 54.90, H 3.76, N 10.65.

Methyl 3-(4-amino-5-nitrothiophen-2-yl)benzoate (6Gg): Yield 0.584 g, 70%, yellow-brown crystals, m.p. 200–201 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.43–8.22 (br.s, 2H, NH₂), 8.13 (s, 1H, C₆H₄), 8.03 (d, *J* = 7.7 Hz, 1H, C₆H₄), 7.95–7.90 (m, 1H, C₆H₄), 7.57–7.49 (m, 1H, C₆H₄), 7.04 (s, 1H, thiophene), 3.92 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 166.6, 157.7, 146.5, 132.6, 131.8, 130.6, 130.2, 130.0, 125.9, 116.5, 115.6, 52.4 ppm. IR (ATR): 3425, 3314, 1721, 1627, 1562, 1448, 1398, 1287, 1222, 1114, 1081, 998, 831, 811, 789, 751, 669, 638, 622, 502, 456 cm⁻¹. MS (ESI): *m/z* = 279.0 [M+H]⁺. Elemental analysis calcd for C₁₂H₁₀N₂O₄S: C 51.79, H 3.62, N 10.07, found: C 51.86, H 3.60, N 10.11.

3-Amino-5-phenylthiophene-2-carbonitrile (6Ha): Yield 0.378 g, 63%, cream colored crystals, m.p. 179–180 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.62–7.58 (m, 2H, C₆H₅), 7.48–7.36 (m, 3H, C₆H₅), 6.94 (s, 1H, thiophene), 6.57 (s, 2H, NH₂) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 158.4, 148.6, 132.6, 129.9, 129.8, 126.1, 116.3, 116.2, 75.3 ppm. Spectroscopic data are in agreement with those reported in the literature.^[29b]

3-Amino-5-(3-methylphenyl)thiophene-2-carbonitrile (6Hb): Yield 0.417 g, 65%, buff colored crystals, m.p. 163–164 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.44–7.37 (m, 2H, C₆H₄), 7.30 (t, *J* = 7.4 Hz, 1H, C₆H₄), 7.17 (d, *J* = 7.0 Hz, 1H, C₆H₄), 7.00 (s, 1H, thiophene), 6.61 (s, 2H, NH₂), 2.36 (s, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 156.7, 147.0, 134.1, 131.4, 129.3, 126.4, 123.2, 120.4, 117.3, 115.5, 75.6, 20.8 ppm. IR (ATR): 3428, 3347, 2187, 1632, 1559, 1521, 1464, 1438, 1407, 1380, 1217, 1181, 1101, 1017, 800, 717, 676, 634, 557, 538, 511, 465 cm⁻¹. MS (ESI): *m/z* = 215.1 [M+H]⁺. Elemental analysis calcd for C₁₂H₁₀N₂S: C 67.26, H 4.70, N 13.07, found: C 67.07, H 4.62, N 13.10.

3-Amino-5-(4-methylphenyl)thiophene-2-carbonitrile (6Hc): Yield 0.507 g, 79%, buff colored crystals, m.p. 184–185 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.48 (d, *J* = 8.2 Hz, 2H, C₆H₄), 7.25 (d, *J* = 7.9 Hz, 2H, C₆H₄), 6.88 (s, 1H, thiophene), 6.53 (s, 2H, NH₂), 2.32 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 157.9 (o), 148.3 (o), 139.1 (o), 129.8 (+), 129.4 (o), 125.5 (+), 115.8 (o), 115.2 (+), 74.2 (o), 20.8 (+) ppm. IR (ATR): 3428, 3347, 2187, 1632, 1559, 1521, 1464, 1438, 1407, 1380, 1217, 1181, 1101, 1017, 800, 717, 676, 634, 557, 538, 511, 465 cm⁻¹. Spectroscopic data are in agreement with those reported in the literature.^[27g]

3-Amino-5-(4-fluorophenyl)thiophene-2-carbonitrile (6Hd): Yield 0.536 g, 82%, cream colored crystals, m.p. 200–201 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.79–7.60 (m, 2H, C₆H₄), 7.30 (t, *J* = 8.7 Hz, 2H, C₆H₄), 6.91 (s, 1H, thiophene), 6.58 (s, 2H, NH₂) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 163.1 (d, *J* = 247.3 Hz), 157.8, 147.3, 131.2 (d, *J* = 3.0 Hz), 129.8 (d, *J* = 8.5 Hz), 118.1 (d, *J* = 22.0 Hz), 116.4, 108.6, 75.1. IR (ATR): 3240, 2194, 1628, 1570, 1502, 1444, 1328, 1233, 1164, 1073, 1018, 968, 917, 890, 815, 723, 688, 514, 453 cm⁻¹. MS (ESI): *m/z* = 219.0 [M+H]⁺. Elemental analysis calcd for C₁₁H₇FN₂S: C 60.54, H 3.23, N 12.84, found: C 60.45, H 3.20, N 12.87.

3-Amino-5-(3-(trifluoromethyl)phenyl)thiophene-2-carbonitrile (6He): Yield 0.482 g, 60%, cream colored crystals, m.p. 160–161 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.96–7.86 (m, 2H, C₆H₄), 7.82–7.74 (m, 1H, C₆H₄), 7.72–7.66 (m, 1H, C₆H₄), 7.10 (s, 1H, thiophene), 6.61 (s, 2H, NH₂) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 157.8 (o), 145.9 (o), 133.1 (o), 130.6 (+), 130.0 (q, *J* = 31.8 Hz, o), 129.7 (+), 125.8 (q, *J* = 3.3 Hz, +), 123.8 (q, *J* = 272.5 Hz, o), 121.9 (q, *J* = 3.7 Hz, +), 117.3 (+), 115.4 (o), 75.6 (o) ppm. IR (ATR): 3452, 3355, 3241, 2191, 1628, 1568, 1500, 1481, 1443, 1417, 1329, 1233, 1168, 1119, 1073, 1020, 1000, 968, 917, 890, 819, 796, 723, 688, 654, 511, 497 cm⁻¹. MS (ESI): *m/z* = 269.0 [M+H]⁺. Elemental analysis calcd for C₁₂H₇FN₂S: C 53.73, H 2.63, N 10.44, found: C 53.65, H 2.70, N 10.40.

3-Amino-5-(3-acetylphenyl)thiophene-2-carbonitrile (6Hf): Yield 0.457 g, 63%, cream colored crystals, m.p. 154–155 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.08 (s, 1H, C₆H₄), 7.99 (d, *J* = 7.9 Hz, 1H, C₆H₄), 7.86 (d, *J* = 8.5 Hz, 1H, C₆H₄), 7.63–7.60 (m, 1H, C₆H₄), 7.06 (s, 1H, thiophene), 6.61 (s, 2H, NH₂), 2.64 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 197.5 (o), 157.9 (o), 146.9 (o), 137.5 (o), 132.5 (o), 130.0 (+), 129.8 (+), 129.0 (+), 124.8 (+), 116.6 (+), 115.5 (o), 75.2 (o), 26.8 (+) ppm. IR (ATR): 3445, 3352, 3240, 2926, 2191, 1685, 1628, 1600, 1568, 1495, 1442, 1409, 1353, 1291, 1277, 1210, 1177, 1102, 1080, 1021, 946, 892, 814, 791, 703, 677, 632, 585, 507, 422 cm⁻¹. MS (ESI): *m/z* = 243.1 [M+H]⁺.

FULL PAPER Accepted Article

Elemental analysis calcd for C₁₃H₁₀N₂OS: C 64.44, H 4.16, N 11.56, found: C 64.38, H 4.13, N 11.57.

Methyl 3-(4-amino-5-cyanothiophen-2-yl)benzoate (6Hg): Yield 0.503 g, 65%, cream colored crystals, m.p. 180–181 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.07 (s, 1H, C₆H₄), 7.97 (d, J = 7.7 Hz, 1H, C₆H₄), 7.87 (t, J = 9.0 Hz, 1H, C₆H₄), 7.64–7.56 (m, 1H, C₆H₄), 7.05 (s, 1H, thiophene), 6.61 (s, 2H, NH₂), 3.89 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 165.6 (o), 157.9 (o), 146.5 (o), 132.5 (o), 130.6 (o), 130.2 (+), 130.0 (+), 129.7 (+), 125.7 (+), 116.6 (+), 115.5 (o), 75.3 (o), 52.4 (+) ppm. IR (ATR): 3474, 3359, 3214, 2982, 1687, 1675, 1596, 1554, 1509, 1465, 1437, 1404, 1368, 1317, 1234, 1181, 1159, 1097, 1023, 972, 952, 835, 815, 716, 691, 576, 561, 516, 494 cm⁻¹. MS (ESI): m/z = 292.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₄FNO₂S: C 61.84, H 4.84, N 4.81, found: C 61.76, H 4.77, N 4.71.

Ethyl (2E)-3-(3-amino-5-phenylthiophen-2-yl)prop-2-enoate (6la): Yield 0.606 g, 74%, yellow crystals, m.p. 156–157 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.98 (d, J = 15.0 Hz, 1H, CH=), 7.59 (d, J = 7.4 Hz, 2H, C₆H₅), 7.45–7.35 (m, 3H, C₆H₅), 6.92 (s, 1H, thiophene), 6.33 (s, 2H, NH₂), 5.63 (d, J = 15.0 Hz, 1H, CH=), 4.13 (q, J = 7.0 Hz, 2H, CH₂), 1.24 (t, J = 7.1 Hz, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 166.7 (o), 152.4 (o), 144.6 (o), 135.8 (+), 132.9 (o), 129.1 (+), 128.7 (+), 125.3 (+), 117.3 (+), 107.7 (o), 107.2 (+), 59.2 (-), 14.4 (+) ppm. IR (ATR): 3372, 3203, 2975, 2901, 1687, 1674, 1590, 1552, 1498, 1463, 1423, 1412, 1368, 1319, 1224, 1178, 1074, 1026, 972, 954, 908, 839, 759, 723, 688, 634, 519, 460 cm⁻¹. MS (ESI): m/z = 274.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₅NO₂S: C 65.91, H 5.53, N 5.12, found: C 65.79, H 5.44, N 5.05.

Ethyl (2E)-3-(3-amino-5-(3-methylphenyl)thiophen-2-yl)prop-2-enoate (6lb): Yield 0.560 g, 65%, yellow crystals, m.p. 145–146 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.97 (d, J = 15.0 Hz, 1H, CH=), 7.45–7.34 (m, 2H, C₆H₄), 7.30 (t, J = 7.4 Hz, 1H, C₆H₄), 7.17 (d, J = 7.0 Hz, 1H, C₆H₄), 6.85 (s, 1H, thiophene), 6.29 (s, 2H, NH₂), 5.59 (d, J = 15.0 Hz, 1H, CH=), 4.12 (q, J = 7.1 Hz, 2H, CH₂), 2.27 (s, 3H, CH₃), 1.22 (t, J = 7.1 Hz, 3H, OCH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 166.8, 152.4, 144.9, 138.4, 135.8, 130.2, 129.4, 129.1, 125.9, 122.6, 116.8, 107.3, 106.8, 59.2, 20.9, 14.4 ppm. IR (ATR): 2978, 2920, 1979, 1709, 1603, 1456, 1373, 1175, 1093, 1035, 823, 783, 695, 594, 512, 435 cm⁻¹. MS (ESI): m/z = 288.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₇NO₂S: C 66.87, H 5.96, N 4.84, found: C 66.87, H 5.84, N 4.79.

Ethyl (2E/Z)-3-(3-amino-5-(4-methylphenyl)thiophen-2-yl)prop-2-enoate (6lc): Yield 0.491 g, 57%, yellow crystals, m.p. 174–175 °C. ¹H NMR (400 MHz, [D₆]DMSO) (E:Z = 0.94:0.06): δ = 7.97 (d, J = 15.0 Hz, 1H, CH=), 7.47 (d, J = 8.2 Hz, 1.9H, C₆H₄), 7.23 (d, J = 7.9 Hz, 2H, C₆H₄), 7.15 (d, J = 8.1 Hz, 0.15H, C₆H₄), 6.86 (s, 1H, thiophene), 6.30 (s, 1.9H, NH₂), 6.15 (s, 0.12H, NH₂), 5.59 (d, J = 15.0 Hz, 0.92H, CH=), 5.26 (d, J = 12.2 Hz, 0.06H, CH=), 4.12 (q, J = 7.1 Hz, 2H, CH₂), 2.31 (s, 2.84H, CH₃), 2.28 (s, 0.2H, CH₃), 1.23 (t, J = 7.1 Hz, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 166.8 (o), 166.5 (o), 155.6 (o), 152.5 (o), 144.9 (o), 141.3 (o), 138.4 (o), 135.8 (+), 132.4 (o), 130.2 (o), 129.7 (+), 129.3 (+), 127.5 (+), 125.2 (+), 116.8 (+), 116.4 (o), 107.3 (o), 106.8 (+), 101.3 (+), 59.2 (-), 20.8 (+), 14.4 (+) ppm. IR (ATR): 3383, 3323, 1684, 1589, 1550, 1510, 1459, 1367, 1318, 1236, 1169, 1113, 1026, 972, 954, 834, 805, 715, 692, 636, 540, 517, 476 cm⁻¹. MS (ESI): m/z = 288.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₇NO₂S: C 66.87, H 5.96, N 4.87, found: C 66.75, H 5.89, N 4.92.

Ethyl (2E)-3-(3-amino-5-(4-fluorophenyl)thiophen-2-yl)prop-2-enoate (6ld): Yield 0.750 g, 86%, yellow crystals, m.p. 148–149 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 7.98 (d, J = 15.0 Hz, 1H, CH=), 7.63 (dd, J_{HH} = 8.8 Hz, J_{HF} = 5.3 Hz, 2H, 3,5H-C₆H₄), 7.32–7.17 (m, 2H, C₆H₄), 6.88 (s, 1H, thiophene), 6.33 (s, 2H, NH₂), 5.62 (d, J = 15.0 Hz, 1H, CH=), 4.13 (q, J = 7.1 Hz, 2H, CH₂), 1.23 (t, J = 7.1 Hz, 3H, CH₃) ppm. ¹³C NMR (101 MHz,

[D₆]DMSO): δ = 166.7 (o), 162.2 (d, J = 246.5 Hz, o), 152.4 (o), 143.5 (o), 135.7 (+), 129.54 (d, J = 3.1 Hz, o), 127.4 (d, J = 8.3 Hz, +), 117.5 (+), 116.1 (d, J = 22.0 Hz, +), 107.7 (o), 107.2 (+), 59.2 (-), 14.3 (+) ppm. IR (ATR): 3474, 3359, 3214, 2982, 1687, 1675, 1596, 1554, 1509, 1465, 1437, 1404, 1368, 1317, 1234, 1181, 1159, 1097, 1023, 972, 952, 835, 815, 716, 691, 576, 561, 516, 494 cm⁻¹. MS (ESI): m/z = 292.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₄FNO₂S: C 61.84, H 4.84, N 4.81, found: C 61.76, H 4.77, N 4.71.

Ethyl (2E)-3-(3-amino-5-(3-(trifluoromethyl)phenyl)thiophen-2-yl)prop-2-enoate (6le): Yield 0.583 g, 57%, yellow crystals, m.p. 120–121 °C. ¹H NMR (500 MHz, [D₆]DMSO): δ = 7.97 (d, J = 15.0 Hz, 1H, CH=), 7.74 (d, J = 7.2 Hz, 1H, C₆H₄), 7.71 (s, 1H, C₆H₄), 7.63–7.43 (m, 2H, C₆H₄), 6.91 (s, 1H, thiophene), 6.32 (s, 2H, NH₂), 5.62 (d, J = 15.0 Hz, 1H, CH=), 4.12 (q, J = 7.0 Hz, 2H, CH₂), 1.23 (t, J = 7.1 Hz, 3H, CH₃) ppm. ¹³C NMR (126 MHz, [D₆]DMSO): δ = 166.9, 152.6, 144.8, 134.3, 130.7, 130.2 (q, J = 33.6 Hz), 129.6, 126.6, 125.3 (q, J = 3.4 Hz), 124.2 (q, J = 276.4 Hz), 121.8 (q, J = 4.3 Hz), 118.5, 107.7, 107.2, 59.2, 14.4 ppm. IR (ATR): 3296, 1593, 1558, 1532, 1489, 1332, 1311, 1111, 999, 973, 960, 915, 894, 859, 839, 793, 753, 715, 683, 613, 493, 436 cm⁻¹. MS (ESI): m/z = 342.1 [M+H]⁺. Elemental analysis calcd for C₁₆H₁₄F₃NO₂S: C 56.30, H 4.13, N 4.10, found: C 56.14, H 4.09, N 4.01.

Ethyl (2E/Z)-3-(5-(3-acetylphenyl)-3-aminothiophen-2-yl)prop-2-enoate (6lf): Yield 0.642 g, 68%, yellow crystals, m.p. 127–128 °C. ¹H NMR (400 MHz, [D₆]DMSO) (E:Z* = 0.75:0.25): δ = 8.10 (s, 0.25H, 2H-C₆H₄), 8.08 (s, 0.75H, 2H-C₆H₄), 7.99 (d, J = 15.1 Hz, 0.75H, =CH), 7.94 (d, J = 7.7 Hz, 1H, C₆H₄), 7.84 (d, J = 7.9 Hz, 1H, C₆H₄), 7.59 (t, J = 7.8 Hz, 1H, C₆H₄), 7.24 (d, J = 12.2 Hz, 0.25H, =CH), 7.04 (s, 0.75H, thiophene), 7.03 (s, 0.25H, thiophene), 6.34 (s, 1.5H, NH₂), 6.21 (s, 0.5H, NH₂), 5.67 (d, J = 15.1 Hz, 0.75H, =CH), 5.33 (d, J = 12.2 Hz, 0.25H, =CH), 4.13 (q, J = 7.1 Hz, 2H, CH₂), 2.63 (s, 3H, CH₃), 1.24 (t, J = 7.1 Hz, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 197.6 (o), 166.7 (o), 166.5* (o), 154.8* (o), 152.3 (o), 145.9* (o), 143.4 (o), 137.5 (o), 137.4* (o), 135.6 (+), 134.2* (+), 133.9* (o), 133.3 (o), 129.7* (+), 129.6 (+), 128.35 (+), 128.30* (+), 124.4 (+), 124.2* (+), 118.2 (+), 116.1* (+), 108.2 (o), 108.0* (o), 107.7 (+), 104.3* (+), 59.3 (-), 59.0* (-), 26.8 (+), 14.4 (+), 14.3* (+) ppm. IR (ATR): 3446, 3349, 3228, 2988, 1678, 1586, 1557, 1494, 1464, 1428, 1367, 1324, 1272, 1233, 1170, 1110, 1030, 980, 940, 841, 786, 716, 677, 588, 521, 468 cm⁻¹. MS (ESI): m/z = 316.1 [M+H]⁺. Elemental analysis calcd for C₁₇H₁₇NO₃S: C 64.74, H 5.43, N 4.44, found: C 64.80, H 5.45, N 4.42.

Methyl 3-(4-amino-5-((1E)-3-ethoxy-3-oxoprop-1-en-1-yl)thiophen-2-yl)benzoate (6lg): Yield 0.605 g, 61%, yellow crystals, m.p. 79–80 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 8.09 (s, 1H, 2H-C₆H₄), 8.05–7.75 (m, 3H, C₆H₄, CH=), 7.64–7.54 (br.s, 1H, C₆H₄), 7.02 (s, 1H, thiophene), 6.35 (s, 2H, NH₂), 5.68 (d, J = 14.7 Hz, 1H, CH=), 4.15–4.12 (m, 2H, CH₂), 3.89 (s, 3H, OCH₃), 1.28–1.20 (br.s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 166.7 (o), 165.7 (o), 152.3 (o), 143.0 (o), 135.6 (+), 133.3 (o), 130.5 (o), 129.8 (+), 129.7 (+), 129.0 (+), 125.4 (+), 118.2 (+), 108.2 (o), 107.8 (+), 59.3 (-), 52.3 (+), 14.3 (+) ppm. IR (ATR): 3435, 3344, 3231, 2981, 2950, 2905, 1703, 1679, 1592, 1558, 1466, 1427, 1366, 1324, 1282, 1239, 1178, 1109, 964, 837, 806, 750, 730, 677, 637, 587, 523, 470 cm⁻¹. MS (ESI): m/z = 332.1 [M+H]⁺. Elemental analysis calcd for C₁₇H₁₇NO₄S: C 61.62, H 5.17, N 4.23, found: C 61.53, H 5.09, N 4.18.

2-Phenylthieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Ja): Yield 0.620 g, 85%, white crystals, m.p. >355 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 11.91–11.48 (br.s, 2H), 7.76–7.66 (m, 2H, C₆H₅), 7.51–7.44 (m, 2H, C₆H₅), 7.43–7.37 (m, 1H, C₆H₅), 7.28 (s, 1H, thiophene), 5.64 (s, 1H, CH) ppm. ¹³C NMR (151 MHz, [D₆]DMSO): δ = 146.2, 129.3, 125.7 ppm. IR (ATR): 3017, 3002, 2807, 1668, 1622, 1586, 1519, 1472, 1436, 1386, 1351, 1319, 1224, 1184, 1161, 1136, 1107, 1015, 998, 861, 808, 748, 719, 706, 676,

FULL PAPER Accepted Article

599, 534, 443, 428 cm⁻¹. MS (ESI): *m/z* = 244.0 [M+H]⁺. Elemental analysis calcd for C₁₃H₉NO₂S: C 64.18, H 3.73, N 5.76, found: C 64.05, H 3.62, N 5.60. ¹H NMR spectrum is in agreement with those reported in the literature.^[58]

2-(3-Methylphenyl)thieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Jb): Yield 0.439 g, 57%, cream colored crystals, m.p. >348 °C. ¹H NMR (400 MHz, [D₆]DMSO): δ = 11.61 (s, 1H, NH), 11.47 (s, 1H, OH), 7.52 (s, 1H, C₆H₄), 7.49 (d, *J* = 7.9 Hz, 1H, C₆H₄), 7.35 (t, *J* = 7.6 Hz, 1H, C₆H₄), 7.27–7.17 (m, 2H, C₆H₄, thiophene), 5.59 (s, 1H, CH) ppm. ¹³C NMR (151 MHz, [D₆]DMSO): δ = 164.2, 146.3, 138.7, 132.7, 129.7, 129.2, 126.1, 122.9, 119.4, 112.7, 95.6, 20.9 ppm. IR (ATR): 3020, 2857, 1662, 1616, 1581, 1476, 1447, 1374, 1344, 1311, 1218, 1122, 943, 800, 743, 696, 637, 557, 522, 469, 445, 428 cm⁻¹. MS (ESI): *m/z* = 258.1 [M+H]⁺. Elemental analysis calcd for C₁₄H₁₁NO₂S: C 65.35, H 4.31, N 5.44, found: C 65.20, H 4.22, N 5.36.

2-(4-Methylphenyl)thieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Jc): Yield 0.532 g, 69%, cream colored crystals, m.p. 250–252 °C (dec.). ¹H NMR (600 MHz, [D₆]DMSO): δ = 11.58 (s, 1H), 11.45 (s, 1H), 7.59 (d, *J* = 8.1 Hz, 2H, C₆H₄), 7.28 (d, *J* = 7.8 Hz, 2H, C₆H₄), 7.20 (s, 1H, thiophene), 5.58 (s, 1H, CH), 2.34 (s, 3H, CH₃) ppm. ¹³C NMR (151 MHz, [D₆]DMSO): δ = 164.7, 146.8, 139.2, 130.6, 130.5, 130.4, 128.2, 127.0, 126.1, 112.8, 96.0, 21.3 ppm. IR (ATR): 2852, 1662, 1615, 1575, 1475, 1447, 1343, 1309, 1217, 1121, 1019, 942, 799, 742, 695, 636, 557, 523, 469, 427 cm⁻¹. MS (ESI): *m/z* = 258.1 [M+H]⁺. Elemental analysis calcd for C₁₄H₁₁NO₂S: C 65.35, H 4.31, N 5.44, found: C 65.20, H 4.24, N 5.31.

2-(4-Fluorophenyl)thieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Jd): Yield 0.603 g, 77%, white crystals, m.p. 240–243 °C (dec.). ¹H NMR (400 MHz, [D₆]DMSO): δ = 11.63 (s, 1H), 11.51 (s, 1H), 7.77 (dd, *J*_{HH} = 8.9 Hz, *J*_{HF} = 5.3 Hz, 2H, C₆H₄), 7.31 (t, *J* = 8.9 Hz, 2H, C₆H₄), 7.23 (s, 1H, thiophene), 5.60 (s, 1H, CH) ppm. IR (ATR): 2803, 1661, 1585, 1471, 1411, 1381, 1347, 1318, 1223, 1159, 1098, 1012, 946, 813, 740, 696, 659, 557, 523, 489, 448, 433, 417 cm⁻¹. MS (ESI): *m/z* = 262.0 [M+H]⁺. Elemental analysis calcd for C₁₃H₈FNO₂S: C 59.76, H 3.09, N 5.36, found: C 59.62, H 2.99, N 5.31.

2-(3-(Trifluoromethyl)phenyl)thieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Je): Yield 0.532 g, 57%, cream colored crystals, m.p. 260–262 °C (dec.). ¹H NMR (500 MHz, [D₆]DMSO): δ = 11.81 (s, 1H, NH), 11.72 (s, 1H, OH), 8.03–7.99 (m, 2H, C₆H₄), 7.76 (d, *J* = 7.7 Hz, 1H, C₆H₄), 7.71 (t, *J* = 7.9 Hz, 1H, C₆H₄), 7.46 (s, 1H, thiophene), 5.76 (s, 1H, CH) ppm. ¹³C NMR (151 MHz, [D₆]DMSO): δ = 164.8 (o), 161.7 (o), 144.4 (o), 144.2 (o), 134.4 (o), 131.0 (+), 130.6 (q, *J* = 32.0 Hz, o), 130.3 (+), 125.8 (q, *J* = 3.6 Hz, +), 124.4 (q, *J* = 272.6 Hz, o), 122.5 (q, *J* = 3.6 Hz, +), 115.0 (+), 111.5 (q, *J* = 3.3 Hz, o), 96.6 (+) ppm. IR (ATR): 2802, 1661, 1615, 1585, 1512, 1475, 1439, 1422, 1380, 1327, 1251, 1225, 1167, 1116, 1096, 1071, 1000, 968, 889, 793, 740, 711, 686, 656, 533, 500, 451, 436, 421, 410 cm⁻¹. MS (ESI): *m/z* = 312.0 [M+H]⁺. Elemental analysis calcd for C₁₄H₈F₃NO₂S: C 54.02, H 2.59, N 4.50, found: C 53.93, H 2.52, N 4.39.

2-(3-Acetylphenyl)thieno(3,2-b)pyridine-5,7(4H,6H)-dione (6Jf): Yield 0.633 g, 74%, cream colored crystals, m.p. 230–232 °C (dec.). ¹H NMR (400 MHz, [D₆]DMSO): δ = 12.00–11.22 (m, 2H, NH, OH), 8.17 (s, 1H, C₆H₄), 8.09–7.82 (m, 2H, C₆H₄), 7.62 (t, *J* = 7.8 Hz, 1H, C₆H₄), 7.38 (s, 1H, thiophene), 5.63 (s, 1H, CH), 2.65 (s, 3H, CH₃) ppm. ¹³C NMR (101 MHz, [D₆]DMSO): δ = 197.6, 164.2, 161.2, 144.9, 143.7, 137.6, 133.2, 130.1, 129.8, 128.6, 124.9, 113.8, 95.9, 26.9 ppm. IR (ATR): 3020, 2857, 1662, 1616, 1581, 1476, 1447, 1374, 1344, 1311, 1218, 1122, 943, 800, 743, 696, 637, 557, 522, 469, 445, 428 cm⁻¹. MS (ESI): *m/z* = 286.1 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₁NO₃S: C 63.14, H 3.89, N 4.91, found: C 62.99, H 3.80, N 4.80.

Methyl 3-(5,7-dioxo-4,5,6,7-tetrahydrothieno(3,2-b)pyridin-2-yl)benzoate (6Jg): Yield 0.596 g, 66%, cream colored crystals, m.p. 345–348 °C (dec.). ¹H NMR (600 MHz, [D₆]DMSO): δ = 11.82–11.42 (br.s, 2H, NH, OH), 8.17 (t, *J* = 1.6 Hz, 1H, C₆H₄), 8.08–7.98 (m, 1H, C₆H₄), 7.98–7.94 (m, 1H, C₆H₄), 7.75–7.54 (m, 1H, C₆H₄), 7.34 (s, 1H, thiophene), 5.61 (s, 1H, CH), 3.90 (s, 3H, CH₃) ppm. ¹³C NMR (151 MHz, [D₆]DMSO): δ = 165.71, 164.3, 144.5, 133.3, 130.7, 130.3, 130.0, 129.4, 125.8, 113.8, 96.0, 52.5 ppm. IR (ATR): 2951, 2809, 1723, 1668, 1622, 1580, 1511, 1479, 1375, 1346, 1287, 1219, 1134, 996, 951, 897, 796, 748, 736, 685, 620, 532, 445, 425 cm⁻¹. MS (ESI): *m/z* = 302.0 [M+H]⁺. Elemental analysis calcd for C₁₅H₁₁NO₄S: C 59.79, H 3.68, N 4.65, found: C 59.60, H 3.52, N 4.53.

Keywords: Meerwein arylation • Arenediazonium salts • Thiophene • Heterocycle synthesis • Thorpe-Ziegler cyclization

- [1] a) J. Schatz, in *Science of Synthesis*, G. Maas, Ed., Thieme Verlag, Stuttgart, 2001, Vol. 9, chapter 9.10., pp. 287–393; b) W.-D. Rudorf, in *Houben-Weyl, Methoden der organischen Chemie*, Thieme Verlag, Stuttgart, 1994, Vol. E6a, p. 186–555.
- [2] D. K. Russel, in *Comprehensive Heterocyclic Chemistry II*, A. R. Katritzky, C. W. Rees, E. F. V. Scriven, Eds.; Pergamon, Oxford, 1996, Vol. 2, Chapter 2.11, pp. 680–729.
- [3] R. Shah, P. K. Verma, *Chem. Central J.* **2018**, 12 (137), 1–22.
- [4] a) K. Yamada, Y. Abe, Y. Kato, A. Nakuchi, Y. Saito, S. Fuchi, PCT Int. Appl. (2019), WO 2019168140 A1 20190906; b) M. F. El-Shehry, H. M. Hosni, A. E. Amr, A. A. Ibrahim, A. A. Fayed, D. H. Elnaggar, *Russ. J. Gen. Chem.* **2019**, 89, 1528–1534; c) X. Wang, Z. Ren, M. Wang, M. Chen, A. Lu, W. Si, C. Yang, *Chem. Central J.* **2018**, 12, 83; d) T. Liu, H. Wu, H. Jiang, L. Zhang, Y. Zhang, L. Mao, *J. Agricult. Food Chem.* **2019**, 67, 6160–6168.
- [5] a) E. J. Roh, J. Changjoon, S. J. Oh, J. H. Jung, U.S. Pat. Appl. Publ. (2015), US 20150148550 A1 20150528; b) S. Pekert, J. Brendel, H. Hemmerle, H.-W. Kleemann, PCT/EP2001/013958.
- [6] a) K. M. Roque Marques, M. Rodrigues do Desterro, S. M. de Arruda, L. Nascimento de Araújo Neto, M. do Carmo Alves de Lima, S. M. Vitalino de Almeida, E. C. Dantas da Silva, T. Mendonça de Aquino, E. Ferreira da Silva-Júnior, J. X. de Araújo-Júnior, M. de M. Silva, M. Dayanne de A. Dantas, J. Carinhanha C. Santos, I. M. Figueiredo, M.-A. Bazin, P. Marchand, T. Gonçalves da Silva, F. J. Bezerra Mendonça Junior, *Curr. Top. Med. Chem.* **2019**, 19, 1075–1091; b) K. C. Gulipalli, P. Ravula, S. Bodige, S. Endoori, P. K. R. Cherukumallil, J. N. Narendra Sharath Chandra, N. Seelam, *Russ. J. Gen. Chem.* **2019**, 89, 1502–1512.
- [7] a) R. M. B. M. Girard, M. Crispim, I. Stolic, F. S. Damasceno, M. Santos da Silva, E. M. F. Pral, M. C. Elias, M. Bajic, A. M. Silber, *Antimicrob. Agents Chemother.* **2016**, 60, 5867–5877; b) R. Arancibia, A. H. Klahn, G. E. Buono-Core, D. Contreras, G. Barriga, C. Olea-Azar, M. Lapier, J. D. Maya, A. Ibanez, M. T. Garland, *J. Organomet. Chem.* **2013**, 743, 49–54.
- [8] a) Y. Unver, M. Tuluk, N. Kahriman, M. Emirik, E. Bektaş, S. Direkeli, *Russ. J. Gen. Chem.* **2019**, 89, 794–799; b) J. J. Brendle, A. Outlaw, A. Kumar, D. W. Boykin, D. A. Patrick, R. R. Tidwell, K. A. Werbovetz, *Antimicrob. Agents Chemother.* **2002**, 46, 797–807.
- [9] J. Fournier dit Chabert, B. Marquez, L. Neville, L. Joucla, S. Brousseau, P. Bouhours, E. David, S. Pellet-Rostaing, B. Marquet, N. Moreau, M. Lemaire, *Bioorg. Med. Chem.* **2007**, 15, 4482–4497.
- [10] a) A. Oster, S. Hinsberger, R. Werth, S. Marchais-Oberwinkler, M. Frotscher, R. W. Hartmann, *J. Med. Chem.* **2010**, 53, 8176–8186; b) E. Bey, S. Marchais-Oberwinkler, R. Werth, M. Negri, Y. A. Al-Soud, P. Kruchten, A. Oster, M. Frotscher, B. Birk, R. W. Hartmann, *J. Med. Chem.* **2008**, 51, 6725–6739.
- [11] L. Yan, R. Budhu, P. Huo, C. L. Lynch, J. J. Hale, S. G. Mills, R. Hajdu, C. A. Keohane, M. J. Rosenbach, J. A. Milligan, G.-J. Shei, G. Chrebet,

FULL PAPER Accepted Article

- J. Bergstrom, D. Card, S. M. Mandala, *Bioorg. Med. Chem. Lett.* **2006**, 16, 3564–3568.
- [12] a) F. D. Tsourtou, S. D. Peroukidis, V. G. Mavrantzas, *J. Mater. Chem. C* **2019**, 7, 9984–9995; b) I. Wurzbach, C. Rothe, K. Bruchlos, S. Ludwigs, F. Giesselmann, *J. Mater. Chem. C* **2019**, 7, 2615–2624.
- [13] a) T. Higashino, K. Ishida, T. Sakurai, S. Seki, T. Konishi, K. Kamada, H. Imahori, *Chem. Eur. J.* **2019**, 25, 6425–6438; b) J. Podlesny, O. Pytela, M. Klikar, V. Jelinkova, I. V. Kityk, K. Ozga, J. Jedryka, M. Rudysh, F. Bures, *Org. Biomol. Chem.* **2019**, 17, 3623–3634.
- [14] M. D. Levi, D. Aurbach, *ECS Transact.* **2007**, 3, 259–263.
- [15] K. Masui, A. Mori, K. Okano, K. Takamura, M. Kinoshita, T. Ikeda, *Org. Lett.* **2004**, 6, 2011–2014.
- [16] T. Ma, H.-F. Wang, K.-Q. Zhao, B.-Q. Wang, P. Hu, H. Monobe, B. Heinrich, B. Donnio, *ChemPlusChem* **2019**, 84, 1439–1448.
- [17] X.-Q. Ran, J.-K. Feng, Y.-L. Liu, A.-M. Ren, L.-Y. Zou, C.-C. Sun, *J. Phys. Chem. A* **2008**, 112, 10904–10911.
- [18] a) K. Gewald, E. Schinke, H. Böttcher, *Chem. Ber.* **1966**, 99, 94–100; b) Z. Puterová, A. Krutošíková, D. Végh, *ARKIVOC* **2010**, 1, 209–246.
- [19] a) S. O. Vardanyan, A. A. Aghekyan, A. S. Avagyan, S. A. Harutyunyan, H. V. Gasparyan, *Russ. J. Org. Chem.* **2019**, 55, 598–601; b) I. Yahaya, N. Seferoglu, Z. Seferoglu, *Tetrahedron* **2019**, 75, 2143–2154; c) I. S. Luna, R. M. Duarte da Cruz, R. S. Aquino de Araujo, F. J. B. Mendonça-Junior, *Curr. Org. Synth.* **2018**, 15, 1026–1042; d) O. Ya. Shyyka, N. T. Pokhodyo, Y. I. Slyvka, E. A. Goreshnik, M. D. Obushak, *Tetrahedron Lett.* **2018**, 59, 1112–1115.
- [20] a) S. Arshadi, E. Vessally, L. Edjlali, E. Ghorbani-Kalhor, R. Hosseinzadeh-Khanmiri, *RSC Adv.* **2017**, 7, 13198–13211; b) B. V. Lichitsky, R. M. Belyi, A. N. Komogortsev, A. A. Dudinov, M. M. Krayushkin, *Russ. Chem. Bull.* **2009**, 58, 387–391; c) H. Wamhoff, in *Adv. Heterocycl. Chem.*, A. R. Katritzky (Ed.), Academic Press, 1985, 38, 299–368; d) H. Wamhoff, A. Schmidt, *J. Org. Chem.* **1993**, 58, 6976–6984.
- [21] D. Dagoneau, A. Kolleth, A. Lumbroso, G. Tanriver, S. Catak, S. Sulzer-Mossé, A. De Mesmaeker, *Helv. Chim. Acta* **2019**, 102, e1900031.
- [22] P. Böger, *J. Pestic. Sci.* **1997**, 22, 257–262.
- [23] M. Snoeck, *Local Reg. Anesth.* **2012**, 5, 23–33.
- [24] D.-B. Sung, B. Mun, S. Park, H.-S. Lee, J. Lee, Y.-J. Lee, H. J. Shin, J. S. Lee, *J. Org. Chem.* **2019**, 84, 379–391.
- [25] W. Steinkopf, P. J. Müller, *Justus Liebigs Ann. Chem.* **1926**, 448, 210–222.
- [26] M. Watanabe, T. Yamamoto, M. Nishiyama, *Chem. Commun.* **2000**, 133–134.
- [27] a) R. Romagnoli, F. Prencipe, P. Oliva, S. Baraldi, P. G. Baraldi, S. Schiaffino Ortega, M. Chayah, M. Kimatrai Salvador, L. C. Lopez-Cara, A. Brancale, S. Ferla, E. Hamel, R. Ronca, R. Bortolozzi, E. Mariotto, E. Mattiuzzo, G. Viola, *J. Med. Chem.* **2019**, 62, 1274–1290; b) Q. Zhang, Z. Hu, Q. Shen, Y. Chen, W. Lu, *Molecules* **2017**, 22, 788–804; c) N. T. Pokhodyo, O. Y. Shyyka, V. S. Matiychuk, M. D. Obushak, *ACS Comb. Sci.* **2015**, 17, 399–403; d) N. T. Pokhodyo, O. Y. Shyyka, M. D. Obushak, *Synth. Commun.* **2014**, 44, 1002–1006; e) H. J. Sahner, M. Groh, M. Negri, J. Haupenthal, R. W. Hartmann, *Eur. J. Med. Chem.* **2013**, 65, 223–231; f) M. Murai, R. Hatano, S. Kitabata, K. Ohe, *Chem. Comm.* **2011**, 47, 2375–2377; g) D. Thomas, G. Kirsch, P. Seck, *Synthesis* **2007**, 7, 1027–1032; h) D. L. Hertzog, K. A. Al-Barazanji, E. C. Bigham, M. J. Bishop, C. S. Britt, D. L. Carlton, J. P. Cooper, A. J. Daniels, D. M. Garrido, A. S. Goetz, M. K. Grizzle, Y. C. Guo, A. L. Handlon, D. M. Ignar, R. O. Morgan, A. J. Peat, F. X. Tavares, H. Zhou, *Bioorg. Med. Chem. Lett.* **2006**, 16, 4723–4727; i) F. X. Tavares, K. A. Al-Barazanji, E. C. Bigham, M. J. Bishop, C. S. Britt, D. L. Carlton, P. L. Feldman, A. S. Goetz, M. K. Grizzle, Y. C. Guo, A. L. Handlon, D. L. Hertzog, D. M. Ignar, D. G. Lang, R. J. Ott, A. J. Peat, H.-Q. Zhou, *J. Med. Chem.* **2006**, 49, 7095–7107.
- [28] a) A. V. Shastin, I. V. Golubinskii, O. N. Lenkova, E. S. Balenkova, V. G. Nenaidenko, *Russ. J. Org. Chem.* **2006**, 42, 238–240; b) K. Schollberg, H. Schaefer, K. Gewald, *J. Prakt. Chem.* **1983**, 325, 876–879.
- [29] a) A. M. Redman, J. S. Johnson, R. Dally, S. Swartz, H. Wild, H. Paulsen, Y. Caringal, D. Gunn, J. Renick, M. Osterhout, J. Kingery-Wood, R. A. Smith, W. Lee, J. Dumas, S. M. Wilhelm, T. J. Housley, A. Bhargava, G. E. Ranges, A. Shrikhande, D. Young, M. Bombara, W. J. Scott, *Bioorg. Med. Chem. Lett.* **2001**, 11, 9–12; b) W.-Y. Ren, K. V. B. Rao, R. S. Klein, *J. Heterocycl. Chem.* **1986**, 23, 1757–1763.
- [30] a) X. Wang, A. Studer, *J. Am. Chem. Soc.* **2016**, 138, 2977–2980; b) C. E. Stephens, M. B. Price, J. W. Sowell Sr., *J. Heterocycl. Chem.* **1999**, 36, 659–665.
- [31] H. Meerwein, E. Buchner, K. van Emster, *J. Prakt. Chem.* **1939**, 152, 237–266.
- [32] a) M. D. Obushak, M. B. Lyakhovich, M. I. Ganushchak, *Tetrahedron Lett.* **1998**, 39, 9567–9570. b) N. D. Obushak, M. B. Lyakhovich, E. E. Bilaya, *Russ. J. Org. Chem.* **2002**, 38, 38–46; c) N. D. Obushak, *Russ. J. Gen. Chem.* **1998**, 68, 443–445.
- [33] a) S. Kindt, M. R. Heinrich, *Synthesis* **2016**, 48, 1597–1606; b) P. Schroll, D. P. Hari, B. König, *ChemistryOpen* **2012**, 1, 130–133; c) M. R. Heinrich, *Chem. Eur. J.* **2009**, 15, 820–833; d) S. E. Vaillard, B. Schulte, A. Studer in *Modern Arylation Methods* (Ed. L. Ackermann) Wiley-VCH, Weinheim, 2009, pp 475–511; e) S. K. Fehler, M. R. Heinrich, *Synlett* **2015**, 26, 580–603.
- [34] a) S. J. F. Macdonald, T. C. McKenzie, W. D. Hassen, *J. Chem. Soc., Chem. Commun.* **1987**, 1528–1530; b) L.-F. Yu, Y.-Y. Li, M.-B. Su, M. Zhang, W. Zhang, L.-N. Zhang, T. Pang, R.-T. Zhang, B. Liu, J.-Y. Li, J. Li, F.-J. Nan, *ACS Med. Chem. Lett.* **2013**, 4, 475–480; c) C. Shinji, S. Maeda, K. Imai, M. Yoshida, Y. Hashimoto, H. Miyachi, *Bioorg. Med. Chem.* **2006**, 14, 7625–7651; d) T. Taniguchi, A. Ishita, M. Uchiyama, O. Tamura, O. Muraoka, G. Tanabe, H. Ishibashi, *J. Org. Chem.* **2005**, 70, 1922–1925; e) T. Taniguchi, K. Iwasaki, M. Uchiyama, O. Tamura, H. Ishibashi, *Org. Lett.* **2005**, 7, 4389–4390.
- [35] a) S. Batsyts, E. G. Hübner, J. C. Namyslo, M. Gjikaj, A. Schmidt, *Org. Biomol. Chem.* **2019**, 17, 4102–4114; b) S. Batsyts, R. Vedmid, J. C. Namyslo, M. Nieger, A. Schmidt, *Eur. J. Org. Chem.* **2019**, 1301–1310; c) A. Schmidt, S. Batsyts, A. Smeyanov, T. Freese, E. G. Hübner, M. Nieger, *J. Org. Chem.* **2016**, 81, 4202–4209; d) N. T. Pokhodyo, O. Ya. Shyyka, M. D. Obushak, *Synth. Commun.* **2014**, 44, 1002–1006; e) N. T. Pokhodyo, V. S. Matiychuk, M. D. Obushak, *Tetrahedron* **2009**, 65, 2678–2683.
- [36] a) A. Rahimi, J. C. Namyslo, M. Drafz, J. Halm, E. Hübner, M. Nieger, N. Rautenberg, A. Schmidt, *J. Org. Chem.* **2011**, 76, 7316–7325; b) A.-L. Lücke, S. Wiechmann, T. Freese, Z. Guan, A. Schmidt, *Z. Naturforsch.* **2016**, 71b, 643–650; c) A.-L. Lücke, S. Wiechmann, T. Freese, A. Schmidt, *Synlett* **2017**, 28, 1990–1993.
- [37] a) V. S. Matiychuk, R. L. Martyak, N. D. Obushak, Y. V. Ostapiuk, N. I. Pidlypyni, *Chem. Heterocycl. Compd.* **2004**, 40, 1218–1219; b) N. D. Obushak, V. S. Matiichuk, R. Ya. Vasylshyn, V. Ostap'yuk, *Russ. J. Organ. Chem.* **2004**, 40, 383–389; c) Y. V. Ostapiuk, M. D. Obushak, V. S. Matiychuk, M. Naskrent, A. K. Gzella, *Tetrahedron Lett.* **2012**, 53, 543–545; d) Y. V. Ostapiuk, V. S. Matiichuk, N. I. Pidlypyni, N. D. Obushak, *Russ. J. Organ. Chem.* **2012**, 48, 519–522; e) V. S. Matiychuk, N. D. Obushak, N. I. Pidlypyni, Y. V. Ostapiuk, R. M. Voloshchuk, *Chem. Heterocycl. Compd.* **2010**, 46, 495–499; f) M. D. Obushak, V. V. Karpyak, Y. V. Ostapiuk, V. S. Matiychuk, *Phosphorus Sulfur Silicon Relat. Elem.* **2007**, 182, 1437–1445; g) Y. V. Ostapiuk, V. S. Matiichuk, M. D. Obushak, *Russ. J. Organ. Chem.* **2017**, 53, 479–480; h) V. V. Turtytsya, Y. V. Ostapiuk, V. V. Matiychuk, M. D. Obushak, *J. Heterocyclic. Chem.* **2014**, 51, 1898–1901; i) M. Fizer, V. Sidey, A. Tupys, Y. Ostapiuk, O. Tymoshuk, Y. Bazel, *J. Mol. Struct.* **2017**, 1149, 669–682; j) N. S. Finiuk, V. P. Hreniuh, Y. V. Ostapiuk, V. S. Matiychuk, D. A. Frolov, M. D. Obushak, R. S. Stoika, A. M. Babskiy, *Biopolym. Cell.* **2017**, 33, 135–146; k) R. Z. Lytvyn, A. O. Neshchadim, Kh. Y. Pitkovich, Yu. I. Horak, J. V. Grazulevicius, T. Lis, V. Kinzhylbalo, M. D. Obushak, *Tetrahedron Lett.* **2016**, 57, 118–121.
- [38] a) E. Matveeva, A. S. Erin, A. G. Osterov, I. F. Leshcheva, A. L. Kurts, *Russ. J. Org. Chem.* **2006**, 42, 388–392.


FULL PAPER Accepted Article

- [39] S. M. Islam, A. S. Roy, R. C. Dey, S. Paul, *J. Mol. Catal. A* **2014**, *394*, 66–73.
- [40] J. R. Falck, A. Bandyopadhyay, D. K. Barma, D.-S. Shin, A. Kundu, R. V. Krishna Kishore, *Tetrahedron Lett.* **2004**, *45*, 3039–3042.
- [41] Y.-F. Zeng, W.-W. Ji, W.-X. Lv, Y. Chen, D.-H. Tan, Q. Li, H. Wang, *Angew. Chem. Int. Ed.* **2017**, *56*, 14707–14711.
- [42] J. N. Kim, J. S. Son, H. R. Kim, E. K. Ryu, *Bull. Korean Chem. Soc.* **1998**, *19*, 812–813.
- [43] K. Schollberg, H. Schäfer, K. Gewald, *J. Prakt. Chem.* **1983**, *325*, 876–879.
- [44] J. A. Hyatt, P. W. Raynolds, *Org. React.* **1994**, *45*, 159–646.
- [45] R. Romagnoli, M. Kimatrai Salvador, S. Schiaffino Ortega, P. G. Baraldi, P. Oliva, S. Baraldi, L. C. Lopez-Cara, A. Brancale, S. Ferla, E. Hamel, J. Balzarini, S. Liekens, E. Mattiuzzo, G. Basso, G. Viola, *Eur. J. Med. Chem.* **2018**, *143*, 683–698.
- [46] G. R. Gokaraju, R. R. Gokaraju, S. Kasina, T. Golakoti, V. Sompalli, S. Krishnan, K. Bhupathiraju, IN 2008CH02230
- [47] F. X. Tavares, K. A. Al-Barazanji, E. C. Bigham, M. J. Bishop, C. S. Britt, D. L. Carlton, P. L. Feldman, A. S. Goetz, M. K. Grizzle, Y. C. Guo, A. L. Handlon, D. L. Hertzog, D. M. Ignar, D. G. Lang, R. J. Ott, A. J. Peat, H. Q. Zhou, *J. Med. Chem.* **2006**, *49*, 7095–7107.
- [48] J. H. Van Duzer, R. Mazitschek, US 20140128391 A1
- [49] M. H. Norman, N. Chen, N. Han, L. Liu, C. R. Hurt, C. H. Fotsch, T. J. Jenkins, O. A. Moreno, WO 9940091 A1
- [50] W. Y. Ren, K. V. B. Rao, R. S. Klein, *J. Heterocycl. Chem.* **1986**, *23*, 1757–1763.
- [51] Agilent Technologies, CrysAlisPro, Version 1.171.40.45a.
- [52] a) G. M. Sheldrick, *Acta Cryst. C* **2015**, *71*, 3–8; b) G. Sheldrick, *Acta Cryst. A* **2008**, *64*, 112–122; c) O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard, H. Puschmann, *J. Appl. Crystallogr.* **2009**, *42*, 339–341.
- [53] H. W. Moore, F. Mercer, D. Kunert, P. Albaugh, *J. Am. Chem. Soc.* **1979**, *101*, 5435–5436.
- [54] J. H. Sahner, M. Empting, A. Kamal, E. Weidel, M. Groh, C. Börger, R. W. Hartmann, *Eur. J. Med. Chem.* **2015**, *96*, 14–21.
- [55] A. Cohen, P. Suzanne, J.-C. Lancelot, P. Verhaeghe, A. Lesnard, L. Basmaciyan, S. Hutter, M. Laget, A. Dumetre, L. Paloque, E. Deharo, M. D. Crozet, P. Rathelot, P. Dallemande, A. Lorthiois, C. H. Sibley, P. Vanelle, A. Valentin, D. Mazier, S. Rault, N. Azas, *Eur. J. Med. Chem.* **2015**, *95*, 16–28.
- [56] J. H. Sahner, M. Groh, M. Negri, J. Haupenthal, R. W. Hartmann, *Eur. J. Med. Chem.* **2013**, *65*, 223–231.
- [57] D. Thomae, J. C. Rodriguez Dominguez, G. Kirsch, P. Seck, *Tetrahedron* **2008**, *64*, 3232–3235.
- [58] D. Thomae, G. Kirsch, P. Seck, T. Kaminski, *Synthesis* **2007**, *14*, 2153–2156.

Accepted Manuscript

FULL PAPER Accepted Article

FULL PAPER



The title compounds are available by Meerwein reaction and proved to be useful starting materials for the synthesis of thiophenes.

Heterocycle synthesis

S. Batsyts, M. Shehedyn, E. Goreshnik, M. D. Obushak, A. Schmidt*, and Yu. Ostapiuk*

Page No. – Page No.
2-Bromo-2-chloro-3-
arylpropanenitriles as C-3 Synths
for the Synthesis of Functionalized 3-
Aminothiophenes

Accepted Manuscript