

Accepted Manuscript

Synthesis, antibacterial and antitubercular activities of benzimidazole bearing substituted 2-pyridone motifs

N.C. Desai , N.R. Shihory , G.M. Kotadiya , Priyanka Desai

PII: S0223-5234(14)00522-4

DOI: [10.1016/j.ejmech.2014.06.004](https://doi.org/10.1016/j.ejmech.2014.06.004)

Reference: EJMECH 7046

To appear in: *European Journal of Medicinal Chemistry*


Received Date: 3 February 2014

Revised Date: 5 April 2014

Accepted Date: 3 June 2014


Please cite this article as: N. Desai, N. Shihory, G. Kotadiya, P. Desai, Synthesis, antibacterial and antitubercular activities of benzimidazole bearing substituted 2-pyridone motifs, *European Journal of Medicinal Chemistry* (2014), doi: 10.1016/j.ejmech.2014.06.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Synthesis, antibacterial and antitubercular activities of benzimidazole bearing substituted 2-pyridone motifs

N C Desai*, N R Shihory, G M Kotadiya, Priyanka Desai



- A series of benzimidazole bearing substituted 2-pyridones were synthesized.
- All the compounds were subjected for their *in vitro* biological screening.
- Some of the tested compounds exhibited higher potency compared to the standard.
- None of the tested compounds exhibited significant cytotoxic activity.
- Lipophilicity plays an important role in antibacterial effect of tested compounds.

Synthesis, antibacterial and antitubercular activities of benzimidazole bearing substituted 2-pyridone motifs

N C Desai*, N R Shihory, G M Kotadiya, Priyanka Desai
Division of Medicinal Chemistry, Department of Chemistry
(UGC NON-SAP & DST-FIST Sponsored),
Mahatma Gandhi Campus, Maharaja Krishnakumarsinhji Bhavnagar University,
Bhavnagar 364 002, India
E-mail: dnisheeth@rediffmail.com,
Telefax: +91 278 2439852

Abstract

A series of benzimidazole bearing 2-pyridones **5a-r** were synthesized and evaluated for their *in vitro* antibacterial and antitubercular activity. Further, all compounds were examined for their cytotoxic study on VERO cell line and characterized by well-known spectral techniques. It was observed that the compounds **5h**, **5i**, **5k**, **5q** and **5r** were found to possess significant broad spectrum antibacterial activity (12.5-100 µg/mL of MIC), while compounds **5g-5i**, **5k** and **5l** proved to be the most potent antitubercular activity in range of 2.76-20.4 µM of MIC at low level of cytotoxicity, indicating good selectivity. From SAR studies, lipophilic profile of compounds was remarkably vital for antibacterial activity, while MIC values of antitubercular could not be directly correlate with lipophilicity.

Keywords: Antibacterial activity; Antitubercular activity; Benzimidazole; Cytotoxicity; MABA assay; 2-Pyridone

1. Introduction

The current first-line tuberculosis drug treatment is more than 40 years old and consists primarily of rifampicin and isoniazid. These antibiotics are drug-susceptible and require longer time and large number of doses, which are multi-drug resistant (MDR) and extensively drug resistant (XDR) to tuberculosis strains [1-3]. However, the rapid increase of multi-drug-resistant tuberculosis (MDR-TB) (resistant to at least isoniazid and rifampicin) and extensively drug-resistant tuberculosis (XDR-TB) (resistant to isoniazid, rifampicin in addition to fluoroquinolone, kanamycin, amikacin or capreomycin among second line anti-TB drugs) has led to an urgent need for the identification of new drug targets and the growth of novel anti-TB drugs. Furthermore, The most commonly encountered antibiotic-resistant bacteria, methicillin-resistant *Staphylococcus aureus* (MRSA), has a major impact on infections in both hospitals and community settings [4,5]. Unfortunately, as antibiotic resistant organisms have become more commonplace, the pipeline for the discovery of new antimicrobial agents has decreased [6]. Thus, there is a pressing need for new antimicrobial agents that are capable of treating resistant bacterial strains.

Moreover, dihydrofolate reductase (DHFR) catalyzes the NADPH-dependent reduction of dihydrofolate to tetrahydrofolate that is essential for DNA synthesis. Inhibition of its activity leads to arrest of DNA synthesis and hence cell death [7]. *Mycobacterium tuberculosis* DHFR is an attractive novel drug target for developing anti-tuberculosis drugs. To overcome this problem, we have synthesized a series of benzimidazole bearing 2-pyridones by replacing the pyrazole motif in our previously synthesized compounds **NCD₁₋₂₀** [8] and screened them for their antibacterial property. In this attempt, we got excellent antibacterial results. Antibacterial studies

impelled us to inspect **5a-r** for their *in vitro* antitubercular activity as well. Structural relevance of title compounds **5a-r** with previously synthesized compounds is shown in **Figure 1**.

Benzimidazole nucleus is a the key building block for numerous compounds that play beneficial roles in the functioning of biologically important molecules [9] and are remarkably effective both with respect to their inhibitory activity and their favorable selectivity ratio [10-12]. Benzimidazoles are considered a promising class of bioactive heterocyclic compounds encompassing a diverse range of biological activities such as antiulcer [13], antihelminthic [14], antihypertensive [15], anticoagulant [16], anti-inflammatory [17], antimicrobial [18-20] and antiparasitic [21]. The azole group of heterocyclic compounds possesses significant pharmacokinetic profile and lipophilicity that influence the ability of drug to reach the target by transmembrane diffusion and along with promising activity against resistant TB by inhibiting the biosynthesis of lipids [22,23]. Further, 2-pyridones represent a unique class of pharmacophores, which are observed in various therapeutic agents [24]. In recent years, 2-pyridones have assimilated much importance as these compounds exhibit several biological activities such as antitumoral [25], antimalarial [26], analgesic [27] and anti-HIV [28] properties. Moreover, 2-pyridones are a class of recently discovered potent antibacterial agents that are of particular interest due to their *in vitro* and *in vivo* antibacterial potencies against the bacterial type II DNA topoisomerases, which include two highly homologous enzymes-DNA gyrase and topoisomerase IV [29,30]. Moreover, among the pharmacokinetic properties, a low and highly variable bioavailability is indeed the main reason for stopping further development of the drug [31].

Motivated by the above findings and from our previous work [32,33], the main aim of the work is to obtain more active antibacterial and antitubercular agents with plausible novel mechanisms of action. It was thought worthwhile to synthesize some new benzimidazole bearing

2-pyridone derivatives comprising of the above aforementioned moieties in single molecular framework in order to investigate their *in vitro* antibacterial and antitubercular activity. In continuation to this, in our present communication, we have synthesized benzimidazole bearing 2-pyridones **5a-r** and evaluated for their *in vitro* antibacterial and antitubercular activity. In addition, cytotoxicity studies were also conducted in VERO cell lines to evaluate the ability of these compounds to inhibit the cell growth. Most active compounds **5h**, **5q** and **5r** were also screened against MRSA strain.

{Insert Figure 1 here}

2. Results and discussion

2.1. Chemistry

We have synthesized new analogues in which 2-pyridone motif is connected to the benzimidazole system. The synthetic strategies adopted for the synthesis of target benzimidazole bearing 2-pyridone derivatives **5a-r** are depicted in **Schemes 1** and **2**. In **Scheme 1**, condensation of 1-(1*H*-benzo[*d*]imidazol-2-yl)ethanone **1** with equimolar quantity of cyanoacetic acid hydrazide in refluxing 1,4-dioxane afforded a single product, that was identified as *N'*-(1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)-2-cyanoacetohydrazide **2**. The elemental analysis and spectral data were in accordance with the proposed *N'*-(1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)-2-cyanoacetohydrazide **2** structure. The IR spectrum of **2** showed strong absorption bands at 2248 and 1681 cm⁻¹ were due to cyanide and carbonyl group, respectively. Its ¹H NMR spectrum apart from the expected aromatic signals, showed two new singlets at δ 3.38 and 8.92 ppm assigned to the reactive methylene protons and proton of secondary amine attached with carbonyl group respectively. The ¹³C NMR spectrum displayed nine carbon signals, the most important signals

92 appeared at δ 13.4, 27.3, 125.8, 171.3 ppm characteristic of methyl, methylene, cyanide and
 93 carbonyl carbons, respectively. The mass spectrum revealed a molecular ion peak at m/z =
 94 241.11 ($M + 1$), in agreement with its proposed structure. The presence of the reactive methylene
 95 group in the hydrazide **2** makes it a versatile precursor for the Michael type condensation with
 96 Knoevenagel product *p*-nitrobenzaldehyde and malononitrile compound **3** in presence of
 97 catalytic amount of piperidine utilizing ethanol (95%) as a solvent furnished 2-pyridone
 98 derivative identified as 1-((1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)amino)-6-amino-4-(4-
 99 nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile **4**. Structure of the latter product was
 100 confirmed by IR spectra which showed characteristic absorption bands at 1688 and 3446 cm^{-1} for
 101 conjugated $>\text{C}=\text{O}$ and primary amine group respectively. Their ^1H NMR spectra displayed a
 102 broad singlet at δ 8.78 ppm for primary amine protons, besides the disappearance of reactive
 103 methylene group and secondary amine singlets due to its involvement in cyclization.
 104 Condensation of 2-pyridone derivative **4** with appropriate aromatic aldehydes in boiling ethanol
 105 afforded the respective targeted benzimidazole bearing 2-pyridones acknowledged as 1-((1-(1*H*-
 106 benzo[*d*]imidazol-2-yl)ethylidene)amino)-6-((arylbenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-
 107 1,2-dihydropyridine-3,5-dicarbonitriles **5a-r**. The structures of the final analogues **5a-r** were
 108 established by IR spectra which showed disappearance of $-\text{NH}_2$ band in compound **4**. A strong
 109 absorption band appeared between 1679-1688 cm^{-1} was assigned to conjugated $>\text{C}=\text{O}$ group.
 110 Their ^1H NMR spectra revealed a singlet of methine proton at δ 9.40-9.57 ppm involved in
 111 azomethine formation, along with the vanishing of primary amine singlet. The ^{13}C NMR
 112 spectrum of compound **5l**, displayed, besides the expected methyl and aromatic signals, three
 113 characteristic signals at δ 115.9, 160.1 and 163.8 ppm due to the carbons of CN, $\text{C}=\text{O}$ and $\text{CH}=\text{N}$
 114 respectively. The mass spectrum of **5l** showed molecular ion peak at m/z = 540.17 ($M + 1$), in

agreement with its proposed structure. Similarly, the spectral values for all the compounds and C, H, N analysis are presented in the experimental part.

{Insert Scheme 1 here}

A plausible mechanistic pathway for the formation of compounds **5a-r** is suggested in **Scheme 2**. Firstly, hydrazone (**A**) underwent Michael addition with Knoevenagel product (**B**) and furnished the intermediate (**C**), which further experienced intramolecular nucleophilic attack on cyanide carbon followed by annulation to yield intermediate (**D**). The intermediate (**D**) transformed to compound (**E**) by intramolecular electron transfer to nitrogen atom. In the last step, intermediate (**E**) was transformed to targeted compounds by intermolecular nucleophilic attack on carbonyl carbon of different aromatic aldehydes.

{Insert Scheme 2 here}

2.2. Biological Evaluation

2.2.1. Antibacterial activity

The intermediates **2**, **4** and target compounds **5a-r** were investigated for their *in vitro* antibacterial activity against two gram-positive and two gram-negative bacteria and the yeast like pathogenic fungus *Candida albicans* using conventional broth-dilution method [34,35]. Minimum inhibitory concentration (MIC) is defined as the concentration of the compound required to obtain complete inhibition of bacterial growth. MICs of the synthesized compounds were compared with ciprofloxacin and chloramphenicol and the results were depicted in **Table 1**. The antibacterial activity assessed (**Table 1**) for analogues **2**, **4** and **5a-r** against several strains flaunted good activity. From the bioassay, it was observed that the final analogues **5a-r** with the

substitutions of phenyl ring were most active against all the pathogenic strains studied than the intermediate hydrazone **2** and 2-pyridone derivative **4**. Intermediate **2** showed very poor antibacterial activity against all tested bacterial strains and it was observed that compound **2** with low lipophilicity ($Clog P = 0.7220$) displayed very low activity as compared to intermediate **4** and final analogues **5a-r**. Precursor **2** reacted with Knoevenagel compound **3** to generate intermediate **4** which was tested against different bacterial strains, and it showed more inhibitory efficacy than compound **2** due to high lipophilicity ($Clog P = 1.4772$) and less activity compared to all final analogues. Therefore, it was observed that formation of 2-pyridone nucleus was responsible for enhancing antibacterial potency. Now, intermediate **4** was treated with different aromatic aldehydes to afford targeted compounds **5a-r** which were found to have broad spectrum antimicrobial efficacy. This could be correlated to structural variations and different aromatic substitutions in phenyl ring.

Good bioavailability can be achieved by an appropriate balance between solubility and partitioning properties. Membrane permeability and bioavailability is always associated with some basic molecular descriptor such as $Clog P$ (lipophilicity). Thus, prediction of bioavailability-related properties, such as lipophilicity is important before actual synthesis, in order to reduce enormous wastage of expensive chemicals and precious time. The computed $Clog P$ values (P is the partition coefficient of the molecule in the water/octanol system) are shown in **Table 1** and **3**. The ChemBioDraw Ultra, version 12.0, software by Cambridge Soft is a program used to predict lipophilicity of compounds.

From the bioassay, final analogues with electron withdrawing halogen substitution were found to be more active than the remaining final analogues against all the bacterial strains. Among all the final active analogues, compound **5q** (3-Br, $Clog P = 3.9272$) endowed with

159 bromine exerted highest inhibition against all the bacterial strains and showed highest efficacy
 160 against *S. aureus* at 12.5 µg/mL of MIC as compared to standards ciprofloxacin (50 µg/mL, Clog
 161 $P = -0.7252$) and chloramphenicol (50 µg/mL, Clog $P = 1.293$). In addition, compound **5q** (Clog
 162 $P = 3.9272$) with higher lipophilicity flaunted potential inhibitory action against *E. coli* and *S.*
 163 *pyogenes* at MIC of 25 µg/mL. Moreover, the above mentioned derivative **5q** displayed excellent
 164 inhibition against *P. aeruginosa* at 25 µg/mL of MIC. Furthermore, among the second line active
 165 compounds, **5h** (3-Cl, Clog $P = 3.7772$) and **5r** (4-Br, Clog $P = 3.9272$) indicated MIC of 50
 166 µg/mL, in which more lipophilic compound **5r** showed better results against both *E. coli* and *S.*
 167 *pyogenes*. In addition, the above mentioned derivatives **5h** and **5r** were also found to
 168 demonstrate excellent efficacy against *S. aureus* at 25 µg/mL of MIC and analogue **5r** also
 169 demonstrated MIC of 50 µg/mL against *P. aeruginosa*. Derivatives **5e** (3-F, Clog $P = 3.2072$)
 170 with fluoro, **5g** (2-Cl, Clog $P = 3.7772$) with chloro and **5k** (3-CH₃, Clog $P = 3.5632$) with
 171 methyl group to the aromatic nucleus bequeathed appreciable potency towards *E. coli* at 100
 172 µg/mL of MIC, in which compound **5g** showing highest lipophilicity gave better results.
 173 Compound **5h** (3-Cl, Clog $P = 3.7772$) with electron withdrawing chlorine substituent indicated
 174 diminished activity at 62.5 µg/mL of MIC against *P. aeruginosa*. Another analogue **5i** (4-Cl,
 175 Clog $P = 3.7772$) with chlorine and **5k** (3-CH₃, Clog $P = 3.5632$) endowed with electron
 176 donating methyl group exerted MIC of 100 µg/mL against *P. aeruginosa* as compared to
 177 standard ciprofloxacin (25 µg/mL, Clog $P = -0.7252$) and chloramphenicol (50 µg/mL, Clog $P =$
 178 1.293) antibiotics. Derivative **5g** (2-Cl, Clog $P = 3.7772$) displayed more potency against *S.*
 179 *aureus* than *E. coli* at MIC of 50 µg/mL and diminished activity at MIC of 62.5 µg/mL against *S.*
 180 *pyogenes*. Final derivatives **5c** (3-OH, Clog $P = 3.2982$), **5e** (3-F, Clog $P = 3.2072$), **5k** (3-CH₃,
 181 Clog $P = 3.5632$) and **5m** (3-OCH₃, Clog $P = 3.4082$) displayed remarkable inhibitory effects at

MIC of 100 µg/mL against *S. aureus*. Among them compound **5k** contemplated same action against *S. pyogenes*. In addition, compound **5i** (4-Cl, Clog *P* = 3.7772) exhibited activity at MIC of 50 and 62.5 µg/mL against *S. pyogenes* and *S. aureus* respectively. Moreover, compound **5e** possessing fluorine atom at 3rd position of phenyl ring increased the potency and showed inhibition at 62.5 µg/mL of MIC against *S. pyogenes*. Furthermore, the inhibitory activity of all the synthesized compounds against *C. albicans* was rather lower than their antibacterial activity, only compound **5q** (3-Br, MIC = 250 µg/mL) exhibited moderate activity as compared to standard ketoconazole.

{Insert Table 1 here}

Furthermore, the most active compounds **5h**, **5q** and **5r** against *S. aureus* and *S. pyogenes* (MIC = 12.5-50 µg/mL) were also tested against methicillin-resistant *S. aureus* (MRSA isolate ATCC 43300) and results are given in **Table 2**. Compound **5r** exhibited more potent activity than the standard drugs against methicillin-resistant *S. aureus*. Compound **5r**, with MIC value of 6.25 µg/mL against MRSA showed four-fold more potency than ciprofloxacin (MIC = 25 µg/mL) and eight-fold more activity than chloramphenicol (MIC = 50 µg/mL). In addition, compound **5q** endowed with bromo group showed two-fold more inhibition at MIC value of 12.5 µg/mL than ciprofloxacin and four-fold higher potency than chloramphenicol against MRSA.

{Insert Table 2 here}

2.2.2. Antitubercular activity

The encouraging results from the antibacterial studies impelled us to go for the screening of title compounds for their *in vitro* antitubercular activity. Intermediates (**2** and **4**) and all the

final compounds along with the standard drug for comparison were firstly evaluated for their activity against the *Mycobacterium tuberculosis* H₃₇Rv strain in Middlebrook 7H12 medium using Microplate Alamar Blue Assay (MABA) MIC method [36]. The drug in clinical use, isoniazid was used as a reference drug. The results of actual MICs of tested compounds were reported in **Table 3**. It was observed that final analogues **5a-r** displayed superior anti-tubercular activity compared to both the intermediates (**2** and **4**). Among the eighteen synthesized compounds, compounds **5g-5i**, **5k** and **5l** endowed with inductively electron withdrawing chlorine and electron donating methyl groups afforded maximum MICs ranging from 0.85 to 6.28 µg/mL against *Mycobacterium tuberculosis*. For final derivatives, introduction of chlorine substituent in **5g-5i** was tolerated, 3-chloro yielding the best activity (3-Cl > 4-Cl > 2-Cl). Furthermore, among the second line active compounds, **5k** and **5l** brandished MICs in range of 2.07-6.28 µg/mL, in which compound **5k** having methyl in *meta* position showed better result at MIC of 2.07 µg/mL whereas compound **5l** displayed inhibition at MIC of 6.28 µg/mL. Compound **5h** (3-Cl) possessed highest inhibition at MIC of 0.85 µg/mL amongst all the tested derivatives as compared to standard isoniazid, while their *ortho* and *para* derivatives **5i** (4-Cl) and **5g** (2-Cl) showed MICs of 1.79 and 2.65 µg/mL respectively. Here, we have discussed and compared antitubercular activity based on first line drug Isoniazid (0.24 µg/mL, Clog *P* = -0.668). From the antitubercular activity results, it may be concluded that final derivatives **5a-o** were mixdly active and no specific relationship was observed in case of Clog *P* or lipophilicity verses antitubercular activity profiles.

2.2.3. Cytotoxicity study

After having identified numerous active antibacterial and antimycobacterial benzimidazole bearing 2-pyridones, the next step was to examine the toxicity of the drug

contenders. *In vitro* cytotoxicity of compounds **5a-r** was determined against mammalian VERO cell lines [37]. After 72 h of exposure, viability was considered on the basis of cellular conversion of [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium and phenylmethasulfazone] (100:20) into formazan product using Promega Cell Titer 96 non-radioactive cell proliferation assay. The cytotoxicity results presented in **Table 3** are expressed as the concentration inhibiting 50% of the cell growth IC_{50} . The compounds exhibited moderate to low level of cytotoxicity with the IC_{50} values in the range of 13.73 - >39.42 $\mu\text{g/mL}$. It was observed that none of the tested compounds exhibited any significant cytotoxic effects on VERO cells, suggesting a great potential for their *in vivo* use as antibacterial and anti-tubercular agents. As for activity against VERO cell lines, the highest cytotoxic activity was flaunted by compound **5d** which showed percentage viability IC_{50} at 13.73 $\mu\text{g/mL}$, whereas, the second line cytotoxic activity was displayed by compounds **5a**, **5c** and **5p** which showed the percentage viability IC_{50} at 17.61, 17.00 and 16.05 $\mu\text{g/mL}$ respectively. Moreover, final analogues **5g-5i**, **5k**, **5l**, **5q** and **5r** showed no toxicity for the percentage viability $IC_{50} > 39.42 \mu\text{g/mL}$.

{Insert Table 3 here}

2.3. Structure-activity relationship

The results of antibacterial and antitubercular screening demonstrated the following assumptions about the structural activity relationship (SAR): in the present study, we investigated the effects of the substitution pattern of the hybrid benzimidazole and 2-pyridone derivatives were carefully selected to impart different electronic environment on the molecules. By comparing the antibacterial activity of the synthesized compounds, it was found that the tested compounds were more effective against Gram-positive bacteria. In addition, the

antibacterial activity was considerably affected by substitution pattern on the phenyl ring and lipophilic profile of the compounds. It is believed that the strong lipophilic character of the molecule plays an essential role in producing antibacterial effect. These properties are seen as an important parameter related to membrane permeation in biological system. Many processes of drug disposition depend on the capability to cross membranes and hence there is a high correlation with measures of lipophilicity. Lipophilicity plays a major role in determining where drugs are distributed within the body after adsorption and as a consequence how rapidly they are metabolized and excreted. In this context, the presence of lipophilic moiety would be important for such activity. In case of antibacterial activity, some analogues of this series were found to have even more potency than the standard drug 'chloramphenicol' while some of them exhibited comparable potency. Compounds **5h**, **5i**, **5k**, **5q** and **5r** bearing Cl, Br, CH₃ groups were found to be most potent antibacterial agents due to its high lipophilicity. Among them, compound **5q** endowed with inductively electron withdrawing bromine at *meta* position emerged as the most effective antibacterial agent at MIC of 12.5 µg/mL. The MIC values of these novel compounds confirmed that the presence of bromine, chlorine and methyl substituent at *meta* position gave rise to better pharmacological potency rather than the same substituent present at *ortho* or *para* positions. Whereas, the opposite trend was observed for antitubercular activity results and there was no specific relationship observed between MIC profiles versus lipophilicity. Among the eighteen synthesized compounds, **5g-5i**, **5k** and **5l** endowed with inductively electron withdrawing chlorine and electron donating methyl group illustrated maximum MICs in the range of 0.85 to 6.28 µg/mL against *Mycobacterium tuberculosis*. Out of them, 3-chloro yielded the best activity at MIC of 0.85 µg/mL as compared to first line drug isoniazid (0.24 µg/mL).

3. Conclusion

The focal point of the current work was on the development of new bioactive based on benzimidazole clubbed 2-pyridone compact system with the hope of generating new bioactive chemical entities that could be useful as potent antibacterial and antitubercular agents. Many of the synthesized motifs (**5h**, **5i**, **5k**, **5q** and **5r**), possessing atom/group such as bromo, chloro and methyl at *meta* or *para* positions were identified as the most compelling antibacterial agents. Albeit, it was observed that the strong lipophilic character of the molecule plays an imperative role in producing antibacterial effect. Compounds **5g-5i**, **5k** and **5l** came out as the most promising antitubercular agents. In addition, the relationship between activity profiles and lipophilicity of the newer analogues was also discussed in which higher lipophilic compounds showed higher bioactivities, while, in case of antitubercular activity, there was no specific relationship observed between MIC profiles versus lipophilicity. Moreover, the potent antibacterial and antitubercular activity of most active compounds **5g-5i**, **5k**, **5l**, **5q** and **5r** were accompanied with relatively low level of cytotoxicity. Consequently, the degree of activity and the encouraging physicochemical parameters displayed by a novel innovative structural combination system of benzimidazole and 2-oxopyridine rings make such compounds a privileged structure to achieve more active derivatives in ongoing studies. Synthesized compounds were screened against *C. albicans*, but it was our observation that none of the compounds exhibited significant activity against this fungi.

4. Experimental

4.1. General Methods

All reactions except those in aqueous media were carried out by standard techniques for the exclusion of moisture. Melting points were determined on an electro thermal melting point

apparatus and were reported uncorrected. TLC on silica gel plates (Merck, 60, F₂₅₄) was used for purity checking and reaction monitoring. Column chromatography on silica gel (Merck, 70-230 mesh and 230-400 mesh ASTH for flash chromatography) was applied when necessary to isolate and purify the reaction products. Elemental analysis (% C, H, N) was carried out by a Perkin-Elmer 2400 CHN analyzer. IR spectra of all compounds were recorded on a Perkin-Elmer FT-IR spectrophotometer in KBr. ¹H NMR and spectra were recorded on Varian Gemini 400 MHz and ¹³C NMR spectra on Varian Mercury-400, 100 MHz in DMSO-*d*₆ as a solvent and tetramethylsilane (TMS) as an internal standard. Mass spectra were scanned on a Shimadzu LC-MS 2010 spectrometer. Anhydrous reactions were carried out in oven-dried glassware in nitrogen atmosphere.

4.2. General procedure for the synthesis of *N'*-(1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)-2-cyanoacetohydrazide (**2**)

A mixture of equimolar amount of acetyl benzimidazole compound **1** (0.01 mol) and cyanoacetic acid hydrazide (0.01 mol) in 1,4-dioxane (50 mL) was refluxed for 3 h. The reaction mixture was concentrated by evaporating to dryness under reduced pressure and then cooled down to room temperature. The separated crystals were filtered, air dried and recrystallized from absolute alcohol. Yield 70%; mp 165 °C. IR (ν_{max}, cm⁻¹, KBr): 3371 (-NH, benzimidazole), 3278 (-NH, -CONH-), 3028 (C-H, aromatic), 2917 (C-H, CH₃), 2248 (CN), 1681 (CO), 1641 (C=C), 1532 (C=N). ¹H NMR (400 MHz, DMSO-*d*₆, δ, ppm): 1.88 (s, 3H, CH₃), 3.38 (s, 2H, -CH₂CN), 7.20 (d, 2H, *J* = 8.0 Hz, Ar-H), 7.61 (d, 2H, *J* = 8.2 Hz, Ar-H), 8.92 (s, 1H, -NHCO- D₂O exch.), 10.32 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ, ppm): 171.3, 155.7, 151.6, 134.9, 125.8, 123.2, 115.1, 27.3, 13.4. LCMS (ESI): *M/Z* = 241.11 [*M*⁺]. Anal. Calcd. for C₁₂H₁₁N₅O: C, 59.74; H, 4.60; N, 29.03. Found: C, 59.69; H, 4.55; N, 29.09.

4.3. General procedure for the synthesis of 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)-amino)-6-amino-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**4**)

A mixture containing compound **2** (0.01 mol) and Knoevenagel compound 2-(4-nitrobenzylidene)malononitrile **3** (0.01 mol) in ethanol (50 mL) was refluxed for 3 h using 2 drops of piperidine as catalyst. The excess of solvent was distilled out and the mixture was then cooled down to room temperature. The crystals formed were filtered, air dried and recrystallized from aqueous DMF. Yield 71%; mp 208-210 °C. IR (ν_{\max} , cm^{-1} , KBr): 3446 (NH_2), 3376 (-NH , benzimidazole), 3030 (C-H, aromatic), 2921 (C-H, CH_3), 2232 (CN), 1688 (CO), 1642 (C=C), 1536 (C=N), 1516, 1332 (N=O , Ar- NO_2). ^1H NMR (400 MHz, $\text{DMSO-}d_6$, δ , ppm): 1.91 (s, 3H, CH_3), 7.22 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.44 (d, 2H, $J = 7.6$ Hz, Ar-H), 7.62 (d, 2H, $J = 8.4$ Hz, Ar-H), 8.21 (d, 2H, $J = 7.7$ Hz, Ar-H), 8.78 (s, 2H, NH_2 D_2O exch.), 10.36 (s, 1H, -NH benzimidazole D_2O exch.). ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$, δ , ppm): 169.5, 160.2, 159.4, 155.8, 151.7, 147.2, 138.7, 134.9, 130.2, 123.1, 123.8, 115.1, 115.5, 115.9, 76.6, 13.8. LCMS (ESI): $\text{M/Z} = 438.11$ [M^+]. Anal. Calcd. for $\text{C}_{22}\text{H}_{14}\text{N}_8\text{O}_3$: C, 60.27; H, 3.22; N, 25.56. Found: C, 60.34; H, 3.25; N, 25.51.

4.4. General procedure for the synthesis of 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((arylbenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydro-pyridine-3,5-dicarbonitriles (**5a-r**)

Compound **4** (0.01 mol), different substituted aromatic aldehydes (0.01 mol) and ethanol (50 mL) were taken in a round bottom flask and refluxed for 5 h. After 5 h, the reaction mass was poured onto crushed ice and separated solid was filtered, dried and recrystallized from DMSO.

4.4.1. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-(benzylideneamino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5a)*

Yield 66%; mp 221 °C. IR (ν_{\max} , cm^{-1} , KBr): 3380 (-NH, benzimidazole), 3034 (C-H, aromatic), 2992 (C-H, CH=N), 2924 (C-H, CH₃), 2228 (CN), 1682 (CO), 1641 (C=C), 1532 (C=N), 1520, 1331 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.90 (s, 3H, CH₃), 7.20 (d, 2H, *J* = 8.1 Hz, Ar-H), 7.41 (d, 2H, *J* = 7.5 Hz, Ar-H), 7.47 (d, 2H, *J* = 7.6 Hz, Ar-H), 7.54 (t, 1H, *J* = 7.4 Hz, Ar-H), 7.63 (d, 2H, *J* = 8.3 Hz, Ar-H), 7.84 (d, 2H, *J* = 7.7 Hz, Ar-H), 8.24 (d, 2H, *J* = 7.6 Hz, Ar-H), 9.51 (s, 1H, CH=N), 10.41 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.8, 160.1, 155.7, 153.3, 151.6, 147.1, 138.6, 134.8, 133.8, 131.2, 130.1, 129.4, 128.9, 123.2, 123.8, 115.0, 115.5, 115.9, 114.7, 13.5. LCMS (ESI): *M/Z* = 526.14 [*M*⁺]. Anal. Calcd. for C₂₉H₁₈N₈O₃: C, 66.16; H, 3.45; N, 21.28. Found: C, 66.22; H, 3.49; N, 21.21.

4.4.2. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((2-hydroxybenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5b)*

Yield 69%; mp 242 °C. IR (ν_{\max} , cm^{-1} , KBr): 3441 (OH), 3382 (-NH, benzimidazole), 3031 (C-H, aromatic), 2990 (C-H, CH=N), 2924 (C-H, CH₃), 2225 (CN), 1681 (CO), 1637 (C=C), 1531 (C=N), 1522, 1330 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.92 (s, 3H, CH₃), 7.01 (d, 1H, *J* = 7.5 Hz, Ar-H), 7.11 (t, 1H, *J* = 7.6 Hz, Ar-H), 7.21 (d, 2H, *J* = 8.2 Hz, Ar-H), 7.44 (d, 2H, *J* = 7.6 Hz, Ar-H), 7.53 (t, 1H, *J* = 7.4 Hz, Ar-H), 7.62 (d, 2H, *J* = 8.4 Hz, Ar-H), 7.71 (d, 1H, *J* = 7.6 Hz, Ar-H), 8.21 (d, 2H, *J* = 7.6 Hz, Ar-H), 8.74 (s, 1H, OH D₂O exch.), 9.54 (s, 1H, CH=N), 10.42 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.6, 163.7, 160.2, 155.6, 153.1, 151.5, 147.0, 138.5, 134.8, 134.1, 133.5, 130.1,

127.9, 127.4, 123.8, 123.1, 115.9, 115.5, 115.0, 115.9, 114.8, 59.4, 13.7, 13.2. LCMS (ESI): M/Z = 542.16 [M⁺]. Anal. Calcd. for C₂₉H₁₈N₈O₄: C, 64.20; H, 3.34; N, 20.65. Found: C, 64.28; H, 3.39; N, 20.71.

4.4.3. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-hydroxybenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5c)

Yield 68%; mp 256 °C. IR (v_{max}, cm⁻¹, KBr): 3440 (OH), 3384 (-NH, benzimidazole), 3034 (C-H, aromatic), 2992 (C-H, CH=N), 2927 (C-H, CH₃), 2223 (CN), 1684 (CO), 1639 (C=C), 1534 (C=N), 1524, 1334 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ, ppm): 1.90 (s, 3H, CH₃), 7.02 (d, 1H, *J* = 7.6 Hz, Ar-H), 7.18 (d, 2H, *J* = 8.0 Hz, Ar-H), 7.27 (t, 1H, *J* = 7.6 Hz, Ar-H), 7.39 (d, 1H, *J* = 7.4 Hz, Ar-H), 7.46 (d, 2H, *J* = 7.6 Hz, Ar-H), 7.53 (s, 1H, Ar-H), 7.63 (d, 2H, *J* = 8.4 Hz, Ar-H), 8.22 (d, 2H, *J* = 7.6 Hz, Ar-H), 8.71 (s, 1H, OH D₂O exch.), 9.55 (s, 1H, CH=N), 10.44 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ, ppm): 169.4, 163.8, 160.1, 158.6, 155.5, 153.2, 151.5, 147.1, 138.7, 135.2, 134.9, 130.5, 130.1, 123.9, 123.1, 121.8, 118.2, 115.9, 115.6, 115.0, 114.8, 114.3, 13.6. LCMS (ESI): M/Z = 542.14 [M⁺]. Anal. Calcd. for C₂₉H₁₈N₈O₄: C, 64.20; H, 3.34; N, 20.65. Found: C, 64.29; H, 3.38; N, 20.72.

4.4.4. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-hydroxybenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5d)

Yield 63%; mp 236 °C. IR (v_{max}, cm⁻¹, KBr): 3444 (OH), 3386 (-NH, benzimidazole), 3036 (C-H, aromatic), 2992 (C-H, CH=N), 2928 (C-H, CH₃), 2224 (CN), 1685 (CO), 1637 (C=C), 1531 (C=N), 1522, 1332 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ, ppm): 1.89 (s, 3H, CH₃), 6.88 (d, 2H, *J* = 7.8 Hz, Ar-H), 7.20 (d, 2H, *J* = 7.9 Hz, Ar-H), 7.44 (d, 2H, *J* = 7.5 Hz, Ar-H), 7.62 (d, 2H, *J* = 8.3 Hz, Ar-H), 7.82 (d, 2H, *J* = 7.4 Hz, Ar-H), 8.24 (d, 2H, *J* = 7.5 Hz, Ar-H),

8.70 (s, 1H, OH D₂O exch.), 9.57 (s, 1H, CH=N), 10.44 (s, 1H, -NH benzimidazole D₂O exch.).
¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.9, 160.8, 160.0, 155.6, 153.1, 151.6, 147.2,
 138.5, 134.8, 130.7, 130.1, 126.2, 123.8, 123.0, 116.1, 115.9, 115.4, 115.0, 114.8, 13.7. LCMS
 (ESI): M/Z = 542.15 [M⁺]. Anal. Calcd. for C₂₉H₁₈N₈O₄: C, 64.20; H, 3.34; N, 20.65. Found: C,
 64.30; H, 3.38; N, 20.73.

4.4.5. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-fluorobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**5e**)

Yield 62%; mp 192 °C. IR (ν_{\max} , cm⁻¹, KBr): 3388 (-NH, benzimidazole), 3038 (C-H, aromatic),
 2994 (C-H, CH=N), 2930 (C-H, CH₃), 2226 (CN), 1688 (CO), 1638 (C=C), 1532 (C=N), 1524,
 1334 (N=O, Ar-NO₂), 1122 (C-F). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.87 (s, 3H, CH₃),
 7.21 (d, 2H, *J* = 8.0 Hz, Ar-H), 7.37 (d, 1H, *J* = 7.4 Hz, Ar-H), 7.45 (d, 2H, *J* = 7.4 Hz, Ar-H),
 7.58 (d, 2H, *J* = 8.3 Hz, Ar-H), 7.65 (d, 1H, *J* = 7.5 Hz, Ar-H), 7.71 (t, 1H, *J* = 7.6 Hz, Ar-H),
 7.84 (d, 1H, *J* = 7.4 Hz, Ar-H), 8.22 (d, 2H, *J* = 7.5 Hz, Ar-H), 9.54 (s, 1H, CH=N), 10.41 (s, 1H,
 -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.3, 163.8, 163.1,
 160.1, 155.7, 153.1, 151.5, 147.1, 138.5, 135.2, 134.9, 130.5, 130.1, 124.8, 123.9, 123.2, 117.7,
 115.9, 115.5, 115.0, 114.8, 114.1, 13.7. LCMS (ESI): M/Z = 544.12 [M⁺]. Anal. Calcd. for
 C₂₉H₁₇FN₈O₃: C, 63.97; H, 3.15; N, 20.58. Found: C, 63.90; H, 3.19; N, 20.52.

4.4.6. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-fluorobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**5f**)

Yield 64%; mp 208 °C. IR (ν_{\max} , cm⁻¹, KBr): 3386 (-NH, benzimidazole), 3037 (C-H, aromatic),
 2993 (C-H, CH=N), 2931 (C-H, CH₃), 2227 (CN), 1686 (CO), 1638 (C=C), 1531 (C=N), 1522,
 1332 (N=O, Ar-NO₂), 1125 (C-F). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.88 (s, 3H, CH₃),

7.22 (d, 2H, $J = 8.1$ Hz, Ar-H), 7.37 (d, 2H, $J = 7.6$ Hz, Ar-H), 7.44 (d, 2H, $J = 7.5$ Hz, Ar-H),
 7.62 (d, 2H, $J = 8.4$ Hz, Ar-H), 7.83 (d, 2H, $J = 7.7$ Hz, Ar-H), 8.23 (d, 2H, $J = 7.5$ Hz, Ar-H),
 9.51 (s, 1H, CH=N), 10.45 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-
 d_6 , δ , ppm): 169.5, 165.3, 163.7, 160.1, 155.7, 153.3, 151.7, 147.2, 138.7, 134.8, 130.9, 130.2,
 129.2, 123.8, 123.1, 115.9, 115.6, 115.2, 114.9, 114.2, 13.5. LCMS (ESI): $M/Z = 544.12$ [M^+].
 Anal. Calcd. for C₂₉H₁₇FN₈O₃: C, 63.97; H, 3.15; N, 20.58. Found: C, 63.91; H, 3.20; N, 20.51.

4.4.7. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((2-chlorobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5g)

Yield 67%; mp 223 °C. IR (ν_{\max} , cm⁻¹, KBr): 3384 (-NH, benzimidazole), 3034 (C-H, aromatic),
 2991 (C-H, CH=N), 2929 (C-H, CH₃), 2226 (CN), 1684 (CO), 1634 (C=C), 1530 (C=N), 1521,
 1331 (N=O, Ar-NO₂), 756 (C-Cl). ¹H NMR (400 MHz, DMSO- d_6 , δ , ppm): 1.92 (s, 3H, CH₃),
 7.19 (d, 2H, $J = 8.2$ Hz, Ar-H), 7.38 (t, 1H, $J = 7.4$ Hz, Ar-H), 7.44 (d, 2H, $J = 7.6$ Hz, Ar-H),
 7.50 (t, 1H, $J = 7.5$ Hz, Ar-H), 7.57 (d, 1H, $J = 7.5$ Hz, Ar-H), 7.68 (d, 2H, $J = 8.4$ Hz, Ar-H),
 7.81 (d, 1H, $J = 7.6$ Hz, Ar-H), 8.22 (d, 2H, $J = 7.5$ Hz, Ar-H), 9.48 (s, 1H, CH=N), 10.42 (s, 1H,
 -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO- d_6 , δ , ppm): 169.4, 163.6, 160.1,
 155.7, 153.3, 151.6, 147.2, 138.7, 134.8, 133.9, 133.3, 132.5, 130.6, 130.1, 127.3, 126.8, 123.8,
 123.2, 115.9, 115.5, 115.0, 114.9, 13.6. LCMS (ESI): $M/Z = 561.12$ [M^+]. Anal. Calcd. for
 C₂₉H₁₇ClN₈O₃: C, 62.09; H, 3.05; N, 19.98. Found: C, 62.19; H, 3.09; N, 19.92.

4.4.8. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-chlorobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5h)

Yield 64%; mp 199 °C. IR (ν_{\max} , cm⁻¹, KBr): 3384 (-NH, benzimidazole), 3035 (C-H, aromatic),
 2992 (C-H, CH=N), 2930 (C-H, CH₃), 2228 (CN), 1682 (CO), 1632 (C=C), 1528 (C=N), 1520,

1330 (N=O, Ar-NO₂), 758 (C-Cl). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.90 (s, 3H, CH₃), 7.21 (d, 2H, *J* = 8.1 Hz, Ar-H), 7.41 (d, 2H, *J* = 7.4 Hz, Ar-H), 7.47 (t, 1H, *J* = 7.4 Hz, Ar-H), 7.58 (d, 1H, *J* = 7.5 Hz, Ar-H), 7.69 (d, 2H, *J* = 8.3 Hz, Ar-H), 7.74 (d, 1H, *J* = 7.5 Hz, Ar-H), 7.92 (d, 1H, *J* = 7.6 Hz, Ar-H), 8.24 (d, 2H, *J* = 7.7 Hz, Ar-H), 9.44 (s, 1H, CH=N), 10.43 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.8, 160.2, 155.6, 153.4, 151.4, 147.1, 138.6, 135.1, 134.9, 134.3, 131.0, 130.6, 130.0, 127.7, 127.2, 123.9, 123.1, 115.8, 115.5, 115.0, 114.7, 13.5. LCMS (ESI): *M/Z* = 561.11 [*M*⁺]. Anal. Calcd. for C₂₉H₁₇ClN₈O₃: C, 62.09; H, 3.05; N, 19.98. Found: C, 62.18; H, 3.10; N, 19.93.

4.4.9. 1-((1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)amino)-6-((4-chlorobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5i**)**

Yield 67%; mp 229 °C. IR (ν_{\max} , cm⁻¹, KBr): 3386 (-NH, benzimidazole), 3036 (C-H, aromatic), 2994 (C-H, CH=N), 2932 (C-H, CH₃), 2229 (CN), 1682 (CO), 1634 (C=C), 1528 (C=N), 1522, 1331 (N=O, Ar-NO₂), 760 (C-Cl). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.88 (s, 3H, CH₃), 7.22 (d, 2H, *J* = 8.2 Hz, Ar-H), 7.44 (d, 2H, *J* = 7.5 Hz, Ar-H), 7.57 (d, 2H, *J* = 7.6 Hz, Ar-H), 7.68 (d, 2H, *J* = 8.2 Hz, Ar-H), 7.79 (d, 2H, *J* = 7.6 Hz, Ar-H), 8.22 (d, 2H, *J* = 7.7 Hz, Ar-H), 9.49 (s, 1H, CH=N), 10.44 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.6, 163.7, 160.1, 155.5, 153.2, 151.5, 147.0, 138.5, 136.7, 134.8, 131.9, 130.7, 130.0, 128.9, 123.8, 123.2, 115.9, 115.6, 115.2, 114.8, 13.7. LCMS (ESI): *M/Z* = 561.11 [*M*⁺]. Anal. Calcd. for C₂₉H₁₇ClN₈O₃: C, 62.09; H, 3.05; N, 19.98. Found: C, 62.19; H, 3.09; N, 19.92.

4.4.10. 1-((1-(1*H*-benzo[*d*]imidazol-2-yl)ethylidene)amino)-6-((2-methylbenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5j**)**

Yield 63%; mp 244 °C. IR (ν_{\max} , cm^{-1} , KBr): 3381 (-NH, benzimidazole), 3032 (C-H, aromatic), 2990 (C-H, CH=N), 2928 (C-H, CH_3), 2224 (CN), 1679 (CO), 1630 (C=C), 1523 (C=N), 1520, 1328 (N=O, Ar- NO_2). ^1H NMR (400 MHz, $\text{DMSO-}d_6$, δ , ppm): 1.91 (s, 3H, CH_3), 2.51 (s, 3H, CH_3), 7.17 (d, 2H, $J = 8.1$ Hz, Ar-H), 7.24 (t, 1H, $J = 7.4$ Hz, Ar-H), 7.30 (d, 1H, $J = 7.5$ Hz, Ar-H), 7.38 (t, 1H, $J = 7.6$ Hz, Ar-H), 7.46 (d, 2H, $J = 7.9$ Hz, Ar-H), 7.67 (d, 2H, $J = 8.3$ Hz, Ar-H), 7.74 (d, 1H, $J = 7.4$ Hz, Ar-H), 8.23 (d, 2H, $J = 7.6$ Hz, Ar-H), 9.43 (s, 1H, CH=N), 10.45 (s, 1H, -NH benzimidazole D_2O exch.). ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$, δ , ppm): 169.5, 163.6, 160.1, 155.7, 153.3, 151.6, 147.1, 138.6, 135.3, 134.9, 131.2, 130.8, 130.1, 129.1, 126.6, 125.9, 123.9, 123.1, 115.9, 115.5, 115.0, 114.7, 18.8, 13.7. LCMS (ESI): $M/Z = 540.18$ [M^+]. Anal. Calcd. for $\text{C}_{30}\text{H}_{20}\text{N}_8\text{O}_3$: C, 66.66; H, 3.73; N, 20.73. Found: C, 66.60; H, 3.77; N, 20.78.

4.4.11. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-methylbenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5k)

Yield 64%; mp 218 °C. IR (ν_{\max} , cm^{-1} , KBr): 3380 (-NH, benzimidazole), 3031 (C-H, aromatic), 2990 (C-H, CH=N), 2927 (C-H, CH_3), 2222 (CN), 1678 (CO), 1630 (C=C), 1523 (C=N), 1521, 1328 (N=O, Ar- NO_2). ^1H NMR (400 MHz, $\text{DMSO-}d_6$, δ , ppm): 1.88 (s, 3H, CH_3), 2.36 (s, 3H, CH_3), 7.18 (d, 2H, $J = 8.2$ Hz, Ar-H), 7.29 (d, 1H, $J = 7.5$ Hz, Ar-H), 7.39 (t, 1H, $J = 7.6$ Hz, Ar-H), 7.45 (d, 2H, $J = 7.5$ Hz, Ar-H), 7.61 (d, 1H, $J = 7.4$ Hz, Ar-H), 7.67 (d, 2H, $J = 8.4$ Hz, Ar-H), 7.76 (s, 1H, Ar-H), 8.21 (d, 2H, $J = 7.4$ Hz, Ar-H), 9.41 (s, 1H, CH=N), 10.44 (s, 1H, -NH benzimidazole D_2O exch.). ^{13}C NMR (100 MHz, $\text{DMSO-}d_6$, δ , ppm): 169.4, 163.8, 160.0, 155.6, 153.2, 151.7, 147.2, 138.8, 138.2, 134.8, 133.7, 131.4, 130.2, 129.5, 128.8, 126.3, 123.9, 123.2, 115.9, 115.5, 115.0, 114.7, 21.4, 13.6. LCMS (ESI): $M/Z = 540.17$ [M^+]. Anal. Calcd. for $\text{C}_{30}\text{H}_{20}\text{N}_8\text{O}_3$: C, 66.66; H, 3.73; N, 20.73. Found: C, 66.58; H, 3.78; N, 20.77.

4.4.12. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-methylbenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**5l**)

Yield 62%; mp 247 °C. IR (ν_{\max} , cm^{-1} , KBr): 3381 (-NH, benzimidazole), 3030 (C-H, aromatic), 2990 (C-H, CH=N), 2928 (C-H, CH₃), 2224 (CN), 1680 (CO), 1631 (C=C), 1524 (C=N), 1520, 1326 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.87 (s, 3H, CH₃), 2.37 (s, 3H, Ar-CH₃), 7.19 (d, 2H, *J* = 8.1 Hz, Ar-H), 7.29 (d, 2H, *J* = 7.7 Hz, Ar-H), 7.44 (d, 2H, *J* = 7.5 Hz, Ar-H), 7.67 (d, 2H, *J* = 8.3 Hz, Ar-H), 7.74 (d, 2H, *J* = 7.4 Hz, Ar-H), 8.24 (d, 2H, *J* = 7.6 Hz, Ar-H), 9.44 (s, 1H, CH=N), 10.46 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.8, 160.1, 155.7, 153.1, 151.6, 147.0, 140.7, 138.5, 134.8, 130.7, 130.1, 129.7, 129.2, 123.8, 123.1, 115.9, 115.5, 115.1, 114.7, 21.5, 13.7. LCMS (ESI): *M/Z* = 540.17 [*M*⁺]. Anal. Calcd. for C₃₀H₂₀N₈O₃: C, 66.66; H, 3.73; N, 20.73. Found: C, 66.58; H, 3.78; N, 20.78.

4.4.13. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-methoxybenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**5m**)

Yield 66%; mp 201 °C. IR (ν_{\max} , cm^{-1} , KBr): 3380 (-NH, benzimidazole), 3027 (C-H, aromatic), 2988 (C-H, CH=N), 2928 (C-H, CH₃), 2851 (C-H, OCH₃), 2221 (CN), 1678 (CO), 1630 (C=C), 1522 (C=N), 1518, 1322 (N=O, Ar-NO₂). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.87 (s, 3H, CH₃), 3.86 (s, 3H, OCH₃), 7.07 (d, 1H, *J* = 7.5 Hz, Ar-H), 7.18 (d, 2H, *J* = 8.1 Hz, Ar-H), 7.32 (t, 1H, *J* = 7.6 Hz, Ar-H), 7.39 (d, 1H, *J* = 7.4 Hz, Ar-H), 7.45 (d, 2H, *J* = 7.6 Hz, Ar-H), 7.54 (s, 1H, Ar-H), 7.66 (d, 2H, *J* = 8.4 Hz, Ar-H), 8.21 (d, 2H, *J* = 7.5 Hz, Ar-H), 9.41 (s, 1H, CH=N), 10.42 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.7, 160.8, 160.1, 155.7, 153.1, 151.5, 147.1, 138.7, 134.9, 134.3, 130.1, 129.7, 123.9, 123.2,

121.6, 116.7, 115.8, 115.4, 115.0, 114.7, 111.3, 55.6, 13.6. LCMS (ESI): $M/Z = 556.15 [M^+]$.

Anal. Calcd. for $C_{30}H_{20}N_8O_4$: C, 64.74; H, 3.62; N, 20.13. Found: C, 64.68; H, 3.67; N, 20.18.

4.4.14. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-methoxybenzylidene)- amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5n)*

Yield 67%; mp 211 °C. IR (ν_{\max} , cm^{-1} , KBr): 3381 (-NH, benzimidazole), 3028 (C-H, aromatic), 2990 (C-H, CH=N), 2929 (C-H, CH_3), 2854 (C-H, OCH_3), 2223 (CN), 1679 (CO), 1632 (C=C), 1524 (C=N), 1520, 1323 (N=O, Ar- NO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, δ , ppm): 1.86 (s, 3H, CH_3), 3.87 (s, 3H, OCH_3), 7.07 (d, 2H, $J = 7.5$ Hz, Ar-H), 7.19 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.41 (d, 2H, $J = 7.4$ Hz, Ar-H), 7.67 (d, 2H, $J = 8.4$ Hz, Ar-H), 7.84 (d, 2H, $J = 7.3$ Hz, Ar-H), 8.20 (d, 2H, $J = 7.5$ Hz, Ar-H), 9.40 (s, 1H, CH=N), 10.42 (s, 1H, -NH benzimidazole D_2O exch.). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$, δ , ppm): 169.4, 163.9, 162.8, 160.0, 155.7, 153.2, 151.7, 147.2, 138.6, 134.9, 130.8, 130.3, 126.1, 123.8, 123.1, 115.9, 115.5, 115.0, 114.8, 114.3, 55.9, 13.7. LCMS (ESI): $M/Z = 556.14 [M^+]$. Anal. Calcd. for $C_{30}H_{20}N_8O_4$: C, 64.74; H, 3.62; N, 20.13. Found: C, 64.68; H, 3.68; N, 20.19.

4.4.15. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-nitrobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5o)*

Yield 69%; mp 251 °C. IR (ν_{\max} , cm^{-1} , KBr): 3385 (-NH, benzimidazole), 3031 (C-H, aromatic), 2994 (C-H, CH=N), 2932 (C-H, CH_3), 2226 (CN), 1684 (CO), 1634 (C=C), 1527 (C=N), 1524, 1326 (N=O, Ar- NO_2). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, δ , ppm): 1.90 (s, 3H, CH_3), 7.21 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.42 (d, 2H, $J = 7.5$ Hz, Ar-H), 7.69 (d, 2H, $J = 8.3$ Hz, Ar-H), 7.80 (t, 1H, $J = 7.5$ Hz, Ar-H), 8.14 (d, 1H, $J = 7.5$ Hz, Ar-H), 8.20 (d, 2H, $J = 7.6$ Hz, Ar-H), 8.27 (d, 1H, $J = 7.7$ Hz, Ar-H), 8.51 (s, 1H, Ar-H), 9.44 (s, 1H, CH=N), 10.48 (s, 1H, -NH benzimidazole D_2O

exch.). ^{13}C NMR (100 MHz, DMSO- d_6 , δ , ppm): 169.4, 163.6, 160.1, 155.7, 153.2, 151.6, 148.1, 147.0, 138.5, 135.4, 134.8, 134.4, 130.2, 129.6, 126.3, 123.8, 123.2, 121.7, 115.8, 115.5, 115.0, 114.8, 13.7. LCMS (ESI): $M/Z = 571.12$ [M^+]. Anal. Calcd. for $\text{C}_{29}\text{H}_{17}\text{N}_9\text{O}_5$: C, 60.95; H, 3.00; N, 22.06. Found: C, 60.89; H, 3.04; N, 22.11.

4.4.16. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-nitrobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5p)*

Yield 66%; mp 261 °C. IR (ν_{max} , cm^{-1} , KBr): 3387 (-NH, benzimidazole), 3034 (C-H, aromatic), 2996 (C-H, CH=N), 2934 (C-H, CH_3), 2229 (CN), 1688 (CO), 1637 (C=C), 1529 (C=N), 1526, 1327 (N=O, Ar- NO_2). ^1H NMR (400 MHz, DMSO- d_6 , δ , ppm): 1.90 (s, 3H, CH_3), 7.21 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.42 (d, 2H, $J = 7.5$ Hz, Ar-H), 7.69 (d, 2H, $J = 8.3$ Hz, Ar-H), 7.80 (t, 1H, $J = 7.5$ Hz, Ar-H), 8.14 (d, 1H, $J = 7.5$ Hz, Ar-H), 8.20 (d, 2H, $J = 7.6$ Hz, Ar-H), 8.27 (d, 1H, $J = 7.7$ Hz, Ar-H), 8.51 (s, 1H, Ar-H), 9.44 (s, 1H, CH=N), 10.48 (s, 1H, -NH benzimidazole D_2O exch.). ^{13}C NMR (100 MHz, DMSO- d_6 , δ , ppm): 169.6, 163.8, 160.1, 155.7, 153.1, 151.7, 150.4, 147.2, 139.9, 138.6, 134.9, 130.1, 127.8, 124.1, 123.9, 123.3, 115.9, 115.5, 115.0, 114.7, 13.5. LCMS (ESI): $M/Z = 571.14$ [M^+]. Anal. Calcd. for $\text{C}_{29}\text{H}_{17}\text{N}_9\text{O}_5$: C, 60.95; H, 3.00; N, 22.06. Found: C, 60.87; H, 3.06; N, 22.12.

4.4.17. *1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((3-bromobenzylidene)amino)-4-(4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (5q)*

Yield 70%; mp 193 °C. IR (ν_{max} , cm^{-1} , KBr): 3386 (-NH, benzimidazole), 3033 (C-H, aromatic), 2992 (C-H, CH=N), 2931 (C-H, CH_3), 2224 (CN), 1684 (CO), 1635 (C=C), 1525 (C=N), 1526, 1324 (N=O, Ar- NO_2), 543 (C-Br). ^1H NMR (400 MHz, DMSO- d_6 , δ , ppm): 1.89 (s, 3H, CH_3), 7.21 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.39 (t, 1H, $J = 7.5$ Hz, Ar-H), 7.45 (d, 2H, $J = 7.4$ Hz, Ar-H),

7.57 (d, 1H, $J = 7.5$ Hz, Ar-H), 7.67 (d, 2H, $J = 7.6$ Hz, Ar-H), 7.79 (d, 1H, $J = 7.4$ Hz, Ar-H),
 7.85 (s, 1H, Ar-H), 8.21 (d, 2H, $J = 7.6$ Hz, Ar-H), 9.49 (s, 1H, CH=N), 10.43 (s, 1H, -NH
 benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-*d*₆, δ , ppm): 169.5, 163.7, 160.2, 155.7,
 153.3, 151.6, 147.1, 138.9, 135.8, 134.7, 133.7, 132.8, 130.2, 129.8, 128.3, 123.9, 123.5, 123.0,
 115.8, 115.5, 115.1, 114.8, 13.7. LCMS (ESI): $M/Z = 604.08$ [M^+]. Anal. Calcd. for
 C₂₉H₁₇BrN₈O₃: C, 57.53; H, 2.83; N, 18.51. Found: C, 57.60; H, 2.87; N, 18.57.

4.4.18. 1-((1-(1H-benzo[d]imidazol-2-yl)ethylidene)amino)-6-((4-bromobenzylidene)amino)-4-
 (4-nitrophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile (**5r**)

Yield 71%; mp 223 °C. IR (ν_{\max} , cm⁻¹, KBr): 3387 (-NH, benzimidazole), 3034 (C-H, aromatic),
 2994 (C-H, CH=N), 2932 (C-H, CH₃), 2225 (CN), 1686 (CO), 1636 (C=C), 1526 (C=N), 1527,
 1325 (N=O, Ar-NO₂), 546 (C-Br). ¹H NMR (400 MHz, DMSO-*d*₆, δ , ppm): 1.88 (s, 3H, CH₃),
 7.20 (d, 2H, $J = 8.0$ Hz, Ar-H), 7.42 (d, 2H, $J = 7.5$ Hz, Ar-H), 7.58 (d, 2H, $J = 7.7$ Hz, Ar-H),
 7.68 (d, 2H, $J = 7.6$ Hz, Ar-H), 7.75 (d, 2H, $J = 7.6$ Hz, Ar-H), 8.20 (d, 2H, $J = 7.6$ Hz, Ar-H),
 9.46 (s, 1H, CH=N), 10.45 (s, 1H, -NH benzimidazole D₂O exch.). ¹³C NMR (100 MHz, DMSO-
*d*₆, δ , ppm): 169.5, 163.7, 160.2, 155.5, 153.2, 151.6, 147.1, 138.7, 134.8, 132.6, 131.8, 130.2,
 128.6, 125.5, 123.8, 123.0, 115.9, 115.4, 115.0, 114.8, 13.7. LCMS (ESI): $M/Z = 604.07$ [M^+].
 Anal. Calcd. for C₂₉H₁₇BrN₈O₃: C, 57.53; H, 2.83; N, 18.51. Found: C, 57.60; H, 2.87; N, 18.58.

4.5. Biological assay

4.5.1. Antibacterial assay

Antibacterial studies of newly synthesized compounds **5a-r** were carried out against the
 representative panel of bacteria such as *Staphylococcus aureus* MTCC-96, *Streptococcus*

555 *pyogenes* MTCC-442, *Escherichia coli* MTCC-443, *Pseudomonas aeruginosa* MTCC-1688 and
556 methicillin-resistant *S. aureus* (MRSA isolate ATCC 43300). Antifungal activity was carried out
557 against the yeast-like pathogenic fungus *Candida albicans* MTCC 227. All MTCC and ATCC
558 cultures were collected from Institute of Microbial Technology, Chandigarh. The activity of
559 compounds was determined as per National Committee for Clinical Laboratory Standards
560 (NCCLS) protocol using Mueller Hinton Broth (Becton Dickinson, USA). Primary screening
561 was done first for antibacterial activity in six sets against *E. coli*, *S. aureus*, *P. aeruginosa* and *S.*
562 *pyogenes* at different concentrations of 1000, 500, 250 µg/mL. The compounds found to be
563 active in primary screening were similarly diluted to obtain 200, 125, 100, 62.5, 50, 25, 12.5 and
564 6.25 µg/mL concentrations for secondary screening to test in a second set of dilution against all
565 microorganisms. Inoculum size for test strain was adjusted to 10^6 CFU/mL (Colony Forming
566 Unit per milliliter) by comparing the turbidity (turbidimetric method). Mueller Hinton Broth was
567 used as nutrient medium to grow and dilute the compound suspension for test organisms. 2%
568 DMSO was used as a diluent/vehicle to obtain the desired concentration of synthesized
569 compounds and standard drugs to test upon standard microbial strains. Synthesized compounds
570 were diluted to 1000 µg/mL concentration, as stock solution. The control tube containing no
571 antibiotic was immediately subcultured [before inoculation] by spreading a loopful evenly over a
572 quarter of plate in a suitable medium for the growth of test organisms. The culture tubes were
573 then incubated for 24 h at 37 °C and the growth was monitored visually and
574 spectrophotometrically. Ten µg/mL suspensions were further inoculated on an appropriate media
575 and growth was noted after 24 h and 48 h. The lowest concentration (highest dilution) required to
576 arrest the growth of bacteria was regarded as minimum inhibitory concentration (MIC) i.e. the
577 amount of growth from the control tube before incubation (which represents the original

inoculum) was compared. Solvent had no influence on strain growth. The result of this was greatly affected by the size of the inoculums. The test mixture should contain 10^6 CFU/mL organisms. DMSO and sterilized distilled water were used as negative control while chloramphenicol (1 U strength) was used as positive control. Standard drugs used in the present study were ‘ciprofloxacin’ and ‘chloramphenicol’ for evaluating antibacterial activity.

4.5.2. Antitubercular assay

Antitubercular activity was determined using the modified radiometric 7H12 broth (BACTEC 12B system) in which stock solutions as test compounds were prepared in dimethylsulfoxide (DMSO) at a concentration of 12.8 mM and the final test concentrations ranged from 39.42 to 0.15 μ g/mL. Controls received 50 μ L DMSO. INH was included as a positive drug control. INH was solubilized and diluted in DMSO and added to BACTEC-12 broth to achieve a range of concentration for determination of minimum inhibitory concentration (MIC, lowest concentration inhibiting \geq 90% of the inoculums, MIC value of INH is 0.24 μ g/mL at 95% inhibition of H₃₇Rv strain). *M. tuberculosis* H₃₇Rv strain (ATCC 27294) was cultured at 37 °C in 100 ml of Middlebrook 7H9 broth (Difco, Detroit, Mich.) supplemented with 0.2% (vol/vol) glycerol, 10% (vol/vol) OADC (oleic acid, albumin, dextrose, catalase) and 0.05% (vol/vol) Tween 80. The complete medium was referred to as 7H9GC-Tween. Cultures were incubated in 500-ml nephelometer flasks on a rotary shaker at 150 rpm and 37 °C until they reached an optical density of 0.4 to 0.5 at 550 nm. Bacteria were washed and suspended in 20 ml of phosphate-buffered saline and passed through an 8-mm pore size filter to eliminate clumps. The filtrates were aliquoted, stored at 280 °C and used within 30 days. Cultures were prepared and an appropriate dilution performed such that a BACTEC-12B vial inoculated with 0.1 mL would reach a growth index (GI) of 999 in five days. Antimicrobial susceptibility testing was

performed in black, clear-bottomed, 96-well microplates in order to minimize background fluorescence. Outer perimeter wells were filled with sterile water to prevent dehydration in experimental wells. Initial drug dilutions were prepared in either dimethyl sulfoxide or distilled deionized water and subsequent two fold dilutions were performed in 0.1 ml of 7H9GC (no Tween 80) in the microplates. BACTEC 12B-passaged inocula were initially diluted 1:2 in 7H9GC, and 0.1 ml was added to wells. Subsequent determination of bacterial titers yielded 1×10^6 CFU/mL in plate wells for H₃₇Rv. Frozen inocula were initially diluted 1:20 in BACTEC 12B medium followed by a 1:50 dilution in 7H9GC. Addition of 1/10 mL to wells resulted in final bacterial titers of 2.0×10^5 for H₃₇Rv. Wells containing drug only were used to detect auto fluorescence of compounds. Cultures were incubated at 37 °C and the Growth of Inhibition (GI) determined daily until control cultures achieved a GI of 999. Assays were usually completed in 5-8 days. Percent inhibition was defined as $1 - (\text{GI of test sample} / \text{GI of control}) \times 100$. The lowest drug concentration effecting an inhibition of $\geq 90\%$ was considered the MIC.

4.5.3. Cytotoxicity assay

VERO cells were cultured in Dulbecco Modified Eagle Medium (DMEM) containing 2 mM Na₂CO₃ supplemented with 10% (v/v) foetal bovine serum (FBS). The cells were incubated at 37 °C under 5% CO₂ and 95% air in a humidified atmosphere until confluent and then diluted with phosphate-buffered saline to 10^6 cells/mL. Stock solutions were prepared in dimethyl sulfoxide (DMSO) and further dilutions were made with fresh culture medium. The concentration of DMSO in the final culture medium was 1%, which had no effect on the cell viability. In a transparent 96-well plate (Falcon Micro test 96), three fold serial dilutions of the macrolide stock solutions resulted in final concentrations of 31.53 to 0.12 µg/mL in a final volume of 200 µL. After incubation at 37 °C for 72 h, medium was removed and monolayer was

washed twice with 100 μ L of warm Hanks' balanced salt solution (HBSS). One hundred microliters of warm medium and 20 μ L of freshly made MTS-PMS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2*H*-tetrazolium and phenylmethasulfazone] (100:20) (Promega) were added to each well, plates were incubated for 3 h, and absorbance was determined at 490 nm using a plate reader. Each concentration was repeated in three wells and control cell viability was considered as 100%. The same experimental conditions were provided for all compounds and analysis was repeated three times for each cell line.

Acknowledgements

Authors are thankful to the University Grants Commission, New Delhi and Department of Science & Technology, New Delhi for financial support under the NON-SAP and DST-FIST programs respectively. One of the authors, G M Kotadiya is thankful to UGC, New Delhi for providing 'Research Fellowships in Science to Meritorious Students'.

References

- [1] S.D. Lawn, P. Mwaba, M. Bates, A. Piatek, H. Alexander, B.J. Marais, L.E. Cuevas, T.D. McHugh, L. Zijenah, N. Kapata, I. Abubakar, R. McNerney, M. Hoelscher, Z.A. Memish, G. B. Migliori, P. Kim, M. Maeurer, M. Schito, A. Zumla, *Lancet. Infect. Dis.* 13 (2013) 349-361.
- [2] S. Prabowo, M. Gröschel, E.L. Schmidt, A. Skrahina, T. Mihaescu, S. Hastürk, R. Mitrofanov, E. Pimkina, I. Visontai, B. Jong, J. Stanford, P.-J. Cardona, S.E. Kaufmann, T. Werf, *Med. Microbiol. Immunol.* 202 (2013) 95.
- [3] I. Abubakar, M. Zignol, D. Falzon, M. Raviglione, L. Ditiu, S. Masham, I. Adetifa, N. Ford, H. Cox, S.D. Lawn, B.J. Marais, T.D. McHugh, P. Mwaba, M. Bates, M. Lipman,

- L. Zijenah, S. Logan, R. McNerney, A. Zumla, K. Sarda, P. Nahid, M. Hoelscher, M. Pletschette, Z.A. Memish, P. Kim, R. Hafner, S. Cole, G.B. Migliori, M. Maeurer, M. Schito, A. Zumla, *Lancet Infect. Dis.* 13 (2013) 529-539.
- [4] P.C. Appelbaum, MRSA - the tip of the iceberg, *Clin. Microbiol. Infect.* 12 (2006) 3-10.
- [5] W.R. Bishai, N. Zetola, J.S. Francis, E.L. Nuernberger, *Lancet Infect. Dis.* 5 (2005) 275-286.
- [6] C. Walsh, *Nature* 406 (2000) 775-781.
- [7] A. Argyrou, M.W. Vetting, B. Aladegbami, J.S. Blanchard, *Nat. Struct. Biol.* doi:10.1038/nsmb1089.
- [8] N.C. Desai, K.M. Rajpara, V.V. Joshi, *Bioorg. Med. Chem. Lett.* 23 (2013) 2714-2717.
- [9] Y. Bansal, O. Silakari, *Bioorg. Med. Chem.* 20 (2012) 6208-6236.
- [10] K.F. Ansari, C. Lal, *Eur. J. Med. Chem.* 44 (2009) 4028-4033.
- [11] H. Goker, C. Kus, D.W. Boykin, S. Yildiz, N. Altanlar, *Bioorg. Med. Chem.* 10 (2002) 2589-2596.
- [12] Z. Kazimierzuk, J.A. Upcroft, P. Upcroft, A. Gorska, B. Starosciak, A. Laudy, *Acta Biochim. Pol.* 49 (2002) 185-195.
- [13] A. Patil, S. Ganguly, S. Surana, *Rasayan J. Chem.* 1 (2008) 447-460.
- [14] A. K. Dubey, P. K. Sanyal, *Online Vet. J.* 5 (2010) 63.
- [15] P. Naik, P. Murumkar, R. Giridhar, M.R. Yadav, *Bioorg. Med. Chem.* 18 (2010) 8418-8456.
- [16] W.W. Mederski, D. Dorsch, S. Anzali, J. Gleitz, B. Cezanne, C. Tsaklakidis, *Bioorg. Med. Chem. Lett.* 14 (2004) 3763-3769.

- 668 [17] M. Mader, A. de Dios, C. Shih, R. Bonjouklian, T. Li, W. White, B. Lopez de Uralde, C.
 669 Sanchez-Martinez, M. del Prado, C. Jaramillo, E. de Diego, L.M. Martin Cabrejas, C.
 670 Dominguez, C. Montero, T. Shepherd, R. Dally, J.E. Toth, A. Chatterjee, S. Pleite, J.
 671 Blanco-Urgoiti, L. Perez, M. Barberis, M.J. Lorite, E. Jambrina, Jr. C.R. Nevill, P.A. Lee,
 672 R.C. Schultz, J.A. Wolos, L.C. Li, R.M. Campbell, B.D. Anderson, *Bioorg. Med. Chem.*
 673 *Lett.* 18 (2008) 179-183.
- 674 [18] T. Lam, M.T. Hilgers, M.L. Cunningham, B.P. Kwan, K.J. Nelson, V. Brown-Driver, V.
 675 Ong, M. Trzoss, G. Hough, K.J. Shaw, J. Finn, *J. Med. Chem.* 57 (3) (2014) 651-668.
- 676 [19] N.C. Desai, N.R. Shihory, G.M. Kotadiya, *Chin. Chem. Lett.* 25 (2014) 305-307.
- 677 [20] D. Seenaiiah, P.R. Reddy, G.M. Reddy, A. Padmaja, V. Padmavathi, N.S. Krishna, *Eur. J.*
 678 *Med. Chem.* 77 (2014) 1-7.
- 679 [21] M. Alp, H. Goker, R. Brun, S. Yildiz, *Eur. J. Med. Chem.* 44 (2009) 2002-2008.
- 680 [22] A. Andreani, M. Granaiola, A. Leoni, R. Morigi, M. Ramballdi, *Eur. J. Med. Chem.* 36
 681 (2001) 743-746.
- 682 [23] S.G. Kini, A.R. Bhat, B. Bryant, J.S. Williamson, F.E. Dayan, *Eur. J. Med. Chem.* 44
 683 (2009) 492-500.
- 684 [24] A.Y. Saiki, L.L. Shen, C.M. Chen, J. Baranowski, C.G. Lerner, *Antimicrob. Agents*
 685 *Chemother.* 43 (1999) 1574-1577.
- 686 [25] M. T. Cocco, C. Congiu, V. Onnis, *Eur. J. Med. Chem.* 38 (2003) 37-47.
- 687 [26] A. Abadi, O. Al-Deeb, A. Al-Afify, H. El-Kashef, *Farmaco* 54 (1999) 195-201.
- 688 [27] G. Ozturk, D.D. Erol, T. Uzbay, M.D. Aytemir, *Farmaco* 56 (2001) 251-256.
- 689 [28] P. Storck, A. Aubertin, D.S. Grierson, *Tetrahedron Lett.* 46 (2005) 2919-2922.
- 690 [29] Q. Li, L.A. Mitscher, L.L. Shen, *Med. Res. Rev.* 20 (2000) 231-293.

- [30] L.A. Mitscher, Chem Rev. 105(2) (2005) 559-592.
- [31] T. Hou, J. Wang, W. Zhang, X. Xu, J. Chem. Inf. Model. 47 (2007) 460-463.
- [32] N.C. Desai, K.M. Rajpara, V.V. Joshi, H.V. Vaghani, H.M. Satodiya, Anti. Infect. Agents 10 (2012) 75-86.
- [33] N.C. Desai, D.D. Pandya, V.V. Joshi, K.M. Rajypara, H.V. Vaghani, H.M. Satodiya, Med. Chem. Res. 21 (2012) 4463-4472.
- [34] P.C. Hannan, Vet. Res. 31 (2000) 373-395.
- [35] N.C. Desai, N. Bhatt, H. Somani, A. Trivedi, Eur. J. Med. Chem. 67 (2013) 54-59.
- [36] L. Collins, S.G. Franzblau, Antimicrob. Agents Chemother. 41 (1997) 1004-1009.
- [37] K. Falzari, Z. Zhu, D. Pan, H. Liu, P. Hongmanee, S.G. Franzblau, Antimicrob. Agents Chemother. 49 (2005) 1447-1454.

Captions

Figure 1 Structural relevance of title compounds **5a-r** with previously synthesized compounds
NCD₁₋₂₀

Scheme 1 Synthetic route for the preparation of title compounds **5a-r**

Scheme 2 Plausible mechanistic pathway for the synthesis of compounds **5a-r**

Table 1 Results of *in vitro* antibacterial screening of tested compounds

Table 2 Inhibitory activity (MIC, µg/mL) of compounds **5h**, **5q** and **5r** against methicillin
resistant *S. aureus*

Table 3 Results of antitubercular and cytotoxic activities of the tested compounds

Table 1 Results of *in vitro* antibacterial screening of tested compounds

Entry	-R	Clog <i>P</i> ^a	Minimum inhibitory concentration (MIC) µg/mL			
			Gram-negative ^b		Gram-positive ^c	
			Ec	Pa	Sa	Sp
2	--	0.7220	>1000	>1000	1000	1000
4	--	1.4772	500	500	500	500
5a	-H	3.0642	500	500	500	500
5b	-2-OH	3.2982	500	500	500	250
5c	-3-OH	3.2982	500	250	100	500
5d	-4-OH	3.2982	250	500	500	250
5e	-3-F	3.2072	100	250	100	125
5f	-4-F	3.2072	250	250	250	100
5g	-2-Cl	3.7772	100	125	50	62.5
5h	-3-Cl	3.7772	50	62.5	25	50
5i	-4-Cl	3.7772	100	100	62.5	50
5j	-2-CH ₃	3.5632	250	250	125	100
5k	-3-CH ₃	3.5632	100	100	100	100
5l	-4-CH ₃	3.5632	250	250	250	125
5m	-3-OCH ₃	3.4082	250	500	100	250
5n	-4-OCH ₃	3.4082	500	500	250	250
5o	-3-NO ₂	2.8072	250	500	250	500
5p	-4-NO ₂	2.8072	1000	1000	500	500
5q	-3-Br	3.9272	25	25	12.5	25
5r	-4-Br	3.9272	50	50	25	50
Ciprofloxacin		-0.7252	25	25	50	50
Chloramphenicol		1.293	50	50	50	50

^a Clog *P* calculated using the ChemBioDraw Ultra, version 12.0, software by Cambridge Soft;^b Ec: *Escherichia coli* (MTCC-443); Pa: *Pseudomonas aeruginosa* (MTCC-1688);^c Sa: *Staphylococcus aureus* (MTCC-96), Sp: *Streptococcus pyogenes* (MTCC-442).

Table 2 Inhibitory activity (MIC, $\mu\text{g/mL}$) of compounds **5h**, **5q** and **5r** against methicillin-resistant *S. aureus*

Entry	MRSA ^a
5h	>50
5q	12.5
5r	6.25
Ciprofloxacin	25
Chloramphenicol	50

^a Methicillin-resistant *S. aureus* (ATCC 43300)

Table 3 Results of antitubercular and cytotoxic activities of the tested compounds

Entry	-R	Clog <i>P</i> ^a	MIC (μg/mL) ^b	IC ₅₀ (μg/mL) ^c
2	--	0.7220	>39.42	n.d.
4	--	1.4772	>39.42	n.d.
5a	-H	3.0642	35.23	17.61
5b	-2-OH	3.2982	31.62	19.83
5c	-3-OH	3.2982	30.18	17.00
5d	-4-OH	3.2982	30.85	13.73
5e	-3-F	3.2072	25.37	23.65
5f	-4-F	3.2072	27.10	24.76
5g	-2-Cl	3.7772	2.65	>39.42
5h	-3-Cl	3.7772	0.85	>39.42
5i	-4-Cl	3.7772	1.79	>39.42
5j	-2-CH ₃	3.5632	9.17	34.55
5k	-3-CH ₃	3.5632	2.07	>39.42
5l	-4-CH ₃	3.5632	6.28	>39.42
5m	-3-OCH ₃	3.4082	17.98	24.22
5n	-4-OCH ₃	3.4082	23.46	25.39
5o	-3-NO ₂	2.8072	28.33	27.63
5p	-4-NO ₂	2.8072	32.21	16.05
5q	-3-Br	3.9272	10.28	>39.42
5r	-4-Br	3.9272	13.79	>39.42
	Isoniazid	-0.668	0.24	--

^a Clog *P* calculated using the ChemBioDraw Ultra, version 12.0, software by Cambridge Soft;

^b Minimum inhibitory concentration against H₃₇Rv strain of *M. Tuberculosis* (μg/mL);

^c Measurement of cytotoxic activity in VERO cell lines: 50% inhibitory concentrations (μg/mL);

n.d. not determined.






Figure 1. Structural relevance of title compounds **5a-r** with previously synthesized compounds **NCD₁₋₂₀**



Scheme 1. Synthetic route for the preparation of title compounds **5a-r**



Scheme 2. Plausible mechanistic pathway for the synthesis of compounds **5a-r**