


# Cell Chemical Biology

## Chemoproteomics-enabled discovery of covalent RNF114-based degraders that mimic natural product function

### Graphical abstract



### Authors

Mai Luo, Jessica N. Spradlin, Lydia Boike, ..., Markus Schirle, Thomas J. Maimone, Daniel K. Nomura

### Correspondence

maimone@berkeley.edu (T.J.M.),  
dnomura@berkeley.edu (D.K.N.)

### In Brief

Using activity-based protein profiling-enabled covalent ligand-screening approaches, Luo et al. have discovered a fully synthetic RNF114 E3 ligase recruiter that can be used in targeted protein degradation applications.

### Highlights

- We have identified a fully synthetic covalent ligand EN219 that targets RNF114
- We show that EN219 mimics the mode of action exploited by natural product nimbolide
- We show that EN219 can be used as an RNF114 recruiter in degrader applications



## Brief Communication

# Chemoproteomics-enabled discovery of covalent RNF114-based degraders that mimic natural product function

Mai Luo,<sup>1,2,7</sup> Jessica N. Spradlin,<sup>1,2,7</sup> Lydia Boike,<sup>1,2</sup> Bingqi Tong,<sup>1,2</sup> Scott M. Brittain,<sup>2,3</sup> Jeffrey M. McKenna,<sup>2,3</sup> John A. Tallarico,<sup>2,3</sup> Markus Schirle,<sup>2,3</sup> Thomas J. Maimone,<sup>1,2,\*</sup> and Daniel K. Nomura<sup>1,2,4,5,6,8,\*</sup>

<sup>1</sup>Department of Chemistry, University of California, Berkeley, Berkeley, CA 94720, USA

<sup>2</sup>Novartis-Berkeley Center for Proteomics and Chemistry Technologies, Cambridge, MA 02139, USA

<sup>3</sup>Novartis Institutes for BioMedical Research, Cambridge, MA 02139, USA

<sup>4</sup>Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA

<sup>5</sup>Department of Nutritional Sciences and Toxicology, University of California, Berkeley, Berkeley, CA 94720, USA

<sup>6</sup>Innovative Genomics Institute, Berkeley, CA 94720, USA

<sup>7</sup>These authors contributed equally

<sup>8</sup>Lead contact

\*Correspondence: [maimone@berkeley.edu](mailto:maimone@berkeley.edu) (T.J.M.), [dnomura@berkeley.edu](mailto:dnomura@berkeley.edu) (D.K.N.)

<https://doi.org/10.1016/j.chembiol.2021.01.005>

## SUMMARY

The translation of functionally active natural products into fully synthetic small-molecule mimetics has remained an important process in medicinal chemistry. We recently discovered that the terpene natural product nimbolide can be utilized as a covalent recruiter of the E3 ubiquitin ligase RNF114 for use in targeted protein degradation—a powerful therapeutic modality within modern-day drug discovery. Using activity-based protein profiling-enabled covalent ligand-screening approaches, here we report the discovery of fully synthetic RNF114-based recruiter molecules that can also be exploited for PROTAC applications, and demonstrate their utility in degrading therapeutically relevant targets, such as BRD4 and BCR-ABL, in cells. The identification of simple and easily manipulated drug-like scaffolds that can mimic the function of a complex natural product is beneficial in further expanding the toolbox of E3 ligase recruiters, an area of great importance in drug discovery and chemical biology.


## INTRODUCTION

Natural products have remained a cornerstone of drug discovery research for decades resulting in numerous FDA-approved medicines and tools for biomedical research across a wide range of therapeutic areas (Newman and Cragg, 2016). Historically, a large percentage of natural product-inspired medicines have utilized natural products as a starting point, in which the tools of synthetic chemistry are used to fine-tune compound properties (i.e., a semisynthetic approach) to achieve the desired endpoint. Alternatively, the function of natural products can also serve as motivation for medicinal chemistry, where the design of fully synthetic small molecules are less constrained by availability, synthetic manipulation limitations, and physicochemical and metabolic liabilities. Modern-day examples of this approach include the translation of the alkaloid cytisine into the smoking cessation drug varenicline, the development of the blockbuster cardiovascular drug atorvastatin from the polyketide lovastatin, and the discovery of the proteasome inhibitor carfilzomib inspired by the natural polypeptide epoxomicin (Figure 1). Such approaches, however, are greatly facilitated by an under-

standing of the binding mode of the small molecule to its protein target as well as the identification of key pharmacophores within the parent natural product (Figure 1).

We recently discovered that the anti-cancer natural product nimbolide, a limonoid-type triterpene isolated from *Azadirachta indica* (neem), covalently reacts with an N-terminal cysteine (C8) within an intrinsically disordered region of the E3 ubiquitin ligase RNF114 in human breast cancer cells (Spradlin et al., 2019). Covalent targeting of RNF114 by nimbolide led to impaired ubiquitination of its endogenous substrate, the tumor suppressor p21, through a nimbolide-dependent competition of the RNF114 substrate binding interaction, thus providing a potential mechanism for the anti-cancer effects of this natural product. The realization that nimbolide targeted a substrate recognition domain within RNF114 suggested that nimbolide could potentially be used as a recruiter of RNF114 for targeted protein degradation (TPD) applications. Consistent with this premise, we showed that a proteolysis-targeting chimera (PROTAC) formed by linking nimbolide to a bromodomain and extraterminal domain (BET) inhibitor JQ1 led to proteasome- and RNF114-dependent degradation of BRD4 in cells. While TPD





**Figure 1. Translation of natural product function**

Many successful drugs have recapitulated the function of natural products with fully synthetic chemical scaffolds. Key pharmacophores are shown in red.

has arisen as a powerful drug discovery paradigm for tackling the undruggable proteome by targeting intracellular proteins for proteasomal degradation rather than classic inhibition, the lack of a broad range of E3 ligase recruiters represents a known limitation in this arena (Bondeson and Crews, 2017; Chamberlain and Hamann, 2019; Lai and Crews, 2017). Indeed, while over 600 different E3 ligases have been annotated, only a small handful of these potential targets have succumbed to the PROTAC strategy. Discovering additional and more synthetically tractable E3 ligase recruiters is thus an important topic in expanding the scope of TPD and may help to address resistance mechanisms (Bond et al., 2020; Ottis et al., 2019), promote differing selectivity or kinetic profiles of degradation (Bondeson et al., 2018; Huang et al., 2018; Tong et al., 2020), and lead to cell-type or location-specific degradation.

While nimbolide has demonstrated success in TPD applications, its high molecular weight, modest chemical stability, and limited points for synthetic modification have prompted us to continually search for RNF114-based ligands for PROTAC development. Here, we realize the successful translation of the binding site of nimbolide into a simple, easily manipulated covalent small molecule. Because the N-terminal region of RNF114 that includes C8 is intrinsically disordered, structure-guided ligand discovery and optimization was not possible thus


requiring unbiased approaches to ligand discovery. Activity-based protein profiling (ABPP)-enabled covalent ligand screening has been previously used to discover covalent recruiters against E3 ligases RNF4 and DCAF16 (Ward et al., 2019; Zhang et al., 2019) and has also facilitated ligand discovery against cysteines targeted by covalently acting natural products (Grossman et al., 2017). Using this technique, we were able to discover fragments that could be used to replace nimbolide as the covalent E3 ligase recruitment module in fully functional degraders against several oncology targets.

## RESULTS

### Covalent ligand screening against RNF114

To discover a fully synthetic covalent ligand that could access the same cysteine (C8) targeted by nimbolide on RNF114, we screened 318 cysteine-reactive chloroacetamide and acrylamide ligands in a gel-based competitive ABPP assay, in which we competed ligands against binding of a rhodamine-functionalized cysteine-reactive iodoacetamide probe (IA-rhodamine) to pure RNF114 protein (Figures 2A, 2B, and S1; Table S1) (Bachovchin et al., 2010; Grossman et al., 2017; Ward et al., 2019). Through this screen, chloroacetamide EN219 emerged as the top hit, showing the greatest inhibition of IA-rhodamine binding to RNF114 (Figure 2C). Dose-response studies showed that EN219 interacted with RNF114 with a 50% inhibitory concentration of 470 nM (Figures 2D and 2E). EN219 also inhibited RNF114-mediated autoubiquitination and p21 ubiquitination *in vitro*, similarly to our previously observed findings with nimbolide (Figure 2F). Given that EN219 possesses a stereogenic center, we also synthesized enantiomeric (R)- and (S)-EN219 derivatives (with >90% enantiomeric excess) (Mahé et al., 2010) and revealed that these enantioenriched molecules showed similar potency for binding to RNF114 as the racemate we used previously (Figure S2A). We also synthesized a non-reactive acetamide analog of EN219, deschloro-EN219 (JNS 2-229), and showed that this compound no longer interacted with RNF114 (Figure S2A). Mapping the sites of EN219 covalent modification on pure RNF114 protein by liquid chromatography-mass spectrometry (LC-MS/MS) also confirmed C8 as the only site of modification (Figure S2B). While EN219 represents an early lead compound within a drug discovery setting and we recognize that, through medicinal chemistry efforts, it could further be tuned, we nonetheless pursued further characterization of EN219 as a fully synthetic ligand for RNF114.

To characterize the proteome-wide interaction profile and the potential for engagement of RNF114 C8 by EN219 in cells, we next mapped the proteome-wide cysteine reactivity of EN219 *in situ* in 231MFP breast cancer cells using several different approaches. First, we crudely assessed general cysteine reactivity of EN219 compared with a previously identified promiscuous scout ligand YP 1-44 (Ward et al., 2019), and showed that EN219 did not broadly inhibit IA-rhodamine cysteine reactivity in 231MFP breast cancer cell lysate even at 250  $\mu$ M, compared with YP 1-44 (Figure S2C). We next assessed proteome-wide selectivity of EN219 by competitive isotopic tandem orthogonal proteolysis-ABPP using previously well-validated methods (Backus et al., 2016; Grossman et al., 2017; Spradlin et al., 2019; Wang et al., 2014; Weerapana et al., 2010). Cells were



**Figure 2. Covalent ligand screening against RNF114**

(A) Gel-based ABPP assay for screening covalent ligands against an IA-rhodamine probe binding to pure RNF114 protein. Loss of fluorescence indicates covalent ligand binding to a cysteine on RNF114.

(B) Quantified results from a gel-based ABPP screen of 318 cysteine-reactive acrylamides and chloroacetamides against IA-rhodamine labeling of RNF114. DMSO vehicle or covalent ligands (50  $\mu$ M) were pre-incubated with pure RNF114 protein (0.1  $\mu$ g) for 30 min before addition of IA-rhodamine (100 nM) for 30 min at room temperature. Proteins were separated by SDS-PAGE and in-gel fluorescence was quantified. Raw gel-based ABPP data are shown in Figure S1. Structures of compounds screened can be found in Table S1. Data expressed as ligand/control ratio of in-gel fluorescent intensity. Shown in the inset are the ligand/control ratios for the top 10 hits.

(C) Structure of top hit EN219 with the chloroacetamide cysteine-reactive warhead in red.

(D) Dose-response of EN219 interaction with RNF114 by competitive gel-based ABPP. DMSO vehicle or covalent ligands were pre-incubated with pure RNF114 protein (0.1  $\mu$ g) for 30 min before addition of IA-rhodamine (100 nM) for 30 min at room temperature. Proteins were separated by SDS-PAGE and in-gel fluorescence was quantified. Gels were also silver stained as a loading control.

(E) Percent inhibition of IA-rhodamine binding to RNF114 in (D) was quantified and 50% inhibitory concentration ( $IC_{50}$ ) was determined to be 0.47  $\mu$ M.

(F) RNF114-mediated autoubiquitination and p21 ubiquitination *in vitro*. DMSO vehicle or EN219 (50  $\mu$ M) was incubated with pure RNF114 for 30 min before addition of PBS or p21, ATP, and FLAG-ubiquitin (Ub) for 60 min. Proteins were separated by SDS-PAGE and blotted for FLAG. Ubiquitinated-RNF114/p21 is noted.

(G) Isotopic tandem orthogonal proteolysis ABPP (isoTOP-ABPP) analysis of EN219 in 231MFP breast cancer cells. 231MFP cells were treated *in situ* with DMSO vehicle or EN219 (1  $\mu$ M) for 90 min. Control and treated cell lysates were labeled with IA-alkyne (100  $\mu$ M) for 1 h, after which isotopically light (control) or heavy (EN219-treated) biotin-azide bearing a TEV tag was appended by CuAAC. Proteomes were mixed in a 1:1 ratio, probe-labeled proteins were enriched with avidin and digested with trypsin, and probe-modified peptides were eluted by TEV protease and analyzed by LC-MS/MS. Average light-to-heavy (control/EN219) ratios and adjusted p values were quantified for probe-modified peptides that were present in two out of three biological replicates and plotted. Shown in red are protein and modified cysteine sites for peptides that showed >2-fold control/EN219 ratio with adjusted p value <0.05. Full data can be found in Table S2.

(H) Structure of alkyne-functionalized EN219 probe.

(I) EN219-alkyne pull-down of RNF114. 231MFP cells were treated with DMSO vehicle or EN219-alkyne (50  $\mu$ M) for 90 min. Biotin-azide was appended to probe-labeled proteins by CuAAC and probe-labeled proteins were avidin enriched and blotted for RNF114 and negative control actin. An aliquot of input proteome from vehicle- and EN219-treated cell lysates was also subjected to blotting as an input control. Data shown in (E) are average values and individual replicate values. Data shown are from n = 1 in (B) and n = 3 for (D–F and I) biological replicate(s) per group. Gels shown in (D and I) are representative gels from n = 3 biological replicates per group. This figure is related to Figures S1 and S2.

treated *in situ* with vehicle or EN219 and cells were subsequently lysed and labeled with an alkyne-functionalized iodoacetamide probe (IA-alkyne), followed by attachment of isotopically light or heavy, tobacco etch virus (TEV)-cleavable biotin-azide enrichment handles through copper-catalyzed azide-alkyne cycloaddition (CuAAC). Probe-labeled proteins were enriched by avidin, digested with trypsin, and probe-modified peptides were eluted by TEV protease and analyzed by LC-MS/MS. Only four proteins showed >2-fold light versus heavy or control versus EN219-treated probe-modified peptide ratios with adjusted *p* value <0.05 out of 686 probe-modified cysteines quantified. These proteins were RNF114 C8, TUBB1 C201, HSPD1 C442, and HIST1H3A C97, among which RNF114 was the only E3 ligase (Figures S2G; Table S2).

To further investigate the selectivity of EN219 and to confirm EN219 engagement of RNF114 in cells, we also synthesized an alkyne-functionalized EN219 probe (EN219-alkyne) (Figure 2H). Initially, we labeled 231MFP cells *in situ* with this EN219-alkyne probe and visualized labeled proteins by gel-based ABPP and investigated labeled protein bands that were competed by EN219 pre-treatment (Figure S2D). This experiment showed about eight potential EN219-alkyne-specific proteins that were competed by EN219 itself in cells, with several bands that were specific to EN219-alkyne (Figure S2D). In some cases, we observed increased EN219-alkyne protein labeling of specific proteins in cells pre-treated with excess EN219, indicating increased labeling of EN219-alkyne-specific off-targets when EN219-specific sites were occupied (Figure S2D). To further elucidate these targets, we performed tandem mass tagging (TMT)-based quantitative proteomic profiling to identify proteins that were enriched by EN219-alkyne *in situ* labeling of 231MFP cells and were competed by EN219 *in situ* treatment (Figure S2E; Table S3). While we did not identify RNF114 in this proteomics experiment, likely due to its relatively low abundance or other many EN219-alkyne-specific targets that occluded detection of EN219-specific targets, we did identify seven additional potential off-targets of EN219 that showed >3-fold competition with adjusted *p* value <0.05 against EN219-alkyne labeling in cells (Figure S2B; Table S3). We found 10 total targets that showed >3-fold enrichment with EN219-alkyne compared with DMSO-treated cells with adjusted *p* value <0.05 (Table S3). This number of potential EN219 off-targets would be consistent with the approximately eight protein bands observed by gel-based ABPP in Figure S2D. None of these seven to ten potential EN219 off-targets were E3 ligases. Collectively, our results suggested that EN219 was a moderately selective covalent ligand against C8 of RNF114. While we did not observe RNF114 by TMT-based proteomic experiments, RNF114 was clearly enriched from 231MFP cells treated with EN219-alkyne *in situ* compared with vehicle-treated controls after CuAAC-mediated appendage of biotin-azide and subsequent avidin pull-down and RNF114 blotting (Figure 2I).


### EN219-based RNF114 recruiter in TPD applications

We next tested whether EN219 could be exploited for TPD applications. We had previously demonstrated that nimbolide could be linked to the BET inhibitor ligand JQ1 to selectively degrade BRD4. Thus, we benchmarked our EN219 RNF114 recruiter by linking EN219 to JQ1 through three different linkers—ML 2-14,

ML 2-31, and ML 2-32 with C4 alkyl, C7 alkyl, and polyethylene glycol (PEG4) linkers, respectively (Figures 3A and S3A). Similar to effects seen with nimbolide-based degraders, ML 2-14 with the shortest linker showed the most robust degradation of BRD4 in 231MFP breast cancer cells compared with ML 2-31 and ML 2-32 with longer linkers, with 50% degradation concentration values of 36 and 14 nM for the long and short isoforms of BRD4, respectively (Figures 3B 3C, S3B, and S3C) (Spradlin et al., 2019). EN219 treatment alone did not affect BRD4 levels, compared with ML 2-14 (Figure S4A). This ML 2-14-mediated degradation was fully averted by pre-treating cells with the proteasome inhibitor bortezomib as well as the E1-activating enzyme inhibitor TAK-243 (Figures 3D–3F and S4B).

To further validate that ML 2-14 degradation of BRD4 was driven through RNF114, we showed that nimbolide pre-treatment completely attenuated BRD4 degradation by ML 2-14 in 231MFP breast cancer cells (Figures 3G and 3H). BRD4 degradation in HAP1 cells was also significantly attenuated in RNF114 knockout cells compared with wild-type counterparts in two independent experiments (Figures 3I, 3J, S4C, and S4D). We note that these purchased RNF114 knockout cells showed residual RNF114 protein expression, likely indicating a mixed population of cells (Figures 3I–3J). Furthermore, we showed that this loss of BRD4 protein levels was not due to transcriptional downregulation of BRD4 expression since ML 2-14 treatment in 231MFP cells did not alter BRD4 mRNA levels (Figure S4E). TMT-based quantitative proteomic profiling of ML 2-14-mediated protein expression changes showed selective degradation of BRD3 and BRD4, but not BRD2. We also observed stabilization of two known or putative RNF114 substrates, including the tumor suppressor CDKN1A (p21) and CTGF (Figure 3K; Table S4) (Han et al., 2013; Spradlin et al., 2019).

To further demonstrate the utility of our fully synthetic RNF114 recruiter EN219 in degrading other more challenging protein targets, we synthesized a degrader linking EN219 to the BCR-ABL inhibitor dasatinib, ML 2-23 and ML 2-22, bearing a longer PEG3 linker and a shorter C3 alkyl linker, respectively (Figures 4A and S4F). For this particular target, ML 2-23 with the longer linker showed more robust degradation of BCR-ABL in K562 leukemia cells compared with ML 2-22, consistent with previously observed structure-activity relationships of nimbolide-based BCR-ABL degraders (Figures 4B, 4C, and S4G) (Tong et al., 2020). Consistent with ML 2-23 engaging BCR-ABL in cells, we observed inhibition of CRKL phosphorylation, a downstream substrate of BCR-ABL signaling (Figures 4B and 4C). Interestingly, EN219 showed preferential degradation of BCR-ABL compared with c-ABL, compared with several previous BCR-ABL degraders utilizing cereblon or VHL recruiters that showed opposite selectivity (Figures 4B and 4C) (Burslem et al., 2019; Lai et al., 2016). This preferential degradation was also observed with the equivalent nimbolide-based degrader (Tong et al., 2020). We further showed that this loss of BCR-ABL and c-ABL was not due to transcriptional downregulation of these genes, since BCR-ABL mRNA levels remained unchanged and we observed a likely compensatory increase in c-ABL mRNA levels (Figure S4H). Furthermore, we do not believe that the loss of BCR-ABL shown here is because of general cytotoxicity since ML 2-23 treatment at the time points used to assess



**Figure 3. EN219-based BRD4 degrader**

(A) Structure of ML 2-14, an EN219-based BRD4 degrader linking EN219 to BET inhibitor JQ1.  
 (B) Degradation of BRD4 by ML 2-14. 231MFP cells were treated with DMSO vehicle or ML 2-14 for 8 h and the long and short isoforms of BRD4 and loading control actin were detected by western blotting.  
 (C) Percentage of BRD4 degradation quantified from (B) shows 50% degradation concentrations of 36 and 14 nM for the long and short isoforms of BRD4, respectively.  
 (D) Proteasome-dependent degradation of BRD4 by ML 2-14. 231MFP cells were treated with DMSO vehicle or with proteasome inhibitor bortezomib (1  $\mu$ M) 30 min before DMSO vehicle or ML 2-14 (100 nM) treatment for 8 h. BRD4 and loading control actin levels were detected by western blotting.  
 (E) Quantification of BRD4 levels from (D).  
 (F) Ubiquitin-activating enzyme (E1)-dependent degradation of BRD4 by ML 2-14. 231MFP cells were treated with DMSO vehicle or with E1 inhibitor TAK-243 (10  $\mu$ M) 30 min before DMSO vehicle or ML 2-14 (100 nM) treatment for 8 h. BRD4 and loading control actin levels were detected by western blotting. Quantification of blot can be found in [Figure S4B](#).  
 (G) BRD4 degradation by ML 2-14 is attenuated by nimbolide. 231MFP cells were treated with DMSO vehicle or with nimbolide (4  $\mu$ M) 30 min before DMSO vehicle or ML 2-14 (100 nM) treatment for 8 h.

(legend continued on next page)

BCR-ABL degradation only exerted relatively modest cell viability impairments in K562 cells; EN219 compromises K562 viability only at higher concentrations (Figure S4I). While rescue experiments with proteasome inhibitors proved challenging due to the cytotoxicity of proteasome inhibitors at the long time points required for robust BCR-ABL degradation, we observed significant rescue of early-stage ML 2-23-mediated BCR-ABL and c-ABL degradation with pre-treatment of K562 cells with the proteasome inhibitor MG132 at a shorter time point (Figures 4D and 4E).

## DISCUSSION

We previously discovered that the natural product nimbolide targets a predicted intrinsically disordered cysteine in RNF114 at a substrate recognition site (Spradlin et al., 2019). Here, we report EN219 as a moderately selective covalent ligand that exploits this same binding modality. We show that EN219 can be linked to the BET inhibitor JQ1 to degrade BRD4 in a nimbolide-sensitive and proteasome- and RNF114-dependent manner. We further show that EN219 can be linked to the kinase inhibitor dasatinib to selectively degrade BCR-ABL over c-ABL in a proteasome-dependent manner in leukemia cells. These findings also further highlight that moderately selective cysteine-targeting ligands can still lead to robust protein degraders (Ward et al., 2019; Zhang et al., 2019). While EN219 represents a promising initial chemical scaffold for RNF114 recruitment, future medicinal chemistry efforts can further optimize the potency and metabolic stability of RNF114 recruiters; in addition, the facile synthesis of pyrazoline derivatives makes this a much more tractable synthetic problem when compared with the necessary chemistry to realize numerous nimbolide derivatives.

Overall, our results highlight the utility of chemoproteomic platforms for discovering chemical scaffolds that can be used as E3 ligase recruiters for TPD applications. Moreover, these studies further speak to the power of unbiased chemoproteomic approaches for identification of synthetic ligands mimicking the function of covalently acting natural products, particularly those in which structural binding information is unavailable (Nomura and Maimone, 2019).

## SIGNIFICANCE

**We previously discovered that the natural product nimbolide could be used as a unique and covalent RNF114 E3 ubiquitin ligase recruiter for targeted protein degradation**

(TPD) applications. However, the adoption of this natural product in degraders has been hindered by the synthetic difficulty of using a complex natural product as a starting point for degrader synthesis. Here, using chemoproteomics-enabled covalent ligand-screening approaches, we have discovered a more synthetically tractable covalent ligand EN219 that targets RNF114 and mimics nimbolide's mode of action. We demonstrate that EN219, like nimbolide, can be incorporated into bifunctional degrader compounds to degrade neosubstrate proteins in an RNF114-dependent manner. Our study highlights the utility of covalent ligand screening in target-based screening endeavors to expand the toolbox of E3 ligase recruiters for TPD applications—an area of great importance in drug discovery and chemical biology.

## STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:


- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
  - Lead contract
  - Materials availability
  - Data and code availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
  - Cell lines
- METHOD DETAILS
  - Chemicals
  - Cell-based degrader assays
  - Gel-based ABPP
  - EN219-alkyne probe labeling *in situ* and pulldown studies
  - LC-MS/MS analysis of pure RNF114 EN219 modification
  - IsoTOP-ABPP
  - Mass spectrometry analysis
  - TMT-based quantitative proteomic profiling
  - RNF114 ubiquitination assay
  - Western blotting
  - Synthetic methods and characterization
  - ML 2-14
  - ML 2-22
  - ML 2-23
  - EN219-alkyne
  - ML 2-31

(H) Quantification of BRD4 levels from (G).

(I) Degradation of BRD4 in RNF114 wild-type (WT) or knockout (KO) HAP1 cells. RNF114 WT or KO HAP1 cells were treated with DMSO vehicle or ML 2-14 (1  $\mu$ M) for 16 h, and BRD4, RNF114, and loading control actin levels were detected by western blotting.

(J) Quantification of data from (I).

(K) TMT-based quantitative proteomic data showing ML 2-14-mediated protein level changes in 231MFP cells. 231MFP cells were treated with DMSO vehicle or ML 2-14 for 8 h. Shown are average TMT ratios for all tryptic peptides identified with at least two unique peptides. Shown in red are proteins that showed >2-fold increased or decreased levels with ML 2-14 treatment with adjusted p values <0.05. Data in (B–K) are from n = 3 biologically independent replicates/group. Data shown in (C) are average and individual replicate values. Data shown in (E, H, and J) are averages  $\pm$  SEM values and also individual biological replicate values. Blots shown in (B, D, F, G, and I) are representative of n = 3 biologically independent replicates/group. Statistical significance was calculated with unpaired two-tailed Student's t test and is expressed as \*p < 0.05 compared with vehicle-treated controls in (E, H, and J), and #p < 0.05 compared with ML 2-14-treated groups in (E and H) and compared with ML 2-14-treated RNF114 WT cells in (J). This figure is related to Figures S3 and S4.



- ML 2-32
- Deschloro-EN219

# ● QUANTIFICATION AND STATISTICAL ANALYSIS

## SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.chembiol.2021.01.005>.

## Figure 4. EN219-based ABL degrader

(A) Structure of ML 2-23, an EN219-based ABL degrader linking EN219 to ABL inhibitor dasatinib. (B) Degradation of BCR-ABL and c-ABL by ML 2-23. K562 cells were treated with DMSO vehicle or ML 2-23 for 16 h, and BCR-ABL, c-ABL, p-CRKL, and loading control actin were detected by western blotting. (C) Percentage of BCR-ABL, c-ABL, and p-CRKL protein levels quantified from (B). (D) Proteasome-dependent degradation of BCR-ABL and c-ABL by ML 2-23. K562 cells were treated with DMSO vehicle or with proteasome inhibitor MG132 (5 μM) 30 min before DMSO vehicle or ML 2-23 (1 μM) treatment for 12 h. BCR-ABL, c-ABL, and loading control actin levels were detected by western blotting. (E) Quantification of BCR-ABL and c-ABL levels from (D). Data shown in (C and E) are averages ± SEM values and also individual biological replicate values. Blots shown in (B and D) are representative of n = 3 biological replicates/group. Statistical significance was calculated with unpaired two-tailed Student's t test and is expressed as \*p < 0.05 compared with vehicle-treated controls in (C and E) and #p < 0.05 compared with ML 2-23-treated groups in (E). This figure is related to Figure S4.

## ACKNOWLEDGMENTS

We thank the members of the Nomura Research Group and Novartis Institutes for BioMedical Research for critical reading of the manuscript. This work was supported by Novartis Institutes for BioMedical Research and the Novartis-Berkeley Center for Proteomics and Chemistry Technologies (NB-CPACT) for all listed authors. This work was also supported by the Nomura Research Group and the Mark Foundation for Cancer Research ASPIRE Award to D.K.N., M.L., J.N.S., and L.B. This work was also supported by grants from

the NIH (R01CA240981 to D.K.N., T.J.M., M.L., J.N.S., and L.B. and F31CA239327 and F99CA253717 to J.N.S.).

#### AUTHOR CONTRIBUTIONS

M.L., J.N.S., T.J.M., and D.K.N. conceived the project and wrote the paper. M.L., J.N.S., L.B., B.T., D.K.N., J.A.T., M.S., J.M.McK., S.M.B., and T.J.M. provided intellectual contributions and insights into project direction. M.L., J.N.S., L.B., B.T., S.M.B., M.S., T.J.M., and D.K.N. designed the experiments. M.L., J.N.S., L.B., B.T., S.M.B., and D.K.N. performed experiments and analyzed data. M.L., J.N.S., L.B., B.T., J.A.T., J.M.McK., M.S., S.M.B., T.J.M., and D.K.N. edited the paper.

#### DECLARATION OF INTERESTS

J.A.T., J.M.McK., M.S., and S.M.B. are employees of Novartis Institutes for BioMedical Research. This study was funded by the Novartis Institutes for BioMedical Research and the Novartis-Berkeley Center for Proteomics and Chemistry Technologies. D.K.N. is a co-founder, shareholder, and adviser for Frontier Medicines.

Received: July 12, 2020

Revised: December 10, 2020

Accepted: January 5, 2021

Published: January 28, 2021

#### REFERENCES

- Bachovchin, D.A., Ji, T., Li, W., Simon, G.M., Blankman, J.L., Adibekian, A., Hoover, H., Niessen, S., and Cravatt, B.F. (2010). Superfamily-wide portrait of serine hydrolase inhibition achieved by library-versus-library screening. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 20941–20946.
- Backus, K.M., Correia, B.E., Lum, K.M., Forli, S., Horning, B.D., González-Páez, G.E., Chatterjee, S., Lanning, B.R., Teijaro, J.R., Olson, A.J., et al. (2016). Proteome-wide covalent ligand discovery in native biological systems. *Nature* **534**, 570–574.
- Bond, M.J., Chu, L., Nalawansa, D.A., Li, K., and Crews, C.M. (2020). Targeted degradation of oncogenic KRASG12C by VHL-recruiting PROTACs. *ACS Cent. Sci.* **6**, 1367–1375.
- Bondeson, D.P., and Crews, C.M. (2017). Targeted protein degradation by small molecules. *Annu. Rev. Pharmacol. Toxicol.* **57**, 107–123.
- Bondeson, D.P., Smith, B.E., Burslem, G.M., Buhimschi, A.D., Hines, J., Jaime-Figueroa, S., Wang, J., Hamman, B.D., Ishchenko, A., and Crews, C.M. (2018). Lessons in PROTAC design from selective degradation with a promiscuous warhead. *Cell Chem. Biol.* **25**, 78–87.e5.
- Burslem, G.M., Schultz, A.R., Bondeson, D.P., Eide, C.A., Savage Stevens, S.L., Druker, B.J., and Crews, C.M. (2019). Targeting BCR-ABL1 in chronic myeloid leukemia by PROTAC-mediated targeted protein degradation. *Cancer Res.* **79**, 4744–4753.
- Chamberlain, P.P., and Hamann, L.G. (2019). Development of targeted protein degradation therapeutics. *Nat. Chem. Biol.* **15**, 937–944.
- Grossman, E.A., Ward, C.C., Spradlin, J.N., Bateman, L.A., Huffman, T.R., Miyamoto, D.K., Kleinman, J.I., and Nomura, D.K. (2017). Covalent ligand discovery against druggable hotspots targeted by anti-cancer natural products. *Cell Chem. Biol.* **24**, 1368–1376.e4.
- Han, J., Kim, Y.-L., Lee, K.-W., Her, N.-G., Ha, T.-K., Yoon, S., Jeong, S.-I., Lee, J.-H., Kang, M.-J., Lee, M.-G., et al. (2013). ZNF313 is a novel cell cycle activator with an E3 ligase activity inhibiting cellular senescence by destabilizing p21(WAF1). *Cell Death Differ.* **20**, 1055–1067.
- Huang, H.-T., Dobrovolsky, D., Paulk, J., Yang, G., Weisberg, E.L., Doctor, Z.M., Buckley, D.L., Cho, J.-H., Ko, E., Jang, J., et al. (2018). A chemoproteomic approach to query the degradable kinome using a multi-kinase degrader. *Cell Chem. Biol.* **25**, 88–99.e6.
- Jessani, N., Humphrey, M., McDonald, W.H., Niessen, S., Masuda, K., Gangadharan, B., Yates, J.R., Mueller, B.M., and Cravatt, B.F. (2004). Carcinoma and stromal enzyme activity profiles associated with breast tumor growth in vivo. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 13756–13761.
- Lai, A.C., and Crews, C.M. (2017). Induced protein degradation: an emerging drug discovery paradigm. *Nat. Rev. Drug Discov.* **16**, 101–114.
- Lai, A.C., Toure, M., Hellerschmied, D., Salami, J., Jaime-Figueroa, S., Ko, E., Hines, J., and Crews, C.M. (2016). Modular PROTAC design for the degradation of oncogenic BCR-ABL. *Angew. Chem. Int. Ed. Engl.* **55**, 807–810.
- Mahé, O., Dez, I., Levacher, V., and Brière, J.-F. (2010). Enantioselective phase-transfer catalysis: synthesis of pyrazolines. *Angew. Chem. Int. Ed.* **49**, 7072–7075.
- Newman, D.J., and Cragg, G.M. (2016). Natural products as sources of new drugs from 1981 to 2014. *J. Nat. Prod.* **79**, 629–661.
- Nomura, D.K., and Maimone, T.J. (2019). Target identification of bioactive covalently acting natural products. *Curr. Top. Microbiol. Immunol.* **420**, 351–374.
- Ottis, P., Palladino, C., Thienger, P., Britschgi, A., Heichinger, C., Berrera, M., Julien-Laferrère, A., Roudnicky, F., Kam-Thong, T., Bischoff, J.R., et al. (2019). Cellular resistance mechanisms to targeted protein degradation converge toward impairment of the engaged ubiquitin transfer pathway. *ACS Chem. Biol.* **14**, 2215–2223.
- Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675.
- Spradlin, J.N., Hu, X., Ward, C.C., Brittain, S.M., Jones, M.D., Ou, L., To, M., Proudfoot, A., Ornelas, E., Woldegiorgis, M., et al. (2019). Harnessing the anti-cancer natural product nimbolide for targeted protein degradation. *Nat. Chem. Biol.* **15**, 747–755.
- Thomas, J.R., Brittain, S.M., Lipps, J., Llamas, L., Jain, R.K., and Schirle, M. (2017). A photoaffinity labeling-based chemoproteomics strategy for unbiased target deconvolution of small molecule drug candidates. *Methods Mol. Biol.* **1647**, 1–18.
- Tong, B., Spradlin, J.N., Novaes, L.F.T., Zhang, E., Hu, X., Moeller, M., Brittain, S.M., McGregor, L.M., McKenna, J.M., Tallarico, J.A., et al. (2020). A nimbolide-based kinase degrader preferentially degrades oncogenic BCR-ABL. *ACS Chem. Biol.* **15**, 1788–1794.
- Wang, C., Weerapana, E., Blewett, M.M., and Cravatt, B.F. (2014). A chemoproteomic platform to quantitatively map targets of lipid-derived electrophiles. *Nat. Methods* **11**, 79–85.
- Ward, C.C., Kleinman, J.I., Brittain, S.M., Lee, P.S., Chung, C.Y.S., Kim, K., Petri, Y., Thomas, J.R., Tallarico, J.A., McKenna, J.M., et al. (2019). Covalent ligand screening uncovers a RNF4 E3 ligase recruiter for targeted protein degradation applications. *ACS Chem. Biol.* **14**, 2430–2440.
- Weerapana, E., Wang, C., Simon, G.M., Richter, F., Khare, S., Dillon, M.B.D., Bachovchin, D.A., Mowen, K., Baker, D., and Cravatt, B.F. (2010). Quantitative reactivity profiling predicts functional cysteines in proteomes. *Nature* **468**, 790–795.
- Xu, T., Park, S.K., Venable, J.D., Wohlschlegel, J.A., Diedrich, J.K., Cociorva, D., Lu, B., Liao, L., Hewel, J., Han, X., et al. (2015). ProLuCID: an improved SEQUEST-like algorithm with enhanced sensitivity and specificity. *J. Proteomics* **129**, 16–24.
- Zhang, X., Crowley, V.M., Wucherpfennig, T.G., Dix, M.M., and Cravatt, B.F. (2019). Electrophilic PROTACs that degrade nuclear proteins by engaging DCAF16. *Nat. Chem. Biol.* **15**, 737–746.

# STAR★METHODS

## KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
<b>Antibodies</b>		
Rb Anti-RNF114	Sigma-Aldrich	HPA021184; RRID: AB_1859185
Mouse monoclonal c-Abl Antibody	Santa Cruz	sc-23; RRID: AB_626775
Rb Phospho-CrkL (Tyr207) Antibody	Cell Signaling Technology	Cat# 3181S; RRID: AB_331068
Mouse GAPDH Monoclonal Antibody	Proteintech Group Inc	60004-1-Ig; RRID: AB_2107436
Mouse Beta Actin Monoclonal antibody	Proteintech Group Inc	66009-1-Ig; RRID: AB_2687938
Rb Anti-Brd4 antibody	Abcam plc	Ab128874; RRID: AB_11145462
<b>Chemicals, peptides, and recombinant proteins</b>		
Myc-Flag-RNF114 human recombinant protein	Origene	TP309752
Recombinant protein of human cyclin-dependent kinase inhibitor 1A	Origene	TP301765
Ubiquitin Activating Enzyme (UBE1)	BostonBiochem	E-305
Ubiquitin-conjugating Enzyme E2D 1 (UBE2D1)	BostonBiochem	E2-615
FLAG (DYKDDDDK)-Ubiquitin	BostonBiochem	U-120
Dimethyl Sulfoxide	Fisher Chemical	D128-500 (CAS 67-68-5)
Dulbecco's Phosphate Buffered Saline (DPBS)	Sigma-Aldrich	D8537-500mL
L-15 Medium	Sigma-Aldrich	L1518-500mL
Iscove's Modified Dulbecco's Medium	Sigma-Aldrich	51471C-1000mL
Opti-MEM Reduced Serum Media	ThermoFisher Scientific	Cat#31985062
Iodoacetamide 98%	ACROS Organics	Cat# AC122270050
Iodoacetamide- <sup>13</sup> C <sub>2</sub> , 2-d <sub>2</sub>	Sigma-Aldrich	Cat#721328
Urea	Fisher Chemical	U15-500 (CAS 57-13-6)
ProteaseMAX Surfactant, lyophilized	Promega	REF#V2072
Ammonium Bicarbonate	Fisher Chemical	A643-500 (CAS 1066-33-7)
TCEP 99%	Strem Chemicals	Cat#15-7400 (CAS 51805-45-9)
Formic Acid, 99.0+%, Optima LC/MS Grade	Fisher Chemical	A117-50 (CAS 64-18-6)
Sequencing Grade Modified Trypsin, Porcine	Promega	V511A
Covalent Ligand Library	Enamine	N/A
Tetramethylrhodamine-5-Iodoacetamide Dihydroiodide (IA-Rhodamine)	Setareh Biotech	6222 (CAS 114458-99-0)
Laemmli SDS sample buffer, reducing (4x)	Alfa Aesar	J60015
Tris(benzyltriazole methylamine) (TBTA)	Sigma-Aldrich	678937 (CAS 510758-28-8)
Biotin-TEV-azide	<a href="#">Weerapana et al. (2010)</a>	N/A
Copper (II) Sulfate	Sigma-Aldrich	451657
AcTEV Protease	Invitrogen	12575-015
N-Hex-5-ynyl-2-iodo-acetamide (IAyne)	Chess GmbH	3187
Spectra Multicolor Broad Range Protein Ladder	Thermo Scientific	Prod#26634
Pierce Protease inhibitor Mini Tablets, EDTA-free	Thermo Scientific	Prod#A32955
High Capacity Streptavidin Agarose Resin	Thermo Scientific	Ref#20357
Streptavidin Agarose Resin	Thermo Scientific	Ref#20349

(Continued on next page)

### Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
4-hydroxybenzaldehyde	Alfa Aesar	A13580-22 (CAS 123-08-0)
3,4-Dihydro-2H-pyran	Sigma-Aldrich	D106208-100mL (CAS 110-87-2)
4'-Bromoacetophenone	Sigma-Aldrich	B56404-25G (CAS 99-90-1)
4-(Boc-amino) butyl bromide	Sigma-Aldrich	90303-500mg-F (CAS 164365-88-2)
Hydrazine monohydrate	Sigma-Aldrich	207942-5g (CAS 7803-57-8)
Chloroacetyl chloride	Sigma-Aldrich	22880-100mL (CAS 79-04-9)
DIPEA	Sigma-Aldrich	D125806-100mL (CAS 7087-68-5)
JQ1	eNovation Chemicals	D372464 (CAS 1268524-70-4)
HATU	Sigma-Aldrich	445460-5g (CAS 148893-10-1)
Propargylamine	Sigma-Aldrich	P50900-1g (CAS 2450-71-7)
N-Boc-7-bromoheptan-1-amine	ACheckBlock	P41297 (CAS 142356-34-1)
tert-Butyl (2-(2-(2-bromoethoxy)ethoxy)ethoxy)ethyl)carbamate	Synthonix	B37532 (CAS 1076199-21-7)
Pyridinium p-toluenesulfonate	Sigma-Aldrich	232238-5G (CAS 24057-28-1)
(E)-1,3-bis(4-bromophenyl)prop-2-en-1-one	Princeton Bio	PRIH93ED8598 (CAS 5471-96-5)
Nimbolide	Cayman Chemicals	19230 (CAS 25990-37-8)
Critical commercial assays		
TMT10plex Isobaric Label Reagent Set	ThermoFisher	A37725
WST8 reagent	APEXBio	CCK-8
Monarch Total RNA Miniprep Kit	New England BioLabs	T2010S
ProtoScript II reverse transcriptase Kit	New England BioLabs	M0368L
DyNAmo HS SYBR Green Kit	Fisher Scientific	F-410L
Deposited Data		
Experimental models: cell lines		
231MFP	( <a href="#">Jessani et al., 2004</a> )	N/A
K562	ATCC	CCL-243
HAP1	Horizon Discovery	HZGHC55905
HAP1 (RNF114 Knockout)	Horizon Discovery	HZGHC55905
Software and algorithms		
ImageJ	( <a href="#">Schneider et al., 2012</a> )	<a href="https://imagej.nih.gov/ij/">https://imagej.nih.gov/ij/</a>
IP2 proteomics pipeline 5.0.1	Integrated Proteomics Applications	N/A

## RESOURCE AVAILABILITY

### Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Daniel K. Nomura, [dnomura@berkeley.edu](mailto:dnomura@berkeley.edu).

### Materials availability

Plasmids, compounds generated in this study will be made available upon reasonable request.

### Data and code availability

Data generated in this study will be made available upon reasonable request. No code was developed for this study.

## EXPERIMENTAL MODEL AND SUBJECT DETAILS

### Cell lines

The 231MFP cells were obtained from Prof. Benjamin Cravatt and were generated from explanted tumor xenografts of female MDA-MB-231 cells as previously described ([Jessani et al., 2004](#)). The 231MFP cells were cultured in L15 medium containing

10% FBS and maintained at 37°C with 0% CO<sub>2</sub>. K562 chronic myeloid leukemia cell lines of female origin were purchased from ATCC. The K562 cells were cultured in Iscove's Modified Dulbecco's Medium containing 10% FBS and maintained at 37°C with 5% CO<sub>2</sub>. HAP1 RNF114 wild-type and knockout cell lines of female origin were purchased from Horizon Discovery. The RNF114 knockout cell line was generated by CRISPR/Cas9 to contain a frameshift mutation in a coding exon of RNF114. HAP1 cells were grown in Iscove's Modified Dulbecco's Medium in the presence of 10% FBS and penicillin/streptomycin.

## METHOD DETAILS

### Chemicals

Covalent ligands screened against RNF114 were purchased from Enamine LLC, including EN219. Structures of compounds screened can be found in [Table S1](#). See [Supplemental information](#) for synthetic methods and characterization for EN219 and degraders. Nimbolide was purchased from Cayman Chemicals (Item No. 19230).

### Cell-based degrader assays

For assaying degrader activity, cells were seeded (500,000 cells for 231MFP and HAP1 cells, 1,000,000 for K562 cells) into a 6 cm tissue culture dish (Corning) in 2.0–2.5 mL of media and allowed to adhere overnight for 231MFP and HAP1 cells. The following morning, media was replaced with complete media containing the desired concentration of compound diluted from a 1,000 x stock in DMSO. For rescue studies, the cells were pre-treated with proteasome inhibitors, nimbolide, or E1 activating enzyme inhibitors 30 min prior to the addition of DMSO or degrader compounds. At the specified time point, cells were washed once with PBS on ice, before addition of 120 µL of lysis buffer (20 mM Tris-HCl at pH 7.5, 150 mM NaCl, 1 mM Na<sub>2</sub>EDTA, 1 mM EGTA, 1% Triton, 2.5 mM sodium pyrophosphate, 1 mM beta-glycerophosphate, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1 µg/ml leupeptin) with Complete Protease Inhibitor Cocktail (Sigma) was added. The cells were incubated in lysis buffer for 5 min before scraping and transferring to microcentrifuge tubes. The lysates were then frozen at –80°C or immediately processed for Western blotting. To prepare for Western blotting, the lysates were cleared with a 20,000g spin for 10 min and the resulting supernatant protein concentrations were quantified via BCA assay. The lysates were normalized by dilution with PBS to match the lowest concentration lysate, and the appropriate amount of 4 x Laemmli's reducing buffer was added.

### Gel-based ABPP

Gel-Based ABPP methods were performed as previously described ([Ward et al., 2019](#)). Pure recombinant human RNF114 was purchased from Boston Biochem (K-220). RNF114 (0.25 µg) was diluted into 50 µL of PBS and 1 µL of either DMSO (vehicle) or covalently acting small molecule to achieve the desired concentration. After 30 min at room temperature, the samples were treated with 250 nM of tetramethylrhodamine-5-iodoacetamide dihydroiodide (IA-Rhodamine) (Setareh Biotech, 6222, prepared in anhydrous DMSO) for 1 h at room temperature. Incubations were quenched by diluting the incubation with 20 µL of 4 x reducing Laemmli SDS sample loading buffer (Alfa Aesar) and heated at 90°C for 5 min. The samples were separated on precast 4–20% Criterion TGX gels (Bio-Rad Laboratories, Inc.). Fluorescent imaging was performed on a ChemiDoc MP (Bio-Rad Laboratories, Inc.). Inhibition of target labeling was assessed by densitometry using ImageJ.

### EN219-alkyne probe labeling *in situ* and pulldown studies

Experiments were performed following an adaption of a previously described protocol ([Thomas et al., 2017](#)). The 231MFP cells were treated with either DMSO vehicle or 50 µM EN219-alkyne probe for 90 min. Cells were collected in PBS and lysed by sonication. For preparation of Western blotting samples, the lysate (1 mg of protein in 500 µl) was aliquoted per sample and then the following were added: 10 µl of 5 mM biotin picolylazide (900912 Sigma-Aldrich) and 50 µl of click reaction mix (three parts TBTA 5 mM TBTA in butanol:DMSO (4:1, v/v), one part 50 mM Cu(II)SO<sub>4</sub> solution and one part 50 mM TCEP). Samples were incubated for 1 h at room temperature with gentle agitation. After CuAAC, proteomes were precipitated by centrifugation at 6,500 g and washed twice in ice-cold methanol (500 µl). The samples were spun in a prechilled (4°C) centrifuge at 6,500 g for 4 min allowing for aspiration of excess methanol and subsequent reconstitution of protein pellet in 250 µl PBS containing 1.2% SDS by probe sonication. Then the proteome was denatured at 90°C for 5 min, the insoluble components were precipitated by centrifugation at 6,500g and soluble proteome was diluted in 1.2 ml PBS (the final concentration of SDS in the sample was 0.2%) to a total volume of 1450 µl, with 50 µl reserved as input. Then 85 µl of prewashed 50% streptavidin agarose bead slurry was added to each sample and samples were incubated overnight at room temperature with gentle agitation. Supernatant was aspirated from each sample after spinning beads at 6,500 g for 2 min at room temperature. Beads were transferred to spin columns and washed three times with PBS. To elute, beads were boiled 5 min in 50 µl LDS sample buffer. Eluents were collected after centrifugation and analyzed by immunoblotting. The resulting samples were also analyzed as described below for TMT-based quantitative proteomic profiling.

### LC-MS/MS analysis of pure RNF114 EN219 modification

Purified RNF114 (10 µg) in 50 µl PBS was incubated 30 min at room temperature either with DMSO vehicle or EN219 (50 µM). The DMSO control was then treated with light iodoacetamide while the compound treated sample was

incubated with heavy iodoacetamide for 1 h each at room temperature (200  $\mu$ M final concentration, Sigma-Aldrich, 721328). The samples were precipitated by addition of 12.5  $\mu$ l of 100% (w/v) trichloroacetic acid and the treated and control groups were combined pairwise, before cooling to  $-80^{\circ}\text{C}$  for 1 h. The combined sample was then spun for at max speed for 10 min at  $4^{\circ}\text{C}$ , supernatant was carefully removed and the sample was washed with ice-cold 0.01 M HCl/90% acetone solution. The pellet was resuspended in 4 M urea containing 0.1% Protease Max (Promega Corp. V2071) and diluted in 40 mM ammonium bicarbonate buffer. The samples were reduced with 10 mM TCEP at  $60^{\circ}\text{C}$  for 30 min. The sample was then diluted 50% with PBS before sequencing grade trypsin (1  $\mu$ g per sample, Promega Corp, V5111) was added for an overnight incubation at  $37^{\circ}\text{C}$ . The next day, the sample was centrifuged at 13,200 rpm for 30 min. The supernatant was transferred to a new tube and acidified to a final concentration of 5% formic acid and stored at  $-80^{\circ}\text{C}$  until mass spectrometry analysis.

### IsoTOP-ABPP

IsoTOP-ABPP studies were done as previously reported (Spradlin et al., 2019). Cells were lysed by probe sonication in PBS and protein concentrations were measured by BCA assay<sup>35</sup>. For *in situ* experiments, cells were treated for 90 min with either DMSO vehicle or covalently acting small molecule (from 1,000 $\times$  DMSO stock) before cell collection and lysis. Proteomes were subsequently labeled with *N*-5-Hexyn-1-yl-2-iodoacetamide (IA-alkyne) labeling (100  $\mu$ M) for 1 h at room temperature. CuAAC was used by sequential addition of tris(2-carboxyethyl) phosphine (1 mM, Sigma), tris[(1-benzyl-1H-1,2,3-triazol-4-yl) methyl]amine (34  $\mu$ M, Sigma), copper(II) sulfate (1 mM, Sigma) and biotin-linker-azide—the linker functionalized with a tobacco etch virus (TEV) protease recognition sequence as well as an isotopically light or heavy valine for treatment of control or treated proteome, respectively. After CuAAC, proteomes were precipitated by centrifugation at 6,500g, washed in ice-cold methanol, combined in a 1:1 control:treated ratio, washed again, then denatured and resolubilized by heating in 1.2% SDS–PBS to  $80^{\circ}\text{C}$  for 5 min. Insoluble components were precipitated by centrifugation at 6,500g and soluble proteome was diluted in 5 ml 0.2% SDS–PBS. Labeled proteins were bound to avidin-agarose beads (170  $\mu$ l resuspended beads per sample, Thermo Pierce) while rotating overnight at  $4^{\circ}\text{C}$ . Bead-linked proteins were enriched by washing three times each in PBS and water, then resuspended in 6 M urea and PBS (Sigma), and reduced in TCEP (1 mM, Sigma), alkylated with iodoacetamide (18 mM, Sigma), before being washed and resuspended in 2 M urea and trypsinized overnight with 0.5  $\mu$ g  $\mu$ l<sup>-1</sup> sequencing grade trypsin (Promega). Tryptic peptides were eluted off. Beads were washed three times each in PBS and water, washed in TEV buffer solution (water, TEV buffer, 100  $\mu$ M dithiothreitol) and resuspended in buffer with Ac-TEV protease and incubated overnight. Peptides were diluted in water and acidified with formic acid (1.2 M, Spectrum) and prepared for analysis.

### Mass spectrometry analysis

Total peptides from TEV protease digestion for isoTOP-ABPP or tryptic peptides for mapping EN219 site of modification on pure RNF114 were pressure loaded onto 250 mm tubing packed with Aqua C18 reverse phase resin (Phenomenex no. 04A-4299), which was previously equilibrated on an Agilent 600 series high-performance liquid chromatograph using the gradient from 100% buffer A to 100% buffer B over 10 min, followed by a 5 min wash with 100% buffer B and a 5 min wash with 100% buffer A. The samples were then attached using a MicroTee PEEK 360  $\mu$ m fitting (Thermo Fisher Scientific no. p-888) to a 13 cm laser pulled column packed with 10 cm Aqua C18 reverse-phase resin and 3 cm of strong-cation exchange resin for isoTOP-ABPP studies. Samples were analyzed using an Q Exactive Plus mass spectrometer (Thermo Fisher Scientific) using a five-step Multidimensional Protein Identification Technology (MudPIT) program, using 0, 25, 50, 80 and 100% salt bumps of 500 mM aqueous ammonium acetate and using a gradient of 5–55% buffer B in buffer A (buffer A: 95:5 water:acetonitrile, 0.1% formic acid; buffer B 80:20 acetonitrile:water, 0.1% formic acid). Data were collected in datadependent acquisition mode with dynamic exclusion enabled (60 s). One full mass spectrometry ( $\text{MS}^1$ ) scan (400–1,800 mass-to-charge ratio ( $m/z$ )) was followed by 15  $\text{MS}^2$  scans of the *n*th most abundant ions. Heated capillary temperature was set to  $200^{\circ}\text{C}$  and the nanospray voltage was set to 2.75 kV, as previously described.

Data were extracted in the form of  $\text{MS}^1$  and  $\text{MS}^2$  files using Raw Extractor v.1.9.9.2 (Scripps Research Institute) and searched against the Uniprot human database using ProLuCID search methodology in IP2 v.3 (Integrated Proteomics Applications, Inc.) (Xu et al., 2015). Probe-modified cysteine residues were searched with a static modification for carboxyamino-methylation (+57.02146) and up to two differential modifications for methionine oxidation and either the light or heavy TEV tags (+464.28596 or +470.29977, respectively) or the mass of the EN219 adduct. Peptides were required to be fully tryptic peptides and to contain the TEV modification. ProLuCID data was filtered through DTASelect to achieve a peptide false-positive rate below 5%. Only those probe-modified peptides that were evident across two out of three biological replicates were interpreted for their isotopic light to heavy ratios. Light versus heavy isotopic probe-modified peptide ratios are calculated by taking the mean of the ratios of each replicate paired light vs. heavy precursor abundance for all peptide spectral matches (PSM) associated with a peptide. The paired abundances were also used to calculate a paired sample t-test p-value in an effort to estimate constancy within paired abundances and significance in change between treatment and control. P-values were corrected using the Benjamini/Hochberg method.

### TMT-based quantitative proteomic profiling

TMT proteomic profiling was performed as previously described (Spradlin et al., 2019).

### RNF114 ubiquitination assay


Recombinant Myc-Flag-RNF114 proteins were purchased from Origene (Origene Technologies Inc., TP309752) or were purified as described previously (Spradlin et al., 2019). For *in vitro* auto-ubiquitination assay, 0.2  $\mu$ g of RNF114 in 25  $\mu$ l of TBS was pre-incubated with DMSO vehicle or the covalently acting compound for 30 min at room temperature. Subsequently, 0.1  $\mu$ g of UBE1 (Boston Biochem. Inc., E-305), 0.1  $\mu$ g UBE2D1 (Boston Biochem. Inc., e2-615), 5  $\mu$ g of Flag-ubiquitin (Boston Biochem. Inc., u-120) in a total volume of 25  $\mu$ l Tris buffer containing 2 mM ATP, 10 mM DTT and 10 mM  $MgCl_2$  were added to achieve a final volume of 50  $\mu$ l. For substrate-protein ubiquitination assays, 0.1  $\mu$ g of p21 (Origene) was added at this stage. The mixture was incubated at 37°C with agitation for 1.5 h. Then, 20  $\mu$ l of Laemmli's buffer was added to quench the reaction and proteins were analyzed by Western blot assay.

### Western blotting

Antibodies to RNF114 (Millipore Sigma, HPA021184), c-ABL (Santa Cruz, 24-11), p-CRKL (Tyr207, Cell Signaling Technology, 3181), GAPDH (Proteintech Group Inc., 60004-1-Ig), BRD4 (Abcam plc, Ab128874), and beta-actin (Proteintech Group Inc., 6609-1-Ig) were obtained from various commercial sources and dilutions were prepared per the recommended manufacturers' procedures. Proteins were resolved by SDS-PAGE and transferred to nitrocellulose membranes using the iBlot system (Invitrogen). Blots were blocked with 5% BSA in Tris-buffered saline containing Tween 20 (TBST) solution for 1 h at room temperature, washed in TBST and probed with primary antibody diluted in diluent, as recommended by the manufacturer, overnight at 4°C. Following washes with TBST, the blots were incubated in the dark with secondary antibodies purchased from Ly-Cor and used at 1:10,000 dilution in 5% BSA in TBST at room temperature. Blots were visualized using an Odyssey Li-Cor scanner after additional washes. If additional primary antibody incubations were required, the membrane was stripped using ReBlot Plus Strong Antibody Stripping Solution (EMD Millipore, 2504), washed and blocked again before being re-incubated with primary antibody. Blots were quantified and normalized to loading controls using Image J.

### Synthetic methods and characterization

**Scheme 1.** Synthetic route to degrader **ML 2-14**.




### (E)-tert-butyl (4-(4-(3-(4-bromophenyl)-3-oxoprop-1-en-1-yl)phenoxy)butyl)carbamate (**2**)

To a solution of enone **1** (226 mg, 0.75 mmol) in anhydrous DMF (8 mL) was added 4-(Boc-amino)butyl bromide (376 mg, 1.5 mmol) and  $K_2CO_3$  (414 mg, 3 mmol) and the reaction mixture was stirred at 60°C for 5 h under an atmosphere of nitrogen. Upon cooling, the inorganic salts were filtered off, the solution was diluted with EtOAc, washed with water, and the volatiles removed *in vacuo*. The crude material was purified by silica gel chromatography (25% EtOAc/hexanes) to yield 310 mg (87%) of **2**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.96–7.88 (m, 2H), 7.83 (d,  $J$  = 15.6 Hz, 1H), 7.72–7.59 (m, 4H), 7.39 (d,  $J$  = 15.6 Hz, 1H), 7.01–6.91 (m, 2H), 4.66 (s, 1H), 4.07 (t,  $J$  = 6.2 Hz, 2H), 3.25 (q,  $J$  = 6.8 Hz, 2H), 1.94–1.82 (m, 2H), 1.79–1.69 (m, 2H), 1.49 (s, 9H).

**tert-butyl (4-(4-(3-(4-bromophenyl)-1-(2-chloroacetyl)-4,5-dihydro-1H-pyrazol-5-yl)phenoxy)butyl)carbamate (3)**


To a solution of enone **2** (550 mg, 1.16 mmol) in EtOH was added hydrazine monohydrate (116 mg, 2.32 mmol) and the reaction mixture heated at 80°C for 5 h under a nitrogen atmosphere. The reaction was cooled to room temperature, diluted with water, and extracted with DCM. The combined organic phase was dried over anhydrous magnesium sulfate and then concentrated *in vacuo* to ~5 mL. [NOTE: All of the workup procedures should be performed quickly (<1.5 h total time) and the rotovap bath kept cool as the crude product can easily undergo autooxidation]. To the concentrated DCM solution was immediately added chloroacetyl chloride (157 mg, 1.39 mmol) and triethylamine (152 mg, 1.5 mmol) at 0°C. The reaction mixture was stirred in an ice bath for 30 minutes, then warmed to room temperature and stirred overnight under N<sub>2</sub>. Upon completion of the reaction, the reaction mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude was purified by silica gel column chromatography (40% EtOAc/hexanes) to yield 381 mg (58%) of **3**: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.60 (q, *J* = 8.8 Hz, 4H), 7.19 – 7.10 (m, 2H), 6.87 – 6.77 (m, 2H), 5.55 (dd, *J* = 11.7, 4.7 Hz, 1H), 4.55 (s, 2H), 3.94 (t, *J* = 6.2 Hz, 2H), 3.74 (dd, *J* = 17.8, 11.7 Hz, 1H), 3.24 – 3.11 (m, 3H), 1.85 – 1.75 (m, 2H), 1.65 (p, *J* = 7.1 Hz, 2H), 1.44 (s, 9H).

**ML 2-14**

*i.* Chloroacetamide **3** (185 mg, 0.327 mmol) was dissolved in DCM (2.5 mL) and trifluoroacetic acid (2.5 mL) was added dropwise slowly over 20 minutes. After an additional 20 minutes of stirring, the solvent was removed *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The deprotected amine was used directly in the next step without purification.

*ii.* The resulting TFA salt was dissolved in DCM (8 mL) and **JQ1-acid** (157 mg, 0.4 mmol), 1- [Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (202 mg, 0.530 mmol), and *N,N*-Diisopropylethylamine (168 mg, 1.31 mmol) were added. The mixture was stirred overnight with monitoring by TLC (5% MeOH in DCM, 100% EtOAc). Upon completion, the reaction mixture was concentrated *in vacuo* and directly purified by silica gel flash column chromatography (1-5% MeOH/DCM). The eluted fractions were insufficiently pure and those containing product were combined, concentrated and purified again by flash silica chromatography (100-0% EtOAc/DCM followed by 0-5% MeOH/DCM) to afford 88.4 mg (32%) of **ML 2-14**: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.66 – 7.46 (m, 4H), 7.39 (d, *J* = 8.2 Hz, 2H), 7.29 (d, *J* = 8.3 Hz, 2H), 7.13 (d, *J* = 8.3 Hz, 2H), 6.86 (t, *J* = 5.9 Hz, 1H), 6.80 (d, *J* = 8.3 Hz, 2H), 5.54 (dt, *J* = 11.7, 4.1 Hz, 1H), 4.63 (t, *J* = 7.0 Hz, 1H), 4.54 (s, 2H), 3.90 (t, *J* = 6.1 Hz, 2H), 3.72 (dd, *J* = 17.8, 11.7 Hz, 1H), 3.54 (dd, *J* = 14.3, 7.5 Hz, 1H), 3.33 (ddt, *J* = 30.5, 13.3, 6.8 Hz, 3H), 3.17 (dd, *J* = 17.9, 4.7 Hz, 1H), 2.64 (s, 3H), 2.39 (s, 3H), 1.84 – 1.72 (m, 2H), 1.69 (d, *J* = 7.4 Hz, 2H), 1.65 (s, 3H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>): δ 170.6, 164.1, 161.4, 158.8, 155.8, 154.5, 151.4, 150.0, 136.9, 136.7, 132.9, 132.3, 132.2, 131.1, 131.0, 130.6, 130.0, 129.9, 128.9, 128.4, 127.2, 125.3, 120.7, 115.1, 67.6, 60.4, 54.7, 42.3, 42.1, 39.7, 39.4, 31.7, 26.7, 26.4, 22.8, 14.5, 14.3, 13.2, 12.0, 11.6. **HRMS (ESI): calcd.** C<sub>40</sub>H<sub>39</sub>BrCl<sub>2</sub>N<sub>7</sub>O<sub>3</sub>S ([M + H]<sup>+</sup>): *m/z* 846.1390, found: 846.1399.

**Scheme 2.** Synthetic route to degraders **ML 2-22** and **ML 2-23**.



**(E)-tert-butyl 2-(4-(3-(4-bromophenyl)-3-oxoprop-1-en-1-yl)phenoxy)acetate (**4**)**

To a solution of **1** (153 mg, 0.50 mmol) in anhydrous DMF (8 mL) was added *tert*-butyl bromoacetate (146 mg, 0.75 mmol) and  $K_2CO_3$  (138 mg, 1 mmol) and the reaction mixture was heated to 60°C for 5 hours under an atmosphere of nitrogen. Upon cooling, the inorganic salts were filtered off, the solution was diluted with EtOAc, washed with water, and concentrated *in vacuo*. The crude was purified by silica gel chromatography (25% EtOAc/hexanes) to yield 150 mg (72%) of **4**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.97 – 7.89 (m, 2H), 7.82 (d,  $J$  = 15.6 Hz, 1H), 7.73 – 7.60 (m, 4H), 7.41 (d,  $J$  = 15.6 Hz, 1H), 7.01 – 6.93 (m, 2H), 4.61 (s, 2H), 1.54 (s, 9H).

**tert-butyl 2-(4-(3-(4-bromophenyl)-1-(2-chloroacetyl)-4,5-dihydro-1H-pyrazol-5-yl)phenoxy)acetate (**5**)**

To a solution of **4** (417 mg, 1.00 mmol) in EtOH was added hydrazine monohydrate (250 mg, 5.00 mmol) and the reaction mixture was heated at 80°C for 5 hours under an atmosphere of nitrogen. The reaction was cooled to room temperature, diluted with water, and extracted with DCM. The combined organic phase was dried over anhydrous magnesium sulfate and then concentrated *in vacuo* to ~5 mL [NOTE: All of the workup procedures should be performed quickly (<1.5 h total time) and the rotovap bath kept cool as the crude product can easily undergo autooxidation]. To the concentrated DCM solution was immediately added chloroacetyl chloride (169 mg, 1.50 mmol) and triethylamine (303 mg, 3.00 mmol) at 0°C under an atmosphere of nitrogen. The reaction was maintained at this temperature for 30 minutes and then warmed to room temperature and stirred overnight under  $N_2$ . The reaction mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (35 – 40% EtOAc/hexanes) to yield 140 mg (28%) of **5**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.69 – 7.58 (m, 4H), 7.24 – 7.16 (m, 2H), 6.93 – 6.84 (m, 2H), 5.60 (dd,  $J$  = 11.7, 4.6 Hz, 1H), 4.58 (d,  $J$  = 1.6 Hz, 2H), 4.52 (s, 2H), 3.78 (dd,  $J$  = 17.8, 11.8 Hz, 1H), 3.23 (dd,  $J$  = 17.8, 4.7 Hz, 1H), 1.52 (s, 9H).

**ML 2-22**

*i.* Compound **5** (15.2 mg, 0.030 mmol) was dissolved in DCM (1.5 mL) and trifluoroacetic acid (0.5 mL) added slowly over the course of 20 minutes. After stirring for an additional 20 minutes, the solvent was removed *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The crude material was used directly in the next step.

ii. The aforementioned crude carboxylic acid was dissolved in 2 mL DCM and **amine 6** (15.0 mg, 0.030 mmol), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (17.1 mg, 0.045 mmol), and *N,N*-Diisopropylethylamine (193.5 mg, 1.50 mmol) were added (Tong et al., 2020). The reaction was stirred overnight monitoring by TLC (20% Methanol in DCM). Upon completion, the reaction mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude material was purified by silica gel flash column chromatography (1-5% MeOH/DCM). The eluted fractions were insufficiently pure and those containing product were combined, concentrated and purified again by flash silica chromatography (100-0% EtOAc/DCM followed by 10-35% MeOH/DCM) to afford 17.1 mg (60%) of **ML 2-22**:  $^1\text{H NMR}$  (600 MHz, DMSO- $d_6$ )  $\delta$  11.47 (s, 1H), 9.88 (s, 1H), 8.23 (s, 1H), 8.11 (d,  $J$  = 18.2 Hz, 1H), 7.78 – 7.71 (m, 2H), 7.71 – 7.64 (m, 2H), 7.40 (dd,  $J$  = 7.8, 1.7 Hz, 1H), 7.34 – 7.23 (m, 2H), 7.18 – 7.10 (m, 2H), 6.97 – 6.88 (m, 2H), 6.06 (s, 1H), 5.54 (dd,  $J$  = 11.7, 4.7 Hz, 1H), 4.78 – 4.64 (m, 2H), 4.46 (s, 2H), 3.86 (dd,  $J$  = 18.2, 11.8 Hz, 1H), 3.50 (s, 4H), 3.31 – 3.08 (m, 4H), 2.41 (s, 6H), 2.31 (s, 2H), 2.25 (s, 3H), 1.64 (s, 2H);  $^{13}\text{C NMR}$  (151 MHz, DMSO)  $\delta$  167.0, 164.7, 162.8, 162.1, 161.9, 159.5, 156.6, 156.5, 154.3, 140.4, 138.4, 133.8, 133.1, 132.0, 131.4, 129.5, 129.0, 128.6, 128.4, 127.7, 126.5, 125.3, 123.7, 114.4, 82.2, 66.7, 59.2, 55.1, 54.5, 51.9, 43.1, 42.0, 41.4, 36.5, 29.5, 28.0, 25.1, 23.0, 22.0, 17.9, 13.5, 10.5; **HRMS (ESI)**: *calcd.*  $\text{C}_{42}\text{H}_{44}\text{BrCl}_2\text{N}_{10}\text{O}_4\text{S}$  ( $[\text{M} + \text{H}]^+$ ): *m/z* 933.1823, found: 933.1814.

### ML 2-23

i. Compound **5** (16.2 mg, 0.032 mmol) was dissolved in DCM (1.5 mL) and trifluoroacetic acid (0.5 mL) added slowly over the course of 20 minutes. After stirring for an additional 20 minutes, the solvent was removed *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The crude material was used directly in the next step.


ii. The aforementioned crude carboxylic acid was dissolved in DCM (2 mL) and **amine 7** (19.8 mg, 0.032 mmol), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (18.2 mg, 0.048 mmol), and *N,N*-Diisopropylethylamine (206.0 mg, 1.60 mmol) were added (Tong et al., 2020). The reaction mixture was stirred overnight with monitoring by TLC (20% Methanol in DCM). Upon completion, the reaction mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude material was purified by silica gel flash column chromatography (1-5% MeOH/DCM). The eluted fractions were insufficiently pure and those containing product were combined, concentrated and purified again by flash silica chromatography (100-0% EtOAc/DCM followed by 10-35% MeOH/DCM) to afford 20.1 mg (59%) of **ML 2-23**:  $^1\text{H NMR}$  (400 MHz, Methanol- $d_4$ )  $\delta$  8.16 (s, 1H), 7.77 – 7.71 (m, 2H), 7.61 (dd,  $J$  = 8.8, 2.3 Hz, 2H), 7.40 – 7.34 (m, 1H), 7.31 – 7.15 (m, 4H), 7.00 – 6.90 (m, 2H), 6.00 (s, 1H), 5.57 (dd,  $J$  = 11.7, 4.7 Hz, 1H), 4.73 (d,  $J$  = 13.8 Hz, 1H), 4.60 (d,  $J$  = 13.9 Hz, 1H), 4.52 (s, 2H), 4.12 (q,  $J$  = 7.1 Hz, 1H), 3.88 (dd,  $J$  = 18.2, 11.7 Hz, 1H), 3.62 (tdd,  $J$  = 13.9, 10.2, 6.7 Hz, 16H), 3.48 (t,  $J$  = 5.5 Hz, 2H), 3.19 (dd,  $J$  = 18.1, 4.8 Hz, 1H), 2.66 (dd,  $J$  = 11.8, 5.7 Hz, 6H), 2.47 (s, 3H), 2.34 (s, 3H);  $^{13}\text{C NMR}$  (151 MHz, DMSO)  $\delta$  167.2, 164.6, 162.8, 161.9, 159.5, 156.6, 154.4, 140.4, 137.6, 133.3, 133.1, 132.0, 131.4, 131.3, 129.1, 128.6, 128.4, 127.3, 126.5, 114.4, 69.3, 69.2, 68.4, 66.7, 59.3, 53.6, 42.0, 41.4, 41.3, 37.8, 35.3, 30.3, 29.5, 27.4, 25.1, 23.0, 22.0, 17.9, 17.6, 15.6, 13.5, 11.9, 10.5; **HRMS (ESI)**: *calcd.*  $\text{C}_{47}\text{H}_{54}\text{BrCl}_2\text{N}_{10}\text{O}_4\text{S}$  ( $[\text{M} + \text{H}]^+$ ): *m/z* 1051.2453, found: 1051.2445.

### EN219-alkyne

i. Compound **5** (17.6 mg, 0.035 mmol) was dissolved in DCM (1.5 mL) and trifluoroacetic acid (0.5 mL) added slowly over the course of 20 minutes. After stirring for an additional 20 minutes, the solvent was removed *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The crude material was used directly in the next step.


ii. The aforementioned crude carboxylic acid was dissolved in DCM (2 mL) and propargylamine (2.3 mg, 0.042 mmol), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (19.8 mg, 0.052 mmol), and *N,N*-Diisopropylethylamine (223.9 mg, 1.750 mmol) were added. The reaction mixture was stirred overnight with monitoring by TLC (60% EtOAc in hexane). Upon completion, the reaction mixture was diluted with DCM, washed with brine, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (25 – 70% EtOAc/hexanes) to yield 14.6 mg (88%) of **EN219-alkyne**:  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.68 – 7.60 (m, 4H), 7.27 – 7.21 (m, 2H), 6.95 – 6.89 (m, 2H), 6.81 (s, 1H), 5.60 (dd,  $J$  = 11.8, 4.8 Hz, 1H), 4.64 – 4.54 (m, 2H), 4.52 (s, 2H), 4.18 (dd,  $J$  = 5.6, 2.6 Hz, 2H), 3.81 (dd,  $J$  = 17.8, 11.8 Hz, 1H), 3.22 (dd,  $J$  = 17.9, 4.8 Hz, 1H), 2.31 (t,  $J$  = 2.6 Hz, 1H);  $^{13}\text{C NMR}$  (101 MHz,  $\text{CDCl}_3$ )  $\delta$  167.7, 163.9, 156.7, 154.3, 134.5, 132.1, 129.6, 128.1, 127.4, 125.2, 115.1, 78.9, 71.8, 67.3, 60.3, 60.0, 42.0, 41.9, 29.6, 28.7, 21.0, 14.1; **HRMS (ESI)**: *calcd.*  $\text{C}_{22}\text{H}_{19}\text{BrCl}_1\text{N}_3\text{O}_3\text{Na}$  ( $[\text{M} + \text{Na}]^+$ ): *m/z* 510.0191, found: 510.0186.

**Scheme 3.** Synthetic route to degrader **ML 2-31**.



**(E)-tert-butyl (7-(4-(3-(4-bromophenyl)-3-oxoprop-1-en-1-yl)phenoxy)heptyl)carbamate (**8**)**


To a solution of **1** (153 mg, 0.50 mmol) in anhydrous DMF (8 mL) was added *N*-Boc-7-bromoheptan-1-amine (300 mg, 1.00 mmol) and  $K_2CO_3$  (276 mg, 2.00 mmol). The reaction mixture was heated at 60 °C for 5 hours under an atmosphere of nitrogen. Upon cooling to room temperature, the inorganic salts were filtered off, the solution was diluted with EtOAc, washed with water, and the volatiles removed *in vacuo*. The crude material was purified by silica gel chromatography (25% EtOAc/hexanes) to yield 238 mg (92%) of **8**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.97 – 7.89 (m, 2H), 7.83 (d,  $J$  = 15.6 Hz, 1H), 7.73 – 7.59 (m, 4H), 7.39 (d,  $J$  = 15.6 Hz, 1H), 7.00 – 6.92 (m, 2H), 4.55 (s, 1H), 4.04 (t,  $J$  = 6.5 Hz, 2H), 3.16 (q,  $J$  = 6.8 Hz, 2H), 1.94 – 1.76 (m, 2H), 1.57 – 1.50 (m, 4H), 1.49 (s, 9H), 1.44 – 1.37 (m, 4H).



**tert-butyl (7-(4-(3-(4-bromophenyl)-1-(2-chloroacetyl)-4,5-dihydro-1H-pyrazol-5-yl)phenoxy)heptyl)carbamate (**9**)**

To a solution of **8** (223 mg, 0.43 mmol) in EtOH was added hydrazine monohydrate (43 mg, 0.86 mmol) and the reaction mixture heated at 80 °C for 3 h under an atmosphere of nitrogen. The reaction mixture was cooled to room temperature, diluted with water and extracted with DCM. The combined organic phase was dried over anhydrous magnesium sulfate and then concentrated *in vacuo* to ~5 mL [NOTE: All of the workup procedures should be performed quickly (<1.5 h total time) and the rotovap bath kept cool as the crude product can easily undergo autooxidation]. To the concentrated solution was quickly added chloroacetyl chloride (58 mg, 0.52 mmol) and triethylamine (57 mg, 0.56 mmol) at 0 °C. The reaction mixture was stirred in an ice bath for 30 minutes, then warmed to room temperature and stirred overnight under  $N_2$ . The reaction mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude was purified by silica gel chromatography (25 – 40% EtOAc/hexanes) to yield 104 mg (40%) of **9**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.69 – 7.53 (m, 4H), 7.23 – 7.12 (m, 2H), 6.89 – 6.82 (m, 2H), 5.57 (dd,  $J$  = 11.7, 4.7 Hz, 1H), 4.57

(s, 2H), 3.93 (t,  $J = 6.5$  Hz, 2H), 3.76 (dd,  $J = 17.8, 11.7$  Hz, 1H), 3.21 (dd,  $J = 17.9, 4.7$  Hz, 1H), 3.13 (q,  $J = 6.8$  Hz, 2H), 1.85 – 1.70 (m, 2H), 1.55 – 1.46 (m, 4H), 1.47 (s, 9H), 1.41 – 1.32 (m, 4H).





### ML 2-31

*i.* Compound **9** (104 mg, 0.171 mmol) was dissolved in DCM (2.5 mL) and TFA (2.5 mL) was added slowly over 20 minutes followed by stirring for an additional 20 minutes at which point the reaction mixture was concentrated *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The crude material was used without further purification for amide coupling.

*ii.* The aforementioned TFA salt was dissolved in DCM (8 mL) and **JQ1-acid** (82 mg, 0.205 mmol), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (97.5 mg, 0.257 mmol), and *N,N*-Diisopropylethylamine (552 mg, 4.28 mmol) were added. The reaction mixture was stirred overnight with monitoring by TLC (5% MeOH in DCM, 100% EtOAc). Upon completion, the reaction mixture was directly concentrated *in vacuo* and the crude material purified by silica gel flash chromatography (1–5% MeOH/DCM). The eluted fractions were insufficiently pure and those containing product were combined, concentrated and purified again by flash silica chromatography (100–0% EtOAc/DCM followed by 0–5% MeOH/DCM) to afford 66 mg (43%) of **ML 2-31**:  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.68 – 7.57 (m, 4H), 7.48 – 7.39 (m, 2H), 7.39 – 7.32 (m, 2H), 7.22 – 7.11 (m, 2H), 6.90 – 6.81 (m, 2H), 6.67 (t,  $J = 5.8$  Hz, 1H), 5.58 (ddd,  $J = 11.8, 4.7, 2.1$  Hz, 1H), 4.69 – 4.61 (m, 1H), 4.59 (d,  $J = 2.5$  Hz, 2H), 3.93 (t,  $J = 6.4$  Hz, 2H), 3.84 – 3.67 (m, 2H), 3.56 (dd,  $J = 14.3, 7.4$  Hz, 1H), 3.42 – 3.30 (m, 2H), 3.28 – 3.16 (m, 3H), 2.70 (s, 3H), 2.47 – 2.35 (m, 3H), 1.82 – 1.71 (m, 2H), 1.71 – 1.67 (m, 3H), 1.56 (p,  $J = 7.0$  Hz, 2H), 1.45 – 1.43 (m, 4H);  $^{13}\text{C NMR}$  (151 MHz,  $\text{CDCl}_3$ ):  $\delta$  170.4, 163.9, 158.9, 155.7, 154.6, 149.9, 136.8, 136.6, 132.6, 132.1, 132.1, 130.9, 130.9, 130.5, 129.8, 128.7, 128.2, 127.0, 125.2, 115.1, 115.1, 114.9, 67.9, 60.2, 54.9, 54.5, 43.0, 42.2, 42.1, 39.6, 39.4, 29.7, 29.4, 29.1, 29.0, 26.8, 25.9, 18.5, 17.2, 14.4, 13.1, 12.4, 11.8; **HRMS (ESI)**: *calcd.*  $\text{C}_{43}\text{H}_{45}\text{BrCl}_2\text{N}_7\text{O}_3\text{S}$  ( $[\text{M} + \text{H}]^+$ ):  $m/z$  888.1865, found: 888.1858.


**Scheme 4.** Synthetic route to degrader **ML 2-32**.





**(E)-tert-butyl (2-(2-(2-(2-(4-(3-(4-bromophenyl)-3-oxoprop-1-en-1-yl)phenoxy)ethoxy)ethoxy)ethoxy)ethyl) carbamate (10)**


To a solution of **1** (153 mg, 0.50 mmol) in anhydrous DMF (8 mL) was added *tert*-Butyl (2-(2-(2-(2-bromoethoxy)ethoxy)ethoxy)ethyl) carbamate (344 mg, 0.97 mmol) and  $K_2CO_3$  (276 mg, 2.00 mmol), and the reaction mixture was stirred at 60°C for 5 hours under an atmosphere of nitrogen. Upon cooling, the inorganic salts were filtered off, the solution was diluted with EtOAc, washed with water, and the volatiles removed *in vacuo*. The crude was purified by silica gel chromatography (50% EtOAc/hexanes) to yield 251 mg (88%) of **10**:  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.97–7.89 (m, 2H), 7.89–7.78 (m, 1H), 7.72–7.55 (m, 4H), 7.40 (d,  $J$  = 15.6 Hz, 1H), 7.04–6.95 (m, 2H), 5.07 (s, 1H), 4.23 (dd,  $J$  = 5.7, 4.0 Hz, 2H), 3.97–3.84 (m, 2H), 3.83–3.77 (m, 2H), 3.76–3.72 (m, 2H), 3.71–3.64 (m, 4H), 3.58 (t,  $J$  = 5.1 Hz, 2H), 3.35 (d,  $J$  = 6.3 Hz, 2H), 1.48 (s, 9H).



**tert-butyl (2-(2-(2-(2-(4-(3-(4-bromophenyl)-1-(2-chloroacetyl)-4,5-dihydro-1H-pyrazol-5-yl)phenoxy)ethoxy)ethoxy)ethoxy)ethyl)carbamate (11)**

To a solution of **10** (244 mg, 0.42 mmol) in EtOH was added hydrazine monohydrate (42 mg, 0.84 mmol) and the reaction solution was heated at 80°C for 5 h under an atmosphere of nitrogen. Upon cooling to room temperature, the solution was diluted with water, extracted with DCM, and dried over anhydrous magnesium sulfate. The combined organic phase was concentrated *in vacuo* to ~5 mL [NOTE: All of the workup procedures should be performed quickly (<1.5 h total time) and the rotovap bath kept cool as the crude product can easily undergo autooxidation]. To the concentrated solution was quickly added chloroacetyl chloride (58 mg, 0.52 mmol) and triethylamine (57 mg, 0.56 mmol) at 0°C. The reaction mixture was stirred for 30 minutes in an ice bath and then warmed to room temperature and stirred overnight under  $N_2$ . Upon completion of the reaction, the mixture was diluted with EtOAc, washed with brine, and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (30–65% EtOAc/hexanes) to yield 82 mg (30%) of **11**:

$^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta$  7.67–7.58 (m, 4H), 7.21–7.14 (m, 2H), 6.92–6.85 (m, 2H), 5.57 (dd,  $J$  = 11.7, 4.7 Hz, 1H), 5.11 (s, 1H), 4.57 (d,  $J$  = 1.1 Hz, 2H), 4.12 (dd,  $J$  = 5.6, 4.2 Hz, 2H), 3.88–3.83 (m, 2H), 3.75–3.73 (m, 2H), 3.72–3.69 (m, 2H), 3.67–3.62 (m, 4H), 3.55 (t,  $J$  = 5.1 Hz, 3H), 3.32 (q,  $J$  = 4.3, 3.4 Hz, 2H), 3.21 (dd,  $J$  = 17.8, 4.7 Hz, 1H), 1.46 (s, 9H).





**ML 2-32**

i. Compound **11** (81.7 mg, 0.120 mmol) was dissolved in DCM (2.5 mL) and trifluoroacetic acid (2.5 mL) was added slowly over 20 minutes followed by stirring for an additional 20 minutes at which point the reaction mixture was concentrated *in vacuo*. To remove residual TFA, the crude material was dissolved in 3 mL of DCM and concentrated *in vacuo* and this process repeated two additional times. The crude material was used directly in the next step.

ii. The aforementioned crude material was dissolved in DCM (8 mL) and **JQ1-acid** (57.6 mg, 0.150 mmol), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate (HATU) (68.4 mg, 0.180 mmol), and *N,N*-diisopropylethylamine (774 mg, 6.00 mmol) were added. The reaction was stirred overnight with monitoring by TLC (10% MeOH in DCM, 100% EtOAc). Upon completion, the reaction mixture was concentrated *in vacuo* and purified by silica gel flash chromatography (1-10% MeOH/DCM). The eluted fractions were insufficiently pure and those containing product were combined, concentrated and purified again by flash silica chromatography (100-0% EtOAc/DCM followed by 0-10% MeOH/DCM) to afford 46 mg (40%) of **ML 2-32**: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.66 – 7.57 (m, 4H), 7.45 – 7.41 (m, 2H), 7.34 (d, *J* = 8.5 Hz, 2H), 7.16 (dq, *J* = 7.9, 3.1 Hz, 2H), 6.98 (d, *J* = 5.1 Hz, 1H), 6.92 – 6.83 (m, 2H), 5.57 (ddd, *J* = 11.8, 7.3, 4.7 Hz, 1H), 4.68 (t, *J* = 6.9 Hz, 1H), 4.57 (s, 2H), 4.12 (dd, *J* = 5.7, 4.0 Hz, 2H), 3.90 – 3.84 (m, 2H), 3.78 – 3.66 (m, 10H), 3.65 – 3.58 (m, 2H), 3.54 – 3.50 (m, 2H), 3.44 – 3.37 (m, 1H), 3.20 (dd, *J* = 17.8, 4.7 Hz, 1H), 2.68 (d, *J* = 1.5 Hz, 3H), 2.42 (s, 3H), 1.69 (s, 3H); <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>) δ 170.4, 163.8, 163.7, 158.4, 155.6, 154.3, 149.7, 145.2, 136.6, 136.6, 132.9, 132.1, 131.9, 131.7, 130.8, 130.6, 130.3, 130.2, 129.9, 129.8, 128.6, 128.1, 126.9, 125.0, 114.9, 70.7, 70.5, 70.5, 70.3, 69.7, 69.6, 67.4, 60.1, 54.3, 54.3, 42.1, 41.9, 39.3, 39.0, 29.6, 14.3, 13.0, 11.7; HRMS (ESI): *calcd.* C<sub>44</sub>H<sub>47</sub>BrCl<sub>2</sub>N<sub>7</sub>O<sub>6</sub>S ([*M* + *H*]<sup>+</sup>): *m/z* 950.1863, found: 950.1852.


**Scheme 5.** Synthetic route to enantioenriched **EN219**.






**tert-butyl (R)-3,5-bis(4-bromophenyl)-4,5-dihydro-1H-pyrazole-1-carboxylate ((R)-12)**

(*E*)-1,3-bis(4-bromophenyl)prop-2-en-1-one (109.8 mg, 0.3 mmol, 1.0 eq.), *tert*-butyl carbazate (43.6 mg, 0.33 mmol, 1.1 eq.), potassium phosphate (82.8 mg, 0.39 mmol, 1.3 eq.), and **catalyst A** (13.3 mg, 0.03 mmol, 10 mol %) were combined in THF (0.6 mL, 0.5 M) under nitrogen and stirred at 0°C for 18 h (Mahé et al., 2010). The mixture was diluted with EtOAc, filtered to remove salts, concentrated *in vacuo* and purified by silica gel chromatography (0–15% EtOH/Hex) to provide 94 mg (65%) of (*R*)-12: <sup>1</sup>H NMR (400 MHz, Chloroform-*d*): δ 7.61 (d, *J* = 8.6 Hz, 2H), 7.55 – 7.43 (m, 4H), 7.14 – 7.08 (m, 2H), 5.31 (dd, *J* = 11.9, 5.1 Hz, 1H), 3.73 (dd, *J* = 17.5, 12.1 Hz, 1H), 3.08 (dd, *J* = 17.5, 5.5 Hz, 1H), 1.35 (s, 9H).




**(R)-1-(3,5-bis(4-bromophenyl)-4,5-dihydro-1H-pyrazol-1-yl)-2-chloroethan-1-one ((R)-EN219)**

(*R*)-12 (45 mg, 0.094 mmol, 1.0 eq) was dissolved in DCM (0.47 mL, 0.2 M), cooled to 0°C, and 4.0 M HCl in dioxane (0.234 mL, 0.94 mmol, 10.0 eq.) was added under nitrogen. The mixture was stirred at room temperature until the starting material was consumed (1.5 h), then additional DCM was added (0.47 mL, reducing to 0.1 M) and the mixture cooled to 0°C. Chloroacetyl chloride (22 μL, 0.28 mmol, 3.0 eq.) was added followed by TEA (0.17 mL, 1.22 mmol, 13.0 eq.), and the reaction allowed to warm to room temperature and stirred for 1 h. The reaction mixture was concentrated *in vacuo* and the crude material purified by silica gel chromatography (15% EtOAc/Hex) to yield 36 mg (85%) of (*R*)-EN219 as a waxy solid. The enantiomeric excess of (*R*)-EN219 was determined by chiral HPLC (CHIRALCEL® OD-H column, *i*-PrOH/hexane (10:90), flow rate 1 mL/min) and found to be 91% ee. <sup>1</sup>H NMR (400 MHz, Chloroform-*d*): δ 7.63 – 7.55 (m, 4H), 7.49 – 7.43 (m, 2H), 7.11 (d, *J* = 8.5 Hz, 2H), 5.55 (dd, *J* = 11.8, 4.9 Hz, 1H), 4.60 – 4.48 (m, 2H), 3.77 (dd, *J* = 17.9, 11.8 Hz, 1H), 3.16 (dd, *J* = 17.9, 4.9 Hz, 1H); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>): δ 164.2, 154.4, 139.8, 132.4, 132.3, 129.7, 128.4, 127.7, 125.6, 122.2, 60.3, 42.1, 42.1; HRMS (ESI): *calcd.* C<sub>17</sub>H<sub>14</sub>ON<sub>2</sub><sup>79</sup>Br<sub>2</sub><sup>35</sup>Cl ([M + H]<sup>+</sup>): *m/z* 454.9156, found: 454.9160.



***tert-butyl (S)-3,5-bis(4-bromophenyl)-4,5-dihydro-1H-pyrazole-1-carboxylate ((S)-12)***


(*E*)-1,3-bis(4-bromophenyl)prop-2-en-1-one (116 mg, 0.32 mmol, 1.0 eq.), *tert*-butyl carbazate (46 mg, 0.35 mmol, 1.1 eq.), potassium phosphate (88 mg, 0.41 mmol, 1.3 eq.), and **catalyst B** (56 mg, 0.13 mmol, 0.4 eq) were combined in THF (2mL) under nitrogen and stirred at 0°C for 18h (Mahé et al., 2010). The mixture was diluted with EtOAc, filtered to remove salts, concentrated and purified by silica gel chromatography (0-35% EtOAc/Hex) to provide 71 mg of (*S*)-**12** intermediate as a yellow oil which was immediately carried forward into (*S*)-**EN219**.



***(S)-1-(3,5-bis(4-bromophenyl)-4,5-dihydro-1H-pyrazol-1-yl)-2-chloroethan-1-one ((S)-EN219)***

(*S*)-**12** (71 mg, 0.148mmol, 1.0 eq) was dissolved in DCM (4mL), cooled to 0°C, and 4.0 M HCl in dioxane (0.370 mL, 1.48mmol mmol, 10.0 eq.) was added under nitrogen. The mixture was stirred at room temperature until the starting material was consumed (2h), then the mixture was cooled again to 0°C. Chloroacetyl chloride (36  $\mu$ L, 0.44 mmol, 3.0 eq.) was added followed by TEA (0.268 mL, 1.92 mmol, 13.0 eq.), and the reaction allowed to warm to room temperature and stirred for 30 minutes. The reaction mixture was extracted in DCM and washed with water 3x. The reaction mixture was then concentrated *in vacuo* and the crude material purified by silica gel chromatography (15% EtOAc/Hex) to yield 35 mg (52%) of (*S*)-**EN219** as a light yellow solid. The enantiomeric excess of (*S*)-**EN219** was determined by chiral HPLC (CHIRALCEL® OD-H column, *i*-PrOH/hexane (10:90), flow rate 1 mL/min) and found to be 91% ee. **<sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)**  $\delta$  7.63–7.56 (m, 4H), 7.48–7.44 (m, 2H), 7.14–7.09 (m, 2H), 5.55 (dd, *J* = 11.8, 4.9 Hz, 1H), 4.56 (d, *J* = 13.5 Hz, 1H), 4.51 (d, *J* = 13.4 Hz, 1H), 3.77 (dd, *J* = 17.8, 11.8 Hz, 1H), 3.16 (dd, *J* = 17.8, 4.9 Hz, 1H). **<sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)**  $\delta$  164.2, 154.4, 139.8, 132.4, 132.3, 129.7, 128.4, 127.7, 125.6, 122.2, 60.3, 42.1, 42.1; **HRMS (ESI): calcd.** C<sub>17</sub>H<sub>13</sub>ON<sub>2</sub><sup>79-</sup>Br<sub>2</sub><sup>35</sup>ClNa ([M + Na]<sup>+</sup>): *m/z* 476.8975, found: 476.8976.

**Scheme 6.** Synthesis of deschloro-**EN219**.



### Deschloro-**EN219**

Hydrazine monohydrate (50% conc., 80.16  $\mu\text{L}$ , 0.8 mmol, 2.0 eq.) was added to a suspension of (E)-1,3-bis(4-bromophenyl)prop-2-en-1-one (146.4 mg, 0.4 mmol, 1.0 eq.) in EtOH (1.33 mL, 0.3 M). The resulting reaction mixture was stirred at reflux temperature for 4 h before it was concentrated under reduced pressure. The crude pyrazoline was then dissolved in DCM (2 mL, 0.2 M) and cooled to 0 °C. Triethylamine (167  $\mu\text{L}$ , 1.2 mmol, 3.0 eq.) was added dropwise, followed by chloroacetyl chloride (42.6  $\mu\text{L}$ , 0.6 mmol, 1.5 eq.). The resulting reaction mixture was stirred at ambient temperature for 1 hour before it was diluted with DCM. The organic phase was sequentially washed with satd. aq.  $\text{NaHCO}_3$  solution, brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated *in vacuo*. Purification by column chromatography (15% EtOAc/Hex) to afford 157 mg (96%) of deschloro-**EN219**:  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.61 (q,  $J$  = 8.6 Hz, 4H), 7.49 (d,  $J$  = 8.4 Hz, 2H), 7.24–7.10 (m, 2H), 5.58 (dd,  $J$  = 12.3, 4.9 Hz, 1H), 3.77 (dd,  $J$  = 17.6, 12.0 Hz, 1H), 3.24–3.06 (m, 1H), 2.44 (s, 3H);  $^{13}\text{C NMR}$  (101 MHz,  $\text{CDCl}_3$ ):  $\delta$  168.9, 152.7, 140.7, 132.10, 132.1, 130.2, 128.0, 127.4, 124.8, 121.7, 59.6, 42.0, 21.9; **HRMS (ESI)**: calcd.  $\text{C}_{17}\text{H}_{15}\text{ON}_2^{79}\text{Br}_2^{35}$  ( $[\text{M} + \text{H}]^+$ ):  $m/z$  420.9546, found: 420.9550.

### QUANTIFICATION AND STATISTICAL ANALYSIS

For quantification of Western blots, bands were quantified using Image J and normalized to protein loading controls. Statistical analysis was performed using a Student's unpaired two-tailed t-test.

For isoTOP-ABPP data analysis, data were extracted in the form of MS1 and MS2 files using Raw Extractor v.1.9.9.2 (Scripps Research Institute) and searched against the Uniprot human database using ProLuCID search methodology in IP2 v.3 (Integrated Proteomics Applications, Inc.) (Xu et al., 2015). Cysteine residues were searched with a static modification for carboxyamino-methylation (+57.02146) and up to two differential modifications for methionine oxidation and either the light or heavy TEV tags (+464.28596 or +470.29977, respectively). Peptides were required to be fully tryptic peptides and to contain the TEV modification. ProLuCID data were filtered through DTASelect to achieve a peptide false-positive rate below 5%. Only those probe-modified peptides that were evident across two out of three biological replicates were interpreted for their isotopic light to heavy ratios. For those probe-modified peptides that showed ratios greater than two, we only interpreted those targets that were present across all three biological replicates, were statistically significant and showed good quality MS1 peak shapes across all biological replicates. Light versus heavy isotopic probe-modified peptide ratios are calculated by taking the mean of the ratios of each replicate paired light versus heavy precursor abundance for all peptide-spectral matches associated with a peptide. The paired abundances were also used to calculate a paired sample t-test P value in an effort to estimate constancy in paired abundances and significance in change between treatment and control. P values were corrected using the Benjamini–Hochberg method.