



Scheme II^a

^aReagents: (a) TsOH-H₂O, CH₃COCH₃, CH₂Cl₂; (b) DAST, CH₂Cl₂; (c) KOtBu, DMSO; (d) EtOH, NH₃; (e) CF₃COOH-H₂O, 4:1. Bz = Benzoyl.

as 3-deazaneplanocin¹⁷ and two analogues of neplanocin A, which lack the 5'-hydroxyl group,¹⁸ are poor substrates for adenosine deaminase and potent inhibitors of SAH hydrolase.

Palmer and Abeles^{7a} reported that 4',5'-didehydro-5'-deoxyadenosine (5) is converted to both adenosine and SAH by SAH hydrolase, presumably via ketone 2. This observation led us to design the corresponding vinyl fluorides 6 and 13 as potential mechanism-based inhibitors of SAH hydrolase.^{19,20} As depicted in Scheme I for 6, enzymatic oxidation would afford the potent electrophile 3, which would be susceptible to addition by an enzyme nucleophile, followed by elimination of fluoride ion to yield

Scheme III^a

^aReagents: (a) (4-CH₃O-C₆H₄-S)₂, Bu₃P, pyridine; (b) MCPBA, CH₂Cl₂; (c) DAST (Et₃NSF₃), CHCl₃; (d) (CH₃OCH₂CH₂)₂O, EtN(iPr)₂, 145 °C; (e) CF₃CO₂H-H₂O, 4:1; Bio-Rad AG1-X2(OH-), MeOH; (f) hν, acetone-CD₃OD.

inactivated enzyme. Alternatively, if the nucleophile were water, inactivation could still take place, since the product 4 (Nu = OH, equivalent to an aldehyde) may bind tightly to the NADH form of the enzyme or covalently attach to an active site residue through the 5'-aldehyde functionality. The latter type of binding has been suggested for adenosine dialdehyde.²¹ In addition, we examined the possibility that 5'-deoxy-5',5'-difluoroadenosine (14, Scheme I) may be a mechanism-based inhibitor of SAH hydrolase, forming the potent electrophile 3, or its geometric isomer. In designing nucleosides 6, 13, and 14, the lack of a 5'-hydroxyl group was anticipated to make these compounds poor substrates for adenosine deaminase.²² Also, the 5'-fluorovinyl ether functional group was expected to impart greater acid stability relative to 5.²³


Synthesis

The first route to the target nucleosides, 6, 13, and 14, is depicted in Scheme II. The starting material 8 was obtained from benzoylation of a known compound.²⁴ Deprotection yielded aldehyde 9,²⁵ which was treated with (diethylamido)sulfur trifluoride (DAST) to provide di-fluoride 10 in 18% yield. Dehydrofluorination with KOtBu in DMSO yielded vinyl fluorides 11 as a 2:1 mixture of *Z*- and *E*-isomers. Concomitant loss of one benzoyl group was observed. Deprotection to diols 12 afforded separable isomers. These were individually converted to nucleosides 6 and 13. Intermediate 10 was also deprotected to nucleoside 14 with trifluoroacetic acid followed by ammonia.

In view of the poor overall yield of this route, a new general approach to the vinyl fluoride nucleosides that could be applied to the synthesis of 34 and 35 was needed. Criteria for the new route were the lack of strong basic conditions and an improved yield in the fluorination step. Because the 4',5'-didehydro nucleoside 5 has been synthesized from 5'-(phenylseleno)-5'-deoxyadenosine by oxidation and subsequent thermolysis at 100 °C,²⁶ we con-


- (17) (a) Christopher, K. H. T.; Marquez, V. E.; Fuller, W. R.; Goldstein, B. M.; Haines, D. R.; McPherson, H.; Parsons, J. L.; Shannon, W. M.; Arnett, G.; Hollingshead, M.; Driscoll, J. S. *J. Med. Chem.* **1989**, *32*, 1442. (b) Glazer, R. I.; Knode, M. C.; Tseng, C. K. H.; Haines, D. R.; Marquez, V. E. *Biochem. Pharmacol.* **1986**, *35*, 4523.
 (18) Narayanan, S. R.; Keller, B. T.; Borcharding, D. R.; Scholtz, S. A.; Borchardt, R. T. *J. Med. Chem.* **1988**, *31*, 500.
 (19) A similar mechanism was recently reported for an inhibitor of D-xylose isomerase: Carrell, H. L.; Glusker, J. P.; Burger, V.; Manfre, F.; Tritsch, D.; Biellmann, J.-F. *Proc. Natl Acad. Sci. U.S.A.* **1989**, *86*, 4440.
 (20) A portion of this work has been communicated: McCarthy, J. R.; Jarvi, E. T.; Matthews, D. P.; Edwards, M. L.; Prakash, N. J.; Bowlin, T. L.; Mehdi, S.; Sunkara, P. S.; Bey, P. *J. Am. Chem. Soc.* **1989**, *111*, 1127.

- (21) Patel-Thombre, U.; Borchardt, R. T. *Biochemistry* **1985**, *24*, 1130.
 (22) Bloch, A.; Robins, M. J.; McCarthy, J. R. *J. Med. Chem.* **1967**, *10*, 908.
 (23) McCarthy, J. R.; Robins, R. K.; Robins, M. J. *J. Am. Chem. Soc.* **1968**, *90*, 4993.
 (24) Ranganathan, R. S.; Jones, G. H.; Moffatt, J. G. *J. Org. Chem.* **1974**, *39*, 290.
 (25) Aldehyde 9 was reported as an intermediate in Vrudhula et al.: Vrudhula, V. M.; Kappler, F.; Afshar, C.; Ginell, S. L.; Lesinger, L.; Hampton, A. *J. Med. Chem.* **1989**, *32*, 885.
 (26) Zylber, N.; Zylber, J.; Gaudemer, A. *J. Chem. Soc. Chem. Comm.* **1978**, 1084.

Scheme IV^a

^a Reagents: (a) Benzoyl chloride, pyridine; (b) MCPBA, CH₂Cl₂; (c) DAST, ClCH₂CH₂Cl; (d) xylene (for 32) or diglyme (for 33); (e) EtOH, NH₃.

sidered extending our fluoro-Pummerer method²⁷ to similar α -fluoro selenoxides. However, preliminary studies with phenyl methyl selenoxide (15) indicated that DAST exclusively deoxygenated 15 to phenyl methyl selenide (16) under the usual conditions.²⁷



Therefore, we turned to the sulfoxide approach (Scheme III), with some concern for the high temperatures that might be required for the thermal elimination step. Isopropylideneadenosine was treated with *p*-anisylidene disulfide under the conditions of Nakagawa et al.²⁸ to provide thioether 17 in 96% yield as the crystalline acetic acid complex. No cyclonucleoside was observed, which is a major side reaction with dialkyl disulfides.²⁹ Oxidation of 17 with *m*-chloroperbenzoic acid (MCPBA) afforded the sulfoxides 18. Either sulfoxide or the diastereomeric mixture was transformed to the same mixture of fluoro thioethers 19 with DAST. Further oxidation with MCPBA yielded sulfoxides 20 as a mixture of isomers. A recent paper by Robins and Wnuk³⁰ described similar transformations on nucleosides using SbCl₃ as a catalyst. In our case the *p*-anisylidene group allows the fluoro-Pummerer reaction to proceed without a catalyst. The *p*-anisylidene-containing intermediates were usually crystalline.

For the thermolysis of 20, diglyme was the solvent of choice, providing good solubility and a high boiling point. The reaction was carried out in the presence of diiso-

propylethylamine³¹ and gave a 77% yield of 21 as a 2.7:1 mixture of *Z*- and *E*-isomers. Deprotection of 21 provided the desired products 6 and 13. These isomers were separated on a Dekker column³² and were identical with products obtained by the method outlined in Scheme II. The *E*-geometric isomer 13 was irradiated with a sun lamp in acetone-methanol, which resulted in a 60–70% conversion to *Z*-isomer 6. The configurations of 6 and 13 were proven by NOE experiments between H-3' and the vinyl proton (see Experimental Section).

The combined overall yield of 6 and 13 in Scheme III was 18%, and this general route was applied to deoxy and arabino analogues (Scheme IV). For these transformations benzoyl-protecting groups were used for both hydroxy and amino groups, and the intermediates were more soluble in organic solvents. The thermal syn elimination of sulfoxides 29 and 31 resulted in concomitant loss of one benzoyl group, as was observed for 10 (in Scheme II). In the arabino series (starting with 23), the *E*- and *Z*-isomers of vinyl fluoride 33 were separated on silica gel and converted individually to 36 and 37. In the deoxy series, the *Z*-isomer 34 was crystalline, whereas isomer 35 was an oil. The initial sample of 34 was isolated as an acetic acid complex.

For the synthesis of chlorovinyl nucleoside 44 (Scheme V), the acetic acid complex 17 was converted to free 38, to avoid side reactions in the benzoylation step. Dibenzoate 39 was oxidized to sulfoxide 40 and chlorinated with sulfur chloride³³ to 41. It should be noted that the acetonide-protecting group was required, as the desired reaction did not take place with benzoate 45.

Sulfoxide elimination from 41 proceeded in the same manner as examples in Scheme III and IV (20, 29, and 31), affording 42 as a 5:1 mixture of *Z*- and *E*-isomers, re-

(27) McCarthy, J. R.; Peet, N. P.; LeTourneau, M. E.; Inbasekaran, M. *J. Am. Chem. Soc.* **1985**, *107*, 735.

(28) Nakagawa, I.; Aki, K.; Hata, T. *J. Chem. Soc. Perkin 1* **1983**, 1315.


(29) Craig, G. W.; Moffatt, J. G. *Nucleosides Nucleotides* **1986**, *5*, 399.

(30) Robins, M. J.; Wnuk, S. F. *Tetrahedron Lett.* **1989**, *29*, 5729.

(31) Amine has been added in a sulfoxide elimination. For example, Babler, J. H.; Haack, R. A. *J. Org. Chem.* **1982**, *47*, 4801.

(32) Dekker, C. A. *J. Am. Chem. Soc.* **1965**, *87*, 4027.

(33) Tsuchihashi, G.; Ogura, K.; Iriuchijima, S.; Tomisawa, S. *Synthesis* **1971**, 89.



spectively. Deprotection of **42** followed by crystallization afforded pure *Z*-isomer **43**. Ammonolysis of **43** provided the desired **44**.

Fluorination of **40** with DAST followed by MCPBA oxidation (Scheme V) afforded fluoro sulfoxide **47**. Chlorination of **47** with sulfur chloride provided chloro fluoro sulfoxide **48**. Thermal elimination of the sulfoxide formed *E*-isomer **49** exclusively. Deprotection of **49** yielded the desired **51** and photolysis of **51** gave a small sample of isomer **52**. With both geometric isomers in hand, structural assignments were made for **51** and **52** on the basis of the carbon-fluorine and hydrogen-fluorine coupling constants (see Experimental Section).

Enzyme Inhibition: Results and Discussion

Initial studies on compounds **6**, **13**, and **14** were performed with partially purified SAH hydrolase from mouse liver. Enzyme activity was assayed in the synthetic direction as described by Ragione and Pegg.³⁴ The pseudo-first-order loss of activity observed for *Z*-isomer **6** is plotted in Figure 1. Kinetic constants were determined according to the method of Kitz and Wilson³⁵ and are listed in Table I. With the mouse liver enzyme, **6** and **13** had similar values of k_{inact} (Table I), but the *Z*-isomer **6** is the more potent inactivator by comparison of k_{inact}/K_I ($5.0 \times 10^5 \text{ min}^{-1} \text{ M}^{-1}$ for **6** and $2.2 \times 10^5 \text{ min}^{-1} \text{ M}^{-1}$ for **13**). Adenosine protected the mouse liver enzyme from inhibition by **6**. Inhibition by **6** was not reversed by extensive dialysis or by incubation of the inactivated enzyme at 37 °C for 90 min with 0.5 mM NAD⁺. The difluoromethyl compound **14** was a less potent time-dependent inhibitor than **6** or **13**, exhibiting $k_{\text{inact}}/K_I = 6 \times 10^3 \text{ min}^{-1} \text{ M}^{-1}$. This was in contrast to Palmer and Abeles' results^{2a} where 5'-deoxy-5'-fluoroadenosine was not a substrate or an inactivator of beef liver SAH hydrolase. As **14** showed nonlinear kinetics with this partially purified enzyme, its K_I was estimated from the initial slope. This nonlinearity may be the result of metabolism of the inhibitor, since linear kinetics were obtained for **14** with the purified rat liver enzyme (Table I). No time-dependent inhibition was observed with **17**, **34**, **35**,³⁶ **36**, **37**, or **51** with mouse liver SAH hydrolase.

To confirm the above results and to perform further mechanistic studies, SAH hydrolase from rat liver was purified.³⁷ This enzyme was assayed in the hydrolytic direction spectrophotometrically.³⁸ The kinetic constants (Table I) show that **6** was a slightly more potent inactivator ($k_{\text{inact}}/K_I = 1.2 \times 10^5 \text{ min}^{-1} \text{ M}^{-1}$) than **13** ($k_{\text{inact}}/K_I = 0.58 \times 10^5 \text{ min}^{-1} \text{ M}^{-1}$), and dramatically more potent than di-




Figure 1. Time- and concentration-dependent inhibition of SAH hydrolase by **6**. Partially purified SAH hydrolase from mouse liver was incubated with various concentrations of **6**. At indicated times, aliquots were withdrawn and tested for remaining enzyme activity as described in the Experimental Section.

fluoronucleoside **14** ($k_{\text{inact}}/K_I = 4.1 \times 10^3 \text{ min}^{-1} \text{ M}^{-1}$).

Compounds **34**, **36**, **37**, **44**, **51**, and **52** were competitive inhibitors and the K_I values were determined for the rat liver enzyme (Table I). Some interesting features of these competitive inhibitors should be pointed out, particularly those relating to the 5'-position. Assuming that K_I reflects binding affinity, the *E*-isomer **37** of the ara-A-derived nucleosides bound more tightly than the *Z*-isomer **36**, which is the order observed with time-dependent inhibitors **6** and **13** with this enzyme. Also, it is worthy of note that the chlorine substituent in **44**, **51**, and **52** does not prevent these nucleosides from binding to the active site of SAH. With the competitive inhibitors no NADH formation was observed during the course of the assay, suggesting that these inhibitors are not rapidly oxidized to the 3'-ketones. This difference in reactivity may have several reasons, but it is clear that substituents on the 5'-carbon play a significant role.

Within the series of inhibitors prepared, the 2'-hydroxyl group appears to be critical for time-dependent inhibition of SAH hydrolase as evidenced by the fact that nucleosides **34**, **35**, and **37** are not oxidized. This finding is of interest since ara-A and 2'-deoxyadenosine are oxidized by SAH hydrolase (and are time-dependent inhibitors)¹ despite the absence or stereochemical change of the 2'-hydroxyl. The 2'-hydroxyl is known to hydrogen bond to the 3'-hydroxyl, affecting its acidity, and has also been reported to affect ring conformation.³⁹ Nucleosides **34**–**37** contain modifications on both sides of the 3'-hydroxyl, which have unknown effects on such factors as the ring conformation and may explain the observation that these compounds are only competitive inhibitors.

Compounds **6** and **14** were deaminated by adenosine deaminase at least 1000-fold slower than adenosine. This slow deamination is consistent with the previously established substrate specificity of adenosine deaminase.²²

Mechanistic Studies

Preliminary results³⁷ indicated that when **6** was mixed with purified (>90%) enzyme from rat liver stoichiometric formation of NADH was observed spectrophotometrically. After complete inactivation of the rat liver enzyme by **6**, the ¹⁹F NMR spectrum of the supernatant obtained after denaturation and removal of the protein indicated that ca. 1 mol of fluoride ion was formed per mole of enzyme

(34) Ragione, F. D.; Pegg, A. E. *Biochem. J.* 1983, 210, 429.


(35) Kitz, R.; Wilson, I. B. *J. Biol. Chem.* 1962, 237, 3245.

(36) Preincubation for 20 min with 5 μM **35** produced no time-dependent inhibition of the mouse enzyme. This compound was not further examined.

(37) Results with the rat liver enzyme have been briefly summarized.²⁰ Full details on mechanistic work with **6**, **13**, and **14** with this enzyme have been reported: Mehdi, S.; Jarvi, E. T.; Koehl, J. R.; McCarthy, J. R.; Bey, P. *J. Enzyme Inh.* 1990, 4, 1.

(38) Wolfson, G.; Chisholm, J.; Tashjian, A. H., Jr.; Fish, S.; Abeles, R. H. *J. Biol. Chem.* 1986, 261, 4492.

(39) Sundaralingam, M. *Ann. N.Y. Acad. Sci.* 1975, 255, 3.

Scheme V^a

^a Reagents: (a) PhCOCl, pyridine; (b) MCPBA, CH₂Cl₂; (c) SO₂Cl₂, CH₂Cl₂, pyridine; (d) diglyme, EtN(iPr)₂; (e) CF₃COOH-H₂O, 4:1; (f) MeOH, NH₃; (g) DAST, CHCl₃; (h) EtOH, NH₃; (i) *hν*, acetone, CH₃OH

Table I. Kinetic Constants for Enzyme Inhibition and IC₅₀ Values against MoLV

compd	mouse liver SAH hydrolase ^a			rat liver SAH hydrolase ^b			IC ₅₀ , μg/mL MoLV ^d
	time-dependent inhibition	K _i , μM	k _{inact} , min ⁻¹	K _i , μM	k _{inact} , min ⁻¹	K _i , ^c μM	
6	yes	0.55	0.28	6.5	0.8	NA	0.05
13	yes	1.04	0.23	2.4	0.14	NA	0.25
14	yes	10 ^e	0.6	9.7	0.04	NA	0.8
34	no	NA ^g	NA	NA	NA	150	10.0
35	no	NA	NA	NA	NA	ND ^f	ND
36	no	NA	NA	NA	NA	11.0	1.2
37	no	NA	NA	NA	NA	3.2	4.4
44	no	NA	NA	NA	NA	1.6	3.3
51	no	NA	NA	NA	NA	3.0	1.3
52	no	NA	NA	NA	NA	1.5	0.1
ara-A	yes	30	ND	ND	ND	NA	1.0

^a Assayed in the synthetic direction (see Experimental Section). ^b Assayed in the hydrolytic direction (see Experimental Section). ^c K_i for competitive inhibition with rat liver SAH. ^d Method: see ref 42. ^e Value from initial slope. ^f ND = not determined. ^g NA = not applicable.

subunit. The stoichiometry indicates that the partition ratio (number of product molecules per each inactivation event) for inhibitor 6 is close to zero.

When 6 (tritiated in the 8-position) was incubated with the enzyme, labeled protein (0.8 mol of inhibitor/mol of enzyme subunit) was isolated. However, recent results³⁷ indicate that the label is lost under three different denaturing conditions.

The above experiments suggest that inhibitor 6 is oxidized to ketone 3, which is then attacked by an enzyme nucleophile or water in the active site of the enzyme. If the displacement (addition-elimination) is rate-limiting in the inactivation, the *Z*-isomer would be predicted to be more potent than the *E*-isomer on the basis of kinetic results reported for the attack of nucleophiles on β-halo-α,β-unsaturated ketones.⁴⁰ It should be noted that the product is not reduced to the 3'-alcohol by the enzyme. If the product 4 contains a covalent link to the enzyme, it

may be stable with respect to further degradation as long as the enzyme is not denatured. Alternatively, product 4 could be the addition product with water and remain tightly bound to the enzyme in the same manner as neplanocin A, which requires enzymatic oxidation. However, while inhibition of the beef liver enzyme by neplanocin A has been reported⁴¹ to be reversed by incubation with NAD⁺, addition of NAD⁺ to SAH hydrolase (rat liver or mouse liver) inactivated by 6 did not reverse the inhibition. This suggests that the mechanism of inhibition by neplanocin A and 6 is different or that there is a difference between the enzymes from different sources.

The mechanism of inactivation by 14 was also investigated further with the rat liver enzyme.³⁷ Incubation of the enzyme with 14 led to formation of NADH as determined spectrophotometrically. When the enzyme was inhibited with an excess of 14, examination of the ¹⁹F spectrum of the supernatant revealed that the ratio of

(40) Modena, G. *Acc. Chem. Res.* 1971, 4, 73.

(41) Matuszewska; Borchardt, R. T. *J. Biol. Chem.* 1987, 262, 265.

moles of fluoride released to moles of enzyme subunits was 1.66, compared to roughly half that number in the case of inhibition by 6. These results are consistent with the mechanism of inactivation of SAH hydrolase by 14 proposed in Scheme I.

Antiviral and Other Biological Activity

The nucleosides were evaluated as antiretroviral agents in the XC-plaque assay using Moloney leukemia virus (MoLV)⁴² as a model for human immunodeficiency virus (HIV). The most potent antiviral activity was observed with time-dependent inhibitors 6 ($IC_{50} = 0.05 \mu\text{g/mL}$) and 13 ($IC_{50} = 0.25 \mu\text{g/mL}$). The less-potent inactivator 14 had antiviral activity in the same range as competitive inhibitors 44, 36, 37, and 51. In general, the results (Table I) indicate that enzyme inhibition correlates with antiviral activity. An exception to this trend is chloro fluoro nucleoside 52 which had antiretroviral activity in the same range as 6 and 13. Aside from enzyme inhibition, other factors, such as membrane transport and cellular metabolism may affect antiviral activity. Among known SAH hydrolase inhibitors, ara-A has an IC_{50} of $1.0 \mu\text{g/mL}$ and neplanocin A has an IC_{50} of $0.008 \mu\text{g/mL}$ against MoLV. However, the latter compound is toxic and is metabolized to other derivatives.¹⁵ The mechanism of antiviral activity was further investigated with 6. MoLV infected fibroblasts (C3H10T1/2) were treated with $1.0 \mu\text{g/mL}$ of 6. A 30% reduction in overall RNA methylation and a 88% reduction in constitutive MoLV production was observed⁴³ compared to controls. These results suggest that the inhibition of RNA-methylation may contribute to the antiretroviral activity at concentrations of 6 not toxic to these cells.

The most potent inactivator, 6, was also evaluated against vaccinia and Herpes I virus. The observed IC_{50} values of $0.03 \mu\text{M}$ for vaccinia⁴⁴ and $10 \mu\text{g/mL}$ for Herpes I are in the range expected for SAH hydrolase inhibitors.⁴⁵ Finally, since SAH hydrolase inhibitors such as neplanocin A⁴⁴ exhibit cytotoxicity, nucleoside 6 was evaluated for HeLa cell antiproliferative activity. An IC_{50} of $1 \mu\text{g/mL}$ was measured, which is well above the antiviral IC_{50} values.

In summary, a new class of mechanism-based inhibitors of SAH hydrolase was designed, on the basis of the enzymatic pathway for the conversion of SAH to adenosine and homocysteine. Evidence was presented supporting the proposed mechanism of action outlined in Scheme I for vinyl fluoride nucleosides 6, 13, and the difluoro nucleoside 14. Other analogues proved to be competitive inhibitors. Antiviral data against MoLV demonstrated that the time-dependent inhibitors 6 and 13 were potent in this model.

Experimental Section

General. All melting points are uncorrected. The IR spectra were recorded with a Perkin-Elmer Model 710B spectrophotometer, ^1H NMR spectra were determined with Varian EM-360A and Varian VXR-300 (multinuclear probe) spectrometers, and ^{19}F NMR spectra were recorded at 282 MHz on the Varian VXR-300 with CFCl_3 as an external standard. Mass spectra were

obtained with a Finnigan MAT Model 4600 and are indicated as either electron impact (EI) or chemical ionization (CI). Exact mass determinations (HRMS) were obtained on a ZAB2-SE high-resolution mass spectrometer with perfluorokerosene as a reference. Combustion analyses for C, H, and N were performed by Merrell Dow Analytical Laboratories, Cincinnati, OH. Thin-layer chromatography (TLC) on silica gel plates was visualized by UV or by staining with alkaline potassium permanganate and heating. Reactions were carried out under nitrogen or argon, except for ammonia reactions and hydrolyses. Silica gel used for chromatography was 40–63 μm size, as described for flash chromatography.⁴⁶ Commercially available (Aldrich, Sure Seal) dry solvents were used as reaction media, unless otherwise indicated.

Caution: (Diethylamido)sulfur trifluoride (DAST) has been reported to decompose violently at temperatures above 50°C .⁴⁷ Suitable safety precautions must be observed with this reagent. Reactions in this paper were carried out in an efficient fume hood in normal pyrex glassware, where some etching was observed.

Preparation and Assay of Mouse Liver SAH Hydrolase.

SAH hydrolase was partially purified from mouse liver as described by Ragione and Pegg.³⁴ The 38–55% ammonium sulfate fraction was stored at -70°C until use. The enzyme was assayed by measuring the radiolabeled S-adenosylhomocysteine formed from $[2\text{-}^3\text{H}]$ adenosine and L-homocysteine. The assay mixture (200 μL) containing 0.025 mM potassium phosphate, 2 mM 2-mercaptoethanol, 0.1 mM adenosine, 0.28 μCi of $[2\text{-}^3\text{H}]$ adenosine (21 Ci/mmol), 10 mM DL-homocysteine, 1.0 μM 2-deoxycoformycin, and enzyme was incubated at 37°C for 10–20 min. The reaction was stopped by the addition of 0.8 mL of 30 mM HCl. A portion of the reaction mixture was applied to a column of Cellex-P (Bio Rad; 0.5 cm \times 0.7 cm; H^+ form). The product, $[2\text{-}^3\text{H}]$ SAH, was retained while adenosine was eluted quantitatively with 10 mL of 10 mM HCl. Radiolabeled SAH was then eluted with 2.0 mL of 0.5 M HCl and the radioactivity in the eluate was measured by liquid scintillation counting. By using this assay, kinetic constants for time-dependent inhibitors were determined by the method of Kitz and Wilson.³⁵

Determination of Kinetic Constants for Rat Liver SAH Hydrolase.

Rat liver SAH hydrolase was assayed by following the formation of homocysteine.³⁸ Assay conditions and preparation of the enzyme are given in the supplementary material. The K_i for a reversible inhibitor was determined from the initial rates of reactions containing varying concentrations of the inhibitor and using a correction for the $[S]$ relative to the K_m . In one case (compound 36), both inhibitor and substrate concentrations were varied, and a double reciprocal plot of the data was used to determine the type of inhibition and the K_i . For time-dependent inhibitors, the inactivation reaction was monitored continuously in the presence of substrate and inhibitor (6). The progress curves were fit to exponentials and the pseudo-first-order rate constants K_{obs} at different inhibitor concentrations were obtained. K_i and k_{inact} were obtained from a plot of $1/k_{\text{obs}}$ versus $1/[I]$. To test for reversibility of inhibition of the rat liver enzyme by compound 6, a sample of inactivated enzyme was gel filtered to remove excess inhibitor, and incubated with NAD^+ (0.5 mM) or without NAD^+ at 37°C for up to 90 min. Portions were removed to test for return of activity. A control sample of enzyme was incubated under the same conditions to monitor any effect on the activity.

^{19}F NMR Experiment with 6 and SAH Hydrolase from Rat Liver. In 0.82 mL of 1 mM EDTA and 10 mM potassium phosphate (pH 7.5), 0.46 mg (9.7 nmol of enzyme subunits) of SAH hydrolase was incubated with 75 nmol of 6. After 10 min at room temperature, less than 2% of original enzyme activity was detected. Ethanol (2 mL) was added, the mixture was kept at 0°C for 5 min and then centrifuged. The pellet was washed once with H_2O –EtOH and the combined supernatant and washing was taken to dryness in vacuo. The residue was dissolved in D_2O . Some insoluble material, possibly denatured protein, was present and was removed by centrifugation. The ^{19}F NMR spectrum (vs ext CFCl_3) showed signals due to fluoride ($\delta - 120.7$) and unreacted inhibitor ($\delta - 162.5$) in a 1:5.9 ratio. The ratio of enzyme to

(42) Bowlin, T. L.; Proffitt, M. R. *J. Interferon Res.* **1983**, *3*, 19.

(43) Full details will be reported in a communication: Prakash, N. J.; Davis, G. F.; Jarvi, E. T.; McCarthy, J. R.; Bowlin, T. L. *Antiviral Res.* **1991**, in press.

(44) Kinney, M., Unpublished observations, Merrell-Dow, Indianapolis, IN.

(45) (a) Hasobe, M.; McKee, J. G.; Borcharding, D. R.; Borchardt, R. T. *Antimicrob. Agents Chemother.* **1987**, *31*, 1849. (b) De Clercq, E.; Cools, M.; Balzarini, J.; Marquez, V. E.; Borcharding, D. R.; Borchardt, R. T. *Antimicrob. Agents Chemother.* **1989**, *33*, 1291.

(46) Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

(47) Cochran, J. *Chem. Eng. News* **1979**, *57*, No. 12, 4.

inhibitor used in the experiment was 1:7.1. Therefore, the amount of fluoride was approximately stoichiometric with the amount of enzyme used. No other signals were observed.

The above experiment was run under routine ^{19}F NMR conditions. The ^{19}F NMR spectrum of a mixture of **6** and fluoride ion was repeated with a 22 s pulse delay. Ratios of integrals were within 10% of those under routine ^{19}F NMR conditions.

Deamination by Adenosine Deaminase. Decrease in absorbance at 265 nm ($-\Delta\epsilon = 8.6 \text{ mM}^{-1} \text{ cm}^{-1}$) due to the deamination of adenosine and of compounds **6** and **14** was monitored spectrophotometrically. The reaction mixture contained adenosine or the adenosine derivative (50 μM) and calf spleen or intestinal mucosa adenosine deaminase in 20 mM potassium phosphate (pH 7.5).

Chemistry. ***N*⁶,*N*⁶-Dibenzoyl-5'-deoxy-2',3'-*O*-isopropylidene-5',5'-(*N,N'*-diphenylethylenediamino)adenosine (8).** *N*⁶-Benzoyl-5'-deoxy-2',3'-*O*-isopropylidene-5',5'-(*N,N'*-diphenylethylenediamino)adenosine²⁴ (2.96 g, 4.9 mmol) in pyridine (10 mL) was cooled in an ice bath, and benzoyl chloride (1.15 mL, 9.9 mmol) was added. The mixture was stirred overnight at room temperature and poured into ice water. The product was extracted with CHCl_3 (100 mL) and dried (MgSO_4). The solution was evaporated and toluene was added. Evaporation was repeated followed by drying in vacuo providing 4.07 g of a yellow foam. The product was percolated through a flash silica gel column (10 cm \times 4 mm) with 4% EtOAc–96% CH_2Cl_2 . Appropriate fractions were combined and evaporated three times from ethanol, providing a solid which was triturated with 50 mL of ethanol and filtered. The solid obtained was dried in vacuo, providing 2.67 g (76%) of **8**: mp 135–138 °C; ^1H NMR (CDCl_3) δ 1.32 (s, 3 H), 1.51 (s, 3 H), 3.48–3.71 (m, 4 H, $\text{NCH}_2\text{CH}_2\text{N}$), 4.61 (m, 1 H, H-4'), 5.14 (m, 2 H, H-2', H-3'), 5.72 (d, 1 H, $J = 2.6$ Hz, N-CH-N), 6.12 (d, 1 H, $J = 2.1$ Hz, H-1'), 6.64 (d, 2 H, $J = 7.9$ Hz), 6.70–6.82 (m, 4 H), 7.13–7.27 (m, 4 H), 7.36 (t, 4 H, $J = 7.6$ Hz), 7.49 (m, 2 H), 7.87 (d, 2 H, $J = 7.0$ Hz), 7.95 (s, 1 H), 8.47 (s, 1 H); MS (CI/CH_4) 708 ($\text{M} + \text{H}$)⁺. Anal. ($\text{C}_{41}\text{H}_{37}\text{N}_7\text{O}_5$) C, H, N.

***N*⁶,*N*⁶-Dibenzoyl-2',3'-*O*-isopropylideneadenosine 5'-Aldehyde (9).** *N*⁶,*N*⁶-Dibenzoyl-5'-deoxy-2',3'-*O*-isopropylidene-5',5'-(*N,N'*-diphenylethylenediamino)adenosine (**8**) (2.64 g, 3.73 mmol) in CH_2Cl_2 (370 mL) at 0 °C was added to a solution of *p*-toluenesulfonic acid monohydrate (1.56 g, 8.2 mmol) in 180 mL of acetone. The mixture was stirred for 1.5 h and filtered. The filtrate was evaporated and partitioned between 200 mL of CH_2Cl_2 and water. The CH_2Cl_2 solution was dried (MgSO_4) and evaporated to a foam. The foam (2.10 g) was dissolved in benzene (200 mL) and water was removed by refluxing with a Dean–Stark trap for 1 h. The solvent was removed in vacuo to provide 2.06 g (100%) of the title compound as a foam.

This material can be used for the following reaction, but the material used below (for best yields) was chromatographed on flash silica gel with 15% EtOAc–85% CH_2Cl_2 and treated in a Dean–Stark setup as above, (80% recovery). The NMR indicated aldehyde (65%) and hydrate (35%) form, resulting in doubling of some NMR signals. Aldehyde **9** has been previously described as an intermediate for which no spectral data was reported.²⁵ ^1H NMR (CDCl_3) δ 1.31 (s, 3 H), 1.52 (s, 3 H), 4.61 (d, 1 H, $J = 1.7$ Hz or two s), 5.25 (d, 1 H, $J = 5.9$ Hz), 5.42 (dd, 1 H, $J = 1.7$ and 7.9 Hz), 6.17 (s, 1 H), 7.24–7.77 (several m, 10 H), 8.07 (s, 1 H), 8.44 (major) and 8.54 (minor, two s, both H-8), 9.26 (s, 0.65 H, CHO); MS (FAB–XENON) m/z 514 ($\text{M} + \text{H}$)⁺. Anal. Calcd ($\text{C}_{27}\text{H}_{23}\text{N}_5\text{O}_6$) C, 63.15; H, 4.51; N, 13.64. Found: C, 64.48; H, 5.08; N, 11.63.

***N*⁶,*N*⁶-Dibenzoyl-5'-deoxy-5',5'-difluoro-2',3'-*O*-isopropylideneadenosine (10).** *N*⁶,*N*⁶-Dibenzoyl-2',3'-*O*-isopropylideneadenosine 5'-aldehyde (**9**) (3.90 g, 7.6 mmol) was dissolved in 25 mL of CH_2Cl_2 (distilled from calcium hydride) and 3.2 mL (22.8 mmol) of (diethylamido)sulfur trifluoride was added. The mixture was stirred for 6 h, diluted with CHCl_3 , and poured into saturated aqueous NaHCO_3 (50 mL). The product was extracted into chloroform (400 mL). The solution was dried (MgSO_4) and evaporated to a foam (3.60 g). The foam was percolated through a silica gel flash column (12 cm \times 40 mm) with 4% EtOAc–96% CH_2Cl_2 . The component of R_f 0.6 (10% EtOAc–90% CH_2Cl_2 TLC solvent) was isolated as a white foam (738 mg, 18%): ^1H NMR (CDCl_3 , 300 MHz) δ 1.42 (s, 3 H), 1.65 (s, 3 H), 4.42–4.53 (m, 1 H), 5.27 (dd, 1 H, $J = 2.7$ and 5.9 Hz), 5.39

(dd, 1 H, $J = 1.7$ and 6.0 Hz), 5.96 (td, 1 H, $J = 55$ and 4.5 Hz), 6.28 (d, 1 H, $J = 1.3$ Hz), 7.34–7.52 (m, 6 H), 7.85 (d, 4 H, $J = 7.2$ Hz), 8.15 (s, 1 H), 8.67 (s, 1 H); ^{19}F NMR (CDCl_3) δ –65.87 (ddd, $J = 12.4$, 55.2 and 299.0 Hz); –50.71 (ddd, $J = 10$, 55.2 and 299.1 Hz); MS (FAB–XENON) m/z 536 ($\text{M} + \text{H}$)⁺. Anal. Calcd. ($\text{C}_{27}\text{H}_{23}\text{F}_2\text{N}_5\text{O}_5$) C, 60.56; H, 4.33; N, 13.08. Found C, 60.26; H, 4.44; N, 12.03.

5'-Deoxy-5',5'-difluoroadenosine (14). The *N*⁶,*N*⁶-dibenzoylated **10** (407 mg, 0.76 mmol) was dissolved in absolute ethanol and evaporated to dryness. The residue was treated with ethanolic ammonia (15 mL, saturated at 0 °C) in a Carius tube for 24 h and evaporated in vacuo. The residue was purified by flash chromatography (1:1 EtOAc– CH_2Cl_2 followed by EtOAc), providing 195 mg (78%) of 5'-deoxy-5',5'-difluoro-2'-3'-*O*-isopropylideneadenosine, ($R_f = 0.21$, EtOAc, TLC). A sample was triturated with CH_2Cl_2 –cyclohexane to afford crystals, mp 161–163 °C. The crude product was used directly in the next step: ^1H NMR (CDCl_3) δ 1.42 (s, 3 H), 1.64 (s, 3 H), 4.37–4.47 (m, 1 H), 5.30–5.36 (m, 1 H), 5.42–5.44 (m, 1 H), 5.68 (bs, 2 H, NH_2), 6.01 (td, 1 H, $J = 56$ and 5.2 Hz), 6.19 (m, 1 H, covered by 6.01 td), 7.87 (s, 1 H), 8.36 (s, 1 H); ^{19}F NMR –125.77 (ddd, $J = 10.7$, 55.8, and 259 Hz), –130.16 (ddd, $J = 10.7$, 55.8, and 259 Hz); MS (EI) m/z 327 (M^+); HRMS calcd for $\text{C}_{13}\text{H}_{15}\text{N}_5\text{O}_3\text{F}_2$ 327.1143, found 327.1140. Anal. ($\text{C}_{13}\text{H}_{15}\text{N}_5\text{O}_3\text{F}_2$) C, H, N.

To 5'-deoxy-5',5'-difluoro-2',3'-*O*-isopropylideneadenosine (165 mg, 0.76 mmol) was added 2 mL of 4:1 CF_3COOH –water. The mixture was stirred for 1 h and evaporated. The residue was purified by flash chromatography (200 mL of EtOAc followed by 400 mL of 5% MeOH–95% EtOAc). The appropriate fractions were combined and evaporated to afford 153 mg (100%) of a white foam ($R_f = 0.35$, 10% MeOH–90% EtOAc). The foam was stirred in 4 mL of ethanol and the resulting crystals were filtered to give 52 mg (36%) of **14**, mp 198–200 °C after drying in vacuo. Thermogravimetric analysis indicated 6% weight loss up to 200 °C and the only impurity found in the assayed sample was ethanol: ^1H NMR ($\text{DMSO}-d_6$, ethanol peaks excluded) δ 4.03–4.12 (m, 1 H), 4.36 (m, 1 H), 4.77 (q, 1 H, $J = 6$ Hz), 5.67 (d, 1 H, $J = 5.1$ Hz, OH), 5.70 (d, 1 H, $J = 6.3$ Hz, OH), 6.13 (d, 1 H, $J = 6.3$ Hz), 6.31 (td, 1 H, $J = 53$ and 4.7 Hz), 7.34 (b s, 2 H, NH_2), 8.16 (s, 1 H), 8.32 (s, 1 H); ^{19}F NMR ($\text{DMSO}-d_6$) –125.9 (ddd, $J = 9.3$, 55.1, and 291.2 Hz), –127.2 (ddd, $J = 9.9$, 54.9, and 291.2 Hz); MS (CI/CH_4) m/z 288 ($\text{M} + \text{H}$)⁺; HRMS calcd for $\text{C}_{10}\text{H}_{11}\text{O}_3\text{N}_5\text{F}_2$ 287.0830, found 287.0850. The ethanol filtrate which contained $\text{CF}_3\text{CO}_2\text{H}$ was evaporated and dissolved in water. A Dowex 50 (H^+) resin (H_2O , then 2 M NH_4OH) provided an additional 47 mg (total yield 68%) of **14**.

***N*⁶-Benzoyl-9-(5-deoxy-5-fluoro-2,3-*O*-isopropylidene- β -D-erythro-pent-4-enofuranosyl)adenine (11).** To 401 mg (0.75 mmol) of crushed *N*⁶,*N*⁶-dibenzoyl-5'-deoxy-5',5'-difluoro-2',3'-*O*-isopropylideneadenosine (**10**) and 335 mg (4 equiv) of potassium *tert*-butoxide under nitrogen was added 2 mL of dimethyl sulfide (distilled from CaH_2 and stored over molecular sieves). The mixture was stirred under nitrogen for 21 h, quenched with 4 mL of saturated ammonium chloride and extracted with ethyl acetate. The ethyl acetate layer afforded 274 mg of yellow oil which was percolated through a flash column (20 mm \times 15 cm) with 30% EtOAc–70% CH_2Cl_2 . TLC revealed two spots very close together at $R_f = 0.55$ (EtOAc solvent). Combined fractions yielded 183 mg (59%) of a foam containing the *Z*- and *E*-isomers in a 2:1 ratio: ^1H NMR (CDCl_3 , 300 MHz) δ 1.34 (major) and 1.37 (minor, 3 H together, 2 s), 1.49 (s, 3 H), 5.35–5.38 (m, 1 H), 5.56 and 5.90 (1 H together; d, $J = 4$ Hz and m, respectively), 6.23 (br s, minor) and 6.25 (1 H together), 6.43 and 6.81 (d, $J = 74$ Hz, major and d, $J = 77$ Hz, minor; 1 H together), 7.39–7.98 (m, 6 H), 8.65 (major) and 8.653 (minor; 2 s, 1 H together), 9.05 (1 H, br, NH); ^{19}F NMR (CDCl_3 , 282 MHz) δ (from external CFCl_3) –158.94 (d, $J = 74$ Hz, major), 174.4 (d, $J = 77$ Hz, minor); MS (CI/CH_4) m/z 412 ($\text{M} + \text{H}$)⁺; HRMS calcd for $\text{C}_{20}\text{H}_{18}\text{FN}_5\text{O}_4$ 411.1343, found 411.1330.

***N*⁶-Benzoyl-9-(5-deoxy-5-fluoro- β -D-erythro-pent-4-enofuranosyl)adenine (12).** Cold 4:1 CF_3COOH –water (10 mL) was used to dissolve 1.28 g (3.1 mmol) of *N*⁶-benzoyl-9-(5-deoxy-5-fluoro-2,3-*O*-isopropylidene- β -D-erythro-pent-4-enofuranosyl)adenine (**11**), and the mixture was stirred for 1 h at room temperature. The reaction was evaporated in vacuo and methanol was added and evaporated, giving 1.60 g of a foam. The foam was percolated through a 40-mm-diameter 17-cm flash silica gel

column consecutively with EtOAc, 5% MeOH–95% EtOAc, and 10% MeOH–90% EtOAc, affording 394 mg of the *E*-isomer (R_f = 0.47 with 10% MeOH–90% EtOAc), 80 mg of a mixture of *E*- and *Z*-isomers and 762 mg of the *Z*-isomer (R_f = 0.37 with 10% MeOH–90% EtOAc). Each of these fractions contained CF₃COOH by ¹⁹F NMR. The 762 mg of *Z*-isomer was stirred in 3 mL of 10% MeOH–90% EtOAc, filtered, and dried to give 491 mg (42%) of the *Z*-isomer of 12 free of CF₃COOH. The partially purified *E*-isomer from above (394 mg) was similarly treated, affording 230 mg (20% yield) of the *E*-isomer of 12 free of CF₃COOH.

(*Z*)-12: mp 150–158 °C; ¹H NMR (CD₃OD, 90 MHz) δ 5.1 (m, 2 H), 6.35 (d, 1 H, J = 6 Hz), 7.5–8.2 (m, 5 H), 8.63 (s, 1 H), 8.72 (s, 1 H); HRMS calcd for C₁₇H₁₄FN₅O₄ 371.1030, found 371.1015.

(*E*)-12: mp 155–160 °C; ¹H NMR (CD₃OD, 90 MHz) δ 5.00–5.10 (m, 2 H), 6.37 (1 H, d, J = 7 Hz), 6.48 (1 H, d, J = 7 Hz), 6.48 (1 H, Cl, J = 75 Hz), 7.54–8.19 (m, 5 H), 8.53 (s, 1 H), 8.62 (s, 1 H); HRMS calcd for C₁₇H₁₄FN₅O₄ 371.1030, found 371.1028.

(*Z*)-9-(5-Deoxy-5-fluoro-β-D-erythro-pent-4-enofuranosyl)adenine (6). **Method A.** A suspension of 491 mg (1.6 mmol) of (*Z*)-12 in (ca. 10 mL) EtOH was evaporated. The residue in EtOH (35 mL) was saturated with anhydrous ammonia at 0 °C. The resulting solution was capped (Carius tube) and stirred at room temperature for 18 h. The mixture was evaporated and the residue recrystallized from methanol (6 mL) to afford 217 mg (61% yield) of 6 (dried under high vacuum at 80 °C). The filtrates were chromatographed (30 mm × 17 cm flash silica gel) with 10% MeOH–90% EtOAc to give an additional 83 mg (84% combined yield) of 6, identical in all respects with the material prepared by Method B from 21.

(*E*)-9-(5-Deoxy-5-fluoro-β-D-erythro-pent-4-enofuranosyl)adenosine (13). **Method A.** A suspension of (*E*)-12 (103 mg, 0.33 mmol) in 10 mL of EtOH was evaporated and the residue treated with ethanolic ammonia as described for (*Z*)-12. The residue from evaporation of the ethanolic ammonia was stirred in 1 mL of MeOH and filtered to provide 51 mg of 13 as a white solid (68% yield), identical with the product prepared by Method B from 21.

5'-Deoxy-5'-[(4-methoxyphenyl)thio]-2',3'-O-isopropylideneadenosine (17) (AcOH Complex). To a mixture of 2',3'-O-isopropylideneadenosine (16.4 g, 0.053 mol) and 4-methoxyphenyl disulfide (26.0 g, 0.094 mol) (Fairfield Chemical) in dry pyridine (125 mL) was added tributylphosphine (23.3 mL, 0.094 mol) via syringe. After 15 min at room temperature, a solution formed. The reaction was stirred overnight under argon, diluted with an equal volume of ethyl acetate, and washed with cold 0.5 N NaOH (2 × 50 mL), dilute acetic acid (2 × 50 mL), and brine (50 mL). The organic layer was dried (MgSO₄) and evaporated to an oil that crystallized under high vacuum. The white crystals were triturated with hexane (2 × 500 mL) and dried in a vacuum oven at 50 °C, providing 17 as an analytically pure acetic acid complex (24.9 g, 96%): mp 70 °C to a wax that clears at 97 °C; ¹H NMR (DMSO-*d*₆, 300 MHz) δ 1.32 (s, 3 H), 1.50 (s, 3 H), 1.91 (s, 3 H, HOAc), 3.07 (dd, 1 H, J = 13.7 and 6.5 Hz), 3.15 (dd, 1 H, J = 13.9 and 8.2 Hz), 3.74 (s, 3 H), 4.14 (ddd, 1 H, J = 2.3, 6.9, and 6.4 Hz), 5.04 (dd, 1 H, J = 6.1 and 2.5 Hz), 5.52 (dd, 1 H, J = 6.1 and 2.2 Hz), 6.16 (d, 1 H, J = 2.0 Hz), 6.88 (d, 2 H, J = 8.8 Hz), 7.31 (d, 2 H, J = 8.6 Hz), 8.17 (s, 1 H), 8.33 (s, 1 H); MS (CI/CH₄) m/z 430 (M + H)⁺. Anal. (C₂₀H₂₃N₅O₄S·CH₃CO₂H) C, H, N.

5'-Deoxy-5'-[(4-methoxyphenyl)sulfinyl]-2',3'-O-isopropylideneadenosine (18). To a solution of 17 (23.1 g, 0.054 mol) in CH₂Cl₂ (225 mL) cooled to –40 °C was added a solution of 85% 3-chloroperbenzoic acid (11.1 g, 0.054 mol) in CH₂Cl₂ (225 mL) dropwise. The reaction was allowed to warm to room temperature, washed with ice cold saturated aqueous NaHCO₃, dried (MgSO₄), and evaporated in vacuo. The resulting gel was purified by flash silica gel chromatography (1 L). Elution with ethanol–ethyl acetate (5:95) provided sulfone (2.2 g), IR (KBr) 1140 cm^{–1}. Further elution with ethanol–EtOAc (1:9) provided the diastereomeric sulfoxides, eluting as two bands. The fractions containing product were combined and evaporated to a white solid (20.1 g, 84%). In another run, the sulfoxides were isolated separately in a 10:9.2 ratio (faster moving:slower moving by TLC). Faster moving sulfoxide: IR (KBr) 1090 cm^{–1}; ¹H NMR (CDCl₃,

300 MHz) δ 1.37 (s, 3 H), 1.61 (s, 3 H), 3.16 (dd, 1 H, J = 13.1 and 4.0 Hz), 3.33 (dd, 1 H, J = 13.1 and 10.2 Hz), 3.83 (s, 3 H), 4.80 (dd, 1 H, J = 4.5 and 3.3 Hz), 5.09 (dd, 1 H, J = 6.2 and 3.3 Hz), 5.55 (dd, 1 H, J = 6.2 and 1.8 Hz), 5.78 (br s, 2 H), 6.08 (d, 1 H, J = 1.8 Hz), 6.92 (d, 2 H, J = 8.9 Hz), 7.37 (d, 2 H, J = 9.0 Hz), 7.87 (s, 1 H), 8.26 (s, 1 H); MS (CI/CH₄) m/z 446 (M + H)⁺; HRMS calcd for C₂₀H₂₃N₅O₅S 445.1420, found 445.143. Slower moving sulfoxide: IR (KBr) 1089 cm^{–1}; ¹H NMR (CDCl₃, 300 MHz) δ 1.37 (s, 3 H), 1.57 (s, 3 H), 3.27 (dd, 1 H, J = 13.3 and 5.7 Hz), 3.38 (dd, 1 H, J = 13.6 and 7.1 Hz), 3.84 (s, 3 H), 4.55 (ddd, 1 H, J = 2.8, 6.4 and 6.4 Hz), 5.27 (dd, 1 H, J = 6.3 and 2.9 Hz), 5.35 (dd, 1 H, J = 6.5 and 2.3 Hz), 5.51 (dd, 1 H, J = 6.3 and 2.4 Hz), 5.73 (br s, 2 H), 6.05 (d, 1 H, J = 2.6 Hz), 6.93 (d, 2 H, J = 8.7 Hz), 7.48 (d, 2 H, J = 8.8 Hz), 7.90 (s, 1 H), 8.33 (s, 1 H); HRMS calcd for C₂₀H₂₃N₅O₅S 445.1420, found 445.1416.

5'-Deoxy-5'-fluoro-5'-[(4-methoxyphenyl)sulfinyl]-2',3'-O-isopropylideneadenosine (20). Sulfoxide 18 (as a mixture of diastereomers) (10.5 g, 0.0236 mol) was dissolved in CH₂Cl₂ (150 mL) and (diethylamido)sulfur trifluoride (13.2 mL, 0.1 mol) was added all at once. The reaction was immediately placed in a preheated oil bath (45 °C) and stirred under an argon atmosphere. The progress of the reaction was followed by TLC (CHCl₃–acetone, 1:1) and after 2.5 h, the solution was washed with ice cold saturated aqueous NaHCO₃, dried (K₂CO₃), and cooled to 10 °C with an ice bath. [A small aliquot of 19 was evaporated in vacuo for ¹⁹F NMR (CDCl₃, 282 MHz) δ –159.14 (dd, J = 53.3 and 14.4 Hz), –159.79 (dd, J = 53.0 and 13.8 Hz). The integrated peak ratios were 10:4.2 for the two fluoro sulfides and the product ratios were the same starting with either isomer of the sulfoxide or the diastereoisomeric mixture.] The solution was stirred and treated portionwise with solid 85% 3-chloroperbenzoic acid (8.5 g) until TLC (CHCl₃–acetone, 1:1) indicated complete conversion to sulfoxide. The reaction solution was washed with ice cold saturated aqueous NaHCO₃, dried (K₂CO₃), and evaporated to a tan foam. The foam was purified by flash chromatography (700 mL silica gel) (CHCl₃–acetone 3:2) providing 20 as a light tan solid (5.9 g, 54%) as a mixture of diastereomers: mp 106–114 °C; IR (KBr) 1180 cm^{–1}; ¹H NMR (CDCl₃) δ 1.4–1.65 (several s, 6 H), 3.81–3.86 (singlets, 3 H, MeO), 4.4–5.8 (several m, 4 H), 6.0–6.3 (several m, 3 H), 6.65–8.36 (several m and s, 6 H); ¹⁹F NMR (CDCl₃) δ –193.11 (dd, J = 47.7 and 26.8 Hz) and –195.05 (dd, J = 47.7 and 19.5 Hz) for two major diastereoisomers; MS (CI/CH₄) m/z 464 (M + H)⁺; HRMS calcd for C₂₀H₂₂FN₅O₅S 463.13257, found 463.1317.

(*E*)- and (*Z*)-4',5'-Didehydro-5'-deoxy-5'-fluoro-2',3'-O-isopropylideneadenosine (21). A solution of fluoro sulfoxide 20 (6.5 g, 0.014 mol), diglyme (120 mL), and diisopropylethylamine (9 g, 0.07 mol) was purged with nitrogen for 20 min and then heated at 145 °C (bath temperature). The progress of the reaction was followed by TLC (EtOAc–MeOH 9:1). After 24 h an additional 6 g of diisopropylethylamine was added and the reaction was heated an additional 24 h. The solvent was removed by Kugelrohr distillation under high vacuum at ca. 70–80 °C, and the residue was purified by flash chromatography (700 mL of silica gel; EtOAc–MeOH 96:4) to provide 3.3 g (77%) of 21 as a white solid, which was a 2.7:1 mixture of *Z*- and *E*-isomers: ¹H NMR (CDCl₃, 300 MHz) δ 1.44 and 1.47 (2 s, 3 H, *E*- and *Z*-isomer), 1.59 (s, 3 H), 5.49 (t, 1 H, J = 6.4 Hz), 5.66 (m, 3 H, includes NH₂), 6.27 and 6.28 (2 d, 1 H, J = 1.4 and 0.9 Hz, *E*- and *Z*-isomer), 6.51 and 6.91 (2 d, J = 74.2 and 77.2 Hz, *Z*- and *E*-isomers, respectively), 7.81 and 7.85 (2 s, 1 H, *E*- and *Z*-isomer), 8.31 and 8.33 (2 s, 1 H, *Z*- and *E*-isomer); ¹⁹F NMR (CDCl₃, 282 MHz) δ (from CFCl₃) –160.21 (d, J = 74.5 Hz), –175.66 (d, J = 77.3 Hz); integrated ratio 3.4 to 1.0; ms (CI/CH₄) m/z 308 (M + H)⁺; HRMS calcd for C₁₃H₁₄FN₅O₃ 308.1159, found 308.1161.

(*Z*)- and (*E*)-5-Deoxy-5-fluoro-β-D-erythro-pent-4-enofuranosyladenine (6) and (13). **Method B.** The protected vinyl fluoride 21 (3.2 g, 0.01 mol) was dissolved in CF₃COOH–water (75:25) (130 mL) and stirred at room temperature for 16 h. The solution was evaporated to dryness in vacuo at room temperature and reevaporated several times with ethanol. The residue was dried under high vacuum at room temperature (4.77 g) and dissolved in methanol (100 mL). The solution was divided into four 25-mL aliquots and applied to four BIO-RAD AD 1-X2 (OH[–] form) (200–400 mesh) columns (200 mL of resin each) packed with methanol.³² The columns were eluted with methanol

at a rate of 4 mL/min and fractions were monitored by a UV detector. After 3 days, the faster moving *E*-isomer (13) started to elute from the column. After 14 h, the peak had completely eluted and the *Z*-isomer (6) started to elute and was collected in ca. 1.4 L of eluant. The two separate bands were concentrated to white crystalline solids in vacuo providing 0.50 g (19%) of the faster moving 13 and 1.30 g (48%) of the slower moving 6. A small sample of 6 was recrystallized from EtOAc: mp 218–221 °C dec; ¹H NMR (DMSO-*d*₆, 300 MHz) δ 4.60 (t, 1 H, *J* = 4.6 Hz), 5.05 (dd, 1 H, *J* = 6.9 and 4.8 Hz), 5.64 (d, 1 H, *J* = 4.6 Hz, OH), 5.79 (d, 1 H, *J* = 6.6 Hz, OH), 6.22 (d, 1 H, *J* = 7.1 Hz), 6.67 (d, 1 H, *J* = 75.8 Hz), 7.38 (s, 2 H, NH₂), 8.17 (s, 1 H), 8.46 (s, 1 H); ¹⁹F NMR (CD₃OD, 282 MHz) δ (from external CFCl₃) -166.76 (d, *J* = 75.2 Hz); MS (CI/CH₄) *m/z* 268 (M + H)⁺ [α]_D²⁰ = -59.4° (MeOH). Anal. (C₁₀H₁₀FN₅O₃) C, H, N.

A small sample of 13 was recrystallized from acetone: mp 233–234 °C dec; ¹H NMR (DMSO-*d*₆, 300 MHz) δ 4.68 (m, 1 H), 5.05 (m, 1 H), 5.78 (m, 1 H, OH), 7.15 (d, 1 H, *J* = 78.5 Hz), 7.38 (m, 2, NH₂), 8.16 (s, 1 H), 8.45 (s, 1 H); ¹⁹F NMR (DMSO-*d*₆) δ (from external CFCl₃) -181.46 (d, *J* = 78.7 Hz); MS (CI/CH₄) *m/z* 268 (M + H)⁺. Anal. (C₁₀H₁₀FN₅O₃) C, H, N.

NOE difference spectroscopy (DMSO-*d*₆) on 6 and 13 showed a 5% enhancement in the vinyl proton on 6 (δ 6.67) when the 3'-methine proton (δ 4.60) was irradiated and no enhancement on the vinyl proton on 13 (δ 7.15) when the 3'-methine proton (δ 4.68) was irradiated. The coupling constants from C-3' to the fluorine were also measured, and *J*_{C-F} is 4.8 Hz for 6 and 0 for 13.

Conversion of 13 to 6. Vinyl fluoride 13 (20 mg, 0.075 mmol) was dissolved in 3 mL of acetone-CD₃OD (2:1) and placed in an NMR tube. The solution was irradiated with a 275-W sunlamp placed 19 cm from the tube. The progress of the conversion was followed by ¹⁹F NMR. After 4.5 h, conversion was greater than 60%. Prolonged irradiation caused decomposition. The solution was concentrated, redissolved in 5 mL of CH₃OH and purified by passage through 5 mL of BIO-RAD AG 1-X2 (OH form) (200–400 mesh) (CH₃OH) to give 12 mg (60%) of 6 and 8 mg (40%) of 13.

Experimental details for compounds 22, 24, 26, 28, 29, and 32 are given in the supplementary material.

(Z)- and (E)-4',5'-Didehydro-2',5'-dideoxy-5'-fluoro-adenosine (34 and 35, Respectively). The mixture 32 from the sulfoxide elimination is treated with ethanolic ammonia to give 34 and 35 (an oil), from which 34 is crystallized (EtOAc).

A lengthier procedure to yield 35 is described in the supplementary material, along with additional spectral data.

34: ¹H NMR (DMSO-*d*₆) δ 2.40–2.48 (m, 1 H, H-2'-down), 3.16 (dt, *J* ≈ 6–7 on 13, H-2'-up) 4.98 (m, 1 H, H-3'), 6.65 (d, 1 H, *J* = 76.5 Hz, vinyl H), 6.70 (t, *J* = 6.5 Hz, H-1'), 7.35 (s, 2 H, NH₂), 8.17 (s, 1 H, H-3), 8.39 (s, 1 H, H-8).

35: ¹H NMR (DMSO-*d*₆) δ 2.40–2.49 (m, 1 H, H-2'-down), 3.13 (dt, 1 H, *J* = 6–7.13 Hz, H-2'-up), 5.24 (m, 1 H, H-3'), 5.79 (d, 1 H, *J* = 5.1 Hz, OH), 6.68 (t, 1 H, *J* = 7.0 Hz, H-1'), 7.08 (d, 1 H, *J* = 78.1 Hz, vinyl H), 7.34 (s, 2 H, NH₂), 8.16 (s, 1 H, H-3), 8.37 (s, 1 H, H-8).

Experimental details for compounds 23, 25, 27, 30, 31, and 33 are given in the supplementary material.

(Z)-9-(5-Deoxy-5-fluoro-β-D-threo-pent-4-enofuranosyl)-9H-purin-6-amine (36). Nucleoside (Z)-33 (1.070 g, 1.8 mmol) was treated with ethanolic saturated ammonia (room temperature, overnight). The mixture was evaporated and the residue was purified by flash chromatography (EtOAc, then 10% MeOH–90% EtOAc) to afford 344 mg (70%) of solid 36. Trituration with 5 mL of boiling EtOAc afforded 190 mg (30%) of 36, mp 203–205 °C dec, after drying at 100 °C in a drying pistol in vacuo. Thermogravimetric analysis revealed that the sample retained EtOAc (7.6% by weight; loss at 100–125 °C): ¹H NMR (excluding EtOAc, DMSO-*d*₆) δ 4.27 (br s, 1 H), 4.63 (br s, 1 H), 5.90 (br s, 1 H, OH), 6.13 (br s, 1 H, OH), 6.61 (d, 1 H, *J* = 4.1 Hz), 6.67 (d, 1 H, *J* = 75.9 Hz), 7.34 (br s, 2 H, NH₂), 8.17 (s, 2 H, H-3, H-8) (D₂O exchange sharpened the δ 4.63 signal to a doublet. Irradiation at δ 4.63 (H-3') produced a NOE on the δ 6.67 (vinyl H) signal as well as the OH signals and H-8.); ¹⁹F NMR (DMSO-*d*₆) -164.89 (d, *J* = 76.6 Hz); MS (CI/CH₄) *m/z* 268 (M + H)⁺; HRMS (M + H)⁺ calcd for C₁₀H₁₁FN₅O₃ 268.0845, found 268.0828. Anal. Calcd (C₁₀H₁₀FN₅O₃) C, 44.95; H, 3.77; N, 26.21.

Found C, 44.78, H, 4.09; N, 23.50.

(E)-9-(5-Deoxy-5-fluoro-β-D-threo-pent-4-enofuranosyl)-9H-purin-6-amine (37). Nucleoside (E)-33 (414 mg) was treated with ammonia as described for (Z)-33. The product was crystallized from methanol to afford 99 mg (52%) of 37: mp 255–260 °C dec; ¹H NMR (DMSO-*d*₆) δ 4.17 (m, 1 H), 4.85 (m, 1 H), 6.05 (d, 1 H, *J* = 4.9 Hz, OH), 6.15 (d, 1 H, *J* = 4.4 Hz, OH), 6.60 (d, 1 H, *J* = 3.5 Hz, H-1'), 7.26 (d, 1 H, *J* = 79.1 Hz), 8.15 (s, 1 H, H-3), 8.17 (s, 1 H, H-8). (Irradiation at δ 4.85 (H-3') produced a NOE on the OH-signals and at δ 8.17.); ¹⁹F NMR (DMSO-*d*₆) -181.48 (d, *J* = 79.2 Hz); HRMS (M + H)⁺ calcd for C₁₀H₁₁FN₅O₃ 268.0845, found 268.0848.

Experimental details for compounds 38, 39, 40, 42, and 43 are given in the supplementary material. As an example of the sulfonyl chloride reaction, the synthesis of 41 is described below.

N⁶,N⁶-Dibenzoyl-5'-deoxy-5'-chloro-5'-[(4-methoxyphenyl)sulfinyl]-2',3'-O-isopropylideneadenosine (41). To nucleoside 40 (1.13 g, 1.7 mmol) in 6.5 mL of dry CH₂Cl₂ was added 0.32 mL (4.0 mmol) of pyridine and the mixture cooled in an ice bath under nitrogen. Then 0.18 mL (1.9 mmol) of sulfonyl chloride was added. The mixture was stirred for 20 min and evaporated in vacuo (aspirator, then 1 mm) to afford a foam. The crude product was percolated through a silica gel column (CH₂Cl₂, then 20% EtOAc–80% CH₂Cl₂) followed by a second column to isolate 41 (*R*_f = 0.22, TLC with 20% EtOAc–CH₂Cl₂) as a foam, 0.72 g (60%).

A sample was triturated in CH₂Cl₂–cyclohexane to give a solid, mp 220–228 °C. The remainder of the foam was evaporated from CH₂Cl₂–cyclohexane to give a solid, which was used for the preparation of 42: ¹H NMR (DMSO-*d*₆) δ 1.35 (s, 3 H), 1.58 (s, 3 H), 3.80 (s, 3 H), 2 m, 1 H, large *J* = 7.6 Hz), 5.14 (d, 1 H, *J* = 9.8 Hz), 5.22–5.24 (m, 1 H), 5.60 (d, 1 H, *J* = 6.3 Hz), 6.57 (s, 1 H), 7.03 (m, 2 H), 7.16 (m, 2 H), 7.46 (m, 4 H), 7.60 (m, 2 H), 7.80 (m, 4 H), 8.716 and 8.722 (2 s, 1 H together), 8.882 and 8.887 (2 s, 1 H, together) (there are at least two isomers whose NMR is almost identical); MS (CI/CH₄) *m/z* 688 (M + H)⁺. Anal. (C₃₄H₃₀ClN₅O₇) C, H, N, Cl.

(Z)-4',5'-Didehydro-5'-chloro-5'-deoxyadenosine (44). Nucleoside 43 (98 mg, 0.25 mmol) was treated with saturated methanolic ammonia in a capped tube for 15 h. Solvents were evaporated, and the residue was chromatographed (silica gel, EtOAc, then 10% MeOH–90% EtOAc). Fractions with 44 (*R*_f = 0.26, 10% MeOH–90% EtOAc) were evaporated and EtOAc was added. Evaporation followed by drying in vacuo at 100 °C afforded 47 mg (65%) of 44, mp 202–207 °C. Thermogravimetric analysis and NMR indicated that EtOAc (0.05 mol per mol of nucleoside) was retained: ¹H NMR (DMSO-*d*₆) δ 4.73 (t, 1 H, *J* = 5 Hz), 5.06 (m, 1 H), 5.76 (d, 1 H, *J* = 7 Hz, OH), 5.81 (d, 1 H, *J* = 6 Hz, OH), 5.90 (s, 1 H), 6.22 (d, 1 H, *J* = 8 Hz), 7.36 (br s, 2 H, NH₂), 8.15 (s, 1 H), 8.45 (s, 1 H). (Irradiation at δ 4.73 (H-3') resulted in a NOE at δ 5.06 (H-2'), 5.81 (3'-OH), and 5.90 (vinyl H)); MS (CI/CH₄) *m/z* 284 (M + H)⁺; HRMS (EI) calcd for C₁₀H₁₀ClN₅O₃ 283.0472; found 283.0470. Anal. C, 42.34; H, 3.55; N, 24.69; found C, 42.56; H, 3.59; N, 23.83.

Experimental details for compounds 46, 47, and 48 are given in the supplementary material.

(E)-N⁶-Benzoyl-4',5'-didehydro-5'-chloro-5'-fluoro-2',3'-O-isopropylideneadenosine (49). In a 250-mL round-bottom flask a mixture of 48 (2.2 g, 3.12 mmol), 4 mL of diisopropylethylamine, and 80 mL of diglyme was purged with N₂ for 15 min. A condenser was attached and the reaction heated at 145 °C under argon for 18 h. The solvent was removed via Kugelrohr vacuum distillation. The residue was chromatographed on 300 g of silica gel (acetone–CHCl₃, 1:3) to afford 622 mg (45%) of 49 as a tan oil: ¹H NMR (CDCl₃) δ 1.48 (s, 3 H), 1.61 (s, 3 H), 5.48 (dd, 1 H, *J* = 1.2 and 5.9 Hz), 5.84 (d, 1 H, *J* = 6.0 Hz), 6.37 (s, 1 H), 7.27–8.78 (m, 7 H); ¹⁹F NMR (CDCl₃) δ -104.03 (s); ¹³C NMR (CDCl₃) 79.47 (d, *J*_{C-F} = 2.3 Hz); MS (CI/CH₄) *m/z* 446 (M + H)⁺.

(E)-4',5'-Didehydro-5'-deoxy-5'-chloro-5'-fluoro-adenosine (51). Compound 49 (610 mg, 1.37 mmol) was stirred with 15 mL of 75% trifluoroacetic acid and 25% water for 22 h at ambient temperature. The reaction was concentrated in vacuo and the residue azeotroped twice with EtOH. The residue (50) was dissolved in 10 mL of EtOH, cooled to 0 °C and saturated with NH₃. The capped tube was allowed to stand at room temperature overnight. The reaction was concentrated and then purified by

flash chromatography (20% CH₃OH-CHCl₃ with 1% NH₄OH) to give 420 mg (100%) **51** as a pale tan oil. The tan oil was dissolved in 15 mL of acetone and placed in a freezer for 4 days. The resulting crystals (170 mg) were collected: mp 232-234 °C; ¹H NMR (CD₃OD) δ 4.78 (d, 1 H, *J* = 4.2 Hz), 5.14 (dd, 1 H, *J* = 5.1 and 7.8 Hz), 6.30 (d, 1 H, *J* = 7.8 Hz), 8.21 (s, 1 H), 8.34 (s, 1 H); ¹⁹F NMR (CD₃OD, 282 MHz) δ -110.355 (s); ¹³C NMR (CD₃OD) δ 74.33 (d, *J*_{C-F} = 1.95 Hz); MS (CI/CH₄) *m/z* 302 (M + H)⁺. Anal. (C₁₀H₉ClFN₅O₃·1/3 C₂H₆O) C, H, N.

(*Z*)-4',5'-Didehydro-5'-deoxy-5'-chloro-5'-fluoroadenosine (**52**). Nucleoside **51** in acetone-CH₃OH (2:1) was irradiated for 4 h. The ¹⁹F NMR of the reaction indicated a 2:1 mixture of **51** and **52**. Nucleoside **52** was isolated by reverse phase HPLC (Zorbax R_x 250 × 4.6 mm column, mobile phase methanol-water) and characterized by NMR and mass spectra.

The coupling constant of C-3' to the fluorine was measured for **51** (2.3 Hz) and **52** (0). Comparison to the values recorded for **6** and **13** allowed assignment of olefin geometry: MS (CI/CH₄) *m/z* 302 (M + H)⁺; ¹H NMR (CD₃OD) δ 4.80 (d, 1 H, *J* = 6 Hz), 5.18 (dd, 1 H, *J* = 5.4 and 7.5 Hz), 6.28 (d, 1 H, *J* = 7.5 Hz), 8.21 (s, 1 H), 8.34 (s, 1 H); ¹⁹F NMR (CD₃OD) δ -126.63 (s).

Acknowledgment. The authors thank Dr. Edward Huber for obtaining NMR spectra and for assistance in interpretation.

Supplementary Material Available: Details of the preparation and assay of rat liver SAH hydrolase and details of the preparation of and spectra for compounds **22-27**, **29**, **31-35**, **38-43**, **47**, and **48** (13 pages). Ordering information is given on any current masthead page.

Quinolone Antibacterials: Preparation and Activity of Bridged Bicyclic Analogues of the C₇-Piperazine

John S. Kiely,* Marland P. Hutt, Townley P. Culbertson, Ruth A. Bucsh, Donald F. Worth, Lawrence E. Lesheski, Rocco D. Gogliotti, Josephine C. Sesnie, Marjorie Solomon, and Thomas F. Mich

Parke-Davis Pharmaceutical Research Division, Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, Michigan 48105.
Received May 8, 1990


A series of quinolone and naphthyridine antibacterial agents possessing as the C₇-heterocycle bicyclic 2,5-diaza-bicyclo[*n.2.m*]alkanes, where *n* = 2, 3 and *m* = 1, 2, and a series including 4-aminopiperidine and 3-amino-8-azabicyclo[3.2.1]octanes have been prepared and evaluated in vitro and in vivo for antibacterial activity against a variety of Gram-negative and Gram-positive organisms. These compounds were also tested against the target enzyme bacterial DNA gyrase. All the examples investigated are nearly equipotent with the parent 7-piperazinyl analogues. Only *endo*-7-(3-amino-8-azabicyclo[3.2.1]oct-8-yl)-1-cyclopropyl-6,8-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid displays activity that surpasses that of the piperazine parent.

Introduction

Quinolone antibacterial agents are a major class of anti-infectives with significant potential for continued development.¹ In virtually all modern examples of this anti-infectives class, a nitrogen heterocycle is attached at the C₇-position of the quinolone or naphthyridine nucleus.² This moiety plays a significant role in determining antibacterial spectrum and potency and represents a site amenable to significant modification. We report our results from a study replacing the frequently employed 7-piperazinyl substituent with various bicyclic analogues.³ We were interested in determining the effect these bicyclic C₇-substituents would have on minimum inhibitory concentrations, inhibition of the gyrase enzyme, and in vivo protective doses in mice.

Ciprofloxacin, which contains a 7-piperazinyl moiety, served as the beginning point for our studies. The SAR of the C₇-side chain has been extensively explored^{4,5} and the importance of proper side-chain length and ring size

Chart I



- (1) *The Quinolones*; Andriole, V. T., Ed.; Academic Press: New York, 1988.
- (2) Heck, J. V. *Annu. Rep. Med. Chem.* 1989, 24, 101. Fernandes, P. B.; Chu, D. T. W. *Ibid.* 1988, 23, 133. Fernandes, P. B.; Chu, D. T. W. *Ibid.* 1987, 22, 117.
- (3) Hutt, M. P.; Mich, T. F.; Culbertson, T. P. U.S. Pat. 4,571,396, 1986.
- (4) (a) Egawa, H.; Miyamoto, T.; Minamida, A.; Nishimura, Y.; Okada, H.; Uno, H.; Matsumoto, J. *J. Med. Chem.* 1984, 27, 1543. (b) Domagala, J. M.; Hanna, L. D.; Heifetz, C. L.; Hutt, M. P.; Mich, T. F.; Sanchez, J. P.; Solomon, M. *J. Med. Chem.* 1986, 29, 394.
- (5) Culbertson, T. P.; Domagala, J. M.; Hagen, S. E.; Hutt, M. P.; Nichols, J. B.; Mich, T. F.; Sanchez, J. P.; Schroeder, M. C.; Solomon, M.; Worth, D. F. *International Telesymposium on Quinolones*; Fernandes, P. B., Ed.; J. R. Prous Science: Barcelona, Spain, 1989.

has been elucidated.⁴ The optimum size for the C₇-substituent has been defined as a ring of five or six atoms.