

Pharmacological characterization of muscarinic receptors in mouse isolated urinary bladder smooth muscle

*¹A. Choppin & ^{1,2}R.M. Eglén

¹Genitourinary-Pharmacology, Neurobiology Unit, Roche Bioscience, Palo Alto, California, CA 94304, U.S.A.

1 The pharmacological characteristics of muscarinic receptors in the male mice urinary bladder smooth muscle were studied.

2 (+)-Cis-dioxolane, oxotremorine-M, acetylcholine, carbachol and pilocarpine induced concentration-dependent contractions of the urinary bladder smooth muscle ($pEC_{50} = 6.6 \pm 0.1$, 6.9 ± 0.1 , 6.7 ± 0.1 , 5.8 ± 0.1 and 5.8 ± 0.1 , $E_{Max} = 3.2 \pm 0.8$ g, 2.7 ± 0.4 g, 1.0 ± 0.1 g, 2.7 ± 0.3 and 0.9 ± 0.2 g, respectively, $n = 4$). These contractions were competitively antagonized by a range of muscarinic receptor antagonists (pK_B values): atropine (9.22 ± 0.09), pirenzepine (6.85 ± 0.08), 4-DAMP (8.42 ± 0.14), methoctramine (5.96 ± 0.05), p-F-HHSiD (7.48 ± 0.09), tolterodine (8.89 ± 0.13), AQ-RA 741 (7.04 ± 0.12), s-secoverine (8.21 ± 0.09), zamifenacin (8.30 ± 0.17) and darifenacin (8.70 ± 0.09).

3 In this tissue, the pK_B values correlated most favourably with pK_i values for these compounds at human recombinant muscarinic M_3 receptors. A significant correlation was also noted at human recombinant muscarinic $m5$ receptors given the poor discriminative ability of ligands between M_3 and $m5$ receptors.

4 In recontraction studies, in which the muscarinic M_3 receptor population was decreased, and conditions optimized to study M_2 receptor activation, methoctramine exhibited an affinity estimate consistent with muscarinic M_3 receptors ($pK_B = 6.23 \pm 0.14$; $pA_2 = 6.16 \pm 0.03$).

5 Overall, these data suggest that muscarinic M_3 receptors are the predominant, if not the exclusive, subtype mediating contractile responses to muscarinic agonists in male mouse urinary bladder smooth muscle.

British Journal of Pharmacology (2001) **133**, 1035–1040

Keywords: Muscarinic receptors; M_3 -receptor; urinary bladder smooth muscle

Abbreviations: AQ-RA 741, (11-({4-[4-(diethylamino)butyl]-1-piperidinyl}acetyl)-5,11-dihydro-6H-pyrido(2,3-b)(1,4)benzodiazepine-6-one); 4-DAMP, 4-diphenylacetoxy-*N*-methylpiperidine; p-F-HHSiD, para fluoro hexahydrosiladifenidol

Introduction

Muscarinic receptors are pharmacologically classified into four subtypes, M_1 , M_2 , M_3 and M_4 which equate with four of the known muscarinic receptor gene products ($m1$, $m2$, $m3$ and $m4$) (Hulme *et al.*, 1990; Caufield, 1993; Eglén *et al.*, 1996 for reviews). A fifth gene has been identified, $m5$, for which a role has yet to be unambiguously defined (see Eglén & Nahorski, 2000 for review).

Being widely distributed, muscarinic receptors play a key physiological role in peripheral organs, including the urinary bladder. In most smooth muscles, the muscarinic M_2 receptor subtype accounts for 70–80% of the receptor population whereas the M_3 receptor subtype forms only 20–30% (Eglén *et al.*, 1996). In this tissue from rat, it was proposed that muscarinic M_3 receptor activation primarily causes direct contraction of the smooth muscle and the muscarinic M_2 receptor contracts the tissue indirectly, by reversing sympathetically mediated relaxation (Hegde *et al.*, 1997). Pharmacological characterization of muscarinic re-

ceptors mediating contraction of detrusor muscle has been well established in rat (Longhurst *et al.*, 1995; Hegde *et al.*, 1997), rabbit (Tobin & Sjogren, 1995; Choppin *et al.*, 1998), guinea-pig (Noronha-Blob *et al.*, 1989) and human (Newgreen & Naylor, 1996). Several investigations of the muscarinic receptors mediating contractions of mouse bladder have been undertaken (Durant *et al.*, 1991; Paravicini *et al.*, 2000; Stengel *et al.*, 2000; Welsh *et al.*, 2000), and most suggest a major role of the muscarinic M_3 receptor in the contractile response, with the role of the M_2 receptor, if any, being unresolved. Recently, the situation has become clearer with the use of transgenic mice that lack either the muscarinic M_2 (Stengel *et al.*, 2000) or M_3 receptor (Matsui *et al.*, 2000). These data collectively indicate a minimal role for the former and that the latter mediates most of the contractile response. The objective of the present study was therefore to examine, using a range of defining antagonists, the pharmacological characteristics of muscarinic receptors present in male mouse urinary smooth muscle using isolated tissue studies.

A preliminary account of the findings has been presented previously to the 9th international symposium on subtypes of muscarinic receptors (Eglén & Choppin, 2001).

*Author for correspondence at: Deltagen, Inc., 1003 Hamilton Avenue, Menlo Park, CA 94025, U.S.A. E-mail: Achoppin@Deltagen.com

²Current address: DiscoverX Corporation, 42501 Albrae Street, Suite 100, Fremont, CA 94538, USA

Methods

In vitro contractile studies

Male C57BL6 mice (25–30 g) were euthanized by CO₂ asphyxiation. The urinary bladder was isolated, cleared of adhering adipose tissue and placed in oxygenated Krebs solution (composition in mM: NaCl 118.2, KCl 4.6, CaCl₂ 2.5, MgSO₄·7H₂O 1.2, KH₂PO₄ 1.2, NaHCO₃ 24.8 and dextrose 10.0). The physiological solution contained indomethacin (10 µM) in order to reduce prostaglandin-induced spontaneous activity of the tissues. Four strips of urinary bladder smooth muscle were cut from the supratrigonal portion of the bladder (longitudinal section). The tissues were mounted in 10 ml organ baths containing Krebs solution, maintained at 37°C and constantly aerated with 95% O₂/5% CO₂ (pH=7.4). Grass FT03 transducers were used to measure changes in isometric tension of the tissues, which were displayed on a Grass 7E polygraph. The tissues were maintained at a resting tension of 1 g during an equilibration period of 60 min. Tension adjustments were made as necessary. The tissues were washed every 15 min.

The viability of each tissue was assessed by determining the contractile response to KCl (30 mM) at the start of the experimental protocol. After washing, tissues were re-equilibrated for 10 min and allowed to regain baseline tension. Cumulative concentration-effect curves to agonists ((+)-cis-dioxolane, oxotremorine-M, acetylcholine, carbachol and pilocarpine; 1 nM–0.1 mM) were then constructed in each tissue. Thereafter, tissues were equilibrated in either the absence (time control) or presence of antagonist for a 90 min period during which tissues were washed every 10 min. Subsequently, a second concentration-effect curve to the same agonist was constructed.

Recontraction experiments

After an initial concentration-response curve to (+)-cis-dioxolane was established, the tissues were washed and equilibrated with 4-DAMP mustard (40 nM) for 60 min in the presence of methoctramine (0.3 µM). This procedure enabled selective alkylation of M₃ but not M₂ receptors (Hegde *et al.*, 1997). 4-DAMP mustard was then removed from the tissues by overflow with Krebs solution containing methoctramine (0.3 µM) every 10 min for 60 min and subsequently with methoctramine-free Krebs solution every 10 min for 90 min. The tissues were then contracted with 90 mM of KCl and subsequently relaxed with isoproterenol (30 µM). Once the tissues had relaxed to baseline, a cumulative concentration-effect curve to (+)-cis-dioxolane (1 nM–0.1 mM) was constructed.

Effects of an M₂ antagonist (methoctramine) on the recontractile responses to (+)-cis-dioxolane

After constructing two concentration-effect curves to (+)-cis-dioxolane under conditions described above, a third cumulative concentration effect curve to (+)-cis-dioxolane (1 nM–0.3 mM) was constructed after equilibration of tissue in absence (time control) or presence of methoctramine (0.1–1.0 µM) for 90 min.

Data analysis

Contractions were recorded as changes in tension from baseline and expressed as a percentage of the maximum response of the first agonist concentration-effect curve. Agonist concentration-response curves were fitted using a nonlinear iterative fitting program (Origin, Microcal Software, Inc., Northampton, MA, U.S.A.) using the relationship of Parker & Waud (1971). Agonist potencies and maximum response are expressed as pEC₅₀ (– logarithm of the molar concentration of agonist producing 50% of the maximum response) and E_{max}, respectively. Concentration-ratios (CRs) were determined from EC₅₀ values in the presence and absence of antagonist. Antagonist affinity estimates (pK_B values) were determined with the equation described by Furchgott (1972) (pK_B = –log ([antagonist]/CR-1)) or using the method of Arunlakshana & Schild (1959) using at least three concentrations of the antagonist (pA₂ values). In cases where the slope of the linear regression was not significantly different from unity, the slope was constrained to unity and the data expressed as the pK_B value. All data are expressed as mean ± s.e.mean. Pearson correlation coefficients (*r*) and associated *P*-values were calculated using the method described by Dixon & Massey (1983). The sum of squares of differences in affinity estimates for each plot (Σ (y–x)², noted ssq) defines the proximity of the data points to the line of identity (y=x).

Compounds used

Atropine sulphate, indomethacin and oxybutynin chloride were obtained from Sigma Chemical Co (MO, U.S.A.). (+)-Cis-dioxolane, acetylcholine, carbachol, oxotremorine-M, pilocarpine, pirenzepine dihydrochloride, methoctramine hydrochloride, 4-diphenylacetoxy-*N*-methylpiperidine (4-DAMP) methiodide, 4-DAMP mustard and para fluoro hexahydrosiladifenidol (p-F-HHSiD) hydrochloride were obtained from Research Biochemicals Inc. (MA, U.S.A.). Darifenacin hydrobromide and zamifenacin fumerate were generously provided by Pfizer Central Research (Sandwich, Kent, U.K.). AQ-RA 741 (11-({4-[4-(diethylamino)butyl]-1-piperidinyl}acetyl)-5,11-dihydro-6H-pyrido(2,3-b)(1,4)benzodiazepine-6-one) was donated by Boehringer Ingelheim Pharmaceuticals, Inc. (Ridgefield, CT, U.S.A.). Isoproterenol, tolterodine and s-secoverine hydrochloride were synthesized at Roche Bioscience (Palo Alto, U.S.A.).

All compounds were diluted in distilled water except indomethacin which was diluted in polyethylene glycol.

Results

Characterization of muscarinic receptors mediating contractions of the mice isolated urinary bladder smooth muscle

(+)-Cis-dioxolane, oxotremorine-M, acetylcholine, carbachol and pilocarpine induced concentration-dependent contractions of the mice urinary bladder smooth muscle (pEC₅₀=6.6±0.1, 6.9±0.1, 6.7±0.1, 5.8±0.1 and 5.8±0.1, E_{max}=3.2±0.8 g, 2.7±0.4 g, 1.0±0.1 g, 2.7±0.3 and 0.9±0.2 g, respectively, *n*=4). Time-control experiments

showed that two consecutive concentration-effect curves to these agonists could be constructed in the same tissue with no significant temporal change in the agonist potency and maximum response (Figure 1).

Pharmacological characterization of the muscarinic receptor mediating direct contractions was done by determination of antagonist affinities. Several antagonists (atropine, pirenzepine, 4-DAMP, methoctramine, p-F-HHSiD, tolterodine, AQ-RA 741, s-secoverine, zamifenacin and darifenacin) were tested for their ability to inhibit (+)-cis-dioxolane-induced responses and their functional affinity estimates (pK_B) are summarized in Table 1. All these compounds, in a concentration-dependent fashion, with parallel rightward displacements, surmountably antagonized cumulative agonist concentration-response curves. The rank order of antagonist affinities (pK_B) was: atropine (9.22 ± 0.09), tolterodine (8.89 ± 0.13), darifenacin (8.70 ± 0.09), 4-DAMP (8.42 ± 0.14), zamifenacin (8.30 ± 0.17), s-secoverine (8.21 ± 0.09), p-F-HHSiD (7.48 ± 0.09), AQ-RA 741 (7.04 ± 0.12), pirenzepine (6.85 ± 0.08) and methoctramine (5.96 ± 0.05).

Comparison of functional data for mice urinary bladder smooth muscle with binding data at human recombinant muscarinic receptors

Correlation analysis between the affinities of the antagonists at muscarinic receptors in the mice urinary bladder smooth

muscle and the affinities at human recombinant muscarinic receptors showed a significant correlation ($r = 0.94$, $P < 0.0001$, $ssq = 1.42$) at m3 but also at m5 receptors ($r = 0.84$, $P = 0.002$, $ssq = 3.70$). In contrast, poor correlations were observed at m1, m2 and m4 ($r = 0.75$, $ssq = 4.38$; $r = 0.32$, $ssq = 12.59$; $r = 0.67$, $ssq = 5.42$) respectively (Figure 2).

Characterization of muscarinic receptors mediating the reconstrictions in mouse urinary bladder smooth muscle

Under control conditions, (+)-cis-dioxolane produced concentration-dependent contractions of the mouse urinary bladder smooth muscle ($pEC_{50} = 6.57 \pm 0.05$, $n = 4$). After preferential alkylation of M₃ receptor (exposure to 4-DAMP mustard in presence of methoctramine), (+)-cis-dioxolane produced recontractile (reversal of contraction) responses ($pEC_{50} = 6.01 \pm 0.05$, $n = 4$) of KCl-precontracted tissues, which were relaxed with isoproterenol. The maximum recontractile response (expressed as per cent of the control curve) was $26 \pm 2\%$ ($n = 4$). No time-dependent changes in agonist sensitivity were observed during the construction of two consecutive concentration-recontractile effect curves. As shown in Figure 3, methoctramine produced surmountable antagonism of the recontractile response to (+)-cis-dioxolane. The affinity estimate (pA_2) for methoctramine was 6.16 ± 0.03 and the slope of the Schild plot was not significantly different than unity.




Figure 1 Effects of muscarinic agonists on mouse urinary bladder smooth muscle. Contractile effects were expressed as percentages of the maximum response of the control curve. The values shown are means $\pm$ s.e. mean, $n \geq 4$ animals.

Table 1 Affinity estimates for muscarinic antagonists in mouse urinary bladder smooth muscle

Antagonist	m_1 pK_i	m_2 pK_i	m_3 pK_i	m_4 pK_i	m_5 pK_i	Bladder pK_B
Atropine (3 nM)	9.1 ^a	8.9 ^a	9.5 ^a	9.2 ^a	9.1 ^a	9.22 ± 0.09
Pirenzepine (1 µM)	8.0	6.3	6.8	7.1	6.9	6.85 ± 0.08
4-DAMP (10 nM)	9.2 ^a	8.1 ^a	9.3 ^a	8.4 ^a	8.9 ^a	8.42 ± 0.14
Methoctramine (10 µM)	6.7	7.7	6.1	7.0	6.3	5.96 ± 0.05
p-F-HHSiD (0.3 µM)	7.3 ^a	6.6 ^a	7.5 ^a	7.2 ^a	6.7 ^a	7.48 ± 0.09
Tolterodine (10 nM)	8.5 ^b	8.4 ^b	8.5 ^b	8.3 ^b	8.5 ^b	8.89 ± 0.13
AQ-RA 741 (0.3 µM)	7.6 ^c	8.9 ^c	7.5 ^c	7.9 ^c	6.0 ^c	7.04 ± 0.12
s-secoverine (10 nM)	8.5 ^c	8.9 ^c	8.3 ^c	8.6 ^c	7.0 ^c	8.21 ± 0.09
Zamifenacin (0.1 µM)	7.6	7.2	7.9	6.9	7.3	8.30 ± 0.17
Darifenacin (10 nM)	7.8	7.0	8.8	7.7	8.0	8.70 ± 0.09

Values shown are means ± s.e.mean., $n=4$. pK_i values are from Loury *et al.* (1999). ^a pK_i values taken from Hedge *et al.* (1997). ^b pK_i values taken from Nilvebrant *et al.* (1996). ^c pK_i values taken from Choppin *et al.* (1999).




Figure 2 Correlation between the functional affinities (pK_B values) of muscarinic antagonists at muscarinic receptor in mouse isolated urinary bladder smooth muscle and binding affinities (pK_i values) at human recombinant muscarinic receptors (m_1 - m_5 ; a-e respectively). The binding data were taken from Dörje *et al.*, 1991; Eglen *et al.*, 1997; Hegde *et al.*, 1997; Nilvebrant *et al.*, 1996. The broken line is the line of identity ($x=y$) while the solid line is the correlation plot (the inserts give the correlation factors (r) and the sum of squares values (ssq)).

Discussion

Previous studies using mouse urinary bladder (Durant *et al.*, 1991; Lundbeck & Sjögren, 1992) have demonstrated a muscarinic-induced contractile response but did not characterize the receptor subtype(s) involved. The present study has examined in detail the pharmacological characteristics of muscarinic receptors in this tissue.

Mouse urinary bladder smooth muscle

Inspection of the agonist potencies and maximal responses suggest that the muscarinic receptor mediating contraction was associated with a low efficacy. Thus, the partial agonist, pilocarpine, yielded a potency similar to the affinity and gave a lower maximal response than seen with the full agonists. Similar observations have been seen in urinary bladder tissue from rat (Hegde *et al.*, 1997). (+)-Cis-dioxolane produced




Figure 3 Recontraction experiments in mouse urinary bladder smooth muscle: effect of methoctramine on the recontractile concentration-effect to (+)-cis-dioxolane obtained after elevation of adenyl cyclase activity following preferential alkylation of muscarinic M_3 receptors ($n=4$). (a) Time control; (b) $+0.3 \mu\text{M}$ methoctramine; (c) $+1.0 \mu\text{M}$ methoctramine; (d) $+3.0 \mu\text{M}$ methoctramine.

concentration-dependent contractions, which were inhibited in a concentration-dependent and competitive fashion by muscarinic antagonists. The apparent affinity estimates of these antagonists correlated most strikingly with the binding affinities of the antagonists at $m3$ recombinant muscarinic receptors (pK_i are: atropine 9.5; 4-DAMP 9.3; AQ-RA 741, 7.5; darifenacin 8.8; methoctramine 6.1; tolterodine 8.5; pirenzepine 6.8; s-secoverine 8.3; p-F-HHSiD 7.5 and zamifenacin 7.9; $r=0.94$, $ssq=1.42$; Dörje *et al.*, 1991; Eglen *et al.*, 1997; Hegde *et al.*, 1997; Nilvebrant *et al.*, 1996) and are consistent with the exclusive involvement of M_3 muscarinic receptors in the direct contractile response to muscarinic agonists. This accords with findings in the rabbit (Tobin, 1995; Tobin & Sjögren, 1995), rat (Longhurst *et al.*, 1995; Hegde *et al.*, 1997), human (Newgreen & Naylor, 1996) and preliminary results in female mice (Paravicini *et al.*, 2000) bladder. It should be noted, however, that a significantly good correlation ($r=0.84$, $ssq=3.70$) was also obtained with the binding affinities of the antagonists at $m5$ recombinant muscarinic receptors (pK_i values: atropine 9.1; 4-DAMP 8.9; AQ-RA 741, 6.0; darifenacin 8.0; methoctramine 6.3; tolterodine 8.5; pirenzepine 6.9; s-secoverine 7.0; p-F-HHSiD 6.7 and zamifenacin 7.3; Dörje *et al.*, 1991; Eglen *et al.*, 1997; Hegde *et al.*, 1997; Nilvebrant *et al.*, 1996). This is unsurprising since many ligands discriminate poorly between M_3 and $m5$ receptors, and highlights the difficulty of excluding a role for the latter in M_3 mediated responses.

In contrast to the rat urinary bladder and several gastrointestinal smooth muscle tissues, functional studies (recontraction experiments) in the mouse bladder revealed no indirect contractile role of M_2 receptors, and only M_3 receptor activation induced bladder contraction. The low affinity of methoctramine ($pA_2=6.16$) argues against the involvement of an M_2 receptor and the remaining 26% of the maximal response observed with (+)-cis-dioxolane after recontraction are likely due to an incomplete alkylation of muscarinic M_3 receptors. Data from studies performed in M_2 knockout mice (Stengel *et al.*, 2000) suggest that muscarinic M_2 receptors play a minor role in carbachol induced contraction of isolated bladder smooth muscle, since the potency of muscarinic agonists is only modestly reduced, and the maximal response unaffected. Concordantly, data from transgenic mice lacking the muscarinic M_3 receptor also suggest predominant involvement of muscarinic M_3 receptor, as the contraction *in vitro* was virtually abolished in these mice (Matsui *et al.*, 2000). *In vivo*, urinary retention was marked in these animals, suggesting that a dominant, if not exclusive, role of this subtype prevails when voiding reflexes are intact (Matsui *et al.*, 2000). The data obtained in the present study are consistent with these findings. Parasympathetic nerves innervating the urinary bladder are endowed with prejunctional inhibitory muscarinic receptors, which have been classified as muscarinic M_2 receptors in the rabbit (Tobin & Sjögren, 1995) and rat (Somogyi & De Groat,

1992) urinary bladder but M_4 in the guinea-pig urinary bladder. M_2 receptors may also act prejunctionally in the mouse bladder but given the difficulty to distinguish between these two subtypes, this function has not been investigated in the present study.

Conclusions

The present study has shown that the pharmacological antagonist profile of the muscarinic receptors present in the

mouse bladder equates most closely with the M_3 muscarinic receptor. Moreover, these data suggest that only M_3 receptors play a role in both direct and indirect contraction in accord with emerging data from knockout animals. It thus appears that the mouse urinary bladder differs from the mouse ileum and urinary bladder tissue from other species, including rat and possibly human. It therefore remains to be established if mouse tissue represents an optimal species to provide a useful model for disorders of human urinary bladder function.

References

- ARUNLAKSHANA, O. & SCHILD, H.O. (1959). Some quantitative uses of drug antagonists. *Br. J. Pharmacol.*, **14**, 48–58.
- CAUFIELD, M.P. (1993). Muscarinic receptors-characterization, coupling and function. *Pharmacol. Ther.*, **58**, 319–379.
- CHOPPIN, A., EGLIN, R.M. & HEGDE, S.S. (1998). Pharmacological characterisation of muscarinic receptors in rabbit isolated iris sphincter muscle and urinary bladder smooth muscle. *Br. J. Pharmacol.*, **124**, 883–888.
- CHOPPIN, A., LOURY, D.N., WATSON, N., HEGDE, S.S. & EGLIN, R.M. (1999). S-secoverine: a defining ligand in muscarinic M_5 receptors characterization. *Br. J. Pharmacol.*, **128**, 33P.
- DIXON, W.J. & MASSEY, F.J. (1983). Introduction to statistical analysis, 4th edition, New York: McGraw-Hill Publishing Company.
- DÖRJE, F., WESS, J., LAMBRECHT, G., TACKE, R., MUTSCHLER, E. & BRANN, M.R. (1991). Antagonist binding profiles of five cloned human muscarinic receptor subtypes. *J. Pharmacol. Exp. Ther.*, **256**, 727–733.
- DURANT, P.A., SHANKLEY, N.P., WELSH, N.J. & BLACK, J.W. (1991). Pharmacological analysis of agonist-antagonist interactions at acetylcholine muscarinic receptors in a new urinary bladder assay (1991). *Br. J. Pharmacol.*, **104**, 145–150.
- EGLIN, R.M., BONHAUS, D.W., CALIXTO, J.J., CHOPPIN, A., LEUNG, E., LOEB, M., LOURY, D., MOY, T., WILDA, M. & HEGDE, S.S. (1997). Characterization of the interaction of tolterodine at muscarinic receptor subtypes *in vitro* and *in vivo*. *Br. J. Pharmacol.*, **120**, 63P.
- EGLIN, R.M. & CHOPPIN, A. (2001). Pharmacological characterisation of muscarinic receptors in mouse urinary bladder smooth muscle. *Life Sci.*, **68**, 2634.
- EGLIN, R.M., HEGDE, S.S. & WATSON, N. (1996). Muscarinic receptor subtypes and smooth muscle function. *Pharmacol. Rev.*, **48**, 531–565.
- EGLIN, R.M. & NAHORSKI, S.R. (2000). The muscarinic M_5 receptor: a silent or emerging subtype? *Br. J. Pharmacol.*, **130**, 13–21.
- FURCHGOTT, R.F. (1972). The classification of adrenoceptors (adrenergic receptors). An evaluation from the standpoint of receptor theory. In *Catecholamines, Handbook of Experimental Pharmacology*, Vol. 33. Ed. Blaschko, H. & Muscholl, E. pp. 283–335. Berlin, Heidelberg, New York: Springer.
- HEGDE, S.S., CHOPPIN, A., BONHAUS, D., BRIAUD, S., LOEB, M., MOY, T.M., LOURY, D. & EGLIN, R.M. (1997). Functional role of M_2 and M_3 muscarinic receptors in the urinary bladder of rats *in vitro* and *in vivo*. *Br. J. Pharmacol.*, **120**, 1409–1418.
- HULME, E.C., BIRDSALL, N.J.M. & BUCKLEY, N.J. (1990). Muscarinic receptor subtypes. *Ann. Rev. Pharmacol. Toxicol.*, **30**, 633–673.
- LONGHURST, P.A., LEGGETT, R.E. & BRISCOE, J.A.K. (1995). Characterization of functional muscarinic receptors in the rat urinary bladder. *Br. J. Pharmacol.*, **116**, 2279–2285.
- LOURY, D.N., HEGDE, S.S., BONHAUS, D.W. & EGLIN, R.M. (1999). Ionic strength of assay buffers influences antagonist binding affinity estimates at muscarinic M_1 – M_5 cholinergic receptors. *Life Sci.*, **64**, 6P.
- LUNDBECK, F. & SJÖGREN, C. (1992). A pharmacological *in vitro* study of the mouse urinary bladder at the time of acute change in bladder reservoir function after irradiation. *J. Urol.*, **148**, 179–182.
- MATSUI, M., MOTOMURA, D., KARASAWA, H., FUJIKAWA, T., JIANG, J., KOMIYA, Y., TAKAHASHI, S. & TAKETO, M.M. (2000). Multiple functional deficits in peripheral autonomic organs in mice lacking muscarinic acetylcholine receptor gene for the M_3 subtype. *Proc. Natl. Acad. Sci.*, **97**, 9579–9584.
- NEWGREEN, D.T. & NAYLOR, A.M. (1996). Characterization of functional muscarinic receptors in human bladder. *Br. J. Pharmacol.*, **119**, 45P.
- NILVEBRANT, L., SUNDQUIST, S. & GILLBERG, P.-G. (1996). Tolterodine is not subtype ($m1$ – $m5$) selective but exhibits functional bladder selectivity *in vivo*. *Neurourol. Urodyn.*, **15**, 34 (abstract).
- NORONHA-BLOB, L., LOWE, V.C., PATTON, A., CANNING, B., COSTELLO, D. & KINNIE, W.J. (1989). Muscarinic receptors: relationships among phosphoinositide breakdown, adenylate cyclase inhibition, *in vitro* detrusor muscle contractions and *in vivo* cystometrograms studies in guinea-pig bladder. *J. Pharmacol. Exp. Ther.*, **249**, 843–851.
- PARAVICINI, T., PENNEFATHER, J.N., LAU, W.A.K., MA, S. & PATAK, E. (2000). Muscarinic receptors mediating contraction of the urinary bladder from the female mouse. *Proc. Aust. Soc. Clin. Exp. Pharmacol. Toxicol.*, **7**, 1–12P.
- PARKER, R.B. & WAUD, D.R. (1971). Pharmacological estimation of drug-receptor dissociation constants. Statistical evaluation. I. Agonists. *J. Pharmacol. Exp. Ther.*, **177**, 1–12.
- SOMOGYI, G.T. & DE GROAT, W.C. (1992). Evidence for inhibitory nicotinic and facilitatory muscarinic receptors on cholinergic nerve terminals of the rat urinary bladder. *J. Auton. Nerv. Syst.*, **37**, 89.
- STENGEL, P.W., GOMEZA, J., WESS, J. & COHEN, M.L. (2000). M_2 and M_4 receptor knockout mice: muscarinic receptor function in cardiac and smooth muscle *in vitro*. *J. Pharmacol. Exp. Ther.*, **292**, 877–885.
- TOBIN, G. (1995). Muscarinic receptor subtypes in the submandibular gland and the urinary bladder of the rabbit: *in vivo* and *in vitro* functional comparisons of receptor antagonists. *J. Auton. Pharmacol.*, **15**, 451–463.
- TOBIN, G. & SJÖGREN, C. (1995). *In vivo* and *in vitro* effects of muscarinic receptor antagonists on contractions and release of [3 H]-acetylcholine in the rabbit urinary bladder. *Eur. J. Pharmacol.*, **281**, 1–8.
- WELSH, N.J., EGLIN, R.M. & SHANKLEY, N. (2000). Pharmacological comparison of the muscarinic receptors mediating contraction of the guinea-pig left atrium, gastric smooth muscle and mouse urinary bladder. *Br. J. Pharmacol.*, **131**, 57P.

(Received February 19, 2001
Revised May 11, 2001
Accepted May 11, 2001)