

Note

Synthesis and Prior Misidentification of 4-*tert*-butyl-2,6-dinitrobenzaldehyde

Daniel J. Martin, Brandon Q. Mercado, and James M. Mayer

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.9b01737 • Publication Date (Web): 21 Aug 2019

Downloaded from pubs.acs.org on August 23, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Synthesis and Prior Misidentification of 4-*tert*-butyl-2,6-dinitrobenzaldehyde

Daniel J. Martin,* Brandon Q. Mercado, and James M. Mayer*


Department of Chemistry, Yale University, New Haven, CT 06520-8107, USA

*Correspondence to: daniel.j.martin@yale.edu, james.mayer@yale.edu

Abstract. Substituted 2,6-dinitrobenzaldehydes are valuable synthetic precursors and have been prepared by several methods. We report here that one reported synthetic method actually forms the 3,5-dinitro isomer, 4-*tert*-butyl-3,5-dinitrobenzaldehyde, instead of the claimed 2,6-isomer, 4-*tert*-butyl-2,6-dinitrobenzaldehyde. Improved syntheses for the large-scale preparation of both compounds and their single-crystal x-ray structures are described.

Substituted 2,6-dinitrobenzaldehydes are useful reagents for the preparation of dyes, pigments and macrocycles such as porphyrins and corroles.¹⁻⁶ In particular, 2,6-dinitrobenzaldehydes are used in the synthesis of tetrakis-5,10,15,20-(2',6'-dinitrophenyl)porphyrins, precursor molecules for bis-picket fence porphyrin derivatives.⁴ The acid-catalyzed condensation of 2,6-dinitrobenzaldehyde or related derivatives with pyrrole produces substituted 2',6'-dinitrophenylporphyrins, but typically only in low yields, ranging from 0.5% to 9%. The highest yields are for the tetrakis-5,10,15,20-(4'-*tert*-butyl-2',6'-dinitrophenyl)porphyrin, a common derivative.³ In order to make reasonable amounts of this porphyrin target (>100 mg), its synthesis requires sizable amounts of 4-*tert*-butyl-2,6-dinitrobenzaldehyde, the starting benzaldehyde precursor.

Scheme 1. Preparations of 4-*tert*-butyl-2,6-dinitrobenzaldehyde and 4-*tert*-butyl-3,5-dinitrobenzaldehyde, showing incorrect prior report (red), corrected reaction product (blue), previously available route (black), and new—chromium and column-free—procedures (gold).



Method A: reported synthesis in reference 6; actually yields **2** (this work). Method B: high yield synthesis of **2**, this work. Method C: previously reported synthesis of (**1**) requiring chromium reagents and chromatography purification, reference 3. Method D: high yield, high throughput synthesis of (**1**), this work.

The synthesis of 4-*tert*-butyl-2,6-dinitrobenzaldehyde (**1**) has been reported using two different routes, Methods A and C in Scheme 1.^{3,6} Method A reported what seemed to be the most direct synthesis, a one-pot nitration and oxidation of 4-*tert*-butylbenzyl alcohol.⁶ This synthesis was stated to be “noteworthy since it occurs with concomitant chemoselective oxidation of the benzylic alcohol to the corresponding aldehyde and represents the by far most practical route” to this common precursor.⁶ Following this preparation, we obtained a product in similar yields, which had ¹H NMR and ¹³C NMR spectra identical to the reported spectra (Figures S1-S2). Slow evaporation of this product from ethyl acetate yielded crystals suitable for single-crystal x-ray diffraction. The single-crystal data revealed that the product of this reaction was not **1** but was instead the isomeric 3,5-dinitro compound, 4-*tert*-butyl-3,5-dinitrobenzaldehyde (**2**) (Figure 1, Right; Scheme 1, Method A). Presumably the hot HNO₃/H₂SO₄ oxidation of the benzyl alcohol proceeds by initial formation of 4-*tert*-butylbenzaldehyde, which directs nitration to positions *meta* to the aldehyde. Consistent with this possible

intermediate formation, we report the synthesis of the 3,5-dinitro isomer, product **2**, in much higher yields (>95%) from the nitration of 4-*tert*-butylbenzaldehyde (Scheme 1, Method B; procedure in the Experimental Section). After a literature search of articles citing reference 6, we believe that this incorrect assignment may have led to a synthesis and misidentification of what is actually tetrakis-5,10,15,20-(4'-*tert*-butyl-3',5'-dinitrophenyl)porphyrin.⁷

An earlier paper by Rose *et al.* reported a three-step preparation of the target 2,6-dinitro isomer (**1**) from 4-*tert*-butyltoluene (Scheme 1, Method C).³ Following this procedure, including chromatography and recrystallization, we obtained a product with ¹H NMR and ¹³C NMR spectra (Figures S3-S4) that were identical to those reported in reference 3. Crystals were obtained from ethyl acetate/ethanol mixtures and single-crystal x-ray diffraction showed that the product of this reaction was correctly assigned as 4-*tert*-butyl-2,6-dinitrobenzaldehyde (**1**) (Figure 1, Left). The ¹H NMR and ¹³C NMR spectra for **1** and **2** are quite distinct, as are the solid-state structures.




Figure 1. Thermal ellipsoid plots (50% probability level) of X-ray structures for 4-*tert*-butyl-2,6-dinitrobenzaldehyde and 4-*tert*-butyl-3,5-dinitrobenzaldehyde. The unit cell of **1** contained two chemically identical, crystallographically distinct molecules of **1** ($Z' = 2$); only one molecule is shown for clarity.

1 It is unfortunate that the apparently scalable Method A does not produce the desired compound **1**. While
2 Method C is suitable for the synthesis of **1** in small quantities, it does not scale well to larger amounts. After
3 three steps, the overall yield of **1** is only 41% from 4-*tert*-butyltoluene, and the synthesis requires stoichiometric
4 amounts of chromium reagents and multiple chromatography columns. We are pleased to report an improved
5 route to **1**, starting from inexpensive 4-*tert*-butyltoluene, that can be done on much larger scales and without the
6 need for chromatography or chromium reagents.
7
8
9
10
11
12
13

14 Our synthesis of **1** (Method D, Scheme 1) was adapted from the reported syntheses of other ortho-
15 substituted aldehydes, including 2,6-dinitrobenzaldehyde.^{8,9} First, 4-*tert*-butyl-2,6-dinitrotoluene was prepared
16 by nitration of 4-*tert*-butyltoluene. The synthesis of this compound has been reported in several articles^{3,6,10,11}
17 and we have found no reports of hazards such as being explosive or forming explosive species. Bromination
18 using *N*-bromosuccinimide yielded 4-*tert*-butyl-2,6-dinitrobenzylbromide. Heating 4-*tert*-butyl-2,6-
19 dinitrobenzylbromide with pyridine in ethanol resulted in the precipitation of 4-*tert*-butyl-2,6-dinitro-
20 benzylpyridinium bromide as a white, crystalline solid. Base-catalyzed condensation of the pyridinium salt with
21 *N,N'*-dimethyl-4-nitrosoaniline generated the *N*-(4-dimethylaminophenyl)- α -(4-*tert*-butyl-2,6-
22 dinitrophenyl)nitro, which precipitated as an orange solid from cold ethanol/water. Acid-catalyzed hydrolysis
23 of the nitrone precipitated **1** as a yellow solid from acidic water. The ¹H NMR and ¹³C NMR for all the
24 intermediates are reported in the supporting information (Figures S5-S12). While this synthesis has more steps,
25 it gives **1** in 65% overall yield (starting from one mole of the parent toluene) and requires only recrystallization
26 and solubility differences for purification (Method D, Scheme 1). An alternative synthesis via direct oxidation
27 of the benzyl bromide intermediate using DMSO/Et₃N was briefly explored. While this gave the target product
28 in fewer steps than the method detailed here, it was not pursued because it required chromatography to separate
29 the unreacted benzyl bromide and was therefore less attractive at large scale.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 In sum, we report here that a published method⁶ yields 4-*tert*-butyl-3,5-dinitrobenzaldehyde (**2**) rather
52 than the claimed 2,6-dinitro isomer, 4-*tert*-butyl-2,6-dinitrobenzaldehyde (**1**). Since the 2,6-isomer is a valuable
53
54
55

synthetic intermediate, we have developed an improved synthesis without the need for chromatography or chromium reagents. This synthesis has been scaled to yield more than 150 grams of **1** with an overall yield of 65%. We also report a convenient synthesis of **2**. The ^1H NMR and ^{13}C NMR spectra and the x-ray crystal structures of both isomers are reported.

Experimental Section

Instrumentation. ^1H NMR spectra were recorded on an Agilent 500 MHz spectrometer and were referenced to proteo solvent impurities. High resolution mass spectrometry was performed using a Waters Xevo G2-XS QToF mass spectrometer. Gas-chromatography mass spectroscopy was performed using an Agilent 6890N/5973 mass spectrometer. Elemental analyses were performed by Robertson Microlit Laboratories, Ledgewood, NJ.

Materials. Potassium nitrate (Sigma,-Aldrich, *ReagentPlus*®, >99%), sulfuric acid (J.T. Baker, 98%), 4-*tert*-butylbenzylalcohol (Sigma-Aldrich, 95%), 4-*tert*-butylbenzaldehyde (Sigma-Aldrich, 97%), 4-*tert*-butyltoluene (Sigma-Aldrich, 95%), *N*-bromosuccinimide (Sigma-Aldrich, *ReagentPlus*®, 99%), 2,2'-azobis(2-methylpropionitrile (Sigma-Aldrich, 98%), pyridine (Sigma-Aldrich, 99.8%), *N,N*-dimethyl-4-nitrosoaniline (Sigma-Aldrich, 97%), aqueous hydrochloric acid (Macron, 36-38% wt), sodium chloride (Sigma-Aldrich, >99%), sodium bicarbonate (Sigma-Aldrich, >99%), sodium hydroxide (Macon, 95%), chloroform (Sigma-Aldrich, 99.8%), ethyl acetate (Sigma-Aldrich, 99.8%), absolute ethanol (Decon Labs, 200 proof), diethyl-ether (Sigma-Aldrich, 99%), and acetone (Sigma Aldrich, 99.8%) were all used as received. Dichloromethane was degassed with argon and dried using a Pure Process Technology solvent system prior to use.

Single Crystal X-ray diffraction methods. Low-temperature diffraction data (ω -scans) were collected on a Rigaku SCX Mini diffractometer coupled to a Rigaku Mercury275R CCD with Mo $K\alpha$ radiation ($\lambda = 0.71073$ Å). The diffraction images were processed and scaled using Rigaku Oxford Diffraction software (CrysAlisPro; Rigaku OD: The Woodlands, TX, 2015). The structures were solved with SHELXT and were refined against F^2

1 on all data by full-matrix least squares with SHELXL.¹ All non-hydrogen atoms were refined anisotropically.
2 Hydrogen atoms were included in both models at geometrically calculated positions and refined using a riding
3 model. The isotropic displacement parameters of all hydrogen atoms were fixed to 1.2 times the U value of the
4 atoms to which they are linked (1.5 times for methyl groups). For **2**, the dataset was refined as a 2-component
5 twin. The fractional volume contribution of the minor twin component was freely refined to a converged value
6 of 0.3850(19). For **1**, the dataset yielded a unit cell containing two chemically identical, crystallographically
7 distinct molecules.
8
9
10
11
12
13
14
15

16 The full numbering scheme of compounds **1** and **2** can be found in the full details of the X-ray structure
17 determinations (CIFs), which are included as Supporting Information. CCDC number 1923128 (**1**) and 1923129
18 (**2**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge
19 from The Cambridge Crystallographic Data Center via www.ccdc.cam.ac.uk/data_request/cif.
20
21
22
23
24

25 **Mixed acid preparation for nitration reactions. Caution:** *Fuming nitric acid* (FNA) is a corrosive
26 strong acid and a strong oxidant; concentrated sulfuric acid is a corrosive strong acid; and the mixture of the
27 two is considered a stronger acid and oxidant. All of these materials are highly hazardous and should be handled
28 with substantial caution with emphasis on appropriate specialized personal protective equipment (PPE). Our
29 laboratory's standard operating procedure (SOP) for FNA and the mixed acid are included as Supporting
30 Information. Fuming nitric acid (FNA, +98%) was freshly prepared immediately before use via distillation of
31 HNO₃ at 80 °C from a mixture of potassium nitrate and concentrated sulfuric acid, prepared as described in the
32 SOP. FNA (1 part by volume, for instance 50 mL) was cooled to 5 °C in an ice bath before slowly adding
33 concentrated sulfuric acid (1.4 parts by volume, for instance 70 mL). Upon addition, the mixed acid warms to
34 15 °C. The mixed acid solution was then cooled back to 5 °C before being used immediately for nitration
35 reactions.
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **Improved synthesis of 4-tert-butyl-3,5-dinitrobenzaldehyde (2).** Approximately 120 mL of the mixed
51 H₂SO₄/HNO₃ acid (see preparation above) was prepared in a 500 mL round bottom flask with a large stirbar and
52
53
54
55

1 was cooled to 5 °C with an ice bath [**Caution:** hazardous! See above and SOP in Supporting Information].
2 Above the round bottom was suspended a dropping funnel filled with 4-*tert*-butylbenzaldehyde (25 mL, 0.149
3 mmol). The benzaldehyde was added dropwise to the stirring solution and the flow rate was adjusted such that
4 the reaction temperature did not rise above 15 °C. After complete addition, the solution was yellow/orange and
5 the reaction flask was allowed to warm to room temperature before being gently heated to 40 °C for 1-2h.
6 During this time, some off-white precipitate formed. The reaction was then quenched by *carefully* pouring the
7 entire reaction into a large amount of crushed ice (ca. 500 mL). The precipitated product was filtered off and
8 washed with excess water. The dilute, acidic filtrate was neutralized and added to the aqueous waste. After air
9 drying, the crude yellow solid (crude yield by weight: 36g, 97%) was dissolved in chloroform and filtered
10 through Celite to remove any over-oxidized 4-*tert*-butyl-2,6-dinitrobenzoic acid (the white solid formed during
11 synthesis). The filtrate was collected and removed with a rotary evaporator to dryness. The collected solid was
12 then recrystallized from boiling ethyl acetate to yield pale yellow crystals (Figure S18). After filtration and air
13 drying, the isolated yield was 29 g (80% yield). ¹H NMR (CDCl₃, 500 MHz): δ (ppm) = 9.99 (s, 1H, CHO),
14 7.99 (s, 2H, Ar-H), 1.49 (s, 9H, C(CH₃)₃). ¹³C NMR (CDCl₃, 500 MHz): δ (ppm) = 187.0, 153.7, 141.1, 135.3,
15 126.6, 38.2, 30.0. Anal. Calcd for C₁₁H₁₂N₂O₅: C, 52.38; H, 4.80; N, 11.11. Found: C, 52.13; H, 4.65; N, 10.95.
16 MS (EI) *m/z* = 252.1 ([M]⁺), 237.1 ([M]⁺ - CH₃).

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37 **4-*tert*-butyl-2,6-dinitrotoluene.** This synthesis was adapted from the literature.⁴ Approximately 240 mL
38 of the mixed H₂SO₄/HNO₃ acid (see preparation above) was prepared in a 1L round bottom flask with a large
39 stirbar and was cooled to 5 °C with an ice bath [**Caution:** hazardous! See above and SOP in Supporting
40 Information]. Above the round bottom was suspended a dropping funnel filled with 4-*tert*-butyltoluene (173
41 mL, 1.0 mol). The 4-*tert*-butyltoluene was added dropwise to the stirring solution over the course of three hours
42 such that the temperature of the reaction stayed between 5-10 °C. The yellow product formed and floated to the
43 reaction surface during the addition. After complete addition, the reaction flask was allowed to warm to room
44 temperature and was stirred for another 12h. The reaction was quenched by *carefully* pouring the reaction
45
46
47
48
49
50
51
52
53
54
55

1 mixture into 500 mL of crushed ice. After filtration, the collected yellow solid was washed with excess water.
2
3 The dilute, acidic filtrate was neutralized and added to the aqueous waste. The damp solid was then
4
5 recrystallized from boiling EtOH (approx. 750 mL). Upon cooling, large yellow crystals formed (Figure S13).
6
7 The solution was cooled at 0 °C overnight before the crystals were filtered and washed with cold EtOH. After
8
9 drying, the isolated yield was 190 g flaky crystalline solid (80% yield). ¹H NMR (CDCl₃, 500 MHz): δ (ppm) =
10
11 7.97 (s, 2H, Ar-*H*), 2.52 (s, 3H, CH₃), 1.37 (s, 9H, C(CH₃)₃). ¹³C NMR (CDCl₃, 500 MHz): δ (ppm) = 152.2,
12
13 151.6, 124.7, 124.1, 35.3, 30.8, 14.5. Anal. Calcd for C₁₁H₁₄N₂O₄: C, 55.46; H, 5.92; N, 11.76. Found: C, 55.42;
14
15 H, 5.78; N, 11.68. MS (EI): *m/z* = 238.1 ([M]⁺), 223.1 ([M]⁺ - CH₃).

16
17
18 ***4-tert-butyl-2,6-dinitrobenzylbromide.*** To a 3L three-necked round bottom flask was added a large
19
20 stirbar, 4-*tert*-butyl-2,6-dinitrotoluene (190 g, 0.798 mol, 1 eq.), *N*-bromosuccinimide (170 g, 0.998 mol, 1.25
21
22 eq.), and AIBN (1.5 g, 0.009 mol, 0.01 eq). Two of the three ports were stoppered and 1.5 L of dry
23
24 dichloromethane was added to the solids. The solution was sparged with N₂ for 20 minutes before a condenser
25
26 was attached to the third port and the reaction was heated to a gentle reflux (45-50 °C). A white LED light (14
27
28 V dc, 0.7 A from Digi-Key) was clamped a few inches away from the flask and the reaction was loosely
29
30 shrouded in aluminum foil. At the 12h, 48h, 74h, and 82h marks, additional NBS (20 g, 0.117 mol, 0.15 eq.) and
31
32 AIBN (0.5 g, 0.0045 mol, 0.005 eq.) were added. The solid additions were made by funneling the NBS/AIBN
33
34 through an opened port under gently flowing N₂ only after cooling the reaction to room temperature. After four
35
36 days, the reaction was nearly complete (>98% conversion based on the ratio of product to starting material by
37
38 ¹H NMR). The solvent was removed using a rotary evaporator to yield a damp solid. The solid was resuspended
39
40 in chloroform and filtered to remove the bulk of the succinimide. The filtrate was separated with sat. Na₂CO₃
41
42 solution to remove residual succinimide before being dried with MgSO₄ and filtered. The solvent was removed
43
44 and the product was recrystallized from hot diethyl ether to yield extremely large pale-yellow crystals (5-7
45
46 grams apiece, Figure S14). Isolated yield was 235.5 g large yellow crystals and 13.2 g non-crystalline powder
47
48
49 with matching ¹H NMR spectra (97.7%). ¹H NMR (CDCl₃, 500 MHz): δ (ppm) = 8.04 (s, 2H, Ar-*H*), 4.84 (s,
50
51
52
53
54
55

2H, CH_2Br), 1.37 (s, 9H, $C(CH_3)_3$). ^{13}C NMR ($CDCl_3$, 500 MHz): δ (ppm) = 155.3, 150.1, 125.9, 123.6, 35.7, 30.7, 20.4. Anal. Calcd for $C_{11}H_{13}BrN_2O_4$: C, 41.7; H, 4.1; N, 8.8. Found: C, 41.69; H, 4.01; N, 8.71. MS (EI): m/z = 301.0 ($[M]^+ - CH_3$), 237.1 ($[M]^+ - Br$), parent ion not observed.

4-tert-butyl-2,6-dinitrobenzylpyridinium bromide. The crude 4-tert-butyl-2,6-dinitrobenzylbromide (246 g, 0.778 mol, 1 eq.) was added to a 2 L beaker. The solid was dissolved in 1 L EtOH (200 proof) and was heated to 78°C with stirring. Pyridine (130 mL, 1.55 mol, 2 eq.) was added dropwise to the stirring solution, which darkened and eventually precipitated flocculent, white crystals (Figure S15). After complete addition, the solution was stirred for an additional hour at reflux before an additional hour of stirring at room temperature. The crystals were filtered and washed with EtOH and acetone. After drying, the first crop of crystals weighed 258 g. The filtrate was pumped to a solid using a rotary evaporator and was repeatedly recrystallized from EtOH to yield an additional 25 g of off-white solid with 1H NMR that matched the crystalline material (total solid collected: 283.0 g, 91.3% yield. 1H NMR (d_6 -DMSO, 500 MHz): δ (ppm) = 9.06 (d, J = 6.0 Hz, 2H, 2,6-py- H), 8.72 (t, J = 7.8 Hz, 1H, 4-py- H), 8.53 (s, 2H, Ar- H), 8.21 (t, J = 7.1 Hz, 2H, 3,5-py- H), 6.14 (s, 2H, Ar- CH_2 -py), 1.41 (s, 9H, $C(CH_3)_3$). ^{13}C NMR (d_6 -DMSO, 500 MHz): δ (ppm) = 157.9, 152.0, 146.7, 144.5, 128.5, 127.6, 116.5, 56.0, 36.2, 30.5. Anal. Calcd for $C_{16}H_{18}N_3O_4Br$: C, 48.50; H, 4.58; N, 10.60. Found: C, 48.31; H, 4.32; N, 10.38. HRMS (ESI/Q-TOF) m/z : $[M - Br]^+$ calcd for $C_{16}H_{18}N_3O_4^+$ 316.1297; found 316.1296.

***N*-(4-dimethylaminophenyl)- α -(4-tert-butyl-2,6-dinitrophenyl)nitron.** To a large Erlenmeyer flask was added approximately 1/3 of the dry 4-tert-butyl-2,6-dinitrobenzylpyridinium bromide (85.4 g, 0.215 mol, 1 eq.) and *N,N*-dimethyl-4-nitrosoaniline (40.0 g, 0.266 mol, 1.24 eq). The solids were suspended in 1.0 L of ethanol (200 proof) and the solution was cooled to 5 °C with an ice bath. An addition funnel was suspended above the reaction and was filled with 600 mL of 1 *N*-sodium hydroxide solution. With vigorous stirring, the NaOH solution was added dropwise such that the temperature was kept between 5-10 °C. After complete addition, the solution was stirred for another 2h before being diluted with 1 L ice water. The nitron precipitated as an orange solid (Figure S16), which was filtered and washed with excess water. This procedure was repeated until nearly

all of the benzylpyridinium bromide was used (279.3 g, 0.714 mol). All of the orange solids were collected and dried in a vacuum oven at 50°C until reaching a constant mass (2-3 days). After drying, the yield was 257.7 g of orange powder (92.4% yield). ¹H NMR (CDCl₃, 500 MHz): δ (ppm) = 8.44 (s, 1H, Ar-CH=N), 8.30 (s, 2H, Ar-H), 7.67 (d, *J* = 9.1 Hz, 2H), 6.67 (d, *J* = 9.1 Hz, 2H), 3.04 (s, 6H, N(CH₃)₂), 1.42 (s, 9H, C(CH₃)₃). ¹³C NMR (CDCl₃, 500 MHz): δ (ppm) = 155.3, 152.1, 149.6, 136.8, 126.1, 122.8, 122.5, 117.8, 111.4, 40.5, 35.9, 30.9. Anal. Calcd for C₁₉H₂₂N₄O₅: C, 59.06; H, 5.74; N, 14.50. Found: C, 58.92; H, 5.56; N, 14.34. HRMS (ESI/Q-TOF) *m/z*: [M + H⁺] calcd for C₁₉H₂₃N₄O₅⁺ 387.1668; found 387.1682.

4-tert-butyl-2,6-dinitrobenzaldehyde (1). In a large beaker, nearly all of the *N*-(4-dimethylaminophenyl)- α -(4-*tert*-butyl-2,6-dinitrophenyl)nitron (257.2 g) was suspended in 2L of water. Concentrated sulfuric acid was added slowly and the solution warmed to approximately 50 °C. The suspension was stirred for 30 minutes, during which the solution color lightened to a pale yellow-brown. The suspension was filtered and the collected solid was washed with excess water. After drying, 164 g of pale-yellow powder was collected (98% yield). The solid can be recrystallized from hot ethanol or from mixtures of ethanol/ethyl acetate to yield golden flake-like crystals (Figure S18). ¹H NMR (CDCl₃, 500 MHz): δ (ppm) = 10.57 (s, 1H, CHO), 8.45 (s, 2H, Ar-H), 1.45 (s, 9H, C(CH₃)₃). ¹³C NMR (CDCl₃, 500 MHz): δ (ppm) = 186.2, 157.1, 147.7, 129.0, 126.8, 36.1, 30.9. Anal. Calcd for C₁₁H₁₂N₂O₅: C, 52.38; H, 4.80; N, 11.11. Found: C, 52.13; H, 4.61; N, 10.94. MS (EI) *m/z* = 252.1 ([M]⁺), 237.1 ([M]⁺ - CH₃).

Supporting Information Available

i) Copies of ¹H NMR and ¹³C NMR spectra. (ii) Crystallographic information files (CIF) and data for compounds **1** and **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>. (iii) Standard Operating Procedures for working with concentrated nitric and sulfuric acids, with emphasis on safety precautions.”

Acknowledgements

This research was supported as part of the Center for Molecular Electrocatalysis, and Energy Frontier Research Center funded by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences. D.J.M. gratefully acknowledges support by a Graduate Research Fellowship from the National Science Foundation. We thank Dr. Fabian Menges (Yale Chemical and Biophysical Instrumentation Center) for assistance with mass spectroscopy and recognize support from the National Science Foundation grant 1828190 for support of mass spectrometry facilities.

References

- (1) Rose, E.; Soleilhavoup, M.; Christ-Tommasino, L.; Moreau, G.; Collman, J. P.; Quekquejeu, M.; Straumanis, A. Bis-Faced Aminoporphyrin Templates for the Synthesis of Chiral Catalysts and Hemeprotein Analogues. *J. Org. Chem.* **1998**, *63*, 2042–2044.
- (2) Quintana, C. A.; Assink, R. A.; Shelnut, J. A. Synthesis and Spectroscopic Characterization of Bis-Pocket Porphyrins: Tetrakis(2',6'-dinitrophenyl)porphyrin and Catalytic Activity of a Manganese(III) Chloride Derivative in Alkane Oxidation. *Inorg. Chem.* **1989**, *28*, 3421–3425.
- (3) Rose, E.; Kossanyi, A.; Quekquejeu, M.; Soleilhavoup, M.; Duwavran, F.; Bernard, N.; Lecas, A. Synthesis of Biomimetic Heme Precursors: The “Double Picket Fence” 5,10,15,20-Tetrakis(2',6'-dinitro-4'-*tert*-butylphenyl)porphyrin. *J. Am. Chem. Soc.* **1996**, *118*, 1567–1568.
- (4) Rose, E.; Lecas, A.; Quekquejeu, M.; Kossanyi, A.; Boitrel, B. Synthesis of Biomimetic Heme Precursors. *Coord. Chem. Rev.* **1998**, *178–180*, 1407–1431.

- 1 (5) Drain, C. M.; Corden, B. B. Synthesis and Characterization of Copper(II) 5,10,15,20-Tetrakis(2,6-
2 dipivalamidophenyl)porphyrinate: A Bis-Picket-Fence Porphyrin. *Inorg. Chem.* **1989**, *28*, 4374–4376.
3
4 (6) Peters, M. V.; Stoll, R. S.; Goddard, R.; Buth, G.; Hecht, S. On the Illusive Nature of O-
5 Formylazobenzenes: Exploiting the Nucleophilicity of the Azo Group for Cyclization to Indazole
6 Derivatives. *J. Org. Chem.* **2006**, *71*, 7840–7845.
7
8 (7) Goodrich, L. E.; Roy, S.; Alp, E. E.; Zhao, J.; Hu, M. Y.; Lehnert, N. Electronic Structure and
9 Biologically Relevant Reactivity of Low-Spin {FeNO}⁸ Porphyrin Model Complexes: New Insight from
10 a Bis-Picket Fence Porphyrin. *Inorg. Chem.* **2013**, *52*, 7766–7780.
11
12 (8) Kalir, A. *o*-Nitrobenzaldehyde. *Org. Synth.* **1966**, *46*, 81–84.
13
14 (9) Clarke, K. Some Aromatic Aldehydes. *J. Chem. Soc.* **1957**, *0*, 3798–3808.
15
16 (10) Brady, O. L.; Lahiri, J. K. 430. The Nitration of Alkyl Benzenes. Part III. The Orientation of Dinitro-*p*-
17 *tert*-butyltoluene. *J. Chem. Soc.* **1934**, No. 0, 1954–1957.
18
19 (11) Yamato, T.; Arimura, T.; Tashiro, M. Metacyclophanes and Related Compounds. Part 16. Preparation of
20 8-Fluoro-*t*-Butyl[2.2]Metacyclophanes and Their Treatment with Aluminium Chloride–Nitromethane in
21 Benzene. *J. Chem. Soc., Perkin Trans. 1* **1987**, No. 0, 1–7.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 For Table of Contents Only
4

