

Communication

Para-Selective C-H Borylation of Common Arene Building Blocks Enabled by Ion-Pairing with a Bulky Counteranion

Madalina T. Mihai, Benjamin D. Williams, and Robert J Phipps

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.9b07267 • Publication Date (Web): 05 Aug 2019

Downloaded from pubs.acs.org on August 5, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Para-Selective C-H Borylation of Common Arene Building Blocks Enabled by Ion-Pairing with a Bulky Counteranion

Madalina T. Mihai, Benjamin D. Williams and Robert J. Phipps*

Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, United Kingdom.

Supporting Information Placeholder

ABSTRACT: The selective functionalization of C-H bonds at the arene *para* position is highly challenging using transition metal catalysis. Iridium-catalyzed borylation has emerged as a leading technique for arene functionalization, but there are only a handful of strategies for *para*-selective borylation, which operate on specific substrate classes and use bespoke ligands or catalysts. We describe a remarkably general protocol which results in *para*-selectivity on some of the most common arene building blocks (anilines, benzylamines, phenols, benzyl alcohols) and uses standard borylation ligands. Our strategy hinges upon the facile conversion of the substrates into sulfate or sulfamate salts, wherein the anionic arene component is paired with a tetrabutylammonium cation. We hypothesize that the bulk of this cation disfavors *meta*-C-H borylation, thereby promoting the challenging *para*-selective reaction.


Use of transition metals to catalytically functionalize arene C-H bonds has undergone tremendous development. Whilst most advances have dealt with *ortho*-selective functionalization, the last decade has seen many advances in *meta*-selective reactions.¹ Many of these approaches have been based on templates or ligands with extended architectures, or on strategies which proceed *via* metalation at the *ortho* position. Considering this, it is not surprising that methods to direct transition metals to the *para* position remain rather fewer, given the greater distance from existing functionality.² Although *S_EAr* mechanisms give the *para* isomer when electrophiles react with electron rich arenes, it is uncommon for transition metals to perform C-H bond functionalization via *S_EAr* mechanisms.³ Exceptions are several examples using gold catalysis⁴ and a smaller number of examples using high-valent copper⁵ or palladium⁶. Yet such selectively metallated intermediates are extremely versatile and new methods to access them are urgently required.^{7,8}

Iridium-catalyzed borylation has emerged to become one of the most versatile and intensively used of modern arene functionalization methods.^{9,10} Whilst regioselectivity is typically determined by steric considerations, numerous modifications can direct borylation to the *ortho* position¹¹ and, to a lesser extent, the *meta* position.¹² In contrast, there are only three instances where *para*-selective iridium catalyzed C-H borylation has been achieved (Figure 1a).^{13,14} Saito, Segawa and Itami used a bulky, chiral phosphine ligand to create a highly sterically congested environment at iridium.¹⁵ However, high *para*-selectivity was limited to substrates possessing a very bulky substituent, and 1,2-disubstituted arenes were unselective. Nakao and co-workers used a bulky Lewis acid catalyst to complex aromatic amides, both activating the substrate and forcing borylation to occur at the most



remote position due to steric interactions.¹⁶ Most recently, Chattopadhyay and co-workers disclosed an 'L-shaped' bipyridine ligand proposed to direct borylation to the *para* position by virtue of interaction between the potassium salt of the ligand and an ester in the substrate.¹⁷

Despite the inventiveness of these approaches, two aspects limit wider application. Firstly, they are restricted to relatively narrow substrate classes. Secondly, they require bespoke or non-commercially available ligands or catalysts, presenting a barrier to those looking to use "off-the-shelf" reagents, such as end-users in the pharmaceutical industry. At the outset of this work, we sought to develop a general and practical strategy that would enable *para*-selective borylation of a range of common substrate classes using readily available catalysts and ligands. We have previously utilized attractive ion-pairing interactions between catalyst and substrate to direct the iridium metal to the arene *meta* position (Figure 1b).^{12c, 12f, 12h} We were concerned that adopting a similar strategy to target the *para* isomer would be very challenging due to the distance that the ligand would be required to reach. A design able to achieve this would likely have a complex, extended structure necessitating lengthy synthesis and reduced practical utility. We envisaged an alternative ion-pairing strategy in which the counterion of the substrate does not deliver the reactive catalyst, but is unfunctionalised and bulky, acting as a 'steric shield' to obstruct borylation at the *meta* position, thereby resulting in *para* selectivity with standard 'off-the-shelf' borylation catalysts (Figure 1c).

We envisaged that the ubiquitous tetrabutylammonium cation may constitute an ideal 'steric shield'; the alkyl chains project outwards at all angles from the tetrahedral nitrogen, occupying a large area. This bulky cation could be paired with a variety of common arene building blocks (anilines, benzylamines, phenols, benzyl alcohols) which can all be rendered temporarily anionic in a single simple step through conversion to the corresponding sulfate (X=O) or sulfamate (X=N) salts (Figure 1d). Herein, we demonstrate the realization of this approach, which we believe represents the most general strategy for *para*-selective borylation developed to date and indeed provides complementarity to existing strategies. In contrast to our earlier work, the key ion-pairing interaction in this system is not being exploited in an attractive sense between substrate and catalyst, but rather to temporarily anchor the 'steric shield' to the substrate. The resulting repulsive steric interactions with the incoming catalyst thereby guide the latter to the remote position that would be the most challenging to reach using a strategy invoking attractive catalyst-substrate interactions.

a) Previous strategies for *para*-selective arene borylation using Ir catalysis:

b) Our previous work: Meta-selective borylation using "attractive" Substrate-Catalyst ion-pairing interaction

c) This work: attractive Substrate-Counterion interaction anchors 'steric shield', forcing catalyst *para*

d) This work:





Figure 1. Previous *para*-selective Ir-catalyzed borylation approaches and that taken in this work.

We began by converting 2-chloroaniline (**1a**) to the corresponding tetrabutylammonium sulfamate salt **2a** by treatment with chlorosulfonic acid followed by cation exchange using $\text{Bu}_4\text{NH}_4\text{SO}_4$ (Scheme 1a). To our delight, iridium-catalyzed borylation of **2a** under standard conditions with *dtbpy* as ligand resulted in >20:1 selectivity for the *para* isomer. Simply treating the crude reaction mixture with HCl in methanol revealed the parent aniline, which give 85% yield of *para*-borylated aniline **3** over two steps and only a single purification procedure. A similar outcome was achieved for the electronically different 2-trifluoromethylaniline (Scheme 1a, **1b** to **3b**). In contrast, direct borylation of parent anilines **1a** and **1b** results in *m:p* ratios of 1:3.9 and 1:1 respectively (Scheme 1b). The moderate *para* selectivity in direct borylation of **1a** echoes a similar observation made by Krska, Maleczka, Smith and co-workers, the origins of which are not obvious.¹⁸ Accordingly, we examined **4a** and **4b**, control substrates much closer in electronic nature to **2a** and **2b**, and found both of these give poor selectivity (Scheme 1c). This outcome suggests that the high *para* selectivity obtained with sulfamate salts **2a** and **2b** does not have electronic origins and that the salt formation and association with the tetrabutylammonium cation are crucial factors.

With these initial results in hand, we next examined a selection of 2-substituted anilines and were pleased to find that the high levels of *para* selectivity were observed for a range of useful building blocks (Scheme 2A). Bromide and iodide substituents were well tolerated (**3c**, **3d**), both giving >20:1 *para* selectivity. The latter is

notable given the incompatibility of many other C-H activation methods with iodide substituents. Arenes bearing trifluoromethoxy (**3e**) and difluoromethoxy (**3g**) substituents are also compatible, as

Scheme 1. a) Initial results. (b), (c) Control reactions.




is the difluorinated **3h**. Interestingly, the *para* selectivity was slightly reduced with a resonance withdrawing substituent (2-CN, **3f**), in-line with the precedent that this substituent has a small *para*-directing (relative to itself) electronic effect in borylation.¹⁹ Substrates bearing electron-donating substituents (isopropyl, methoxy) gave low conversion, albeit with excellent *para*-selectivity (see SI).

Given the ubiquity of benzylamines as aromatic building blocks, we next sought to test our strategy on benzylamine-derived sulfamates. As with the aniline variant, these tetrabutylammonium salts are easily synthesized from cheap materials. This class proved highly amenable, in most cases giving >20:1 regioselectivity (Scheme 2B). A range of useful halide substituents was tolerated, including Cl (**7a**), Br (**7b**), I (**7c**) and F (**7g**). Trifluoromethyl (**7d**), trifluoromethoxy (**7e**) and difluoromethoxy (**7f**) substituted variants also performed well. A methoxy-substituted variant gave excellent selectivity but with poor conversion (**7h**). It was necessary to derivatize the free amine products to enable purification on silica; isolated yields are following borylation/hydrolysis/trifluoroacetylation.

Encouraged by the effectiveness of the strategy thus far, we next extrapolated this to the ubiquitous oxygen-containing building blocks, phenols and benzyl alcohols.²⁰ The tetrabutylammonium sulfate salts of these compounds are also easily obtained and cleaved. Gratifyingly, high *para*-selectivity was observed for a number of *ortho*-substituted phenol-derived sulfate salts, encompassing Cl (**9a**), Br (**9b**), I (**9c**), CF₃ (**9d**), OCF₃ (**9e**), *i*Pr (**9f**) and a difluorinated analogue (**9h**) (Scheme 2C). Control borylations on 2-chlorophenol and on a 2-chlorophenol sulfate ester gave poor selectivity.²¹ Interestingly, selectivity (by crude ¹H-NMR analysis following borylation) was somewhat reduced compared with the aniline-derived sulfamates. Particularly, substrates possessing conjugatively withdrawing groups such as nitro (**9g**) exhibited only

Scheme 2. Scope of *para* C-H borylation of aniline (A), benzylamine (B), phenol (C) and benzyl alcohol (D) derivatives, and aryl and benzylsulfonates (E).^{a,b}



^a Typically: Substrate (0.25 mmol), B₂Pin₂ (0.25–0.5 mmol), [Ir(COD)OMe]₂ (1.5–2.5 mol%), dtbpy (3–5 mol%), 1,4-dioxane (0.33–0.5 M), 70 °C (See SI for full details). ^b Isomeric ratios are *para:meta* taken from analysis of crude ¹H-NMR spectra after borylation, unless otherwise stated. Yields shown are isolated and typically include *meta* and *para* isomers which are generally inseparable. ^c Selectivity assessed after HCl step. ^d Selectivity assessed after TFA protection step. ^e Tmpen used instead of dtbpy. ^f Oxidized to corresponding phenol for isolation.

moderate *para*-selectivity. This is likely due to the substituent electronic influence on the desired *para* selectivity (*vide supra*).

Remarkably, the extension to a fourth common substrate class, benzyl alcohols, was well tolerated (Scheme 2D, **11a–11h**). As in the phenol-derived sulfates, regioselectivity was adversely affected by resonance withdrawing groups (**11h**), but positively reinforced by resonance donors (**11f**). These results suggest that in substrate classes when the *para*-directing influence from the associated cation is weaker, substituents electronic effects can have greater impact on selectivity.


Finally, to further reinforce the generality and practicality of our strategy for the selective synthesis of complex, multifunctional building blocks, we targeted aryl sulfonates and benzyl sulfonates, substrates which inherently bear a negative charge (Scheme 2E). The former are readily accessed from commercially available aryl sulfonyl chlorides by treatment with tetrabutylammonium hydroxide, and the latter by the action of sodium sulfite on benzyl halides,

followed by cation exchange. Furthermore, through treatment with POCl₃, both sulfonates can be easily converted to sulfonyl chlorides, valuable precursors to sulfonamides. Both of these substrate classes result in *para*-selective borylation, in line with our hypothesis. Aryl sulfonates bearing *ortho* Cl (**13a**), Br (**13b**), CF₃ (**13c**) and OCF₃ (**13d**) gave excellent *para*-selectivity, as did the analogously-substituted benzyl sulfonates (**13e–13h**). Following borylation, the crude borylated sulfonates were either converted through to the corresponding piperidine sulfonamides (**13a–13d**) or were isolated as sulfonyl chlorides (**13e–13h**).

At the outset we advanced the tentative hypothesis that the bulk of the associated tetrabutylammonium cation may occupy sufficient space around the arene such that borylation at the normally accessible *meta* positions becomes disfavored (Figure 1C). To probe this hypothesis, we synthesized the 2-Cl member of the first four substrate classes with the smaller tetraethylammonium cation (synthesis of the tetramethyl variant for all members proved problematic)

to gauge the impact on selectivity (Scheme 3a). These experiments showed that for three of the four classes, *para* selectivity was significantly reduced, in line with our hypothesis (in the fourth, selectivity remained excellent at >20:1). We next selected the benzyl sulfonate substrate class to carry out a complete evaluation of sequential modulation of cation size on a single substrate. These experiments showed a clear and consistent trend to higher *para*-selectivity as the cation grows larger (Scheme 3b).

Scheme 3. Experiments to probe origin of selectivity.



Finally, in order to visualize the steric effect of the associated tetrabutylammonium cation, we were able to obtain X-ray structures of the starting materials for three substrate classes (Figure 2). These give a visual impression of the bulk provided by the cation and support our central tenet that it could feasibly act as a non-covalent 'shield' to block the arene *meta*-position. Accordingly, one might expect that removal of the *ortho* substituent incorporated into all substrates thus far may reduce *para*-selectivity as both *meta* positions would be open to borylation and the cation would not be able to block both at any one time. Indeed, this is the case; the *para* selectivity was drastically reduced for all unsubstituted variants providing further support for our hypothesis.²¹ We also extended the chain length of all three classes of substrate as the 2-Cl variant but found that with two methylenes between the arene and the anionic group, selectivity was reduced to around 3:1 *p:m*, presumably due to the much greater flexibility.²¹




Figure 2. X-Ray crystal structures of starting materials allowing visualization of the bulk of the tetrabutylammonium cation.

In conclusion, we have developed a general method for the *para*-selective C-H borylation of a broad range of the most useful arene building blocks encompassing anilines, benzylamines, phenols and benzyl alcohols as well as several classes of arenesulfonate, valuable precursors to sulfonamides. Our method produces a variety of complex trisubstituted arene cores typically with three functional handles for further elaboration that in many cases would require numerous steps to obtain by other methods. We provide evidence to suggest that this is due to ion-pairing of a bulky tetrabutylammonium cation with the substrate, which is rendered temporarily anionic. The bulky cation blocks the *meta* position, promoting borylation at the most remote and typically most challenging *para* position. Our approach provides a practical and general solution to the challenge of *para*-selective C-H borylation and demonstrates a new strategy for the utilization of ion-pairing interactions in catalysis.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures and spectral data is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*rjp71@cam.ac.uk

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENTS

We are grateful to AstraZeneca for a PhD studentship through the AZ-Cambridge PhD program (M.T.M.), the Cambridge Trust and Sidney Sussex College, Cambridge for a PhD studentship (B.D.W.), the Royal Society for a University Research Fellowship (R.J.P.), the EPSRC (EP/N005422/1) and the ERC (StG 757381). We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University. We thank Dr. Andrew Bond acquiring and solving the X-ray data. Thanks to Iain Cumming (AstraZeneca) for useful discussion.

REFERENCES

- (a) Li, J.; De Sarkar, S.; Ackermann, L., meta- and para-Selective C-H Functionalization by C-H Activation. In *C-H Bond Activation and Catalytic Functionalization I*, Dixneuf, P. H.; Doucet, H., Eds. Springer International Publishing: Cham, 2016; pp 217-257; (b) Dey, A.; Agasti, S.; Maiti, D., Palladium catalysed meta-C-H functionalization reactions. *Org. Biomol. Chem.* **2016**, *14*, 5440-5453; (c) Mihai, M. T.; Genov, G. R.; Phipps, R. J., Access to the meta position of arenes through transition metal catalysed C-H bond functionalisation: a focus on metals other than palladium. *Chem. Soc. Rev.* **2018**, *47*, 149-171.
- Dey, A.; Maity, S.; Maiti, D., Reaching the south: metal-catalyzed transformation of the aromatic para-position. *Chem. Commun.* **2016**, *52*, 12398-12414.
- For selected recent examples or para-selective functionalization via radical mechanisms, see: (a) Romero, N. A.; Margrey, K. A.; Tay, N. E.; Nicewicz, D. A., Site-selective arene C-H amination via photoredox catalysis. *Science* **2015**, *349*, 1326-1330; (b) Boursalian, G. B.; Ham, W. S.; Mazzotti, A. R.; Ritter, T., Charge-transfer-directed radical substitution enables para-selective C-H functionalization. *Nature Chemistry* **2016**, *8*, 810; (d) Berger, F.; Plutschack, M. B.; Riegger, J.; Yu, W.; Speicher, S.; Ho, M.; Frank, N.; Ritter, T., Site-selective and versatile aromatic C-H functionalization by thianthrene. *Nature* **2019**, *567*, 223-228.
- For selected examples, see: (a) Ball, L. T.; Lloyd-Jones, G. C.; Russell, C. A., Gold-Catalyzed Direct Arylation. *Science* **2012**, *337*, 1644-1648; (b) Brand, J. P.; Waser, J., Para-Selective Gold-Catalyzed Direct Alkynylation of Anilines. *Org. Lett.* **2012**, *14*, 744-747; (c) Cambeiro, X.

C.; Boorman, T. C.; Lu, P.; Larrosa, I., Redox-Controlled Selectivity of C-H Activation in the Oxidative Cross-Coupling of Arenes. *Angew. Chem. Int. Ed.* **2013**, *52*, 1781-1784; (d) Yu, Z.; Ma, B.; Chen, M.; Wu, H.-H.; Liu, L.; Zhang, J., Highly Site-Selective Direct C-H Bond Functionalization of Phenols with α -Aryl- α -diazoacetates and Diazooxindoles via Gold Catalysis. *J. Am. Chem. Soc.* **2014**, *136*, 6904-6907; (e) Cambeiro, X. C.; Ahlsten, N.; Larrosa, I., Au-Catalyzed Cross-Coupling of Arenes via Double C-H Activation. *J. Am. Chem. Soc.* **2015**, *137*, 15636-15639.

5. Ciana, C.-L.; Phipps, R. J.; Brandt, J. R.; Meyer, F.-M.; Gaunt, M. J., A Highly Para-Selective Copper(II)-Catalyzed Direct Arylation of Aniline and Phenol Derivatives. *Angew. Chem. Int. Ed.* **2011**, *50*, 458-462.

6. For selected examples, see: (a) Wang, X.; Leow, D.; Yu, J.-Q., Pd(II)-Catalyzed para-Selective C-H Arylation of Monosubstituted Arenes. *J. Am. Chem. Soc.* **2011**, *133*, 13864-13867; (b) Luan, Y.-X.; Zhang, T.; Yao, W.-W.; Lu, K.; Kong, L.-Y.; Lin, Y.-T.; Ye, M., Amide-Ligand-Controlled Highly para-Selective Arylation of Monosubstituted Simple Arenes with Arylboronic Acids. *J. Am. Chem. Soc.* **2017**, *139*, 1786-1789; (c) Naksomboon, K.; Poater, J.; Bickelhaupt, F. M.; Fernandez-Ibanez, M. A., para-Selective C-H Olefination of Aniline Derivatives via Pd/S₂O-Ligand Catalysis. *J. Am. Chem. Soc.* **2019**, *141*, 6719-6725.

7. For examples of Pd-catalyzed para-functionalization, see: (a) Bag, S.; Patra, T.; Modak, A.; Deb, A.; Maity, S.; Dutta, U.; Dey, A.; Kancherla, R.; Maji, A.; Hazra, A.; Bera, M.; Maiti, D., Remote para-C-H Functionalization of Arenes by a D-Shaped Biphenyl Template-Based Assembly. *J. Am. Chem. Soc.* **2015**, *137*, 11888-11891; (b) Patra, T.; Bag, S.; Kancherla, R.; Mondal, A.; Dey, A.; Pimparkar, S.; Agasti, S.; Modak, A.; Maiti, D., Palladium-Catalyzed Directed para C-H Functionalization of Phenols. *Angew. Chem. Int. Ed.* **2016**, *55*, 7751-7755; (c) Maji, A.; Guin, S.; Feng, S.; Dahiya, A.; Singh, V. K.; Liu, P.; Maiti, D., Experimental and Computational Exploration of para-Selective Silylation with a Hydrogen-Bonded Template. *Angew. Chem. Int. Ed.* **2017**, *56*, 14903-14907; (d) Maji, A.; Dahiya, A.; Lu, G.; Bhattacharya, T.; Brochetta, M.; Zanon, G.; Liu, P.; Maiti, D., H-bonded reusable template assisted para-selective ketonisation using soft electrophilic vinyl ethers. *Nature Commun.* **2018**, *9*, 3582.

8. For examples of Ru-catalyzed para-functionalization, see: (a) Leitch, J. A.; McMullin, C. L.; Paterson, A. J.; Mahon, M. F.; Bhonoah, Y.; Frost, C. G., Ruthenium-Catalyzed para-Selective C-H Alkylation of Aniline Derivatives. *Angew. Chem. Int. Ed.* **2017**, *56*, 15131-15135; (b) Yuan, C.; Zhu, L.; Chen, C.; Chen, X.; Yang, Y.; Lan, Y.; Zhao, Y., Ruthenium(II)-enabled para-selective C-H difluoromethylation of anilides and their derivatives. *Nature Commun.* **2018**, *9*, 1189; (c) Yuan, C.; Zhu, L.; Zeng, R.; Lan, Y.; Zhao, Y., Ruthenium(II)-Catalyzed C-H Difluoromethylation of Ketoximes: Tuning the Regioselectivity from the meta to the para Position. *Angew. Chem. Int. Ed.* **2018**, *57*, 1277-1281.

9. (a) Cho, J.-Y.; Tse, M. K.; Holmes, D.; Maleczka, R. E.; Smith, M. R., Remarkably Selective Iridium Catalysts for the Elaboration of Aromatic C-H Bonds. *Science* **2002**, *295*, 305-308; (b) Ishiyama, T.; Takagi, J.; Ishida, K.; Miyaura, N.; Anastasi, N. R.; Hartwig, J. F., Mild Iridium-Catalyzed Borylation of Arenes. High Turnover Numbers, Room Temperature Reactions, and Isolation of a Potential Intermediate. *J. Am. Chem. Soc.* **2002**, *124*, 390-391; (c) Mkhalid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F., C-H Activation for the Construction of C-B Bonds. *Chem. Rev.* **2010**, *110*, 890-931.

10. (a) Hartwig, J. F., Regioselectivity of the borylation of alkanes and arenes. *Chem. Soc. Rev.* **2011**, *40*, 1992-2002; (b) Preshlock, S. M.; Ghaffari, B.; Maligres, P. E.; Krska, S. W.; Maleczka, R. E.; Smith, M. R., High-Throughput Optimization of Ir-Catalyzed C-H Borylation: A Tutorial for Practical Applications. *J. Am. Chem. Soc.* **2013**, *135*, 7572-7582.

11. Ros, A.; Fernandez, R.; Lassaletta, J. M., Functional group directed C-H borylation. *Chem. Soc. Rev.* **2014**, *43*, 3229-3243.

12. (a) Kuninobu, Y.; Ida, H.; Nishi, M.; Kanai, M., A meta-selective C-H borylation directed by a secondary interaction between ligand and substrate. *Nature Chem.* **2015**, *7*, 712-717; (b) Bisht, R.; Chattopadhyay, B., Formal Ir-Catalyzed Ligand-Enabled Ortho and Meta Borylation of Aromatic Aldehydes via in Situ-Generated Imines. *J. Am. Chem. Soc.* **2016**, *138*, 84-87; (c) Davis, H. J.; Mihai, M. T.; Phipps, R. J., Ion Pair-Directed Regiocontrol in Transition-Metal Catalysis: A Meta-Selective C-H Borylation of Aromatic Quaternary Ammonium Salts. *J. Am. Chem. Soc.* **2016**, *138*, 12759-12762; (d) Davis, H. J.; Genov, G. R.; Phipps, R. J., meta-Selective C-H Borylation of Benzylamine-, Phenethylamine-, and Phenylpropylamine-Derived Amides Enabled by a Single Anionic Ligand. *Angew. Chem. Int. Ed.* **2017**, *56*, 13351-13355; (e) Bisht, R.; Hoque, M. E.; Chattopadhyay, B., Amide Effects in C-H Activation: Noncovalent Interactions with L-Shaped Ligand for meta Borylation of Aromatic Amides.

Angew. Chem. Int. Ed. **2018**, *57*, 15762-15766; (f) Mihai, M. T.; Davis, H. J.; Genov, G. R.; Phipps, R. J., Ion Pair-Directed C-H Activation on Flexible Ammonium Salts: meta-Selective Borylation of Quaternized Phenethylamines and Phenylpropylamines. *ACS Catal.* **2018**, *8*, 3764-3769; (g) Yang, L.; Uemura, N.; Nakao, Y., meta-Selective C-H Borylation of Benzamides and Pyridines by an Iridium-Lewis Acid Bifunctional Catalyst. *J. Am. Chem. Soc.* **2019**, *141*, 7972-7979; (h) Lee, B.; Mihai, M. T.; Stojalninkova, V.; Phipps, R. J., Ion-Pair-Directed Borylation of Aromatic Phosphonium Salts. *J. Org. Chem.* **2019**, doi: 10.1021/acs.joc.9b00878.

13. For leading examples of non-catalyzed electrophilic borylation, see: (a) Bagutski, V.; Del Grosso, A.; Carrillo, J. A.; Cade, I. A.; Helm, M. D.; Lawson, J. R.; Singleton, P. J.; Solomon, S. A.; Marcelli, T.; Ingleson, M. J., Mechanistic Studies into Amine-Mediated Electrophilic Arene Borylation and Its Application in MIDA Boronate Synthesis. *J. Am. Chem. Soc.* **2013**, *135*, 474-487; (b) Del Grosso, A.; Ayuso Carrillo, J.; Ingleson, M. J., Regioselective electrophilic borylation of haloarenes. *Chem. Commun.* **2015**, *51*, 2878-2881; (c) Yin, Q.; Klare, H. F. T.; Oestreich, M., Catalytic Friedel-Crafts C-H Borylation of Electron-Rich Arenes: Dramatic Rate Acceleration by Added Alkenes. *Angew. Chem. Int. Ed.* **2017**, *56*, 3712-3717.

14. For rhodium-catalyzed silylation which has enhanced sensitivity to existing sterics, see: Cheng, C.; Hartwig, J. F., Rhodium-Catalyzed Inter-molecular C-H Silylation of Arenes with High Steric Regiocontrol. *Science* **2014**, *343*, 853-857.

15. (a) Saito, Y.; Segawa, Y.; Itami, K., para-C-H Borylation of Benzene Derivatives by a Bulky Iridium Catalyst. *J. Am. Chem. Soc.* **2015**, *137*, 5193-5198; (b) Haines, B. E.; Saito, Y.; Segawa, Y.; Itami, K.; Musaev, D. G., Flexible Reaction Pocket on Bulky Diphosphine-Ir Complex Controls Regioselectivity in para-Selective C-H Borylation of Arenes. *ACS Catal.* **2016**, *6*, 7536-7546.

16. Yang, L.; Semba, K.; Nakao, Y., para-Selective C-H Borylation of (Hetero)Arenes by Cooperative Iridium/Aluminum Catalysis. *Angew. Chem. Int. Ed.* **2017**, *56*, 4853-4857.

17. Hoque, M. E.; Bisht, R.; Haldar, C.; Chattopadhyay, B., Noncovalent Interactions in Ir-Catalyzed C-H Activation: L-Shaped Ligand for Para-Selective Borylation of Aromatic Esters. *J. Am. Chem. Soc.* **2017**, *139*, 7745-7748.

18. (a) Preshlock, S. M.; Plattner, D. L.; Maligres, P. E.; Krska, S. W.; Maleczka, R. E.; Smith, M. R., A Traceless Directing Group for C-H Borylation. *Angew. Chem. Int. Ed.* **2013**, *52*, 12915-12919; (b) Smith, M. R.; Bisht, R.; Haldar, C.; Pandey, G.; Dannatt, J. E.; Ghaffari, B.; Maleczka, R. E.; Chattopadhyay, B., Achieving High Ortho Selectivity in Aniline C-H Borylations by Modifying Boron Substituents. *ACS Catal.* **2018**, *8*, 6216-6223.

19. Tajuddin, H.; Harrison, P.; Bitterlich, B.; Collings, J. C.; Sim, N.; Batsanov, A. S.; Cheung, M. S.; Kawamorita, S.; Maxwell, A. C.; Shukla, L.; Morris, J.; Lin, Z.; Marder, T. B.; Steel, P. G., Iridium-catalyzed C-H borylation of quinolines and unsymmetrical 1,2-disubstituted benzenes: insights into steric and electronic effects on selectivity. *Chem. Sci.* **2012**, *3*, 3505-3515.

20. (a) Boebel, T. A.; Hartwig, J. F., Silyl-Directed, Iridium-Catalyzed ortho-Borylation of Arenes. A One-Pot ortho-Borylation of Phenols, Arylamines, and Alkylarenes. *J. Am. Chem. Soc.* **2008**, *130*, 7534-7535; (b) Chattopadhyay, B.; Dannatt, J. E.; Andujar-De Sanctis, I. L.; Gore, K. A.; Maleczka, R. E.; Singleton, D. A.; Smith, M. R., Ir-Catalyzed ortho-Borylation of Phenols Directed by Substrate-Ligand Electrostatic Interactions: A Combined Experimental/in Silico Strategy for Optimizing Weak Interactions. *J. Am. Chem. Soc.* **2017**, *139*, 7864-7871; (c) Li, H.-L.; Kanai, M.; Kuninobu, Y., Iridium/Bipyridine-Catalyzed ortho-Selective C-H Borylation of Phenol and Aniline Derivatives. *Org. Lett.* **2017**, *19*, 5944-5947.

21. See supporting information for full details.

For Table of Contents Only:

