

Communication

Para-Selective, Iridium-Catalyzed C–H Borylations of Sulfated Phenols, Benzyl Alcohols, and Anilines Directed by Ion-Pair Electrostatic Interactions

Jose R. Montero Bastidas, Thomas J Oleskey, Susanne L. Miller, Milton R. Smith, and Robert E. Maleczka

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.9b08464 • Publication Date (Web): 16 Sep 2019

Downloaded from pubs.acs.org on September 16, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Para-Selective, Iridium-Catalyzed C–H Borylations of Sulfated Phenols, Benzyl Alcohols, and Anilines Directed by Ion-Pair Electrostatic Interactions

Jose R. Montero Bastidas, Thomas J. Oleskey, Susanne L. Miller, Milton R. Smith, III,* and Robert E. Maleczka, Jr.*

Department of Chemistry, Michigan State University, 578 South Shaw Lane, East Lansing, Michigan 48824, United States

Supporting Information Placeholder

ABSTRACT: *Para* C–H borylation (CHB) of tetraalkylammonium sulfates and sulfamates have been achieved using bipyridine-ligated Ir boryl catalysts. Selectivities can be modulated both by the length of the alkyl groups in the tetraalkylammonium cations and the substituents on the bipyridine ligands. Ion-pairing, where the alkyl groups of the cation shield the meta C–H bonds in the counter-anions, is proposed to account for para selectivity. The 4,4'-dimethoxy-2,2'-bipyridine ligand gave superior selectivities.

For two competing pathways, a difference in barrier heights of 2.5 kcal·mol⁻¹ is sufficient for 99% of the reactants to follow the favored pathway in a chemical reaction. This is lower than the barrier for converting the anti-conformer of butane to the gauche form.¹ In transition-metal mediated reactions, a classic mode for selectively functionalizing bonds in a substrate relies on the coordination of an atom in a reactant functional group to the metal center of a compound or catalytic intermediate. The magnitudes of the ligand-metal interactions are at least an order of magnitude greater than the difference in barrier heights necessary for 99:1 selectivity. Consequently, design of catalysts where selectivity is conferred by weakly coordinating groups,² as well as catalysts that leverage even weaker interactions (e.g. hydrogen-bonding, ion-pairing, dipole-dipole, etc.) for selective transformations,^{3,4} is attracting significant attention.

C–H functionalizations offer both atom and step economical means of converting ubiquitous C–H bonds to a range of functional groups.^{5,6} C–H borylations (CHBs) convert C–H bonds to C–B bonds and are mediated by both metal and metal-free catalysts.^{7–9} CHB reactions are valuable due to (i) facile substitution of the boron moiety by numerous functional groups, and (ii) functional group tolerance of CHB catalysts, particularly those containing Ir.

The first Ir CHB catalysts enabled C(sp²)–H functionalizations with high selectivity for the most sterically accessible C–H bonds.^{10–12} In substrates where multiple C–H bonds are sterically accessible, early generation catalysts often give isomer mixtures, as well as multiply borylated products. To overcome

these limitations, more selective Ir catalysts have been designed. The first examples were ortho-selective, relying on strongly coordinating functional groups in the substrate,^{13,14} while later reports exploited weaker interactions for ortho selectivity.^{15,16}




Figure 1. Para C–H borylations.

By comparison, meta and para CHBs pose different challenges because their C–H bonds are further from the functional group. One meta-selective CHB has been ascribed to a classical


chelate-directed mechanism,¹⁷ while others rely on Ir ligands bearing groups that engage in non-covalent interactions with substrate functional groups to effect meta CHB.^{18–20}

Figure 1 depicts approaches for para-selective CHB. The first CHBs with high para selectivity involved electrophilic additions of borenium cations to arenes bearing ortho,para-directors.²¹ Sterically directed CHBs rely on hindered phosphine ligands and substrates with large substituents.²² More recently, para borylations of esters have been achieved through non-covalent interactions with K ions or coordination of the ester oxygen to hindered Lewis acids.^{23,24}

Our inspiration was based on the Phipps' group ion-pair directed CHBs with one key difference.¹⁹ Instead of using *oppositely* charged groups on the ligand and substrate, combinations where groups on the ligand and substrate had the *same* charge were surveyed with the expectation that para borylation would be favored because electrostatic repulsions between the ligand and substrate would be minimized. However, a control experiment where tetrabutylammonium 2-chlorophenyl sulfate (**1a'**) was subjected to standard borylation conditions with a *neutral* bipyridine ligand (sealed tube with B₂pin₂ as the boron source, 1.5 mol % [Ir(cod)OMe]₂ as the precatalyst and 3 mol % dtbpy as the ligand, in THF at 80 °C) showed promising results with 6:1 para to meta regioselectivity (Table 1, entry 1). We hypothesized that substrate ion-pairing interactions, where the *n*-butyl groups of the cation shield the meta C–H bonds of the counter-anions, accounted for the para selectivity.

In Nakao's study, ligand geometry played a key role in enhancement of the para selectivity.²⁴ Similarly, we have observed that ligand choice can impact CHB regiochemistry where there is little steric differentiation between different arene C–H bonds.²⁵

Table 1. Ligand Effect on Para CHB of 1a'



entry	ligand (R =)	solvent	T °C	conv. ^a	p:m ^a
1	<i>t</i> -Bu (L1)	THF	80	100	6:1
2	CN (L2)	THF	80	0	ND
3	H (L3)	THF	80	89	7:1
4	Me (L4)	THF	80	96	10:1
5	OMe (L5)	THF	80	100	13:1
6	tmphen (L6)	THF	80	100	6:1
7	phen (L7)	THF	80	45	9:1
8	NMe ₂ (L8)	dioxane	80	100	12:1
9	OMe (L5)	dioxane	80	100	14:1
10	OMe (L5)	dioxane	60	100	16:1
11	OMe (L5)	dioxane	40	100	17:1
12 ^b	OMe (L5)	dioxane	rt	83	21:1

^aDetermined by ¹H NMR analysis. ^bReaction carried out for 50 h





Figure 2. Lowest energy conformation geometry of 1a'.


Therefore, we tested commercially available substituted bipyridine and phenanthroline ligands (Table 1) in CHB reactions run at 80 °C. The reactivity of the ligands is in accordance with previously noted electronic effects,²⁶ with electron rich ligands affording a more active system relative to electron poor ligands. The borylation in THF with 4,4'-dimethoxy-2,2'-bipyridine (**L5**) as the ligand went to 100% conversion and afforded the best para selectivity (13:1). This observation is notable in that **L5** is a non-traditional CHB ligand. Switching the solvent to dioxane slightly increased the para selectivity, whereas other apolar solvents worsened regioselectivity (see SI for details). Running the reactions at lower temperature (60 °C (entry 10) and 40 °C (entry 11)) further improved the para selectivity while still allowing for full conversion. The reaction at room temperature (entry 12) afforded 21:1 para selectivity, but starting material remained even after 50 h.

With the experiments in Table 1 establishing 4,4'-dimethoxy-2,2'-bipyridine and dioxane as our ligand and solvent of choice, we next investigated the effect of the tetraalkylammonium salt on para selectivity. DFT geometry optimization of 1a' (Figure 2), using B3LYP functional and 6-31G* basis set for all the atoms suggested to us that a slightly shorter alkyl chain would still block the meta position but leave the para position more exposed, thus potentially leading to improved para selectivity. As shown in Scheme 1, the CHB of tetrapropylammonium 2-chlorophenyl sulfate (**1a**) validated this hypothesis, as running the reaction with ligand **L5** in dioxane at 40 °C pushed the para selectivity to 22:1. We also examined tetraethylammonium 2-chlorophenyl sulfate (**1a''**) as a substrate. In terms of chain shortening, clearly diminishing returns had set in as the para selectivity decreased to 6:1.

Based on our results, we chose to test a series of phenol derived sulfates with *n*-Pr₄N⁺ as the counterion to determine substrate scope. As illustrated in Scheme 2, borylations of a series of 2-substituted phenol derived sulfates produced the para regioisomer as the major isomer, often with >20:1 selectivity. Most isolated yields were in the 70–80% range. Notably upon

Scheme 1. Effect of alkyl chain length on regioselectivity




Scheme 2. Borylation of phenol derived sulfates^a

^aPara/meta ratios were measured by ¹H-NMR on crude reaction mixtures of the borylated sulfates. Yields refer to isolated material with the para/meta ratio of the isolated material given in parentheses (**2o** and **2p** were not isolated). See the SI for experimental details, including preparations of the starting sulfates.

isolation the para to meta isomer was enhanced, in some cases to >50:1.

Given that CHB ortho to small substituents is common,²⁷ borylation at the C-3 and C-5 meta CH bonds of 2-fluoro- (**1e**) and 2-cyanophenol sulfate (**1i**) are possible. Indeed, analysis of the crude reaction mixture indicated that **1i** gave a mixture of the para to 5-Bpin to 3,5-diBpin products in a ratio of approximately 7.5:1:0.4. For substrates **1e** and **1i**, the observed minor isomer was that with the Bpin ortho to the fluoro group and no diborylation was observed. Given this preference, it was perhaps somewhat surprising that 3-fluorophenol sulfate (**1d**) produced a relatively large amount of the meta regioisomer.

Not surprising was that the CHB of 3-substituted phenol sulfates (**1n-p**) gave the meta isomer as the major product, showing that such ion-pair interactions are limited in their ability to overcome steric crowding of the para CH position. Last, we

Scheme 3. Borylation of aniline derived sulfamates^a


^aPara/meta ratios were measured by ¹H-NMR (or ¹⁹F-NMR for **4c'**) on crude reaction mixtures of the borylated sulfates. Yields refer to isolated material with the para/meta ratio of the isolated material given in parentheses. See the SI for experimental details, including preparations of the starting sulfamates. ^bRun with the $n-Pr_4N^+$ counterion. ^cProduct isolated as the acetamide.

borylated the sulfate of phenol and observed the para, meta, and 3,5-dimeta borylated products in a ratio of 4.4:1.1.8, or a para/meta ratio of 1.6:1. This result is consistent with the assumption that the ion-pairing can only block one meta site and thus the reactions need a 2-substituent to sterically block the second meta CH bond.

Turning to anilines (Scheme 3), we subjected tetrapropylammonium 2-chlorophenylsulfamate (**3a**) to our now standard conditions. The para selectivity (40:1) was even better than that observed for **1a**. Questioning if the chain length of the tetraalkylammonium salt would also impact the para selectivity for aniline derivatives, we prepared and reacted the tetrabutylammonium salt (**3a'**). In contrast to the phenol sulfates, employing this counterion met with 43:1 para selectivity and a higher isolated yield. Owing to this result and that the tetrabutylammonium salts are somewhat easier to prepare and isolate, we chose $n-Bu_4N^+$ as the counterion for CHB's of a series of aniline sulfamates. The para selectivities for **3b'** and **3d'** were excellent, while again selectivity for a 3-fluoro substrate (**3c'**) suffered, giving only a 2.4:1 para/meta ratio. Isolation of the hydrolyzed aniline following borylation of **3c'** proved difficult. Therefore, the reaction was quenched with acetyl chloride, facilitating the isolation of **4c'**.

Finally, we surveyed three benzyl alcohol derived sulfates (Scheme 4). Generally, these substrates reacted with somewhat diminished para selectivity relative to their phenol and aniline counterparts. Products **6b** and **6d** were generated in lower yields owing in part to lower conversions and for **6d**, loss of meta isomer upon isolation. Again, borylation of a substrate with fluorine in the 2-position (**5g**) afforded a significant amount of product with the Bpin ortho to the fluorine. Applying the CHB conditions to the $n-Bu_4N^+$ counterion, **5a-5c** revealed that the counterion has a similar influence on the regioselectivity as observed for the phenols.

Scheme 4. Borylation of benzyl alcohol derived sulfates^a



^aPara/meta ratios were measured by ¹H-NMR on crude reaction mixtures of the borylated sulfates. Yields refer to isolated material with the para/meta ratio of the isolated material given in parentheses. See the SI for experimental details, including preparations of the starting sulfates. ^bRun with the *n*-Bu₄N⁺ counterion.

In summary, ion-pair electrostatic interactions can be used to direct Ir-catalyzed borylation to the para position of sulfates and sulfamates derived from phenols, anilines, and benzyl alcohols. We hypothesize that the source of the para selectivity is a steric block created by the carbon chain of the tetrabutylammonium counterion. For sulfates derived from phenols and benzyl alcohols, *n*-Pr₄N⁺ salts gave better selectivity than their *n*-Bu₄N⁺ counter parts. The chain length of tetrabutylammonium salt was not as influential on the borylation of the sulfamates derived from anilines. Notably, optimal results were observed with the non-traditional CHB ligand 4,4'-dimethoxy-2,2'-bipyridine. This serves to remind the community to look beyond dtbpy or tmphen when optimizing CHB reactions.²⁸

ASSOCIATED CONTENT

Supporting Information. Experimental procedures, including preparation of starting materials, and compound characterization data (PDF). The Supporting Information is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Author

* Email: smithmil@msu.edu

* Email: maleczka@chemistry.msu.edu

Notes

Conflict of Interest: S.L.M., M.R.S., and R.E.M. own a percentage of BoroPharm, Inc.

ACKNOWLEDGMENT

We thank Drs. Daniel Holmes and Li Xie of the Max. T. Rogers NMR Facility and the MSU RTSF Mass Spectrometry Core for their assistance in compound characterization. We thank the NIH (GM63188) for financial support and BoroPharm, Inc. for a gift of B₂pin₂.

REFERENCES

- (1) Anslyn, E. V.; Dougherty, D. A. *Modern Physical Organic Chemistry*; University Science Books, 2006.
- (2) Engle, K. M.; Mei, T.-S.; Wasa, M.; Yu, J.-Q. Weak Coordination as a Powerful Means for Developing Broadly Useful C-H Functionalization Reactions. *Acc. Chem. Res.* **2012**, *45*, 788–802.
- (3) Raynal, M.; Ballester, P.; Vidal-Ferran, A.; van Leeuwen, P. W. N. M. Supramolecular Catalysis. Part 1: Non-Covalent Interactions as a Tool for Building and Modifying Homogeneous Catalysts. *Chem. Soc. Rev.* **2014**, *43*, 1660–1733.
- (4) Davis, H. J.; Phipps, R. J. Harnessing Non-Covalent Interactions to Exert Control over Regioselectivity and Site-Selectivity in Catalytic Reactions. *Chem. Sci.* **2017**, *8*, 864–877.
- (5) Dixneuf, P. H.; Doucet, H. C-H Bond Activation and Catalytic Functionalization I; Springer, 2015.
- (6) Dixneuf, P. H.; Doucet, H. C-H Bond Activation and Catalytic Functionalization II; Springer, 2016.
- (7) Mkhalid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. C-H Activation for the Construction of C-B Bonds. *Chem. Rev.* **2010**, *110*, 890–931.
- (8) Xu, L.; Wang, G.; Zhang, S.; Wang, H.; Wang, L.; Liu, L.; Jiao, J.; Li, P. Recent Advances in Catalytic C-H Borylation Reactions. *Tetrahedron* **2017**, *73*, 7123–7157.
- (9) Fontaine, F.-G.; Rochette, É. Ambiphilic Molecules: From Organometallic Curiosity to Metal-Free Catalysts. *Acc. Chem. Res.* **2018**, *51*, 454–464.
- (10) Cho, J. Y.; Iverson, C. N.; Smith, M. R., III. Steric and Chelate Directing Effects in Aromatic Borylation. *J. Am. Chem. Soc.* **2000**, *122*, 12868–12869.
- (11) Cho, J.-Y.; Tse, M. K.; Holmes, D.; Maleczka, R. E., Jr; Smith, M. R., III. Remarkably Selective Iridium Catalysts for the Elaboration of Aromatic C-H Bonds. *Science* **2002**, *295*, 305–308.
- (12) Ishiyama, T.; Takagi, J.; Ishida, K.; Miyaura, N.; Anastasi, N. R.; Hartwig, J. F. Mild Iridium-Catalyzed Borylation of Arenes. High Turnover Numbers, Room Temperature Reactions, and Isolation of a Potential Intermediate. *J. Am. Chem. Soc.* **2002**, *124*, 390–391.
- (13) Kawamori, S.; Ohmiya, H.; Hara, K.; Fukuoka, A.; Sawamura, M. Directed Ortho Borylation of Functionalized Arenes Catalyzed by a Silica-Supported Compact Phosphine-Iridium System. *J. Am. Chem. Soc.* **2009**, *131*, 5058–5059.
- (14) Ishiyama, T.; Isou, H.; Kikuchi, T.; Miyaura, N. Ortho-C-H Borylation of Benzoate Esters with Bis(pinacolato)diboron Catalyzed by Iridium-Phosphine Complexes. *Chem. Commun.* **2010**, *46*, 159–161.
- (15) Roosen, P. C.; Kallepalli, V. A.; Chattopadhyay, B.; Singleton, D. A.; Maleczka, R. E., Jr; Smith, M. R., III. Outer-Sphere Direction in Iridium C-H Borylation. *J. Am. Chem. Soc.* **2012**, *134*, 11350–11353.
- (16) Chattopadhyay, B.; Dannatt, J. E.; Andujar-De Sanctis, I. L.; Gore, K. A.; Maleczka, R. E., Jr; Singleton, D. A.; Smith, M. R., III. Ir-Catalyzed Ortho-Borylation of Phenols Directed by Substrate-Ligand Electrostatic Interactions: A Combined Experimental/in Silico Strategy for Optimizing Weak Interactions. *J. Am. Chem. Soc.* **2017**, *139*, 7864–7871.

1 (17) Bisht, R.; Chattopadhyay, B. Formal Ir-Catalyzed Ligand-Ena-
2 bled Ortho and Meta Borylation of Aromatic Aldehydes via in Situ-
3 Generated Imines. *J. Am. Chem. Soc.* **2016**, *138*, 84–87.

4 (18) Kuninobu, Y.; Ida, H.; Nishi, M.; Kanai, M. A Meta-Selective C-
5 H Borylation Directed by a Secondary Interaction between Ligand and
6 Substrate. *Nat. Chem.* **2015**, *7*, 712–717.

7 (19) Davis, H. J.; Mihai, M. T.; Phipps, R. J. Ion Pair-Directed Regio-
8 control in Transition-Metal Catalysis: A Meta-Selective C-H Borylation
9 of Aromatic Quaternary Ammonium Salts. *J. Am. Chem. Soc.* **2016**,
10 *138*, 12759–12762.

11 (20) Yang, L.; Uemura, N.; Nakao, Y. Meta-Selective C–H Borylation
12 of Benzamides and Pyridines by an Iridium–Lewis Acid Bifunctional
13 Catalyst. *J. Am. Chem. Soc.* **2019**, *141*, 7972–7979.

14 (21) Del Grosso, A.; Singleton, P. J.; Muryn, C. A.; Ingleson, M. J.
15 Pinacol Boronates by Direct Arene Borylation with Borenum Cations.
16 *Angew. Chem. Int. Ed.* **2011**, *50*, 2102–2106.

17 (22) Saito, Y.; Segawa, Y.; Itami, K. Para-C–H Borylation of Benzene
18 Derivatives by a Bulky Iridium Catalyst. *J. Am. Chem. Soc.* **2015**, *137*,
19 5193–5198.

20 (23) Hoque, M. E.; Bisht, R.; Haldar, C.; Chattopadhyay, B. Noncova-
21 lent Interactions in Ir-Catalyzed C-H Activation: L-Shaped Ligand for
22 Para-Selective Borylation of Aromatic Esters. *J. Am. Chem. Soc.* **2017**,
23 *139*, 7745–7748.

24 (24) Yang, L.; Semba, K.; Nakao, Y. Para-Selective C- H Borylation of
25 (Hetero) Arenes by Cooperative Iridium/Aluminum Catalysis. *Angew.*
26 *Chem. Int. Ed.* **2017**, *56*, 4853–4857.

27 (25) Miller, S. L.; Chotana, G. A.; Fritz, J. A.; Chattopadhyay, B.;
28 Maleczka, R. E., Jr; Smith, M. R., III. C-H Borylation Catalysts That Dis-
29 tinguish Between Similarly Sized Substituents Like Fluorine and Hydro-
30 gen. *Org. Lett.* **2019**, *21*, 6388–6392.

31 (26) Ishiyama, T.; Takagi, J.; Hartwig, J. F.; Miyaura, N. A Stochio-
32 metric Aromatic C-H Borylation Catalyzed by iridium(I)/2,2'-Bipyridine
33 Complexes at Room Temperature. *Angew. Chem. Int. Ed.* **2002**, *41*,
34 3056–3058.

35 (27) Chotana, G. A.; Rak, M. A.; Smith, M. R., III. Sterically Directed
36 Functionalization of Aromatic C-H Bonds: Selective Borylation Ortho
37 to Cyano Groups in Arenes and Heterocycles. *J. Am. Chem. Soc.* **2005**,
38 *127*, 10539–10544.

39 (28) During preparation of our manuscript we became aware that
40 the Phipps group had developed a similar approach to para-selective
41 borylation. We are grateful to them for agreeing to publish their re-
42 sults in a back-to-back fashion with our own.

1 SYNOPSIS TOC (Word Style "SN_Synopsis_TOC"). If you are submitting your paper to a journal that requires a synopsis graphic
2 and/or synopsis paragraph, see the Instructions for Authors on the journal's homepage for a description of what needs to be
3 provided and for the size requirements of the artwork.
4

5 Authors are required to submit a graphic entry for the Table of Contents (TOC) that, in conjunction with the manuscript title,
6 should give the reader a representative idea of one of the following: A key structure, reaction, equation, concept, or theorem,
7 etc., that is discussed in the manuscript. Consult the journal's Instructions for Authors for TOC graphic specifications.
8

