

0040-4020(94)01051-X

ONE- AND TWO-STEP METAL ION TEMPLATED SYNTHESES OF THE CRYPTANDS

Krzysztof E. Krakowiak^a, Jerald S. Bradshaw^{b*}, Xiaolan Kou^b, and N. Kent Dalley^b

IBC Advanced Technologies, Inc., P.O. Box 565, Provo, UT 84603-0565 and Department of Chemistry, Brigham Young University, Provo, UT 84602-4670

Abstract. New cryptands [3.2.1], [4.2.1] and [4.2.2] were prepared in good yields by treating the appropriate diazacrown with the ditosylate esters of triethylene, tetraethylene or pentaethylene glycol as required. Treatment of 4,10-diaza-15-crown-5 with α, α' -dibromo-*m*-xylene gave a mixture of the cryptand (1:1 cyclization) and the tricyclic bisdiaza-15-crown-5 connected by two *m*-xylylene units (2:2 cyclization) when K₂CO₃ was used as the base. When Li₂CO₃ was used, the same reaction gave only the 1:1 cyclization. An ¹H NMR spectrum of the *m*-xylylene-containing cryptand exhibited a peak at δ 8.85 which suggests that the proton on benzene position 2 is close to the diaza-15-crown-5 cavity. The X-ray crystal structure of that compound showed that to be the case. A metal ion template effect was also observed in the preparation of [3.3.3] from 3,6,9-trioxa-1,11-diiodoundecane and its diamine analog. Only [3.3.3] was observed with K₂CO₃. When Cs₂CO₃ was used, mainly a bis(aza-12-crown-4) was isolated. When the diiodide was changed to the corresponding ditosylate, both [3.3.3] and the bis(aza-12-crown-4) were isolated.

INTRODUCTION

Cryptands, first discovered by Lehn and coworkers,^{1,2} are superior and selective complexing agents in water for the alkali and alkaline earth cations.^{3,4} The spherands,⁵ calixspherands,⁶ and torands⁷ also have been reported to form strong complexes with these cations but only in chloroform saturated with water. These latter complexing agents are not soluble in water so that no comparative thermodynamic measurements can be carried out in aqueous solutions. Since most of the practical applications of the macrocyclic complexing agents occur in aqueous solution and require ligands with high affinity and selectivity for the metal cations, cryptands are important ligands to study.

These macrocyclic ligands have been attached to solid supports for the removal of cations from aqueous solutions⁸⁻¹⁰ and in clinical chemistry for the determination of lithium, sodium, and potassium in body fluids.¹¹⁻¹⁵

Cryptands are of increasing interest because they now can be prepared in convenient one- and two-step syntheses from inexpensive starting materials.¹⁶⁻²³ Only 4 cryptands can be purchased. In view of the importance

* Department of Chemistry

^b IBC Advanced Technologies, Inc.

of the cryptands, it is important to publish all simple methods for their preparation.

The simplest one-step method for the synthesis of cryptands containing two identical bridges requires ditosylate or diiodide derivatives of the oligoethylene glycols, the appropriate diamine, and a template cation.^{16,17} Indeed, our one-step process produced four cryptands in good yields from the appropriate oligoethylene glycol ditosylates and diamines.²¹ To make cryptands containing an aromatic subcyclic unit, the appropriate diazacrown, the aromatic-containing ditosylate or dihalide and the template cation are reacted together.^{16,17} Many researchers have stressed that the template cation is very important in these reactions but no real data have been presented. In this paper, we extend our one- and two-step methods²⁰⁻²³ to prepare additional cryptands and to study the effect of metal template cations on the yields of these macrocyclization reactions. A crystal structure of the *m*-xylylene-containing cryptand (1) is also included.

RESULTS AND DISCUSSION

Cryptands 2, 3, and 5-7 (see Figure 1) were prepared by a one-step method from the appropriate diamine and ditosylate, diiodide, or dimesylate as shown in Scheme I. Cryptand 7 was in one instance accompanied by a bis(aza-12-crown-4) by-product as will be discussed later. The structure of the 7-KI complex was recently published²⁴ without complete details of the template effect for different cations. This template effect will be discussed later. The synthesis of cryptands 2, 3, 5, and 6 have been reported in communication form without complete synthetic details.²¹



Figure 1. Cryptands mentioned in this study

Cryptands 4, 8, and 9 were prepared from the appropriate diazacrown and ditosylate as shown in Scheme II. Cryptand 1 and the 2:2 tricyclic adduct 10 were prepared from diaza-15-crown-5, α, α' -dibromo-*m*-xylene and K₂CO₃ in acetonitrile. We observed similar results when diaza-18-crown-6 was treated with α, α' -dibromo-*m*-xylene to give the corresponding cryptand and the 2:2 tricyclic adduct similar to 10.¹⁶ Other researchers have reported only 10 in a 30% yield in this reaction.^{25,26} In our hands, potassium, rubidium or cesium ion templates gave both 1 and 10. However with Li₂CO₃, only 1 was formed. We believe that the metal template must be smaller than the cryptand cavity to effectively catalyze its synthesis. So, Li⁺ is smaller than the cavity of 1.

Cryptand 1 does not form the typical teardrop-shaped spot on silica gel TLC using NH_4OH and methanol as we previously reported for the aliphatic cryptands.^{18,20} Many macrocyclic ligands other than the cryptands, such as 18-crown-6 and 24-crown-8, also form the teardrop-shaped TLC spots. It is important to note that the shape of the spot changes with a water/methanol eluant which indicates that NH_4OH is essential for producing the

Scheme I. One-step Method to Prepare Cryptands 2, 3, and 5-7

H₂N O 11 (n 12 (n 13 (n	$H_2 + 2 \times (0) \times $	$\frac{CH_3CN}{M_2CO_3}$ 2, 3, and 5-7
11	+ 14 (X = OTs)	Na ₂ CO ₃ 2 [2.2.1] (11%)
11	+ 14 (X = I)	Na ₂ CO ₃ 2 [2.2.1] (10%)
12	+ 14 (X = OTs)	K ₂ CO ₃ 3 [2.2.2] (36%)
13	+ 15 (X = I)	K₂CO₃ 5 [3.3.1] (20%)
13	+ 14 (X = OTs, I, OMs)	K ₂ CO ₃ 6 [3.2.2] (50, 45, 23%)
13	+ 15 (X = I, OTs)	K ₂ CO ₃ 7 [3.3.3] (56, 12%)





distinctive spot, probably a result of an ammonium complex with the macrocyclic ligand. In the case of 1, the short bridge containing the benzene ring could cause sufficient steric hindrance to prohibit ammonium ion complexation by 1. Indeed, the ¹H NMR signal attributed to the proton on position 2 of the benzene ring in 1 is shifted downfield to 8.85 δ in comparison to 8.45 for the analagous cryptand prepared from diaza-18-crown-6 and 7.9 δ for the cryptand from 24-crown-8. This result indicates that the proton on position 2 of the benzene ring of 1 is very close to the cavity of the 15-crown-5 moiety. The X-ray study of 1, discussed below, also shows that the proton is very close to the diaza-15-crown-5 cavity. The ¹H NMR spectrum of biscrown 10 is very different from that of 1.

Interesting template effects were also observed for the formation of 7 (cryptand [3.3.3]) (see Scheme III). When 3,6,9-trioxa-1,11-diiodoundecane (**15a**) was treated with diamine **13** and K_2CO_3 as the base, only cryptand 7 was isolated in a 40% yield. The same reactants, in the presence of Cs_2CO_3 , gave bis(aza-12-crown-4) (**22**) in a 33% yield and cryptand 7 in only a 13% yield. Using ditosylate **15b** instead of diiodide **15a** resulted in lower yields of the products.

Scheme III. Preparation of Cryptand 7 and Biscrown 22



The first cryptands were prepared in 1968.^{1,2} Only aliphatic cryptands with cavity sizes from [1.1.1] to [3.3.3] have been reported. Cryptands [3.2.1] (4), [3.3.1] (5), [4.2.1] (8) and [4.2.2] (9) are new macrotricycles. It is not possible to use the simple one-step reaction of 1 mole of a diamine with 2 moles of a dihalide (ditosylate) to prepare a cryptand with 3 different bridges between the nitrogen atoms. For those types of cryptands, one needs to condense the appropriate diazacrown with the appropriate dihalide or ditosylate (see Scheme II). Cryptands 4, 8, and 9 were prepared by this process as mentioned above, since the starting diaza-15-crown-5 (17) and two diaza-18-crown-6 reactants (16 and 18) are readily available. Other cryptands could be prepared in the same manner if the diazacrowns were available. Diazacrowns such as 16 can be prepared by a convenient one-step method from the appropriate dihalide (ditosylate) and a diamine as reported.²⁷ Cryptand 4 was prepared from both 16 and 17, giving about the same yields. Thus, either method could be used, but 4,10-diaza-15-crown-5 is more available than 4,10-diaza-18-crown-6, although the latter compound can easily be prepared.²⁷

An X-ray study of 1 was initiated in order to determine if the hydrogen on the 2-position of the benzene (H18A) is close to the cavity of the crown 15 ring. A computer drawing showing the conformation of 1 and giving

the atom labels is presented in Figure 2.²⁸ The figure shows that H18A is directed into the cavity. Because there is no given point that fixes the location of the cavity, it is difficult to show the relationship of an atom to the cavity of the molecule, but the following indicators are helpful. H18A is 1.46 Å from the least-squares plane



Figure 2. Computer drawing of 1 with O13' and hydrogens, except for H18A, omitted for clarity.

calculated for the hetero atoms of the crown portion of the molecule. The distances of H18A from N1, O4, N7, O10, and O13 are 2.511, 2.652, 2.547, 3.468 and 3.154 Å, respectively, and it is 0.43 Å above the least-squares plane of N1, O4 and N7. By way of comparison, in the diaza-18-crown-6 analogue of 1,²⁰ the hydrogen in the 2-position of the benzene ring is 1.76 Å from the least-square plane of the hetero atoms. The distances of that hydrogen to the two nitrogens are 2.44 and 2.36 Å but the distances to the four oxygens are 3.656, 3.735, 3.931, and 3.984 Å. Clearly the structural study indicates that H18A more closely approaches the cavity of 1 than does the similar H in the diaza-18-crown-6 analogue.

EXPERIMENTAL

Cryptands 3^{21} 5^{21} and 7^{24} were prepared as reported. Other cryptands, such as 2 and 6, were obtained in a similar manner and their physical and spectral properties were the same as those reported.²⁹

Preparation of Cryptand 1 and Macropolycycle 10. Diaza-15-crown-5 (0.55 g, 2.52 mmol), 0.67 g (2.53 mmol) of α , α' -dibromo-*m*-xylene and 10.0 g of anhydrous K₂CO₃ were refluxed in 200 mL of CH₃CN for 20 h. After cooling, the mixture was filtered and the solvent was evaporated under reduced pressure. CH₂Cl₂ (30 mL) was added to the residue and the mixture was stirred and filtered. The solvent was evaporated from the filtrate and the residue was purified on an alumina column using CH₂Cl₂/THF: 5/1 as eluant to give 0.36 g (54%)

of 1, ¹H NMR (δ); 2.5 (m, 2 H), 2.7 (m, 2 H), 2.9 (m, 6 H), 3.35 (m, 2 H) 3.65 (m, 10 H), 4.0 (s, 1 H), 4.08 (s, 1 H), 6.85-7.15 (m, 3 H), 8.85 (s, 1 H). *Anal.* Calcd for C₁₈H₂₈N₂O₃: C, 67.47; H, 8.81. found: C, 67.48; H, 8.79; MS (m/e): 320 (100%), MS (CI): 321.

The above alumina column was further eluted with CH_2Cl_2/THF : 3/1 and 1/1 to give 0.20 g (30%) of **10**, ¹H NMR (δ): 2.7-2.9 (m, 16 H), 4.5-4.7 (m, 32 H), 7.1-7.3 (m, 6 H), 7.6 (s, 2 H); *Anal.* Calcd for $C_{36}H_{56}N_4O_6$: C, 67.47; H, 8.81. Found: C, 67.40; H, 8.63; MS (m/e) 640 (100%); MS (CI) 641.

Preparation of Bis(aza-12-crown-4) (22) (Scheme III). Diamine **13** (0.96 g, 5 mmol), 4.35 g (10.5 mmol) of diiodide **15a** and 10.0 g of Cs_2CO_3 were stirred under reflux in 200 mL of CH_3CN for 24 h. After cooling the reaction mixture was filtered and evaporated under reduced pressure. The residue was stirred in 50 mL of CH_2Cl_2 and filtered. The filtrate was evaporated and the residue was purified on an alumina column using THF/C₈H₅OH: 50:1, 20:1 and 10:1 as eluants and on silica gel using CH_3OH/NH_4OH : 20/1, 10/1, 5/1 and 2/1 as eluants. The appropriate fractions of the second main spot were combined, evaporated, mixed with $C_6H_5CH_3$ and filtered to give 0.75 g (33%) of biscrown **22**; ¹H NMR (δ): 2.75 (m, 12 H), 3.65 (m, 36 H); ¹³C NMR (δ): 56.21, 56.60, 69.78, 70.23, 70.84, 71.01, 71.21, 71.77 *Anal.* Calcd for $C_{24}H_{48}N_2O_9$: C, 56.68; H, 9.50. Found: C, 56.81; H, 9.27; MS (M/E) 508; MS (CI) 509.

Preparation of Cryptands 4,8, and 9 (Scheme II). Diazacrowns **16**, **17** or **18** (0.39 mmol), 4.1 mmole of **19**, **20** or **21** and 10.0 g of Na₂CO₃ were stirred under reflux in 300 mL of CH₃CN for 7 days. The reaction mixtures were cooled and evaporated. Water (40-50 mL) was added and the resulting mixture was extracted 4 times with 150 mL potions of CH₂Cl₂. The combined CH₂Cl₂ extracts were dried (MgSO₄) and evaporated. The reside was chromatographed on alumina to decompose the complex using THF/C₂H₅OH: 100/1 as the eluant. The solvents were evaporated and the residue was chromatographed on silica gel using CH₃OH/NH₄OH: 40/1, 20/1 and 10/1 as eluants. The residue was dissolved in C₆H₅CH₃ and filtered through a glass filter to remove silica gel. The products were obtained as oils follows:

<u>Cryptand 4</u> (46%), ¹H NMR (δ): 2.7 (m, 12 H), 3.6 (m, 24 H); *Anal*. Calcd for C₁₈H₃₆N₂O₆: C, 57.36; H, 9.63. Found: C, 57.76; H, 9.53; MS (m/e) 376 (82%); MS (CI) 377.

<u>Cryptand 8</u> (30%), ¹H NMR (δ): 2.75 (m, 12 H), 3.6 (m, 28 H); *Anal*. Calcd for C₂₀H₄₀N₂O₇: C, 57.12; H, 9.58. Found: C, 57.35; H, 9.38; MS (m/e), 420 (86%); MS (CI), 421.

<u>Cryptand 9</u> (30%), ¹H NMR (δ): 2.8 (m, 12 H), 3.65 (m, 32 H); *Anal*. Calcd for C₂₂H₄₄N₂O₈: C, 56.87; H, 9.54. Found: C,56.69; H, 9.29; MS (m/e) 464 (96%); MS (CI) 465.

X-ray Crystal Study of 1. Crystallographic and single crystal intensity data for 1 were obtained using a Siemens R3m/V automated diffractometer which utilized graphite monocromated Mo K α radiation ($\lambda =$ 0.71073Å). The lattice parameters and orientation matrix were obtained using a least-square procedure involving 32 carefully centered reflections ($8.9 \le 20 \le 32.7^{\circ}$). Single crystal data were collected using a variable rate 20- θ scanning procedure. The crystal data and a summary of the experimental conditions are contained in Table 1. The structure was solved using direct methods. It was possible to locate all non-hydrogen atoms in an Emap. A difference map revealed the presence of a small peak near O13 which was a disordered O13'. The ratio of the population parameters for O13:O13'are 0.86:0.14. The approximate ratio was suggested by the peak heights in difference maps and the final values were obtained in the refinement process. The large thermal parameters for C12 and C14 suggested that these atoms were also disordered but the disorder could not be resolved. This resulted in unusual bond angles involving O13'. Positions for all hydrogens were calculated and the hydrogens were assigned isotropic thermal parameters. These H atoms were allowed to ride on their neighboring carbons during the refinement. In the calculations of H positions, O13' was omitted so realistic bond angles could be calculated. It is also omitted from Figure 2 for clarity. In the final stages of refinement all non-hydrogen atoms except O13' were refined anisotropically. O13' was refined isotropically and the hydrogen thermal parameters were not refined. Atomic scattering factors were obtained from the International Tables for X-ray crystallography.³⁰ All computer programs used in the solution, refinement and display of this structure are contained in SHLEXTL-PLUS^{TM_31}

Formula	$C_{18}H_{28}N_2O_3$
Formula weight	320.4
F(000)	696
Crystal size, mm	0.24 x 0.42 x 0.44
μ , mm ⁻¹	0.082
Crystal system	Monoclinic
Space group	$P2_1/c$
a, Å	12.843(3)
b, Å	15.145(3)
c, Å	9.604(4)
β, deg	108.80(2)
V	1768.4
Z	4
Dx,g/cc	1.203
20 range, deg	4.00 to 50.00
Independent data	$3129 (R_{int} = 1.64\%)$
Observed data	1469 (F > 3.5 σ (F))
Data/parameter ratio	6.9:1
R	7.72%
Rw	8.23%
Weighing scheme	$w^{-1} = \sigma^2(F) + 0.0009F^2$
Goodness of fit	1.68
Largest peak in difference map e A-3	0.27
Largest hole in difference map e A ⁻³	-0.28

TABLE 1. Crystal Data and Summary of Experimental Conditions for 1

Acknowledgement. Department of Energy, Basic Energy Sciences Grant No. DE-FG02-86 ER 13463 is greatfully acknowledged.

REFERENCES

- 1. Dietrich, B.; Lehn, J.-M.; Sauvage, J.-P.; Blanzat, J. Tetrahedron Lett. 1969, 2885.
- 2. Dietrich, B. Cryptate Complexes in Inclusion Compounds, Vol. 2, Atwood, J.L.; Davies, J.E.D.; McNicol, D.D. Eds, Academic Press, London, 1984.
- 3. Izatt, R.M.; Bradshaw, J.S.; Nielsen, S.A.; Lamb, J.D.; Christensen, J.J.; Sen, D. Chem Rev. 1985, 85, 271.
- 4. Izatt, R.M.; Pawlak, K.; Bradshaw, J.S.; Bruening, R.L. Chem. Rev. 1991, 91, 1721.
- 5. Cram, D.J.; Lein, G.M. J. Am. Chem. Soc. 1985, 107, 3657.
- 6. Dijkstra, P.J.; Brunink, J.A.J; Bugge, K.E.; Reinhoudt, D.N.; Harkema, S.; Ungaro, R.; Vgorzoli, F.; Ghiclini, E.J. J. Am. Chem. Soc. 1989, 111, 7567.
- 7. Bell, T.W.; Sahni, S.K. Bridged Pyridine Hosts in Inclusion Compounds Vol. 4, Atwood, J.L.; Davies, J.E.D.; MacNicol, D.D.; Eds, Oxford University Press, Oxford 1991.
- 8. Izatt, R.M.; Bradshaw, J.S., Bruening, R.L.; Krakowiak, K.E.; Tarbet, B.J. *Proceedings at the First Handford Separation Science Workshop* 1993, Pacific Northwest Laboratory Richland, Washington 1993.
- 9. Izatt, R.M.; Bradshaw, J.S.; Bruening, R.L.; Bruening, M.L. American Lab., in press.
- 10. Izatt, R.M.; Bradshaw, J.S; Bruening, R.L.; Tarbet, B.J.; Krakowiak, K.E. in *Emerging Separation Technologies for Metals and Fuels*, Ed. Lakshmanan, V.I.; Bautista, R.G.; Somasundaran. The Minerals, Metals and Material Society 1993.
- Cram, D.J.; Chapoteau, E.; Czech, P.; Gebauer, C.R.; Helgeson, R.C.; Kumar, A. Novel Chromogenic Ionophores for Determination of Sodium and Potassium in Biological Fluids: Synthesis and Application in Proceedings of the Fifth International Symposium on Inclusion Phenomena and Molecular Recognition, J.L. Atwood, Ed. Plenum Press, NY, 1990.
- 12. Czech, B. P.; Chapoteau, E.; Kumar, A. U.S. Patent 5,817,103, 1993.
- 13. Czech, B. P.; Chapoteau, E.; Chimenti, M. Z.; Zazulak, W.; Gebauer, C. R.; Kumar, A. Anal. Chim. Acta 1992, 263, 159.
- 14. Czech, B. P.; Chapoteau, E.; Zazulak, W.; Gebauer, C. R.; Kumar, A. Anal. Chim. Acta 1990, 241, 127.
- 15. Zazulak, W.; Chapoteau, E.; Czech, B. P.; Kumar, A. J. Org. Chem. 1992, 57, 6720.
- 16. Krakowiak, K.E.; Bradshaw, J.S. Isr. J. Chem. 1992, 32, 3.
- 17. Krakowiak, K.E.; Bradshaw, J.S. Synlett. 1993, 611.
- 18. Krakowiak, K.E.; Bradshaw, J.S. J. Org. Chem. 1991, 56, 3723.
- 19. Krakowiak, K.E.; Bradshaw, J.S., An. H.-Y.; Izatt, R.M. Pure and Appl Chem. 1993, 65, 511.
- 20. Krakowiak, K.E.; Bradshaw, J.S.; Dalley, N.K.; Zhu, C.-Y.; Yi, G.-L.; Curtis, J.C.; Li, D.; Izatt, R.M. J. Org. Chem. 1992, 57, 3166.
- 21. Krakowiak, K.E.; Krakowiak, P.A.; Bradshaw, J.S. Tetrahedron Lett. 1993, 36, 777.
- 22. Bradshaw, J.S.; Krakowiak, K.E., An H.-Y.; Wang, T.-M.; Zhu, C.-Y.; Izatt, R.M. Tetrahedron Lett. 1992, 33, 4871.
- 23. Bradshaw, J.S.; An, H.-Y.; Krakowiak, K.E.; Wang, T.-M.; Zhu, C.-Y.; Izatt, R.M. J. Org. Chem. 1992, 57, 6112.
- 24. Dalley, N.K.; Krakowiak, K.E.; Bradshaw, J.S., England, M.M.; Kou, X.; Izatt, R.M. Tetrahedron 1994, 50, 2721.
- 25. Sutherland, I.O. Pure & Appl. Chem. 1989, 61, 1547.
- 26. Cheetham, G.M.T.; Harding, M.M. Acta Cryst. 1991, C47, 2478.
- 27. Krakowiak, K.E.; Mass, G.E.; Bradshaw, J.S.; Hathaway, J.K.; Izatt, R.M. J. Heterocyclic Chem. submitted.
- 28. Tables of positional and thermal parameters and bond lengths and angles for 1 have been deposited in the Cambridge Cryptallographic Data Center. These data can be obtained from the Director, Cambridge Crystallographic Center, 12 Union Road, Cambridge, CB2 IEZ, U.K.
- 29. Dietrich, B.; Lehn, J.-M.; Sauvage, J.-P.; Blanzat, J. Tetrahedron 1973, 29, 1629.
- 30. Ibers, J.A.; Hamilton, W.C., eds, International Tables for X-ray Crystallography, Vol. 4, p. 89, Kynock Press, Birmingham, England, 1974.
- 31. Sheldrick, G.M., SHELXTL-PLUSTM, Siemens Analytical X-ray Instruments, Inc., Madison, Wisconsin, 1990.

(Received in USA 30 August 1994; accepted 17 November 1994)