

INFLUENCE OF NUCLEOPHILIC REAGENTS ON THE REACTIONS OF PHOSPHORUS SULFIDES AND ALKYL HOMOLOGUES OF DAVY'S REAGENT WITH ALKYL HALIDES AND DIALKYL DISULFIDES

Il'Yas S. Nizamov , Gul'Nur G. Sergeenko , Alexey V. Matseevskii & Elvira S. Batyeva

To cite this article: Il'Yas S. Nizamov , Gul'Nur G. Sergeenko , Alexey V. Matseevskii & Elvira S. Batyeva (1998) INFLUENCE OF NUCLEOPHILIC REAGENTS ON THE REACTIONS OF PHOSPHORUS SULFIDES AND ALKYL HOMOLOGUES OF DAVY'S REAGENT WITH ALKYL HALIDES AND DIALKYL DISULFIDES, Phosphorus, Sulfur, and Silicon and the Related Elements, 132:1, 85-100, DOI: [10.1080/10426509808036977](https://doi.org/10.1080/10426509808036977)

To link to this article: <http://dx.doi.org/10.1080/10426509808036977>



Published online: 24 Sep 2006.



Submit your article to this journal [↗](#)



Article views: 21



View related articles [↗](#)



Citing articles: 1 View citing articles [↗](#)

INFLUENCE OF NUCLEOPHILIC REAGENTS ON THE REACTIONS OF PHOSPHORUS SULFIDES AND ALKYL HOMOLOGUES OF DAVY'S REAGENT WITH ALKYL HALIDES AND DIALKYL DISULFIDES

IL'YAS S. NIZAMOV*, GUL'NUR G. SERGEENKO,
ALEXEY V. MATSEEVSKII and ELVIRA S. BATYEVA

*A. E. Arbuzov Institute of Organic and Physical Chemistry, Russian Academy of
Sciences, Arbuzov Str. 8, 420088 Kazan, Russia*

(Received 18 December 1997)

Use of potassium carbonate, sodium sulfide and sodium alkylthiolates results in the reactivity enhancement of phosphorus sulfides (P_4S_{10} , P_4S_7 and P_4S_5) and alkyl homologues of Davy's reagent in their reactions with alkyl halides and dialkyl disulfides.

Keywords: Phosphorus sulfides; homologues of Davy's reagent; potassium carbonate; sodium sulfide; sodium alkylthiolates; alkyl halides; dialkyl disulfides; organothiophosphorus compounds

INTRODUCTION

During the past few years we have become interested in the problem of reactivity enhancement of lower and higher phosphorus sulfides (P_4S_{10} , P_4S_7 , P_4S_5 and P_4S_3) and their organic derivatives of 2,4-bis(substituted) 2,4-dithio-1,3,2 λ^5 ,4 λ^5 -dithiadiphosphetane structure (Lawesson's and Davy's-like reagents) in their reactions with organic compounds. Thus, the reactions of tetraphosphorus trisulfide and tetraphosphorus decasulfide with alkyl chlorides proceed under severe conditions (160–330°C) to form alkyl dichlorodithiophosphates and dialkyl chlorotrithiophosphates in the absence of any additives. [1, 2]


*Author to whom correspondence should be addressed. Tel.: (8432) 76-16-35 Fax: (8432) 75-22-53 Telex: 224864 PTB SU E-mail: nizamov@glass.ksu.ras.ru

The reactions of tetraphosphorus trisulfide, tetraphosphorus pentasulfide (P_4S_5), Lawesson's and Davy's-like reagents with dialkyl disulfides and thioacetals occurred sometimes at 100-200°C (up to 22 h) and are accompanied by resinous materials. [3-6] In order to increase reactivity of phosphorus sulfides and Lawesson's and Davy's-like reagents in the reactions with disulfides, thioacetals, aminals and sulfenamides we have previously developed efficient methods of facilitation of these reactions by involvement of organic amines and benzoyl peroxide, [3] and iodine. [7, 8] UV and ultrasonic irradiations have been shown to be effective methods of promoting the reactions of tetraphosphorus trisulfide, tetraphosphorus pentasulfide (P_4S_5), Lawesson's and Davy's-like reagents with organic disulfides, thioacetals, acetals, bis(tributylstannyl)sulfide and (isobutylthio)tributylstannane. [4, 9-11] In continuation of these works we involved potassium carbonate, sodium sulfide and sodium alkylthiolates in the reactions of phosphorus sulfides (P_4S_{10} , P_4S_7 and P_4S_5) and alkyl homologues of Davy's reagent in their reactions with alkyl halides and dialkyl disulfides and present the results in this article.

RESULTS AND DISCUSSION

Additions of inorganic salts such as sodium sulfide, Na_2S , [12] sodium carbonate, Na_2CO_3 , [12-15] sodium hydrogen-carbonate, $NaHCO_3$, [12, 15] and calcium fluoride, CaF_2 , [16] to the reaction mixture of tetraphosphorus decasulfide with carbonyl compounds were reported to accelerate the sulfurization of carbonyl groups. Salts of 1,2,3,5,6,7,4,8-hexathiadiphosphocane-4,8-dithio-4,8-dithiolic acid were obtained in the reaction of tetraphosphorus decasulfide with tetraalkylammonium bromides in the presence of sodium azide, NaN_3 , in glyme. [17] Treatment of Lawesson's reagent (and its homologues) with potassium fluoride, [18] potassium azide, [18] potassium cyanide [19] and alkali thiocyanates [20] in ketones or acetonitrile as solvents results in the formation of the alkali salts of the corresponding phosphonodithioic acids. 4-Methoxyphenyl dithiophosphonate derivatives were prepared by treatment of Lawesson's reagent with sodium and potassium salts of carboxylic and xanthogenic acids [21] or sodium alkoxides [22] in their subsequent interactions with alkyl halides or benzoyl chlorides. Obviously these salts lead to the increase of nucleophilic reactivity of tetraphosphorus decasulfide or Lawesson's reagent via the formation of some reactive and thermal instable organothiophosphorus intermediates.

We assumed that involvement of nucleophilic reagents such as potassium carbonate, sodium sulfide and sodium alkylthiolates in the reactions of phosphorus sulfides and 1,3,2,4-dithiadiphosphetane-2,4-disulfides with alkyl

TABLE I Experimental data and yields of tetrathiophosphates **4**

Initial compounds	Reaction conditions ^a Temp., °C/Time, h	Product (Yield, %)
P ₄ S ₁₀ 1 , PhCH ₂ Cl 2a , K ₂ CO ₃ 3	60-70/2	(PhCH ₂ S) ₃ P(S) 4a (51 ^b /26 ^c)
(EtSPS ₂) ₂ 8a , PhCH ₂ Cl 2a , K ₂ CO ₃ 3	60-70/2	EtSP(S)(SCH ₂ Ph) ₂ 4b (56 ^b /19 ^c)
(EtSPS ₂) ₂ 8a , BuBr 2b , K ₂ CO ₃ 3	60-70/1	EtSP(S)(SBU) ₂ 4c (71 ^b /33 ^c)
(i-BuSPS ₂) ₂ 8b , BuBr 2b , K ₂ CO ₃ 3	60/1	i-BuSP(S)(SBU) ₂ 4d (55 ^c)
(EtSPS ₂) ₂ 8a , EtSSEt 12a , K ₂ CO ₃ 3	70/1	(EtS) ₃ P(S) 4e (51 ^d)
(i-BuSPS ₂) ₂ 8b , i-BuSSBu-i 12b , K ₂ CO ₃ 3	60/1	(i-Bu) ₃ P(S) 4f (66 ^d)
(i-BuSPS ₂) ₂ 8b , PrSSPr 12c , K ₂ CO ₃ 3	60/1	i-BuSP(S)(SPr) ₂ 4g (50 ^c)
P ₄ S ₁₀ 1 , PhCH ₂ Cl 2a , Na ₂ S 15	50-60/2	(PhCH ₂ S) ₃ P(S) 4a (54 ^b /23 ^c)
(EtSPS ₂) ₂ 8a , PhCH ₂ Cl 2a , Na ₂ S 15	20/4.5	EtSP(S)(SCH ₂ Ph) ₂ 4b (21 ^b /9 ^c)
P ₄ S ₅ 16 , PrSSPr 12c , Na ₂ S 15	20/2	(PrS) ₃ P(S) 4h (46 ^b /32 ^d)
P ₄ S ₇ 18 , PrSSPr 12c , Na ₂ S 15	20/6	(PrS) ₃ P(S) 4h (22 ^b /19 ^d)
P ₄ S ₁₀ 1 , i-BuSSBu-i 12b , Na ₂ S 15	20/6	(i-Bu) ₃ P(S) 4f (39 ^b /21 ^d)
(EtSPS ₂) ₂ 8a , PrSSPr 12c , Na ₂ S 15	20/6	EtSP(S)(SPr) ₂ 4i (25 ^d)
P ₄ S ₇ 18 , PrSSPr 12c , NaSPr-i 17b	20/1	i-PrSP(S)(SPr) ₂ 4j (10 ^b /7 ^d)

^aIn MeCN. ^bYield of crude product. ^cYield of crude product purified by column chromatography. ^dYield of crude product purified by distillation.

halides and dialkyl disulfides may result in the formation of organothiophosphorus products under milder reaction conditions. Indeed, we have found that the reaction of tetraphosphorus decasulfide **1** with benzyl chloride **2a** in the presence of dried potassium carbonate **3** in anhydrous acetonitrile proceeds at 60-70°C for 2 h and gives tribenzyl tetrathiophosphate **4a** (Equation (1), and Table (I) and (II)).

The structure of tetrathiophosphate **4a** has been established by IR (Table III), ¹H NMR (Table IV), ³¹P NMR (Table II), mass spectral data (Table V) as well as elemental analyses (Table II) and by comparing its NMR spectral data with literature ones. [23] The ³¹P NMR spectrum of product **4a** (C₆H₆) reveals resonance at δ_p 90.5 ppm (cf. Lit. [23] δ_p 90.2 ppm). The electron impact mass spectrum of **4a** shows a mass peak of the molecular ion [M]⁺ m/e 432.

TABLE II Physical, analytical and the ^{31}P NMR data of tetrathiophosphates RSP(S)(SR')₂

Prod. C	R	R'	R _f Value(C ₆ H ₆)	n _D ²⁰	^{31}P NMR δ_p , ppm(C ₆ H ₆)	Molecular formula(Mol. mass)	FoundCalc., %
	H	P	S				
4a (432.1)	PhCH ₂ 58.33	PhCH ₂ 4.91	0.82 7.17	1.6569 29.60	90.5	C ₂₁ H ₂₁ PS ₄	58.08 5.60 7.01 29.13
4b^a (370.0)	Et 51.89	PhCH ₂ 5.19	0.80 8.37	1.6478 34.56	92.1	C ₁₆ H ₁₉ PS ₄	51.52 5.05 8.20 34.44
4c (302.1)	Et 39.72	Bu 7.69	0.84 10.25	1.5859 42.33	91.2	C ₁₀ H ₂₃ PS ₄	39.50 7.80 10.53 42.44
4d (330.1)	i-Bu 43.62	Bu 8.26	0.86 9.38	- 38.74	94.0	C ₁₂ H ₂₇ PS ₄	43.96 8.01 9.56 38.98
4g (302.1)	i-Bu 39.72	Pr 7.69	0.89 10.25	1.5802 42.33	93.9	C ₁₀ H ₂₃ PS ₄	39.50 7.24 10.44 42.61
4i^b (274.0)	Et 35.04	Pr 7.00	- 11.30	1.5805 46.67	94.0	C ₈ H ₁₉ PS ₄	35.32 7.21 11.21 46.34
4j^c (288.1)	i-Pr 37.49	Pr 7.36	- 10.75	1.5872 44.39	94.9	C ₉ H ₂₁ PS ₄	37.01 7.48 10.32 44.61

^aB.p. 195-198°C (0.07 mm Hg); ^bB.p. 98-100°C (0.07 mm Hg); ^cB.p. 136-138°C (0.02 mm Hg).

TABLE III IR spectral data (ν , cm^{-1}) of tetrathiophosphates **4**

Prod.	=C-H ^a , Ar	CH ₃ ^a as, s; CH ₂ ^a as, s	C=C ^a , Ar	δ^b (CH ₃ as)	P=S ^a , PS ^a as	PS ₂ ^a s, P-SC ^a
4a	3096, 3060 π , 3040	2985, 2940, 2882	1605, 1500	1450 525, 465	709, 685	570
4b	3090, 3060, 3040	2985, 2940, 2882	1605, 1500	1450	709, 685	530
4c		2970, 2938, 2880		1460	690	532
4d		2965, 2930, 2882		1470	695	540
<P<B4g		2935, 2910, 2880		1468	692	500
4i		2970, 2933, 2910, 2880		1460	690	530
4j		2980, 2940, 2930, 2880		1460	690	530

^aValence vibrations. ^bDeformation vibrations.

TABLE IV ¹H NMR data of tetrathiophosphates **4**

Prod.	CCl ₄ , δ , ppm, J, Hz
4a	4.05 (d, 3H, CH ₂ SP, ³ J _{P-H} 14.0); 6.93-7.27 (m, 15H, C ₆ H ₅)
4b	1.23 (t, 3H, CH ₃ CH ₂ SP, ³ J _{H-H} 7.5); 2.86 (dq, 2H, CH ₃ CH ₂ SP, ³ J _{H-H} 7.5, ³ J _{P-H} 16.5); 3.88 (d, 4H, CH ₂ C ₆ H ₅ , ³ J _{P-H} 14.0); 6.97-7.26 (m, 10H, C ₆ H ₅).
4c	0.98 (t, 3H, CH ₃ CH ₂ SP, ³ J _{H-H} 7.5); 1.46 (t, 6H, CH ₃ CH ₂ CH ₂ CH ₂ S, ³ J _{H-H} 7.5); 1.37-1.90 and 2.69-3.33 (m, 2H, CH ₃ CH ₂ S + m, 12H, CH ₃ CH ₂ CH ₂ CH ₂ S).
4d	0.55-2.02 (m, CH ₃ CHCH ₂ SP + CH ₃ CHCH ₂ SP + C ₄ H ₉ SP); 2.90 (dd, 2H, CH ₃ CHCH ₂ , ³ J _{H-H} 7.0, ³ J _{P-H} 15.0).
4g	1.03 (t, 6H, CH ₃ CH ₂ CH ₂ , ³ J _{H-H} 7.0); 1.09 (d, 6H, CH ₃ CHCH ₂ , ³ J _{H-H} 7.0); 1.40-2.07 (m, 4H, CH ₃ CH ₂ CH ₂ S + m, 1H, CH ₃ CHCH ₂ S); 2.48-3.18 (m, 4H, CH ₃ CH ₂ CH ₂ SP + m, 2H, CH ₃ CHCH ₂ SP).
4i	0.94 (t, 3H, CH ₃ CH ₂ SP, ³ J _{H-H} 7.5); 1.39 (t, 6H, CH ₃ CH ₂ CH ₂ S, ³ J _{H-H} 7.5); 1.28-1.83 (m, 4H, CH ₃ CH ₂ CH ₂ SP); 2.63-3.27 (m, 2H, CH ₃ CH ₂ SP); 2.93 (dt, 4H, CH ₃ CH ₂ CH ₂ SP, ³ J _{H-H} 7.5, ³ J _{P-H} 24.0).
4j	1.02 (t, 6H, CH ₃ CH ₂ CH ₂ SP, ³ J _{H-H} 7.0); 1.43 (d, 6H, CH ₃ CHSP, ³ J _{H-H} 7.0); 1.25-2.05 (m, 4H, CH ₃ CH ₂ CH ₂ SP); 2.42-3.38 (m, 1H, CH ₃ CHSP); 2.93 (dt, 4H, CH ₃ CH ₂ CH ₂ SP, ³ J _{H-H} 7.0, ³ J _{P-H} 17.0).

The reaction (Equation (1)) was accompanied by the evolution of carbon dioxide and formation of a precipitate of potassium chloride **5a**, which was removed by washing with water. The product **4a** was not decomposed by washing with water. Tetrathiophosphate **4a** was consequently purified by column chromatography. It should be noted that unlike with the literature data on the reactions of tetraphosphorus decasulfide with alkyl halides [2] the involvement of potassium carbonate in the reaction studied (Equation (1)) lead not only to decrease in reaction temperature but the change of the reaction route gave a different synthetic result.


TABLE V Mass spectral data of tetrathiophosphates 4

Prod.	<i>i</i> -C ₄ H ₁₀ , <i>m/e</i> (<i>I</i> _{rel.} , %)
4a ^a	433 [M + H] ⁺ (25), 310 [M + H - PhCH ₂ - S] ⁺ (25), 246 [M + H - PhCH ₂ - 3S] ⁺ (100).
4a ^b	432 [M] ⁺ (13), 309 [M - PhCH ₂ - S] ⁺ (16), 277 [M - PhCH ₂ - 2S] ⁺ (25).
4b ^a	371 [M + H] ⁺ (30); 310 [M + H - Et - S] ⁺ (55); 338 [M - S] ⁺ (30).
4b ^b	309 [M - Et - S] ⁺ (28); 280 [M - 2Et - S] ⁺ (10), 247 [M - PhCH ₂ - S] ⁺ (100), 218 [M - Et - S - PhCH ₂] ⁺ (51), 123 [M - Et - 3S - PhCH ₂] ⁺ (71).
4c ^a	303 [M + H] ⁺ (44); 274 [M + H - Et] ⁺ (100).
4c ^b	302 [M] ⁺ (5); 273 [M - Et] ⁺ (41), 245 [M - Bu] ⁺ (44), 241 [M - Et - S] ⁺ (22), 213 [M - Bu - S] ⁺ (66).
4d ^a	331 [M + H] ⁺ (52), 274 [M + H - Bu] ⁺ (100), 242 [M + H - Bu - S] ⁺ (39).
4f ^b	330 [M] ⁺ (10), 273 [M - Bu] ⁺ (70).
4j ^a	289 [M + H] ⁺ (100).

^aChemical ionization, 100 eV. ^bElectron impact, 70 eV.

Some assumptions may be suggested concerning the structures of the intermediates and initial products of the interactions of phosphorus sulfide **1** with chloride **2a** in the presence of carbonate **3**. It would seem to be reasonable to assume that the degradation of the cage-like molecule of tetraphosphorus decasulfide is a multi-step process. A nucleophilic attack of the oxygen atom of the K-O fragment of the carbonate **3** upon a phosphorus atom of **1** would be expected to lead to the rupture of the P-S bond and consequent migration of the potassium ion to the sulfur atom neighbouring the P-S fragment. It seems that this proposed mechanism is similar to that suggested by Roesky and Remmers for the reactions of tetraphosphorus decasulfide with trimethylsilyldimethylamine in which a nucleophilic nitrogen atom attacks a phosphorus atom. [24] Finally, a series of consequent transformations involved the elimination of the carbon dioxide molecule and the formation of tripotassium thiophosphate **7** and tripotassium trithiophosphate **6** (Equation (2)).

The interaction of benzyl chloride **2a** with this reaction mixture (Equation (2)) is assumed to result in alkylation of these tripotassium thiophosphate **7** and tripotassium trithiophosphate **6** and/or some precursor intermediates containing P-S-K and P-O-K bonds formed in earlier reaction steps with formation of




reaction of tetraphosphorus decasulfide **1** with potassium carbonate **3** may be attributed to tripotassium thiophosphate **7**. Other small intensity signals (δ_p 128.4 and 91.9 ppm, Table VI) may be attributed to some intermediates with P-S-K and P-O-K bonds. The treatment of this mixture with benzyl chloride **2a** leads to the appearance of the signal of tetrathiophosphate **4a** (δ_p 90.5 ppm).


Use of inorganic salts has been shown to be an effective means of promoting the reactions of phosphorus sulfides with organic compounds. We extended this approach to the alkyl homologues of Davy's reagent. Thus, the 2,4-bis(alkylthio)-2,4-dihydro-1,3,2 λ^5 ,4 λ^5 -dithiadiphosphetanes **8a,b** react with alkyl chlorides **2a,b** in the presence of carbonate **3** in MeCN at 60-70°C for 1-3 h to form tetrathiophosphates **4b-d** with mixed substituents ($R \neq R'$) (Equation (4), Tables (I)-(V)).

The ^{31}P NMR spectra of the products **4b-d** reveal the signals at δ_p 92.1-94.0 ppm. Most of the known tetrathiophosphates appear in the same region. [25] The chemical ionization mass spectrum of tetrathiophosphate **4b** shows the mass peak of molecular ion $[\text{M} + \text{H}]^+$ m/e 371.

In this case we also assumed an initial nucleophilic attack of the oxygen atom of the K-O fragment of carbonate **3** upon a phosphorus atom of dithiadiphosphetanes **8** with subsequent elimination of carbon dioxide. A subsequent symmetrization of the intermediate O,S-dipotassium alkyltrithiophosphates seems to lead to the formation of dipotassium alkylidithiophosphates **9** and dipotassium alkyltetrathiophosphates **10** (Equation (5)).

Similarly interaction of alkyl chlorides **2a,b** with this reaction mixture (Equation (5)) is likely to occur by alkylation of the salts **9** and **10** and/or some intermediates containing P-S-K and P-O-K bonds with the formation of tetrathiophosphates **4b-d** and S-alkyl-O,O'-dialkylidithiophosphates **B**. The latter seems to undergo S,O-exchange processes in accordance with the Pishchimuka








reaction with subsequent exchange of a phosphoryl group to a thiophosphoryl one under the effect of dithiadiphosphetanes **8** with the formation of final tetrathiophosphates **4b-d** as the only final organothiophosphorus products (Equation (6)).

The ^{31}P NMR spectrum of the crude reaction mixture of dithiadiphosphetanes **8** with potassium carbonate **3** in acetonitrile at 20°C after evolution of carbon dioxide shows two signals at δ_{P} 97.1 and 89.5 ppm in the ratio 3:1 (Table VI) which may be attributed to salts **9** and **1**.

We have previously shown that the reaction of dithiadiphosphetanes **8** with dialkyl disulfides proceeds at $90\text{--}100^\circ\text{C}$ for 1-1.5 h to form S,S'-dialkyl S'',S''-alkyltetrathiolothionophosphates with an S-S bridge bond. [6] This reaction was also facilitated when carbonate **3** was employed. Thus, the reaction of dithiadiphosphetanes **8a,b** with disulfides **12a-c** in the presence of an equimolar amount of carbonate **3** occurs in MeCN at $60\text{--}70^\circ\text{C}$ for 1 h (Equation (7), Tables (I) and (II)).


This reaction (Equation (7)) proceeds via the initial formation of pentathiophosphates **13a-c**. Their intermediate formation was confirmed by the ^{31}P NMR spectra of the crude reaction mixtures indicating the signals at δ_{P} (C_6H_6) 98 and 100 ppm (Reference ⁶: δ_{P} (C_6H_{16}) **13a** 98.0 ppm, and **13b** and 100.5 ppm). The ^{13}P NMR spectrum of pentathiophosphate **13c** reveals resonance at δ_{P} (C_6H_6) 100.7 ppm. The chemical ionization mass spectrum of pentathiophosphate **13a**






shows the mass peak of molecular ion $[\text{M} + \text{H}]^+$ m/e 279. There is the mass peak $[\text{M}]^+$ m/e 362 in the electron impact mass spectrum of compound **13b** which may be attributed to its molecular ion. We have previously shown that the penthiophosphates **13a** with S-S thiol bond are thermally unstable compounds. [4] Products **13a-c** formed in the reaction (7) are unstable during distillation as well and tetrathiophosphates **4e-g** were isolated in pure form (distilled or chromatographed) from the reaction mixtures (Tables (I)-(IV)).

We assumed that tetrathiophosphates may be also formed in the reactions of phosphorus sulfides with alkyl halides in the presence of sodium sulfide. Indeed, the reaction of tetraphosphorus decasulfide **1** with benzyl chloride **2a** in the presence of sodium sulfide **15** in anhydrous acetonitrile at 50-60°C for 2 h gave tribenzyl tetrathiophosphate **4a** (Equation (8), and Table (I)).








Thus sodium sulfide has proved to be a more effective promoter as regards phosphorus sulfides. Thus tetrathiophosphate **4b** was also obtained in the reaction of dithiadiphosphetane **8a** with chloride **2a** under mild conditions (20°C, 4.5 h, MeCN) when sodium sulfide **15** was employed (Equation (9), and Table (I)).

We extended this technique to the reactions of phosphorus sulfides (P₄S₁₀, P₄S₇ and P₄S₅) and dithiadiphosphetanes **8** with dialkyl disulfides because these reactions traditionally proceed under severe conditions (100-160°C). [4, 7, 8] Thus, the formation of tetrathiophosphate **4h** occurred already at 20°C for 2 h in the reaction of tetraphosphorus pentasulfide (P₄S₅) **16** in the presence of an equimolar amount of sodium sulfide **15** in MeCN (Equation (10), and Table (I)).

Similarly tetrathiophosphate **4h** was also formed in the reaction of tetraphosphorus heptasulfide (P₄S₇) **18** with disulfide **12c** and sodium sulfide **15** at 20°C for 6 h in MeCN (Equation (11), and Table (I)).

Both reactions (10) and (11) proceed via the intermediate formation of pentathiophosphate **13d** (R = R' = Pr) (δ_P (C₆H₆) 100.2 ppm). Similar formation of pentathiophosphate **13b** was also observed in the reaction of tetraphosphorus decasulfide **1** with disulfide **12b** in the presence of sodium sulfide **15** at 20°C for






6 h in MeCN. Tetrathiophosphate **4f** was isolated from the reaction mixture by Distillation (Equation (12), and Table (I) and (II)).

Tetrathiophosphate **4i** with mixed substituents was isolated in the reaction of dithiadiphosphetane **8a** with disulfide **12c** in the presence of an equimolar amount of sodium sulfide **15** under mild conditions (20°C, 2 h, MeCN) (Equation (13), and Table (I)-(IV)).

We have recorded the ^{31}P NMR spectra of the crude reaction mixtures of phosphorus sulfides and dithiadiphosphetane **8a** with sodium sulfide **15** in MeCN in the absence of organic reagents at 20°C (Table (VI)). These signals may be attributed to some intermediates with P-S-Na bonds. It seems the treatment of this mixture with alkyl halides or disulfides leads to the formation of the final tetrathiophosphates **4**.

In accordance with the stoichiometry of reactions of phosphorus sulfides (P_4S_{10} , P_4S_7 and P_4S_5) and dithiadiphosphetane **8a** with disulfides **12** in the presence of potassium carbonate and sodium sulfide (Equations (7) and (10)-(13)) they should be accompanied by the formation of potassium or sodium alkylthiolates **14** or **17**. It seemed of interest to study the influence of alkaline alkylthiolates on the reactivity of phosphorus sulfides. We carried out the reaction of tetraphosphorus heptasulfide (P_4S_7) **18** with disulfide **12c** in the presence of sodium isopropylthiolate **17b** in MeCN at 20°C for 1 h and obtained tetrathiophosphate **4j** with mixed substituents (Equation (14), Tables (I)-(V)).

The intermediate formation of pentathiophosphate **13e** was confirmed by the ^{31}P NMR spectrum of the crude reaction mixture (Equation (14)) (signal at δ_{p} (C_6H_6) 104.8 ppm) and by the chemical ionization mass spectrum data of the reaction mixture (molecular ion $[\text{M} + \text{H}]^+$ m/e 321). Tetrathiophosphate **4j** has different alkyl substituents (*n*- and *i*-propyl) that have been established by its ^1H NMR spectrum. The $\text{CH}_3\text{CH}_2\text{CH}_2$ protons appear as a triplet at δ 1.02 ppm ($^3J_{\text{H-H}}$



The mixture was stirred at 100°C for 0.5 h. Chloride **2a** (6.9 g, 54.5 mmol) was added dropwise with stirring at 20°C to the mixture and stirring was continued for 2 h at 60-70°C. The mixture was filtered. The precipitate was extracted with benzene. The extractions and filtrate were combined and evaporated at reduced pressure (0.04 mm Hg) at 70°C for 5 h and gave crude tetrathiophosphate **4a** (2.0 g, 26%). The residue was diluted with 50 mL of benzene and washed with water. The organic layer was dried (MgSO₄) and filtered. The filtrate was evaporated under vacuum (0.1 and 0.02 mm Hg) at 50°C for 2 h. The residue was chromatographed on Al₂O₃ column with benzene as eluant and gave pure **4a** (1.0 g, 13%) (See Table I-V).

Similarly dithiadiphosphetane **8a** (3.0 g, 9.6 mmol), chloride **2a** (4.9 g, 38.7 mmol) and carbonate **3** (2.7 g, 19.5 mmol) (reaction conditions: 60-70°C, 3 h, 10 mL MeCN) gave crude S-ethyl S',S''-dibenzyltetrathiophosphate **4b** (2.0 g, 28%). Pure **4b** (0.7 g, 10%) was isolated from the residue by means of column chromatography (Al₂O₃, benzene).

Similarly dithiadiphosphetane **8a** (2.5 g, 8.0 mmol), bromide **2b** (4.4 g, 32.1 mmol) and carbonate **3** (2.2 g, 15.9 mmol) (reaction conditions: 60-70°C, 1 h, 10 mL MeCN) gave crude S-ethyl S',S''-dibutyltetrathiophosphate **4c** (1.7 g, 35%) and pure (chromatographed) **4c** (0.8 g, 17%).

Similarly dithiadiphosphetane **8b** (5.2 g, 16.7 mmol), bromide **2b** (9.1 g, 66.4 mmol) and carbonate **3** (4.6 g, 33.3 mmol) (reaction conditions: 60°C, 1 h, 10 mL MeCN) gave crude S,S'-dibutyl-S''-isobutyltetrathiophosphate **4d** (3.0 g, 27%).

Similarly dithiadiphosphetane **8a** (4.9 g, 15.7 mmol), disulfide **12a** (5.8 g, 47.5 mmol) and carbonate **3** (2.2 g, 15.9 mmol) (reaction conditions: 70°C, 1 h, 10 mL MeCN) gave distilled triethyltetrathiophosphate **4e** (2.0 g, 26%), b.p. 110°C (0.03 mm Hg), <BOLDn</BOLD_B²⁰ 1.6207. The ³¹P NMR (C₆H₆) δ_p: 91.7 ppm. The mass spectrum (EI, 70 eV), m/e (I_{rel.}, %): 246 [M]⁺. (28) (Reference ^[26]: b.p. 124-125°C (1.5 mm Hg), <BOLDn</BOLD_B²⁰ 1.6201. The ³¹P NMR (δ_p: 91.7 ppm).

Similarly dithiadiphosphetane **8b** (3.2 g, 8.7 mmol), disulfide **12b** (4.6 g, 25.8 mmol) and carbonate **3** (1.2 g, 8.7 mmol) (reaction conditions: 60°C, 1 h, 10 mL MeCN) gave distilled triisobutyltetrathiophosphate **4f** (1.9 g, 33%), b.p. 130°C (0.03 mm Hg), <BOLDn</BOLD_B²⁰ 1.5559. The ³¹P NMR (C₆H₆) δ_p: 93.9 ppm. The mass spectrum (EI, 70 eV), m/e (I_{rel.}, %): 330 [M]⁺. (43) (Reference ^[41]: b.p. 136-137°C (0.02 mm Hg), <BOLDn</BOLD_B²⁰ 1.5549. The ³¹P NMR (C₆H₆) δ_p: 92.8 ppm).

Similarly dithiadiphosphetane **8b** (2.0 g, 5.4 mmol), disulfide **12c** (2.4 g, 16.0 mmol) and carbonate **3** (0.8 g, 5.8 mmol) (reaction conditions: 70°C, 1 h, 10 mL MeCN) gave crude S,S'-dipropyl-S''-isobutyltetrathiophosphate **4g** (2.0 g, 61%)

which was chromatographed (Al_2O_3 column, benzene) and gave pure **4g** (0.8 g, 24%).

Reaction of Tetraphosphorus Decasulfide **1** with Benzyl Chloride **2a** in the Presence of sodium sulfide **15**. Typical Procedure

Dried sulfide **15** (2.4 g, 30.8 mmol) was added portionwise to a stirred suspension of **1** (2.3 g, 5.2 mmol) in 10 mL of anhydrous acetonitrile at 20°C. Chloride **2a** (7.9 g, 62.4 mmol) was added dropwise with stirring at 20°C to the mixture and stirring was continued for 2 h at 50-60°C. The mixture was diluted with 40 mL of MeCN and washed with water. The organic layer was dried (MgSO_4) and filtered. The filtrate evaporated under vacuum (0.1 and 0.07 mm Hg) at 50°C for 2 h and gave crude tetrathiophosphate **4a** (4.9 g, 54%). The residue was chromatographed on Al_2O_3 column with benzene as eluant and gave pure **4a** (2.1 g, 23%) (See Tables I-V).

Similarly dithiadiphosphetane **8a** (4.5 g, 14.4 mmol), chloride **2a** (7.3 g, 57.7 mmol) and sulfide **15** (2.3 g, 29.5 mmol) (reaction conditions: 20°C, 4.5 h, 15 mL MeCN) gave crude tetrathiophosphate **4b** (2.2 g, 21%). Distillation of the residue gave pure **4b** (1.0 g, 9%), b.p. 195-198°C (0.07 mm Hg) (See Tables I-V).

Similarly tetraphosphorus pentasulfide (P_4S_5) **16** (2.7 g, 9.5 mmol), disulfide **12c** (10.0 g, 66.6 mmol) and sulfide **15** (0.74 g, 9.5 mmol) (reaction conditions: 20°C, 2 h, 10 mL MeCN) gave crude tetrathiophosphate **4h** (5.1 g, 46%) and distilled **4h** (3.5 g, 32%), b.p. 128-130°C (0.04 mm Hg), δ_{P}^0 1.5892. The ^{31}P NMR (C_6H_6) δ_{P} : 92.0 ppm. The mass spectrum (CI, 100 eV), m/e (I_{rel} , %): 289 [$\text{M} + \text{H}$] $^+$ (100) (Reference 114): b.p. 131-132°C (0.5 mm Hg), δ_{P}^0 1.5885. The ^{31}P NMR δ_{P} : 92.5 ppm).

Similarly tetraphosphorus heptasulfide (P_4S_7) **18** (4.5 g, 12.9 mmol), disulfide **12c** (13.6 g, 90.6 mmol) and sulfide **15** (1.0 g, 12.8 mmol) (reaction conditions: 20°C, 6 h, 10 mL MeCN) gave crude tetrathiophosphate **4h** (3.3 g, 22%) and distilled **4h** (2.8 g, 19%).

Similarly dithiadiphosphetane **8a** (5.2 g, 16.7 mmol), disulfide **12c** (7.5 g, 50.0 mmol) and sulfide **15** (1.3 g, 16.7 mmol) (reaction conditions: 20°C, 2 h, 10 mL MeCN) gave distilled tetrathiophosphate **4i** (2.3 g, 25%) (See Tables I-IV).

Reaction of Tetraphosphorus heptasulfide **18** with Dipropyl Disulfide **12c** in the Presence of sodium Isopropylthiolate **17b**

Dried thiolate **17b** (3.5 g, 35.7 mmol) was added portionwise to a stirred suspension of **18** (3.1 g, 8.9 mmol) in 10 mL of anhydrous acetonitrile at 20°C. Disulfide **12c** (8.0 g, 53.3 mmol) was added dropwise with stirring at 20°C to the

mixture and stirring was continued for 1 h at 20°C. The mixture was washed with water. The organic layer was dried (MgSO₄) and evaporated under vacuum (0.1 and 0.03 mm Hg) at 50°C for 3 h and gave crude tetrathiosphosphate **4j** (1.0 g, 10%). Distillation of the residue gave pure **4j** (0.7 g, 7%) (See Tables I-V).

References

- [1] A. D. F. Toy and E. H. Uhing, *US Pat.*, 4076746, 1974; *Chem. Abstr.*, **89**, 43771g (1978).
- [2] E. H. Uhing and A. D. F. Toy, *US Pat.*, 3879500, 1973; *Chem. Abstr.*, **83**, 27554j (1978).
- [3] I. S. Nizamov, E. S. Batyeva, V. A. Al'fonsov, R. Z. Musin and A. N. Pudovik, *Phosphorus, Sulfur, and Silicon*, **55**, 229 (1991).
- [4] I. S. Nizamov, L. A. Al'metkina, G. G. Garifzyanova, E. S. Batyeva, V. A. Al'fonsov and A. N. Pudovik, *Phosphorus, Sulfur, and Silicon*, **83**, 191 (1993).
- [5] A. A. El-Barbary, *Monatsh. Chem.*, **115**, 769 (1984).
- [6] I. S. Nizamov, L. A. Al'metkina, E. S. Batyeva, V. A. Al'fonsov and A. N. Pudovik, *Phosphorus, Sulfur, and Silicon*, **72**, 229 (1992).
- [7] I. S. Nizamov, G. G. Garifzyanova, L. A. Al'metkina, G. G. Sergeenko and E. S. Batyeva, *Phosphorus, Sulfur, and Silicon*, **97**, 173 (1994).
- [8] I. S. Nizamov, L. A. Al'metkina, G. G. Garifzyanova, G. G. Sergeenko and E. S. Batyeva, *Phosphorus, Sulfur, and Silicon*, **102**, 71 (1995).
- [9] F. H. Musa, B. W. Tattershall and W. Bruce, *J. Chem. Soc., Dalton Trans.*, 1517 (1984).
- [10] B. W. Tattershall, *J. Chem. Soc., Dalton Trans.*, 1707 (1985).
- [11] I. S. Nizamov, G. G. Garifzyanova, and E. S. Batyeva, *Phosphorus, Sulfur, and Silicon*, **88**, 39 (1994).
- [12] J. W. Scheeren, P. H. J. Ooms and R. J. F. Nivard, *Synthesis*, 149 (1973).
- [13] K. I. Pashkevich, V. I. Saloutin, M. B. Bobrov and D. S. Yufit, *Zh. Org. Khim. (Russ.)*, **24**, 2397 (1988).
- [14] D. Brillon, *Synth. Commun.*, **20**, 3085 (1990).
- [15] B. C. Stahly, *US Pat.*, 4935510, 1990; *Chem. Abst.*, **113**, 191408y (1990).
- [16] B. C. Stahly, *US Pat.*, 4935513, 1989; *Chem. Abst.*, **113**, 212030m (1990).
- [17] H. W. Roesky, *US Pat.*, 3533736, 1966; *Chem. Abst.*, **74**, 5141j (1970).
- [18] G. Schrader, *Ger. Pat.*, 1104506, 1961; *Chem. Abst.*, **56**, 1482h (1962).
- [19] G. Schrader, *Ger. Pat.*, 1124946, 1962; *Chem. Abst.*, **58**, 5724g (1963).
- [20] G. Schrader, *Ger. Pat.*, 1129954, 1962; *Chem. Abst.*, **58**, 5724h (1963).
- [21] H. Hoffman and G. Schumacher, *Tetrahedron Lett.*, 2963 (1967).
- [22] R. Shabana, *Phosphorus Sulfur*, **29**, 293 (1987).
- [23] M. Demarq, *J. Chem. Soc., Dalton Trans.*, 2221 (1988).
- [24] H. W. Roesky and G. Remmers, *Z. anorg. allg. Chem.*, **431**, 221 (1977).
- [25] M. Grayson, E. J. Griffith (eds): *Topics on Phosphorus Chemistry. P³¹ Nuclear Magnetic Resonance*, Interscience Publishers, a Division of John Wiley and Sons, New York, London, Sydney, vol. **5** (1967).
- [26] D. E. Ailman and R. J. Magee, in "Organic Phosphorus Compounds" (Eds. G. M. Kosolapoff and L. Maier (John Wiley and Sons: New York, London, Sydney, Toronto, 1976), Vol. 7, p. 791.