This article was downloaded by: [North Dakota State University] On: 21 April 2014, At: 17:05 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

A Facile One-Pot Microwave-Induced Synthesis of Some Novel Selenolo[2,3-b]quinoline Derivatives under Solvent-Free Conditions

M. Raghavendra^a, Halehatty S. Bhojya Naik^a & Bailure S. Sherigara^a

^a Department of P G Studies and Research in Industrial Chemistry, School of Chemical Sciences, Kuvempu University Shankaraghatta, Karnataka, India

Published online: 04 Jun 2008.

To cite this article: M. Raghavendra , Halehatty S. Bhojya Naik & Bailure S. Sherigara (2008) A Facile One-Pot Microwave-Induced Synthesis of Some Novel Selenolo[2,3-b]quinoline Derivatives under Solvent-Free Conditions, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:6, 1501-1509

To link to this article: http://dx.doi.org/10.1080/10426500701690855

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the

Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



A Facile One-Pot Microwave-Induced Synthesis of Some Novel Selenolo[2,3-*b*]quinoline Derivatives under Solvent-Free Conditions

M. Raghavendra, Halehatty S. Bhojya Naik, and Bailure S. Sherigara

Department of P G Studies and Research in Industrial Chemistry, School of Chemical Sciences, Kuvempu University Shankaraghatta, Karnataka, India

A rapid, solvent free microwave assisted synthetic strategy has been developed for the synthesis of some selenolo[2,3-b]quinoline derivatives using potassium carbonate by the reaction between 2-seleno-3-formyl-quinolines (2a-d) with 2chloroacetamide and phenacylbromide. This method found to be very effective, products obtained in high yield, isolation is just treating the reaction mixture with water.

 ${\bf Keywords}~{\rm Microwave}$ irradiation; quinolines; selenolo
[2,3-b]quinolines; solvent-free conditions

INTRODUCTION

The use of microwave for the synthesis of organic compounds under solvent-free conditions proved to be efficient safe and environmentally benign techniques that with shorter reaction time, high yields, and easier manipulation. Additionally, it can also avoid the use of poisonous and expensive solvent, and as such can be environmentally benign, and make manipulations much easier.^{1–2}

Several polycyclic analogues of natural or synthetic antitumor agents are well known, and have attracted considerable interest because of their significant anticancer activity.³ There is a evidence that the antitumor activity is due to these compounds intercalation between the

Address correspondence to Halehatty S. Bhojya Naik, Department of P G Studies and Research in Industrial Chemistry, School of Chemical Sciences, Kuvempu University, Shankaraghatta, Karnataka, India. E mail: hsb.naik@rediffmail.com

Received 17 January 2007; accepted 8 August 2007.

M. Raghavendra is thankful to the Chairman, Kuvempu University, Shankaraghatta, for providing research facilities and the Head of the Sophisticated Instruments Facility, IISc, Bangalore, for spectral data.

base pairs of DNA and interference with normal functioning of the enzyme topoisomerase II, which is involved in the breaking and releasing of DNA strands.⁴ The intercalative binding of these drugs is due to the presence of planar linearly fused tri and tetra cyclic systems.

Further, the biological and pharmaceutical activities of different selenium compounds are gains special interest because it has been associated with modification of metal toxicity⁵ and with prevention of cancer.⁶ In addition to naturally occurring selenium compounds, certain synthetic organoselenium compounds such as *p*-methoxybenzeneselenol, benzylselenocyanate, 1,4-phenylenebisselenocyanate were found to inhibit azoxymethane-induced hepatocarcinogenesis in female F 344 rats without clinical signs of toxicity,^{7–9}*p*-methoxybenzyl selenocyanate and 1, 4-phenylenebis (methylene) selenocyanate prevented both precancerous cell growth and tumor growth in animals after being treated with a colorectal cancer-inducing agent with no side effects.¹⁰ Moreover, some patented phenylaminoalkyl selenides confirmed antihypertensive properties.

On the other hand, five- and six-membered heterocyclic compounds containing one or two heteroatom fused to quinoline ring in a linear fashion are found in natural products, as well as in synthetic compounds of biological interest have antitumor and anticancer properties.^{11,12} They are also known to exhibit antiallergenic,¹³ antifungal,¹⁴ hypocholesterolemic, hypolemic,¹⁵ antibacterial,¹⁶ and antiviral¹⁷ properties. In addition, various condensed quinolines were studied for their intercalative DNA binding, cytotoxic, and antitumor activities.¹⁸

Due to their great biological properties, compounds containing the quinoline system have been the subject of many synthetic studies. In these directions, there has been increasing interest in the synthesis of quinoline containing Se moieties to obtain improved chemoprevention with lower toxic activity. So, in continuation of our studies on synthesis of condensed quinolines¹⁹ by taking the advantage of microwave, we wish to report on synthesis of novel selenolo[2,3-*b*]quinoline derivatives under solvent free microwave irradiation conditions.

RESULTS AND DISCUSSION

The starting compounds 2-chloro-3-formylquinoline (1a-d), were prepared according to literature method.²⁰ The compounds 2-seleno-3formylquinoline (2a-d) were prepared by the action of sodium hydroselenide on 2-chloro-3-formylquinolines (1a-d) (Scheme 1).

A wide range of substituted selenolo[2,3-*b*]quinolines such as (**3a-d**), and (**4a-d**) were prepared by treating (**2a-d**) with phenacylbromide and 2-chloroacetamide under solvent free microwave irradiation conditions



SCHEME 1

in one pot to furnished the title compounds in good yields as shown in (Scheme 2). The reaction was known²¹ to proceed through replacing halogen of the reactants by selenium moiety, the carbanion results in the intermediate by base, adds on the aldehydic carbon with simultaneous elimination of water resulting in the formation of products with good yields. The structural elucidation of all the newly synthesized compounds was established based on their IR, ¹H NMR and mass spectral data. (which are presented in the experimental section).

As an example, the IR spectrum of (3a) showed an absence of tautomeric SeH, NH, groups and CHO stretching frequency in the region of 3150–3300, and 1650 cm⁻¹, which appeared in the 2-seleno-3-formylquinoline (2a) were found to be absent in the IR spectrum of (3a). The ¹H NMR spectrum of 2a exhibits singlet at δ 5.8 ppm corresponding to tautomeric SeH, NH, group and singlet at δ 10.35 ppm corresponding to -CHO group, the singlet in the intermediate near at δ 4.2 ppm corresponding to an unreacted –SeCH₂ group were found to be absent in (**3a**) indicates the attachment of the reactive partner to quinoline moiety. Further, compound (**3a**) showed signal at δ 8.50 for C4-H proton and signal at δ 8.62 for C3-H proton, this might be due to hydrogen



bonding and anisotropy effect of the adjacent carbonyl groups. The other aromatic protons resonated at δ 7.50–8.16 (4H). Finally, the structure assigned was confirmed by its mass spectrum through the appearance of a molecular ion peak at m/z 334 [M⁺]. The melting points, percentage of yield and elemental data were shown in Table I.

MATERIALS AND METHODS

Melting points were determined in open capillaries and are uncorrected. The FT-IR spectra were recorded on NICOLETAVATAR 360-FTIR instrument by using KBr pellets. The ¹H NMR were recorded on a BRUCKER AMX-400 spectrometer operating at 400 MHz. Mass spectra were recorded on AGILENT LC-MSD-TRAP-XCT mass spectrometer

Compound no.	Microwave irradiation/Reflux				Elemental analysis Calcd./Found (%)		
	M.P. (°C)	Yield (%)	Time/hr/min	Mol. formula	С	н	N
2a	255	91	1hour	C ₁₀ H ₇ NOSe	50.84	2.96	5.93
					50.80	3.00	5.99
2b	247	90	1hour	$C_{11}H_9N0Se$	52.81	3.63	5.60
				11 0	52.81	3.63	3.63
2c	266	92	1hour	C ₁₁ H ₉ N0 ₂ Se	49.64	3.41	5.26
				11 0 1	49.64	3.43	5.28
2d	240	90	1hour	C ₁₁ H ₉ N0Se	52.81	3.63	5.60
				11 0	52.81	3.63	5.60
3a	235	89	7	C ₁₈ H ₁₁ NOSe	64.28	3.27	4.16
				10 11	64.30	3.24	4.12
3b	208	88	8	C ₁₉ H ₁₃ NOSe	65.14	3.71	4.00
				10 10	65.18	3.68	4.06
3c	231	85	7	$C_{19}H_{13}NO_2Se$	62.29	3.55	3.82
				10 10 1	62.32	3.52	3.86
3d	208	88	8	C ₁₉ H ₁₃ NOSe	65.14	3.71	4.00
					65.20	3.73	3.96
4a	237	89	8	$C_{12}H_8N_2OSe$	52.36	2.90	10.18
					52.39	2.94	10.20
4b	217	90	8	$C_{13}H_{10}N_2OSe$	53.97	3.46	9.68
				10 10 2	53.94	3.42	9.64
4c	229	88	8	$C_{13}H_{10}N_2O_2Se$	51.14	3.27	9.18
					51.16	3.30	9.22
4 d	225	90	8	$C_{13}H_{10}N_2OSe$	53.97	3.46	9.68
				10 10 2	53.94	3.42	9.64

 TABLE I Characterization Data of the Newly Synthesized

 Compounds

Elemental analyses were done on Vario EL. CHNOS elemental analyzer.

General Method for the Preparation of 2-Seleno-3-formylquinoline (2a)

Solution of sodium hydrogen selenide [freshly prepared from finely divided selenium powder (1 g, 0.013 mol) and sodium borohydride (1 g, 0.026 mol) in water (50 ml)],²²was added to 2-chloro-3-formylquinoline **(1a)** (1.91 g, 0.01 mol) in ethanol (80 ml). The reaction mixture was refluxed on water bath for one hour, cooled, poured to ice water (100 ml) and neutralized with dilute hydrochloric acid (4N) the resultant solid,

2-seleno-3-formyl quinoline obtained was filtered, washed with water (300 ml) dried and recrystallized from alcohol.

Yield: IR (KBr) cm⁻¹ 3150–3300 (tautomeric form of -NH, SeH), 1650 (CHO) 1180 (C=Se). ¹H NMR (400 MHz, DMSO-d₆) 5.8 (tautomeric form of -NH, SeH), 10.35 (-CHO), 7.25–8.35 (m, 5H, Ar-H). MS m/z: 236 (M⁺).

2-Seleno-6-methylquinoline-3-carbaldehyde (2b)

Prepared from methyl derivative of quinoline 1b.

IR (KBr) cm⁻¹ 3155–3310 (tautomeric form of -NH, SeH), 1650 (CHO) 1180 (C=Se). ¹H NMR (400 MHz, DMSO-d₆) 2.60 (3H, s, CH₃), 5.85 (tautomeric form of -NH, SeH), 10.35 (-CHO) 7.20–8.30 (m, 4H, Ar-H). MS m/z: 250 (M⁺).

2-Seleno-6-methoxy-3-carbaldehyde (2c)

Prepared from methoxy derivative of quinoline 1c.

IR (KBr) cm⁻¹ 3160–3300 tautomeric form of (–NH, SeH) 1650 (CHO) 1180 (C=Se). ¹H NMR (400 MHz, DMSO-d₆) 3.95 (3H, s, OCH₃), 5.90 (tautomeric form of –NH, SeH), 10.40 (–CHO) 7.35–8.40 (m, 4H, Ar–H). MS, m/z 266 (M⁺).

2-Seleno-7-methylquinoline-3-carbaldehyde (2d)

Prepared from methyl derivative of quinoline 1d.

IR (KBr) cm⁻¹ 3155–3310 (tautomeric form of –NH, SeH), 1650 (CHO) 1182 (C=Se). ¹H NMR (400 MHz, DMSO-d₆) 2.65 (3H, s, CH₃), 5.85 (tautomeric form of –NH, SeH), 10.35 (–CHO) 7.25–8.37 (m, 4H, Ar–H). MS m/z: 250 (M⁺).

General Procedure: Synthesis 1-Selenolo[2,3-b]quinolin-2-ylethanone

A mixture of 2a (2.360 g, 10 mmol), phenacyl bromide (1.99 g, 10 mmol) and potassium carbonate (1.380 g, 20 mmol) were ground for uniform mixing. The mixture was then subjected to microwave radiation in a domestic microwave oven for 8 min at an interval of 1 min at 160 W as required to complete the reaction (TLC). The reaction mixture was then poured into water, stirred, filtered and dried. The crude product was recrystallized from aqueous DMF to give 2.990 g (89%) of **3a**. All the other compounds were prepared in a similar way with 85–88% yield.

Selenolo[2,3-*b*]quinolin-2-yl(phenyl)methanone (3a)

Prepared according to the general method of preparation.

Solid; (89%), MS, m/z 336 (M+); IR (KBr) (cm⁻¹): 1640 (CO), ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 8.52 (s, C₃-H), 8.25 (s, C₄-H), 7.50–8.16 (9H, m, Ar-H).

(6-Methylselenolo[2,3-b]quinolin-2-yl)(phenyl)methanone (3b)

Prepared according to the general method of preparation.

Solid; (88%), MS, m/z 350 (M+); IR (KBr) (cm⁻¹): 1643 (CO), ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 2.45 (3H, s, Ar-CH₃), 8.58 (s, C₃-H), 8.24 (s, C₄-H), 7.55–8.13 (8H, m, Ar-H).

(6-Methoxyselenolo[2,3-*b*]quinolin-2-yl)(phenyl)methanone (3c)

Prepared according to the general method of preparation.

Solid; (85%), MS, m/z 366 (M+); IR (KBr) (cm⁻¹): 1640 (CO), ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 3.95 (3H, s, Ar-OCH₃), 8.60 (s, C₃-H), 8.24 (s, C₄-H), 7.58–8.12 (8H, m, Ar-H).

(6-Methylselenolo[2,3-b]quinolin-2-yl)(phenyl)methanone (3d)

Prepared according to the general method of preparation.

Solid; (88%), MS, m/z 350 (M+); IR (KBr) (cm⁻¹): 1645 (CO), ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 2.47 (3H, s, Ar-CH₃), 8.58 (s, C₃-H), 8.26 (s, C₄-H), 7.55–8.15 (8H, m, Ar-H).

Selenolo[2,3-b]quinoline-2-carboxamide (4a)

Prepared according to the general method of preparation.

Solid; (89%), MS, m/z 275 (M+); IR (KBr) (cm⁻¹): 3350-3175 (CONH₂), 1630 (CO). ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 7.42 (2H, b s, CONH₂), 8.50 (s, C₃-H), 8.20 (s, C₄-H), 7.54–8.05 (4H, m, Ar-H).

6-Methylselenolo[2,3-b]quinoline-2-carboxamide (4b)

Prepared according to the general method of preparation.

Solid; (90%), MS, m/z 289 (M+); IR (KBr) (cm⁻¹): 3360–3185 (CONH₂), 1625 (CO). ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 2.40

Downloaded by [North Dakota State University] at 17:05 21 April 2014

(3H, s, Ar-CH₃), 7.42(2H, b s, CONH₂) 8.56 (s, C₃-H), 8.25 (s, C₄-H), 7.62– 8.12 (3H, m, Ar-H).

6-Methoxyselenolo[2,3-b]quinoline-2-carboxamide (4c)

Prepared according to the general method of preparation.

Solid; (88%), MS, m/z 305 (M+); IR (KBr) (cm⁻¹): 3360–3190 (CONH₂), 1625 (CO). ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 3.90 (3H, s, Ar-OCH₃), 7.45(2H, b s, CONH₂) 8.53 (s, C₃-H), 8.26 (s, C₄-H), 7.65–8.14 (3H, m, Ar-H).

6-Methylselenolo[2,3-b]quinoline-2-carboxamide (4d)

Prepared according to the general method of preparation.

Solid; (90%), MS, m/z 289 (M+); IR (KBr) (cm⁻¹): 3370–3195 (CONH₂), 1630 (CO). ¹H NMR (400 MHz) (DMSO-d6) (δ) ppm: 2.45 (3H, s, Ar-CH₃), 7.40(2H, b s, CONH₂) 8.60 (s, C₃-H), 8.25 (s, C₄-H), 7.62–8.15 (3H, m, Ar-H).

REFERENCES

- (a) S. Caddick, *Tetrahedron.* 51, 10403 (1995); (b) A. R. Katritzky and K. S. Sandeep, *Arkivoc.*, (xiii), 68 (2003).
- [2] C. B. Khour, C. B. Darth, A. Lalnger, and M. E. Daves, J. Catal., 149, 195 (1994).
- [3] (a) W. D. Kingsbury, J. C. Boehm, D. R. Jakas, K. G. Holden, S. M. Hecht, G. Gallagher, M. J. Caranfa, F. L. McCabe, L. F. Faucette, R. K. Johnson, and R. P. Hertzberg, J. Med. Chem., 34, 98 (1991); (b) M. C. Woni, A. W. Nicholas, G. Manikumar, and M. E. Wall, J. Med. Chem., 30, 1774 (1987); (c) M. J. Luzzio, J. M. Besterman, D. L. Emerson, M. G. Evans, K. Lackey, P. L. Leitner, G. McIntyre, B. Morton, P. L. Myers, M. Peel, J. M. Sisco, D. D. Sternbach, W. Q. Tong, A. Truesdale, D. E. Uehling, A. Vuong, and J. Yates, J. Med. Chem. 38, 395 (1995); (d) M. E. Wall, M. C. Wani, A. W. Nicholas, G. Manikumar, C. Tele, A. Moore, D. Truesdale, P. Leitner, and J. M. Besterman, J. Med. Chem., 36, 2689 (1993); (e) G. W. Gribble, Advances in Heterocyclic Natural Products Synthesis., 1, 43 (1990).
- [4] B. Gatto, G. Capranico, and M. Palumbo, Curr. Pharacol. Design, 5, 195 (1999).
- [5] L. Magos and M. Webb, Crit. Rev. Toxicol., 8, 1 (1980).
- [6] L. Fishbein, Toxicology and Environmental Chemistry., 12, 1 (1986).
- [7] T. Tanaka, B. S. Reddy, and K. El-Bayoumy, Japanese J. Cancer Research., 76, 462 (1985).
- [8] K. El-Bayoumy, Cancer Res., 45, 3631 (1985).
- [9] B. S. Reddy, T. Tanaka, and K. El-Bayoumy, J. Natl. Cancer Inst., 74, 1325 (1985).
- [10] I. Jibril, E. Abd Alhadi, and Z. Hamadah, Trans Metals in Chemistry., 25, 407 (2000).
- [11] H. Thomys, Ber. Deut. Pharm. Ges., 33, 68 (1923).
- [12] P. R. Loaiza, A. Quintero, R. Rodríguez-Sotres, J. D. Solano, and A. L. Rocha, Eur. J. Med. Chem., 39, 5 (2004).
- [13] T. H. Althuis, S. B. Khadin, L. J. Czuba, P. F. Moore, and H. J. Hess, J. Med. Chem., 23, 262 (1980).

- [14] J. B. Jiang and D. Isaacson.U. S. Patent., 4,656,274 (1987). Chem. Abstr., 107, 39643 (1987).
- [15] R. E. Graeve, J. R. Pociask, and R. G. Stein, U. S. Patent. 3,600,393 (1971).
- [16] A. M. Farghaly, N. S. Habib, M. A. Khalil, O. A. El-Sayed, and A. Alaxandria, J. Pharm. Sci., 3, 90 (1989). Chem. Abstr., 112, 7420 (1990).
- [17] V. Zikan, S. Radl, F. Smejkal, and D. Zelena. Czech. Patent., 233,445 (1986); Chem. Abstr., 106, 138,447 (1987).
- [18] (a) M. Gopal, S. Veeranna, and L. S. Doddamani. Spect Lett., 37, 347 (2004); (b) V. Nithyadevi and S. P. Rajendran, Phosphorus, Sulfur, and Silicon, and their related elem., 180, 1849 (2005); (c) M. Gopal, M. S. Shahabuddin, and S. R. Inamdar, Proc. Indian Acad. Sci. (Chem. Sci)., 114, 687 (2002); (d) M. Gopal, S. Shenoy, and L. S. Doddamani, J. Photochem. Photobiol., 72, 69 (2003); (e) G. J. Atwell, B. F. Cain, and R. N. Seelye, J. Med. Chem., 15, 611 (1972); (f) B. F. Cain, R. N. Seelye, and G. J. Atwell, J. Med. Chem., 17, 922 (1974); (g) B. F. Cain, G. J. Atwell, and W. A. Denny, J. Med. Chem., 18, 1110 (1975); (h) G. W. Rewcastle, G. J. Atwell, D. Chambers, B. C. Gaguley, and W. A. Denny, J. Med. Chem., 29, 472 (1986).
- [19] (a) B. P. Nandeshwarappa, D. B. Arun Kumar, H. S. Bhojya Naik, V. P. Vaidya, and K. M. Mahadevan, J. Sulfur Chem., 26, 373 (2005); (b) B. P. Nandeshwarappa, D. B. Arun Kumar, H. S. Bhojya Naik, V. P. Vaidya, and K. M. Mahadevan, Indian J. Chem., 44B, 2155 (2005); (c) M. Raghavendra, H. S. Bhojya Naik, and B. S. Sherigara, J. Sulfur Chem., 27, 347 (2006); (d) H. S. Bhojya Naik, M. S. Ramesh, B. V. Swetha, and T. R. Roopa, Phosphorous, Sulfur, Silicon and their related elem., 181, 533 (2006).
- [20] O. Meth-Cohn, B. Narine, and B. Tarnowski, J. Chem. Soc. Perkin Trans., 1, 1520 (1981).
- [21] V. P. Litvinov, V.Yu. Mortikov, Yu. A. Sharanin, and A. M. Shestopalov, Synthesis., 98 (1985).
- [22] D. L. Klayman and T. S. Griffin, J. Am. Chem. Soc., 95, 197 (1973).