# Journal of Medicinal Chemistry



# Identification of *N*-[(5-{[(4-Methylphenyl)sulfonyl]amino}-3-(trifluoroacetyl)-1*H*-indol-1-yl)acetyl]-L-leucine (NTRC-824), a Neurotensin-like Nonpeptide Compound Selective for the Neurotensin Receptor Type 2

James B. Thomas,<sup>\*,†</sup> Angela M. Giddings,<sup>†</sup> Robert W. Wiethe,<sup>†</sup> Srinivas Olepu,<sup>†</sup> Keith R. Warner,<sup>†</sup> Philippe Sarret,<sup>‡</sup> Louis Gendron,<sup>‡</sup> Jean-Michel Longpre,<sup>‡</sup> Yanan Zhang,<sup>†</sup> Scott P. Runyon,<sup>†</sup> and Brian P. Gilmour<sup>†</sup>

<sup>†</sup>Center for Organic and Medicinal Chemistry, Research Triangle Institute, P.O. Box 12194, 3040 Cornwallis Road, Research Triangle Park, North Carolina 27709, United States

<sup>‡</sup>Department of Physiology and Biophysics, Faculty of Medicine and Health Sciences, Université de Sherbrooke, 3001, 12th Avenue North, Sherbrooke, Quebec J1H 5N4, Canada

# **Supporting Information**

**ABSTRACT:** Compounds acting via the neurotensin receptor type 2 (NTS2) are known to be active in animal models of acute and chronic pain. To identify novel NTS2 selective analgesics, we searched for NTS2 selective nonpeptide compounds using a FLIPR assay and identified the title compound (NTRC-824, **5**) that, to our knowledge, is the first nonpeptide that is selective for NTS2 versus NTS1 and behaves like the endogenous ligand neurotensin in the functional assay.



# ■ INTRODUCTION

The discovery of novel analgesics has been a major pursuit of medicinal chemists for more than a hundred years. Although opiates and opioids remain the treatment of choice for severe acute pain, they produce serious side effects including addiction, respiratory depression, and tolerance. Furthermore, opioids often fail to provide relief from chronic pain, a persistent form of pain resulting from nerve injury. Thus, identification of nonopioid analgesics remains a key challenge of discovery science.

The tridecapeptide neurotensin (NT, Glu-Leu-Tyr-Glu-Asn-Lys-Pro-Arg-Arg-Pro-Tyr-Ile-Leu), acting via two G-protein coupled receptors NTS1 and NTS2 (NTR1, NTR2, etc.) and the multiligand type-1 receptor sortilin, mediates many biological functions including nonopioid analgesia.<sup>1–5</sup> Compounds acting via neurotensin receptor types 1 and 2 have been reported to be active in animal models of both acute and chronic pain.<sup>6–11</sup> NT mediated analgesia is also synergistic, with opioid analgesia suggesting that NT-based compounds could function alone or as adjuncts to opioids in the management of pain.<sup>12</sup> Together, these findings underpin the rational for identification of NT-based analgesics.

The search for such compounds is decades old, yet to date, nearly all of the NT compounds reported to be active in animal models of pain are peptides, variants of the terminal hexapeptide fragment of NT (NT(8–13), **1a**, Chart 1).<sup>12–16</sup> The vast majority of these compounds produce analgesia that is accompanied by hypothermia and hypotension, a feature ascribed to interaction with the NTS1 receptor.<sup>17,18</sup> Reports

from the study of the NTS2 selective peptide NT79 (1b) support this notion as it possessed activity against visceral pain but lacked the side effects described above.<sup>12,19</sup>

Surprisingly, only one nonpeptide compound has so far been described to possess NT-based analgesic properties, the NTS2 versus NTS1 selective compound levocabastine (2), which shows activity in both visceral and chronic pain models.<sup>10,20,21</sup> Coupled with our desire to identify nonpeptide compounds, the lower side effect profile demonstrated by the NTS2 selective peptide **1b** prompted us to develop methods of identifying NTS2 selective compounds.<sup>22</sup>

Literature reports with a CHO cell line stably expressing rNTS2 indicated that the pyrazole compound **3a** was an agonist in the FLIPR assay and that NT was an antagonist of the calcium release mediated by **3a**.<sup>23</sup> This suggested that we could identify NT-like (antagonist) compounds by first activating NTS2 with **3a** and then screening for compounds that would block this activity. The binding assay versus [<sup>125</sup>I]NT could then be used as a secondary screen for active compounds to verify interaction with NTS2.

We tested this idea using the CHO cell line above and a FLIPR Tetra and studied the peptide NT, the nonpeptide levocabastine (2), and the two well-known nonpeptide pyrazole-based ligands SR48692 (3a) and SR142948a (3b), all of which are known to bind NTS2.<sup>24,25</sup> In line with literature precedent, we found that both pyrazole compounds (3a,b)

Received:
 June 9, 2014

 Published:
 August 26, 2014



behaved as full agonists with **3b** being more potent than **3a** but equally efficacious. Levocabastine showed potent partial agonist activity (14-16% percent of **3b**). NT, on the other hand, did not induce calcium release but was an antagonist of **3b** in the FLIPR assay.

While it seemed counterintuitive, our data showed clearly that NT was an antagonist and levocabastine (2) a very low efficacy potent partial agonist in the FLIPR assay, although both are known to be antinociceptive in animal models of pain. The two pyrazole compounds (3a,b), on the other hand, were found to be agonists even though it is well documented that both antagonize the analgesic action of NT-based compounds in a variety of animal models. Overall, this pilot study suggested that our search for novel NTS2-based analgesics should begin with identification of compounds with in vitro profiles mimicking either NT or 2 versus 3a or 3b.

We recently reported using this assay to drive an SAR study that led to the identification of the NTS2 selective, low efficacy, potent partial agonist 4 (NTRC-739).<sup>22</sup> In this article, we report a parallel study that identified the NTS2 selective nonpeptide compound 5 that, like NT, is an antagonist of 3b in the FLIPR assay. The details of this work are presented herein.

# CHEMISTRY

The chemistry used to prepare target compounds 5 and 13 is illustrated in Scheme 1. Thus, 5-nitroindole (7) was alkylated with methyl bromoacetate to give ester 8 that was subsequently hydrolyzed to the corresponding acid intermediate 9 by treatment with LiOH in dioxanes and water. Compound 9 was then coupled to leucine *tert*-butyl ester using *O*benzotriazol-1-yl-N,N,N,N'-tetramethyluronium hexafluorophosphate (HBTU) and triethylamine to give intermediate 10. This was reduced using hydrogen and palladium on carbon

# Scheme 1. Synthetic Scheme for Compounds 5 and 13<sup>a</sup>



<sup>*a*</sup>Reagents and conditions: (i) acetone, methyl bromoacetate,  $K_2CO_3$ , RT, 4 days; (ii) 2N LiOH, 1,4-dioxanes,  $H_2O$ , RT, 16 h; (iii) L-leucine *tert*-butyl ester hydrochloride, HBTU, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, RT, 4 h; (iv) H<sub>2</sub>, Pd/C, EtOH, RT, 2 h; (v) tosyl chloride, 4-dimethylaminopyridine (DMAP), CH<sub>2</sub>Cl<sub>2</sub>, RT, 16 h; (vi) TFA/CH<sub>2</sub>Cl<sub>2</sub> (1:1), RT, 30 min to 12 h; (vii) trifluoroacetic anhydride/CH<sub>2</sub>Cl<sub>2</sub>, 30 min, RT.

to give intermediate 11 and then coupled to tosyl chloride to give intermediate 12. Compound 13 was obtained from 12 by treatment with trifluoroacetic acid (TFA) in  $CH_2Cl_2$ . Compound 5 was obtained from 12 by treatment with trifluoroacetic anhydride (TFAA) to give the acylated indole derivative 14, which was subsequently deprotected with TFA to give 5.

## BIOLOGY

The binding affinities of the test compounds for the rNTS1 and rNTS2 receptors, listed in Table 1, were determined using

Table 1. Data for Reference Compounds NT and 3b and Test Compounds 5 and 13 in the FLIPR Assay (NTS2) and Binding Assays (NTS1 and NTS2 Receptors)

	FLIPR assay			binding assays <sup>a</sup>	
	NTS2			NTS1	NTS2
	EC <sub>50</sub> <sup>b</sup>	$E_{\max}^{c}$	IC <sub>50</sub>	K <sub>i</sub>	K <sub>i</sub>
NT	$NA^d$		$114 \pm 24$	$1.1\pm0.3$	$28 \pm 6$
3b	$22 \pm 5$	$100 \pm 6$		$1.4 \pm 0.4$	$5.8 \pm 1.1$
5	NA		$38 \pm 11$	$>30 \ \mu M$	$202 \pm 44$
13	NA		$3322 \pm 556$	$>30 \ \mu M$	$1504 \pm 244$
<sup>a</sup> [ <sup>125</sup> I]NT. <sup>b</sup> EC <sub>50</sub> , IC <sub>50</sub> and $K_i$ values are nM ± SEM. <sup>c</sup> $E_{max}$ value is %					

**3b**. <sup>*d*</sup>Not active.

<sup>[125</sup>I]NT in previously reported competitive binding assays.<sup>23</sup> The cells used were CHO-k1 cells (American Type Culture Collection) engineered to overexpress either the rNTS1 or rNTS2 receptor. Measures of functional agonist activity were obtained by measuring changes in intensity of a calciumsensitive fluorescent dye as an indirect measure of changes in internal calcium concentrations. Antagonism was measured as inhibition of calcium mobilization induced by **3b**. These measurements were performed using a FLIPR Tetra (Molecular Devices).

# LIBRARY SCREENING

Fan and co-workers reported that the nonpeptide compound **6a**, Chart 1, possessed partial agonist activity at the NTS1 receptor.<sup>26</sup> With the hope that some variant of this compound might also be active at NTS2, we made a library of 66 compounds with general structure **6b** using commercially available sulfonyl chlorides and amino acid esters by following the chemistry route described for compound **13** in Scheme 1. We screened these for agonist and antagonist activity in parallel at a single concentration (10  $\mu$ M) using **3b** as our NTS2 agonist standard and NT as our antagonist standard. Only one compound **(13)** showed >50% inhibition of **3b** induced calcium mobilization in the NTS2 screen. Compound **13** was also devoid of NTS2 agonist activity and showed no activity at NTS1 (data not shown).

# RESULTS AND DISCUSSION

In confirming the hit 13, an impurity was discovered that represented 25% of the material in the well. Mass spectral and  $^{19}$ F NMR analysis suggested that the desired compound (13) had incorporated a trifluoroacetyl group to give byproduct 5. We postulated that this would have occurred at the indole 3-position during deprotection with TFA and tested this by synthesizing 5 from a different direction as described in Scheme 1. Comparison of the <sup>1</sup>H NMR and mass spectral data of 5 to that of the impurity confirmed that they were identical. Both compounds 5 and 13 were carried forward into dose response testing.

In Table 1, we provide the FLIPR assay (NTS2) and binding assay (NTS1 and NTS2) data collected for the reference compounds NT and **3b** as well as the test compounds **5** and **13**. In the FLIPR assay, the reference compounds performed as expected. NT showed no agonist activity up to 100 mM, but as previously reported, the  $K_e$  experiment with NT at NTS2 versus **3b** demonstrated insurmountable antagonist activity.<sup>22,27,28</sup> For this reason, we determined and report here its IC<sub>50</sub> value of 114 nM. In the binding assay at NTS2, NT provided a  $K_i$  value of 28 nM. Reference compound **3b** gave an EC<sub>50</sub> of 22 nM (100%  $E_{max}$ ) in the FLIPR assay and a  $K_i$  of 5.8 nM in the binding assay.

Neither of the test compounds (5 or 13) showed activity in the NTS1 FLIPR assay (data not shown), but in the NTS2 FLIPR assay, they both behaved like NT. Each compound antagonized the action of 3b in the  $K_e$  experiment in an insurmountable manner. The graph of the  $K_e$  experiment for 5 is illustrated in Figure 1. Both 5 and 13 were tested in the IC<sub>50</sub> experiment and gave IC<sub>50</sub> values of 38 and 3322 nM, respectively. The results from the IC<sub>50</sub> experiment of NT and 5 versus 3b are illustrated in Figure 2.

In the binding assays at NTS2, **5** and **13** showed  $K_i$  values of 202 and 1504 nM, respectively, versus 28 nM for NT. Testing of compounds **5** and **13** at the NTS1 receptor showed no binding activity in the NTS1 assay up to 30  $\mu$ M, making **5** selective for NTS2 in both the binding and functional assays.

A comparison of FLIPR data for compounds 5 and 13 revealed that the impurity 5 was not only 90-fold more active than the target compound 13 but also 3-fold more active than the endogenous ligand NT. This result was unexpected, as the impurity was found to be far more interesting than the intended compound. Indeed, it is unlikely that this discovery would have been made if not by accident as substitution with the trifluoroacetyl group is not a common practice in medicinal



**Figure 1.**  $K_e$  assays were run against dose-response curves of the control agonist **3b**. Compound **5** demonstrated the characteristics of insurmountable antagonist activity versus **3b** in the  $K_e$  experiment. This was also seen for NT as reported earlier.<sup>22</sup>



**Figure 2.**  $IC_{50}$  assays were run against the  $EC_{80}$  of **3b** (73 nM final concentration). The  $IC_{50}$  curves for **5** and NT in the NTS2 FLIPR assay demonstrate that **5** is almost 3-fold more active than the endogenous ligand NT.

chemistry. Overall, however, we have identified a molecular region of **5** that is very important to antagonism of calcium mobilization at NTS2.

The activity of both 5 and 13 in the NTS2 FLIPR and binding assays provides additional evidence that the calcium release mediated by 3b is an NTS2 mediated event. Also, a comparison of their binding data at NTS2 revealed that the trifluoroacetyl substitution of 5 did not increase the binding affinity to the same degree observed for the FLIPR assay. We found a 90-fold boost to antagonist activity for 5 versus 13 in the FLIPR assay and a 7.5-fold boost to binding affinity. Although these need not necessarily be equivalent, we suspect that this could be a consequence of measuring the two end points using two different reference compounds. In the FLIPR assay, 5 is believed to compete directly with 3b, while in the binding assay 5 is competing against NT. In most assay systems, the endogenous ligand (or another more active compound) serves as both the reference compound in the functional assay and the radioligand in the binding assay. In this case, however, the endogenous ligand (NT) does not mobilize calcium. Thus, we would require radioactive 3b in order to

#### Journal of Medicinal Chemistry

achieve parity. While this is possible, radioactive NT is commercially available and far more convenient to obtain.<sup>29</sup> Empirically, however, we observe that every compound so far identified that inhibits **3b** in the FLIPR assay also competes with NT in the binding assay, suggesting that the difference in magnitude of effect does not prevent the detection of NTS2 active compounds.

As noted above, we do not know that 5 is competing directly with 3b for its binding site, this is only a working assumption. There is, however, empirical evidence that supports the assumption that the binding sites for 3b and NT are, at least partially, overlapping. The most important observation is that NT, 3a, 3b, and 5 all rely heavily upon a free carboxylic acid group for binding to NTS2 (and NTS1). In studies of the NTS1 receptor, the ionic interaction established by the binding of this group to an arginine residue in the receptor, formed the principle ligand to receptor stabilizing interaction.<sup>30</sup> Along with this, a partially overlapping scenario has been postulated for NT and 3a binding sites based on evidence obtained from molecular modeling and point mutation studies at the NTS1 receptor.<sup>31</sup> Although not conclusive, these observations do offer support to the notion that these compounds all share partially overlapping binding sites.

#### SUMMARY

The work presented herein provides additional evidence that 3b can be used as an agonist in the NTS2 FLIPR assay to identify novel NTS2 active compounds. Because NT is an antagonist of 3b, novel NT-like antagonist compounds can be identified. To our knowledge, 5 is the first nonpeptide compound that behaves like, and is more active than, the endogenous ligand in this functional assay. This work compliments our discovery of compound 4, providing an additional nonpeptide compound whose in vitro properties align with those of compounds that are known to be active in models of pain. Together, 4 and 5 should be useful in additional studies related to the biology of the NT receptors. We are evaluating these compounds in animal models of pain to establish the relevance of this assay with respect to identification of potential NTS2-based analgesics. Additional SAR studies with 5 are in progress and will be reported in the near future.

#### EXPERIMENTAL SECTION

Reactions were conducted under a nitrogen atmosphere using ovendried glassware as required. All solvents and chemicals used were reagent grade. 5-Nitroindole (7) was purchased from Combi-blocks, Inc. Flash column chromatography was carried out using a Teledyne ISCO Combiflash Rf system and Redisep Rf gold prepacked HP silica columns. Flow rates are automatically determined by column size. Fractions were collected as "peaks only" over 30 min. The purity of target compounds was determined to be  $\geq$ 95% by combustion analysis. NMR spectra were recorded on a Bruker Avance DPX-300 (300 MHz). Low-resolution mass spectra were obtained using a Waters Alliance HT/Micromass ZQ system (ESI) in both positive and negative mode. Optical rotations were measured on an Auto Pol III polarimeter at the sodium D line. Thin layer chromatography (TLC) was performed on EMD precoated silica gel 60 F<sub>254</sub> plates, and spots were visualized with UV light and I2 or phosphomolybdic acid stain. CHO-k1 cells were from the American Type Culture Collection, and [<sup>125</sup>I]NT was from PerkinElmer. Calcium 5 dye was from Molecular Devices. Cell culture reagents were from Life Technologies.

N-[[5-[[(4-Methylphenyl)sulfonyl]amino}-3-(trifluoroacetyl)-1H-indol-1-yl]acetyl}-L-leucine (5). 14 (78.0 mg, 0.14 mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (2 mL), and to this was added TFA (2 mL) and the reaction stirred at RT for 2 h. This mixture was then concentrated to dryness and the residue dissolved in a minimal amount of CH<sub>2</sub>Cl<sub>2</sub>, and then 2 mL each of ethyl ether and hexanes was added to precipitate the acid. This was filtered, washed with hexanes, and then dried under vacuum to provide **5** (67.3 mg, 86%) as an off-white powder. <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  8.71 (d, *J* = 8.0 Hz, 1H), 8.29 (d, *J* = 1.6 Hz, 1H), 7.99 (d, *J* = 2.0 Hz, 1H), 7.63 (d, *J* = 8.4 Hz, 2H), 7.37 (d, *J* = 8.9 Hz, 1H), 7.26 (d, *J* = 8.0 Hz, 2H), 7.20 (dd, *J* = 2.0, 8.9 Hz, 1H), 4.97–5.15 (m, 2H), 4.38–4.53 (m, 1H), 2.35 (s, 3H), 1.58–1.79 (m, 3H), 0.94 (dd, *J* = 5.9, 19.5 Hz, 6H). <sup>13</sup>C NMR (CD<sub>3</sub>OD)  $\delta$  21.39, 21.69, 23.35, 26.11, 41.51, 50.53, 52.35, 103.00, 104.59, 110.56, 112.39, 116.26, 120.93, 128.39, 128.54, 130.51, 135.49, 136.26, 137.95, 142.01, 142.09, 144.94, 168.73, 175.58. <sup>19</sup>F NMR (CD<sub>3</sub>OD)  $\delta$  –74.02. MS (ESI): 552.7 (M – H)<sup>-</sup>; [ $\alpha$ ] <sup>25</sup><sub>D</sub> –17.1 (*c* 0.62, CH<sub>3</sub>OH). Anal. (C<sub>25</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O<sub>6</sub>S): C, H, N.

**Methyl (5-Nitro-1***H***-indol-1-yl)acetate (8).** Into a 250 mL RBF with a magnetic stir bar are placed 5-nitroindole (7, 4.864 g, 0.030 mol) and  $K_2CO_3$  (4.561 g, 0.033 mol) and 90 mL of acetone. The methyl bromoacetate (5.048 g, 0.033 mol, 3.10 mL) was added over a minute via syringe and then stirred 4 days at RT. The reaction was then concentrated, poured into water, and extracted with EtOAc (2×), and the combined organics were washed (H<sub>2</sub>O, brine) and dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give a yellow oil. This was chromatographed on an ISCO silica gel column (80 g) using a 0–60% EtOAc/hexanes gradient and then crystallized from ethyl acetate to give the desired product as a yellow solid (4.38 g, 65% yield). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.60 (d, *J* = 2.1 Hz, 1 H), 8.14 (dd, *J* = 2.2, 9.1 Hz, 1 H), 7.26 (d, *J* = 9.1 Hz, 1 H), 7.25 (d, *J* = 3.3 Hz, 1 H), 6.75 (d, *J* = 3.2 Hz, 1 H), 4.92 (s, 2 H), 3.78 (s, 3 H).

(5-Nitro-1*H*-indol-1-yl)acetic Acid (9). To a solution of 8 (10.10 g, 0.045 mol) in 50 mL of 1,4-dioxane was added 50 mL of 1 N LiOH, and this was stirred overnight at RT. The reaction was concentrated to remove dioxane and then diluted with 100 mL of H<sub>2</sub>O and acidified to pH 2. This was then extracted with 100 mL of ethyl acetate (3×), and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration gave 9 (8.72 g, 88%) as a white solid. <sup>1</sup>H NMR (DMSO- $d_6$ )  $\delta$  13.13 (br s, 1H), 8.58 (d, *J* = 2.3 Hz, 1 H), 8.03 (dd, *J* = 2.3, 9.0 Hz, 1 H), 7.66–7.58 (m, 2 H), 6.76 (d, *J* = 3.2 Hz, 1 H), 5.16 (s, 2 H). MS (ESI): 219.3 (M – H)<sup>-</sup>.

*tert*-Butyl *N*-[(5-Nitro-1*H*-indol-1-yl)acetyl]-L-leucinate (10). Into a 100 mL RBF with stir bar containing 9 (2.202 g, 0.010 mol), (*S*)-(L)-leucine *tert*-butyl ester HCl salt (2.461 g, 0.011 mol), and HBTU (4.172g, 0.011 mol) was added CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and then Et<sub>3</sub>N (3.036 g, 0.030 mol, 4.18 mL), and this was stirred at RT overnight. The reaction mixture was then concentrated and chromato-graphed on an 80 g ISCO silica gel column using a 10–60% gradient of EtOAc in hexanes to give **10** (4.07 g) as a yellow oil which crystallized when dried under vacuum to give a yellow–orange solid (3.859 g, 99%). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  8.59 (d, *J* = 2.2 Hz, 1 H), 8.13 (dd, *J* = 2.2, 9.0 Hz, 1 H), 7.35 (d, *J* = 9.0 Hz, 1 H), 7.31 (d, *J* = 3.3 Hz, 1 H), 6.80 (dd, *J* = 0.5, 3.2 Hz, 1 H), 5.96 (d, *J* = 8.3 Hz, 1 H), 4.88 (s, 2 H), 4.50 (dt, *J* = 5.2, 8.5 Hz, 1 H), 1.59–1.42 (m, 2 H), 1.40 (s, 9 H), 0.85 (dd, *J* = 3.3, 6.2 Hz, 6 H). MS (ESI): 360.5 (M + H)<sup>+</sup>.

*tert*-Butyl *N*-[(5-Amino-1*H*-indol-1-yl)acetyl]-L-leucinate (11). 10 (3.894 g, 0.010 mol) was dissolved in ethanol (200 mL) and reduced using a Parr apparatus with 10% Pd/C (400 mg) under 40 psig of H<sub>2</sub>. The reaction mixture was filtered through Celite, the cake was washed with EtOH, and the filtrate was concentrated under vacuum. Drying under HVAC gave 11 (3.620 g, 100%) as a red oil. [NOTE: This material darkens with exposure to air.] <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.09 (d, J = 8.6 Hz, 1 H), 7.01 (d, J = 3.1 Hz, 1 H), 6.96 (d, J = 2.1 Hz, 1 H), 6.72 (dd, J = 2.2, 8.6 Hz, 1 H), 6.44 (d, J = 2.7 Hz, 1 H), 5.62 (d, J = 8.5 Hz, 1 H), 4.74 (s, 2 H), 4.45 (dt, J = 5.1, 8.7 Hz, 1 H), 1.50–1.41 (m, 2 H), 1.36 (s, 9 H), 1.31–1.20 (m, 1 H), 0.81 (dd, J = 6.4, 7.8 Hz, 6 H). MS (ESI): 358.5 (M – H)<sup>-</sup>, 360.3 (M + H)<sup>+</sup>.

*tert*-Butyl *N*-({5-[(4-Methylphenylsulfonyl)amino]-1*H*-indol-1-yl]acetyl)-L-leucinate (12). 11 (3.595 g, 0.010 mol) was dissolved in  $CH_2Cl_2$  (50 mL) in a 250 mL RBF equipped with a stir bar. DMAP (1.344 g, 0.011 mol) was added, followed by a solution of tosyl chloride (2.002 g, 0.011 mol) in  $CH_2Cl_2$  (20 mL). The reaction was

#### Journal of Medicinal Chemistry

stirred overnight and then concentrated and chromatographed on an 80 g silica gel cartridge with a 0–80% EtOAc/hexanes gradient. The product crystallized from the fractions upon standing to give **12** (4.65g, 91%) as colorless crystals. <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.61 (d, *J* = 8.2 Hz, 2 H), 7.39 (d, *J* = 1.9 Hz, 1 H), 7.18 (s, 2 H), 7.16 (s, 1 H), 7.12–7.09 (m, 1 H), 6.95 (dd, *J* = 1.9, 8.8 Hz, 1 H), 6.52 (d, *J* = 3.2 Hz, 1 H), 5.75 (d, *J* = 8.5 Hz, 1 H), 4.77 (s, 2 H), 4.46 (dt, *J* = 5.1, 8.5 Hz, 1 H), 2.35 (s, 3 H), 1.53–1.41 (m, 3 H), 1.37 (s, 9 H), 0.79 (t, *J* = 6.2 Hz, 6 H). MS (ESI): 514.3 (M + H)<sup>+</sup>.

N-[(5-{[(4-Methylphenyl)sulfonyl]amino}-1H-indol-1-yl)acetyl]-L-leucine (13). 12 (51.4 mg, 0.1 mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (2 mL), and then TFA was added (2 mL) and the reaction stirred at RT for 30 min. This was then concentrated to dryness and the residue dissolved in a minimal amount of CH2Cl2/MeOH and then triturated with 2 mL each of ethyl ether and hexanes to precipitate the acid as an off-white powder. This was collected by filtration and washed with hexanes, and then the solids were dried under vacuum. This was then purified on a 4 g ISCO silica gel column using a gradient of 0-10% MeOH/NH<sub>4</sub>OH (10:1) in CH<sub>2</sub>Cl<sub>2</sub> to provide the 13 (11.7 mg, 26%) as a off-white solid. <sup>1</sup>H NMR  $(CD_3OD) \delta$  7.55 (d, I = 8.3 Hz, 1H), 7.23–7.31 (m, 1H), 7.16–7.23 (m, 1H), 7.02–7.15 (m, 2H), 6.98 (t, J = 8.5 Hz, 1H), 6.89 (dt, J = 2.0, 4.3 Hz, 1H), 6.82 (dd, J = 2.0, 8.8 Hz, 1H), 6.73 (dd, J = 2.0, 8.8 Hz, 1H), 6.40 (d, J = 3.0 Hz, 1H), 4.72–4.86 (m, 2H), 4.40–4.62 (m, 1H), 2.16–2.41 (m, 3H), 1.52–1.79 (m, 4H), 0.81–0.98 (m, 6H). <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ 21.42, 21.75, 23.38, 26.05, 41.57, 50.04, 52.20, 102.88, 110.73, 116.78, 119.64, 128.37, 130.38, 130.42, 130.90, 131.35, 136.16, 138.10, 144.60, 170.62, 175.76. MS (ESI): 456.3 (M - H)<sup>-</sup>;  $[\alpha]_{D}^{25}$  –8.84 (c 0.37, CH<sub>3</sub>OH). Anal. (C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>5</sub>S): C, H, N.

*tert*-Butyl *N*-{[[4-Methylphenyl]sulfonyl]amino}-3-(trifluoroacetyl)-1*H*-indol-1-yl]acetyl}-L-leucinate (14). 12 (128.4 mg, 0.25 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and to this was added trifluoroacetic anhydride (57.8 mg, 0.04 mL, 0.28 mmol) and the reaction was stirred at RT for 2 h. The reaction was then concentrated to a solid film and purified on a 4 g ISCO silica gel column using a 0–60% EtOAc/hexanes gradient. The desired fractions were concentrated to provide 14 (85.9 mg, 56%) as a colorless film. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.92 (d, *J* = 1.5 Hz, 1H), 7.84 (s, 1H), 7.76 (s, 1H), 7.61 (d, *J* = 8.3 Hz, 2H), 7.10–7.19 (m, 4H), 6.83 (d, *J* = 8.3 Hz, 1H), 4.88 (s, 2H), 4.49–4.62 (m, 1H), 2.31 (s, 3H), 1.51–1.68 (m, 3H), 1.46 (s, 9H), 0.90 (d, *J* = 3.9 Hz, 6H). MS (ESI): 609.2 (M – H)<sup>-</sup>.

Calcium Mobilization Assay for NTS2 Receptor. CHO-k1rNTS2 cells were maintained in DMEM/F12 supplemented with 10% FBS, pen/strep, 100  $\mu$ g/mL normocin, and 400  $\mu$ g/mL Geneticin. For calcium mobilization assays, cells were plated at 25000 cells/well in black, clear-bottom 96-well plates 24 h before the assay and incubated at 37 °C, 5% CO2. Then 100 µL of reconstituted Calcium 5 dye (Molecular Devices, diluted 1:20 in assay buffer (1× HBSS, 20 mM HEPES, 2.5 mM Probenicid, pH 7.4)) was added per well and plates were incubated for 45 min at 37 °C, 5% CO<sub>2</sub>. For agonist assays, cells were pretreated with 1:10 addition of 10% DMSO, and the plates were returned to 37 °C, 5% CO2. Full-log serial dilutions of the test compounds were made at 10× the desired final concentration in 1% DMSO assay buffer and warmed to 37 °C. After the pretreatment incubation, fluorescence intensity was measured on a FLIPR Tetra fluorometric imaging plate reader (Molecular Devices). Relative fluorescence units (RFUs) were measured every second for 100 s (14 readings before compound addition to establish baseline fluorescence, 85 after), exposure time 0.53 s, gain = 2000, excitation intensity = 30%, gate open = 10%. For antagonist assays, 1/10 volume of 10× concentration of the test compound dilutions in 10% DMSO in assay buffer was added to cells in lieu of the 10% DMSO only in assay buffer for the pretreatment. Ke assays were run against dose-response curves of the control agonist 3b, and IC<sub>50</sub> assays were run against the EC<sub>80</sub> of **3b** (73 nM final concentration).

**Competitive Binding Assays.** Relative binding affinity was evaluated using  $^{125}\mathrm{I}$  labeled neurotensin and CHO-k1 cell lines overexpressing either the rNTS1 or rNTS2 receptor essentially as described by Gendron.  $^{23}$ 

**Data Analysis.** To determine  $EC_{50}$  and  $IC_{50}$  values, data were fit to a three-parameter logistic equation using GraphPad Prism software.  $K_i$  values for radioligand binding assays were determined from  $IC_{50}$  values using the equation of Cheng and Prusoff. All data are from at least three independent experiments run in duplicate wells.

# ASSOCIATED CONTENT

#### **S** Supporting Information

Elemental analysis results for test compounds 5, 13, and 14. This material is available free of charge via the Internet at http://pubs.acs.org.

# AUTHOR INFORMATION

#### **Corresponding Author**

\*Phone: 919-541-6375. Fax: 919-541-6499. E-mail: jbthomas@rti.org.

#### **Author Contributions**

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

We gratefully acknowledge the NIMH Chemical Synthesis and Drug Supply Program for providing us with the samples of SR142948a (3b), and SR48692 (3a). This research was supported by the National Institute on Drug Abuse, grant DA029961.

#### ABBREVIATIONS USED

NT, neurotensin; NTS1, neurotensin receptor 1; NTS2, neurotensin receptor 2; CNS, central nervous system 5; H1, histamine receptor 1; FLIPR, fluorometric imaging plate reader; CHO, Chinese hamster ovary

#### REFERENCES

(1) Caraway, R.; Leeman, S. E. The isolation of a new hypotensive peptide, neurotensin, from bovine hypothalami. *J. Biol. Chem.* **1973**, 248, 6854–6861.

(2) Clineschmidt, B. V.; McGuffin, J. C. Neurotensin administered intracisternally inhibits responsiveness of mice to noxious stimuli. *Eur. J. Pharmacol.* **1977**, *46*, 395–396.

(3) Kitabgi, P. Targeting neurotensin receptors with agonists and antagonists for therapeutic purposes. *Curr. Opin. Drug Discovery Dev.* **2002**, *5*, 764–776.

(4) Vincent, J. P.; Mazella, J.; Kitabgi, P. Neurotensin and neurotensin receptors. *Trends Pharmacol. Sci.* **1999**, *20*, 302–309.

(5) Mazella, J.; Zsurger, N.; Navarro, V.; Chabry, J.; Kaghad, M.; Caput, D.; Ferrara, P.; Vita, N.; Gully, D.; Maffrand, J. P.; Vincent, J. P. The 100 kDa neurotensin receptor is gp95/sortilin, a non-G-proteincoupled receptor. *J. Biol. Chem.* **1998**, 273, 26273–26276.

(6) Tanaka, K.; Masu, M.; Nakanishi, S. Structure and functional expression of the cloned rat neurotensin receptor. *Neuron* **1990**, *4*, 847–854.

(7) Vita, N.; Laurent, P.; Lefort, S.; Chalon, P.; Dumont, X.; Kaghad, M.; Gully, D.; Le Fur, G.; Ferrara, P.; Caput, D. Cloning and expression of a complementary DNA encoding a high affinity human neurotensin receptor. *FEBS Lett.* **1993**, *317*, 139–142.

(8) Chalon, P.; Vita, N.; Kaghad, M.; Guillemot, M.; Bonnin, J.; Delpech, B.; Le Fur, G.; Ferrara, P.; Caput, D. Molecular cloning of a levocabastine-sensitive neurotensin binding site. *FEBS Lett.* **1996**, *386*, 91–94.

(9) Dobner, P. R. Neurotensin and pain modulation. *Peptides* 2006, 27, 2405–2414.

(10) Tetreault, P.; Beaudet, N.; Perron, A.; Belleville, K.; Rene, A.; Cavelier, F.; Martinez, J.; Stroh, T.; Jacobi, A. M.; Rose, S. D.; Behlke, M. A.; Sarret, P. Spinal NTS2 receptor activation reverses signs of neuropathic pain. *FASEB J.* **2013**, *27*.

(11) Guillemette, A.; Dansereau, M. A.; Beaudet, N.; Richelson, E.; Sarret, P. Intrathecal administration of NTS1 agonists reverses nociceptive behaviors in a rat model of neuropathic pain. *Eur. J. Pain* **2012**, *16*, 473–484.

(12) Boules, M.; Shaw, A.; Liang, Y.; Barbut, D.; Richelson, E. NT69L, a novel analgesic, shows synergy with morphine. *Brain Res.* **2009**, 1294, 22–28.

(13) Hughes, F. M.; Shaner, B. E.; May, L. A.; Zotian, L.; Brower, J. O.; Woods, R. J.; Cash, M.; Morrow, D.; Massa, F.; Mazella, J.; Dix, T. A. Identification and functional characterization of a stable, centrally active derivative of the neurotensin (8–13) fragment as a potential first-in-class analgesic. *J. Med. Chem.* **2010**, *53*, 4623–4632.

(14) Boules, M.; Liang, Y.; Briody, S.; Miura, T.; Fauq, I.; Oliveros, A.; Wilson, M.; Khaniyev, S.; Williams, K.; Li, Z.; Qi, Y.; Katovich, M.; Richelson, E. NT79: A novel neurotensin analog with selective behavioral effects. *Brain Res.* **2010**, *1308*, 35–46.

(15) Akunne, H. C.; Demattos, S. B.; Whetzel, S. Z.; Wustrow, D. J.; Davis, D. M.; Wise, L. D.; Cody, W. L.; Pugsley, T. A.; Heffner, T. G. Agonist properties of a stable hexapeptide analog of neurotensin, N alpha MeArg-Lys-Pro-Trp-tLeu-Leu (NT1). *Biochem. Pharmacol.* **1995**, 49, 1147–1154.

(16) Bredeloux, P.; Cavelier, F.; Dubuc, I.; Vivet, B.; Costentin, J.; Martinez, J. Synthesis and biological effects of c(Lys-Lys-Pro-Tyr-Ile-Leu-Lys-Lys-Pro-Tyr-Ile-Leu) (JMV2012), a new analogue of neurotensin that crosses the blood-brain barrier. *J. Med. Chem.* **2008**, *51*, 1610–1616.

(17) Tyler-McMahon, B. M.; Stewart, J. A.; Farinas, F.; McCormick, D. J.; Richelson, E. Highly potent neurotensin analog that causes hypothermia and antinociception. *Eur. J. Pharmacol.* **2000**, *390*, 107–111.

(18) Fantegrossi, W. E.; Ko, M. C.; Woods, J. H.; Richelson, E. Antinociceptive, hypothermic, hypotensive, and reinforcing effects of a novel neurotensin receptor agonist, NT69L, in rhesus monkeys. *Pharmacol., Biochem. Behav.* **2005**, *80*, 341–349.

(19) Smith, K. E.; Boules, M.; Williams, K.; Richelson, E. NTS1 and NTS2 mediate analgesia following neurotensin analog treatment in a mouse model for visceral pain. *Behav. Brain Res.* **2012**, *232*, 93–97.

(20) Tasaka, K.; Kamei, C.; Tsujimoto, S.; Yoshida, T.; Aoki, I. Central effect of the potent long-acting H1-antihistamine levocabastine. *Arzneimittelforschung* **1990**, *40*, 1295–1299.

(21) Sarret, P.; Esdaile, M. J.; Perron, A.; Martinez, J.; Stroh, T.; Beaudet, A. Potent spinal analgesia elicited through stimulation of NTS2 neurotensin receptors. *J. Neurosci.* 2005, 25, 8188–8196.

(22) Thomas, J. B.; Giddings, A. M.; Wiethe, R. W.; Olepu, S.; Warner, K. R.; Sarret, P.; Gendron, L.; Longpre, J. M.; Zhang, Y.; Runyon, S. P.; Gilmour, B. P. Identification of 1-({[1-(4-fluorophenyl)-5-(2-methoxyphenyl)-1*H*-pyrazol-3-yl]carbonyl}-amino)cyclohexane carboxylic acid as a selective nonpeptide neurotensin receptor type 2 compound. J. Med. Chem. **2014**, *57*, 5318–5332.

(23) Gendron, L.; Perron, A.; Payet, M. D.; Gallo-Payet, N.; Sarret, P.; Beaudet, A. Low-affinity neurotensin receptor (NTS2) signaling: internalization-dependent activation of extracellular signal-regulated kinases 1/2. *Mol. Pharmacol.* **2004**, *66*, 1421–1430.

(24) Gully, D.; Canton, M.; Boigegrain, R.; Jeanjean, F.; Molimard, J. C.; Poncelet, M.; Gueudet, C.; Heaulme, M.; Leyris, R.; Brouard, A. Biochemical and pharmacological profile of a potent and selective nonpeptide antagonist of the neurotensin receptor. *Proc. Natl. Acad. Sci. U. S. A.* **1993**, *90*, 65–69.

(25) Gully, D.; Labeeuw, B.; Boigegrain, R.; Oury-Donat, F.; Bachy, A.; Poncelet, M.; Steinberg, R.; Suaud-Chagny, M. F.; Santucci, V.; Vita, N.; Pecceu, F.; Labbe-Jullie, C.; Kitabgi, P.; Soubrie, P.; Le Fur, G.; Maffrand, J. P. Biochemical and pharmacological activities of SR 142948A, a new potent neurotensin receptor antagonist. *J. Pharmacol. Exp. Ther.* **1997**, *280*, 802–812.

(26) Fan, Y.; Lai, M. H.; Sullivan, K.; Popiolek, M.; Andree, T. H.; Dollings, P.; Pausch, M. H. The identification of neurotensin NTS1 receptor partial agonists through a ligand-based virtual screening approach. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 5789–5791.

(27) Gaddum, J. H. Theories of drug antagonism. *Pharmacol. Rev.* **1957**, *9*, 211–218.

(28) Kenakin, T.; Jenkinson, S.; Watson, C. Determining the potency and molecular mechanism of action of insurmountable antagonists. *J. Pharmacol. Exp. Ther.* **2006**, *319*, 710–723.

(29) Labbe-Jullie, C.; Barroso, S.; Nicolas-Eteve, D.; Reversat, J. L.; Botto, J. M.; Mazella, J.; Bernassau, J. M.; Kitabgi, P. Mutagenesis and modeling of the neurotensin receptor NTR1. Identification of residues that are critical for binding SR 48692, a nonpeptide neurotensin antagonist. J. Biol. Chem. **1998**, 273, 16351–16357.

(30) Henry, J. A.; Horwell, D. C.; Meecham, K. G.; Rees, D. C. A structure–affinity study of the amino acid side chains in neurotensin: N and C terminal deletions and Ala-scan. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 949–952.

(31) Labbe-Jullie, C.; Botto, J. M.; Mas, M. V.; Chabry, J.; Mazella, J.; Vincent, J. P.; Gully, D.; Maffrand, J. P.; Kitabgi, P. [3H]SR 48692, the first nonpeptide neurotensin antagonist radioligand: characterization of binding properties and evidence for distinct agonist and antagonist binding domains on the rat neurotensin receptor. *Mol. Pharmacol.* **1995**, *47*, 1050–1056.