

Isolation and Identification of Insecticidal Components from *Citrus aurantium* Fruit Peel Extract

ELIAS P. SISKOS, BASILIS E. MAZOMENOS,* AND MARIA A. KONSTANTOPOULOU

Chemical Ecology and Natural Products Laboratory, NCSR "Demokritos",
150 13 Aghia Paraskevi, Attikis, Greece

Three active components were identified by bioassay-guided fractionation of bitter orange (*Citrus aurantium* L.) fruit peel petroleum ether extract. Silica gel fractionation of the extract yielded a fraction that inflicted up to 96% mortality to adults of the olive fruit fly *Bactrocera oleae* (Gmelin) three days post-treatment. Subsequent HPLC purification of the active fraction resulted in the isolation of three components, eluted in fractions F₂₂₂, F₂₂₄, and F₂₂₆, that induced adult mortality. Considering the data obtained from UV, FTIR, MS, and ¹H NMR spectra, they were identified as 7-methoxy-8-(3'-methyl-2'-butenyl)-2*H*-1-benzopyran-2-one (osthol), 4-methoxy-7*H*-furo[3,2-*g*]benzopyran-7-one (bergapten), and 4-((*E*)-3'-methyl-5'-(3'',3''-dimethyloxiran-2''-yl)pent-2'-enyloxy)-7*H*-furo[3,2-*g*]benzopyran-7-one (6',7'-epoxybergamottin). Our results are in concordance with those reported in the literature and were further verified by direct comparison to authentic components. 6',7'-Epoxybergamottin was toxic when tested individually, while bergapten and osthol were found to act synergistically to 6',7'-epoxybergamottin.

KEYWORDS: Bitter orange; *Citrus aurantium*; HPLC; coumarin; furanocoumarins; plant insecticides; fruit flies; *Bactrocera oleae*

INTRODUCTION

Peel and seed oils from several citrus plants contain chemicals that exhibit insecticidal and antifungal activity (1–3). Limonoids, extremely bitter chemicals present in citrus seeds, act as antifeedant or exhibit antagonistic to ecdysone action in many insect species (4, 5). Four major components have been isolated from lemon peel oil: 5,7-dimethoxy-2*H*-1-benzopyran-2-one, 9-[(3,7-dimethyl-2,6-octadienyl)oxy]-7*H*-furo[3,2-*g*][1]benzopyran-7-one, 4-[(3,7-dimethyl-2,6-octadienyl)oxy]-7*H*-furo[3,2-*g*][1]benzopyran-7-one, and 5-[(3,7-dimethyl-2,6-octadienyl)oxy]-7-methoxy-2*H*-1-benzopyran-2-one. The blend of those compounds exhibited insecticidal activity against the adults of the beetle species *Sitophilus oryzae* L. and *Callosobruchus maculatus* (F.) (6). Recent studies reported that flavones and some phenolic compounds in certain citrus essential oils exhibit insecticidal activity (7, 8).

Bitter orange (*Citrus aurantium* L.) fruits have been reported to possess antibacterial, antifungal, and pharmacological properties (9). We have recently demonstrated that ripe *C. aurantium* peels homogenized in methanol and separated using solvents of increasing polarity exhibit strong insecticidal activity against the olive fruit fly *Bactrocera oleae* (Gmelin) (10).

The objectives of the present study were (a) to isolate and identify the biologically active chemicals present in the bitter orange peels and (b) to test the effect of the isolated fraction(s) and the synthetic compound(s) on the viability of *B. oleae* adults.

MATERIALS AND METHODS

Chemicals. All chemicals used in the study, such as petroleum ether (PE), dichloromethane (DCM), *n*-hexane (Hex), ethyl acetate (EtOAc), and methanol (MeOH), were analytical grade. Sodium sulfate (Na₂SO₄) was purchased from Merck and 3-chloroperbenzoic acid (MCPBA) from Fluka (Buchs, Switzerland). The deuterated chloroform-*d* (CDCl₃) and the synthetic bergapten were purchased from Sigma-Aldrich Corp. (St. Louis, MO). The synthetic bergamottin was purchased from Carl-Roth, (Karlsruhe, Germany) and the synthetic osthol sample was kindly donated by Dr. M. R. Berenbaum.

Plant Material. Fruits were collected in September 2006 from *C. aurantium* trees located in the Zografos area of Athens, Greece. They were immediately transferred to the laboratory and washed in tap water containing 0.01% detergent using a soft brush, rinsed with tap water, and finally rinsed with distilled water. Peels were excised from 60 fruits using a razor blade, which left the white spongy portion (albedo) on the fruit.

Extraction of Fruit Peels. Fresh peels (300 g) were homogenized with 600 mL of petroleum ether for 5 min in a blender. The homogenate was placed in a 2 L Erlenmeyer flask. Three hundred milliliters of petroleum ether were added in the flask and further extracted three times for 10 min each in an ultrasonic bath at room temperature. The

* Author to whom correspondence should be addressed. Telephone: +30 210 6503556. Fax: +30 210 6511767. E-mail: bmazom@bio.demokritos.gr).




Figure 1. HPLC profile of the F₂₂ fraction from silica gel column of the petroleum ether fruit peel extract of *C. aurantium* and UV absorbance wavelength spectra of (A) the coumarin osthol and the furanocoumarins (B) bergapten and (C) 6',7'-epoxybergamottin, isolated from *C. aurantium* fruit peel extracts.

extract was filtered and evaporated under vacuum at 35 °C to dryness. The residue (2.22 g) was collected in 5 mL of DCM and stored at −20 °C until use.

Flash Chromatography. The petroleum peel extract was fractionated on a 500 × 45 mm glass column, packed with 200 g of silica gel (0.04–0.063 mm, 230–400 mesh; Macherey-Nagel, GmbH & Co. KG, Postfach, Germany) suspended in dichloromethane. A portion of the petroleum ether peel extract equivalent to 2 g in 20 mL of dichloromethane was introduced onto the column. Three fractions were collected by sequentially eluting the column with 1 L each of, dichloromethane (DCM-F₁ fraction, 542 mg), ethyl acetate (EtOAc-F₂ fraction, 1254 mg), and methanol (MeOH-F₃ fraction, 138 mg).

The biologically active (EtOAc-F₂) fraction was further fractionated. The same glass column packed with 100 g of silica gel (suspended in 20% EtOAc in Hex). One gram of the EtOAc-F₂ fraction dissolved in 50 mL of 20% EtOAc in Hex was introduced onto the column. Four fractions were collected by sequentially eluting the column with 500 mL each of, 20% EtOAc in Hex (F₂₁ fraction, 41 mg), 40% EtOAc in Hex (F₂₂ fraction, 243 mg), 60% EtOAc in Hex (F₂₃ fraction, 633 mg), and 100% MeOH (F₂₄ fraction, 80 mg).

HPLC Fractionation. Analyses were performed on an HPLC system equipped with a binary LC pump 250, an LC-235 diode array detector (Perkin-Elmer, Norwalk, CT), and the Simple peak 202 Data System (SRI Instruments, Torrance, CA). The fractionation was carried out on a 250 mm × 10 mm i.d. 10 μm LiChrospher 100 CN column (Merck KGaA, Darmstadt, Germany), supplied with a 10 mm precolumn packed with the same stationary phase. The mobile phase consisted of *n*-hexane (solvent A) and ethyl acetate (solvent B). The gradient program commenced at 90:10 (A:B) for 35 min, followed by a linear gradient from A:B 90:10 to 100% B for 5 min and maintained at 100% B for 20 min. Prior to each sample injection, the column was re-equilibrated for 10 min at the initial conditions. The flow rate was adjusted to 4 mL/min, the column was maintained at ambient temperature, and the absorbance was monitored at 310 nm (optical bandwidth 15 nm). The active F₂₂ fraction was introduced (1 mg/20 μL EtOAc per injection) onto the column via a Rheodyne 7125 injector (Rohnert Park, CA 94928) fitted with a 20 μL sample loop. Seven fractions were collected [F₂₂₁ (18.9 mg), F₂₂₂ (9.1 mg), F₂₂₃ (18 mg), F₂₂₄ (9.1 mg), F₂₂₅ (9.4 mg), F₂₂₆ (22.3 mg), and F₂₂₇ 13 mg)]. One major peak was prominent in each of the fractions F₂₂₂, F₂₂₄, and F₂₂₆, while several minor ones were present in the other four fractions (Figure 1). One hundred

milligrams of the F₂₂ fraction was fractionated in total; the fractions were concentrated and stored at −20 °C until used.

Identification of the Major HPLC Active Fractions: UV Spectroscopy. UV spectra of fractions F₂₂₂, F₂₂₄, and F₂₂₆ were obtained during the HPLC fractionation of the active fraction (F₂₂). They were recorded in the region of 245–365 nm and printed (GP-100 Graphics Printer, Perkin-Elmer) (Figure 1).

GC-FTIR. Analysis of the active fractions was carried out on a Perkin-Elmer Autosystem gas chromatograph interfaced to a Perkin-Elmer System 2000 (Perkin-Elmer Ltd., Buckinghamshire, England) Fourier transform infrared spectrometer equipped with a liquid-nitrogen cooled narrow-band (4000–750 cm^{−1}) infrared detector (mercury cadmium telluride). Helium was used as the carrier gas (2 mL/min) and the effluent from the GC mixed with the sweep gas helium (0.5 mL/min; transfer line temperature 280 °C) and passed through the IR light pipe with KBr windows. All spectra were obtained at 8 cm^{−1} resolution. All samples were chromatographed on a 30 m × 0.32 mm × 1 μm film thickness DB-5 column (J&W Scientific, Folsom, CA). The oven temperature program was 50 °C for 2 min and then 5 °C/min to 280 °C, where it was held for 60 min. Splitless injections were made (1 μL) at an injector temperature of 250 °C and a splitless period of 90 s.

GC-MS. Gas chromatography–mass spectrometry analysis of F₂₂₂ and F₂₂₄ fractions was carried out on a Hewlett-Packard 5890 Series II gas chromatograph interfaced to a Fisons VG Trio 1000 (Manchester M23 9BE, UK) quadrupole mass spectrometer. Electron impact ionization was used at 70 eV and a trap current of 200 μA. The interface was kept at 290 °C. All samples were chromatographed on a 60 m × 0.25 mm × 0.1 μm film thickness DB-5 column (J&W Scientific). The oven temperature was set at 50 °C for 2 min, raised to 250 at 5 °C/min, held for 1 min, raised 2 °C/min to 280 °C, and held there for 50 min. Helium was used as the carrier gas at a flow rate of 1 mL/min. Splitless injections were made (1 μL) at an injector temperature of 250 °C and a splitless period of 90 s. The mass spectrometric data for fraction F₂₂₆ were obtained using the direct contact inlet (DCI) probe of the VG Trio 1000 mass spectrometer. The DCI was set at 0 mA for 2 min and then raised to 200 mA at 100 mA/min, held for 2 min, raised then 300 mA/min to 1000 mA, held for 4 min, and then ramped down at 200 mA/min to 0 mA, where it stayed for 5 min.

¹H NMR. ¹H NMR spectra of F₂₂₂, F₂₂₄, and F₂₂₆ fractions were recorded on Bruker Advance DPX-400 spectrometer. CDCl₃ was used




Figure 2. Chemical structures of osthol (1), bergapten (2), and 6',7'-epoxybergamottin (3).

Table 1. Toxicity of *Citrus aurantium* Flash Chromatography Fractions F_2 and F_{22} ; the Three Major HPLC Fractions F_{222} , F_{224} , F_{226} ; and the Corresponding Synthetic Compounds against Olive Fruit Fly *Bactrocera oleae* Adults in a Petri Dish Exposure Bioassay 72 h Post-Treatment^{a,b}

fractions	I. concentration set to 5 $\mu\text{g}/\text{cm}^2$	II. concentration equivalent to natural presence	
	% mortality	concentration ($\mu\text{g}/\text{cm}^2$)	% mortality
F_2	96 $\pm$ 3.1a	20.0	94 $\pm$ 2.4a
F_{22}	98 $\pm$ 2.0a	4.9	90 $\pm$ 3.2a
F_{222}	0 $\pm$ 0b	0.45	2 $\pm$ 2c
F_{224}	0 $\pm$ 0b	0.45	2 $\pm$ 2c
F_{226}	100 $\pm$ 0a	1.12	46 $\pm$ 6.8b
$F_{222} + F_{224} + F_{226}$		0.45 + 0.45 + 1.12	80 $\pm$ 3.2a
6',7'-epoxybergamottin		1.12	48 $\pm$ 7.3b
osthol + bergapten + 6',7'-epoxybergamottin		0.45 + 0.45 + 1.12	74 $\pm$ 8.1a
control	0 $\pm$ 0b		2 $\pm$ 2c

^a Data were transformed into arcsine $\sqrt{p}$ value before ANOVA and Tukey's studentized range honestly significant difference (HSD) test. Values followed by the same letter within a column are not significantly different at the 0.05 level by HSD test. SE = standard error of mean. ^b Concentrations of all fractions were selected according to their participation in the preceding purification step, respectively. Concentrations of synthetic compounds were equal to those of their corresponding HPLC fractions.

as solvent. Tetramethylsilane or residual solvent peaks (e.g., chloroform) were used as frequency standards. Coupling constants were determined using the computer program Multiplot and are quoted in hertz (Hz). Values were reported to 0.1 Hz, but had an uncertainty of ca. ± 0.3 Hz (at 400 MHz), due to the digital resolution of the FID accumulation and Fourier transformation. ^1H NMR spectra were simulated using RACCOON.

Synthesis of 6',7'-Epoxybergamottin. We synthesized 6',7'-epoxybergamottin using synthetic bergamottin and 3-chloroperbenzoic acid (MCPBA) (11, 12). The synthetic 6',7'-epoxybergamottin was HPLC purified using the same chromatographic conditions as the natural isolate from the peel extract.

Component 1: Osthol ($\text{C}_{15}\text{H}_{16}\text{O}_3$). Component 1 was obtained as a white powder, mp 83–84 $^\circ\text{C}$. UV (λ_{max}): (250, 310). GC–MS m/z (% intensity): 244 (100), 229 (77), 213 (48), 201 (72), 189 (81), 159 (40), 131 (62). IR (cm^{-1}): 3000, 2800, 1770, 1600, 1450, 1300, 1050. ^1H NMR δ_{H} (CDCl_3 , 400 MHz): 7.55 (1H, d, J = 9.4 Hz, 4-H), 7.20 (1H, d, J = 8.6 Hz, 5-H), 6.77 (1H, d, J = 8.6 Hz, 6-H), 6.17 (1H, d, J = 9.4 Hz, 3-H), 5.16 (1H, t, J = 7.2 Hz, 2'-H), 3.85 (3H, s, 6'-H), 3.45 (2H, d, J = 7.2 Hz, 1'-H), 1.78 (3H, s, 4'-H), 1.59 (3H, s, 5'-H).

The UV, IR, and GC–MS spectra were similar to that reported for osthol isolated from the wild celery *Apium graveolens* and the nonvolatile fraction of the grapefruit essential oil (13, 14).

Component 2: Bergapten ($\text{C}_{12}\text{H}_8\text{O}_4$). Component 2 was obtained as white needles, mp 188–191 $^\circ\text{C}$. UV (λ_{max}): (250, 305). GC–MS m/z (% intensity): 216 (100), 201 (35), 188 (15), 173 (64), 145 (33). IR (cm^{-1}): 3000, 2800, 1770, 1600, 1450, 1300, 1050. ^1H NMR δ_{H} (CDCl_3 , 400 MHz): 8.14 (1H, d, J = 9.9 Hz, 5-H), 7.57 (1H, d, J = 2.4 Hz, 2-H), 7.11 (1H, s, br, 9-H), 7.01 (1H, dd, J = 2.4 and 0.8 Hz, 3-H), 6.27 (1H, d, J = 9.9 Hz, 6-H), 4.27 (3H, s, 1'-H). The ^1H NMR and the GC–MS spectra was consistent with that reported for bergapten isolated from *Glehnia littoralis* (Umbelliferae) and from *Euodia borbonica* (Rutaceae) (15, 16).

Component 3: 6',7'-Epoxybergamottin ($\text{C}_{21}\text{H}_{22}\text{O}_5$). Component 3 was obtained as white needles, mp 55–56 $^\circ\text{C}$. UV (λ_{max}): (250, 305). GC–MS m/z (% intensity): 202 (69), 174 (42), 153 (30), 145 (27), 71 (100). IR (cm^{-1}): 3000, 2800, 1770, 1600, 1450, 1300, 1050. ^1H NMR δ_{H} (CDCl_3 , 400 MHz): 8.09 (1H, d, J = 9.9 Hz, 5-H), 7.53 (1H, d, J = 2.5 Hz, 2-H), 7.09 (1H, s, 9-H), 6.88 (1H, dd, J = 2.5 and 0.9 Hz, 3-H), 6.21 (1H, d, J = 9.9 Hz, 6-H), 5.52 (1H, m, 2'-H), 4.89 (2H, d, J = 6.9 Hz, 1'-H₂), 2.64 (1H, dd, J = 7.0 and 5.4 Hz, 6'-H), 2.15 (2H, m, 4'-H), 1.67 (3H, s, 10'-H), 1.55 (2H, m, 5'-H), 1.25 (3H, s, 9'-H), 1.23 (3H, s, 8'-H). The ^1H NMR spectrum of 6',7'-epoxybergamottin was consistent to that reported for 6',7'-epoxybergamottin isolated from *Citrus macroptera* and that from grapefruit juice (11, 17).

Insects. Adult olive fruit flies *B. oleae*, were obtained from an artificially reared colony maintained in our laboratory at NCSR "Demokritos" for many generations. All the insects' stages were kept under the same rearing conditions: 25 $\pm$ 2 $^\circ\text{C}$, 65 $\pm$ 5% relative humidity, and 12:12 light:dark regime.

Toxicity Studies. The Petri dish exposure bioassay described elsewhere (10) with slight modifications was used to evaluate the insecticidal activity of the fractions eluted from the silica gel and the HPLC columns and their blends, plus the three synthetic active components. The tested concentrated samples diluted in 1 mL of dichloromethane plus olive oil (0.65% v/v) were prepared. Groups of 5 pairs of 2–3-day-old flies of *B. oleae* were chill-immobilized for 5 min and subsequently introduced into each of the treated Petri dish and exposed for 12 h. Control Petri dishes were treated in a similar manner with dichloromethane plus olive oil only. Five replicates of 10 flies were employed for each treatment and the control.

Statistics. Mortality data were analyzed by one-way analysis of variance (ANOVA). Raw mortality data were transformed to arcsine $\sqrt{p}$ before ANOVA analysis, where p is the proportion of dead insects, in order to equalize the variance among treatments. Comparisons and separations of means were performed by Tukey's honestly significant difference (Tukey's HSD) test (18) at P = 0.05. SPSS 8.0 software (SPSS Inc., Chicago, IL) was used for our statistical analyses.

RESULTS AND DISCUSSION

Isolation and Identification of the Active Compounds. The fractions EtOAc- F_2 and 40% EtOAc in Hex (F_{22}) from the two flash chromatographic steps induced a substantial mortality level to the olive fruit flies. Further HPLC fractionation of the F_{22} fraction yielded three active components (fractions F_{222} , F_{224} , and F_{226}) (Figure 1). When these three fractions were tested as a blend on the flies (in the same proportions as they were in fraction F_{22} : 0.45 + 0.45 + 1.12 $\mu\text{g}/\text{cm}^2$) high mortality rates were recorded. The chemical structures of the compounds in F_{222} , F_{224} , and F_{226} were assigned as 7-methoxy-8-(3'-methyl-2'-butenyl)-2H-1-benzopyran-2-one (osthol) in F_{222} , 4-methoxy-7H-furo[3,2-g]benzopyran-7-one (bergapten) in F_{224} , and 4-((E)-3'-methyl-5'-(3'',3''-dimethyloxiran-2''-yl)pent-2'-enyloxy)-7H-furo[3,2-g][1]benzopyran-7-one (6',7'-epoxybergamottin) in F_{226} (Figure 2).

Toxicity Studies. The active fractions F_2 and F_{22} eluted from the silica gel columns were tested and attained a 96% (EtOAc- F_2) and 98% (F_{22}) mortality 72 h after treatment at

a concentration of 5 $\mu\text{g}/\text{cm}^2$ (Table 1.I). When 5 $\mu\text{g}/\text{cm}^2$ of F₂₂₆ was tested, the same insect mortality was attained. However, when F₂₂₆ was tested at concentrations equivalent to that of its natural presence in fraction F₂₂, a lesser degree in insect mortality was achieved (Table 1.II). 6',7'-Epoxybergamottin (fraction F₂₂₆) exhibited substantial toxicity when tested individually, while the other major components, osthol in fraction F₂₂₂ and bergapten in F₂₂₄, were inactive. This is the first report of the insecticidal activity of 6',7'-epoxybergamottin. However, 6',7'-epoxybergamottin has been reported to inhibit the activity of the enzyme cytochrome P450 3A4 (CYP3A4) (19, 20). Comparison of the insecticidal activity of 6',7'-epoxybergamottin and its precursor, bergamottin, revealed that the epoxide group is essential for the insecticidal properties of 6',7'-epoxybergamottin, since bergamottin was inactive even at high concentrations (data not shown). Although bergamottin did not show any significant effect on the olive fruit flies when tested individually in the Petri dish exposure bioassay, it has been reported to exert antifeedant activity to several species of *Lepidoptera* (21, 22). The coumarin osthol when tested alone did not exhibit any activity on *B. oleae* flies. However, osthol is known as an antibacterial, antimalarial, antimutagenic, cytostatic, and cytotoxic agent (23, 24). Also the antifeedant activity of bergapten is well-documented (21, 22, 25).

Mixing of the three components (osthol, bergapten and 6',7'-epoxybergamottin) increases the adults' mortality to the level of the initial fraction (F₂₂), indicating the occurrence of a synergistic effect (Table 1.II). Furanocoumarins have been reported to elicit behavioral responses and physiological effects in herbivorous insects. They are repellents and feeding deterrents (5) as well as toxins (26). In addition to their inherent toxicity, they may act synergistically (21) or antagonistically (27) when mixed with other coumarins or furanocoumarins. Toxicity in most instances is greatly enhanced in the presence of UV light, suggesting that photoactivation, possibly involving DNA, is a major mechanism of toxicity (28). However, there are also toxic effects of furanocoumarins, presently known, that are independent of UV light (26).

Considering the advantages of using botanical insecticides for pest management, it can be concluded that *C. aurantium* peel extracts could have great potential for the control of insect pests, particularly taking into account the effect of synergism present in this case. It is well-established that the evolution of insect resistance to plant extracts is extremely slow, in contrast to resistance to pure compounds, which develops rapidly (29).

ACKNOWLEDGMENT

We gratefully acknowledge Dr. David Kelly (Department of Chemistry, Cardiff University) for his help on the ¹H NMR analyses. We also thank Dr. M. Berenbaum (Department of Entomology, University of Illinois) for providing authentic osthol samples.

LITERATURE CITED

- (1) Ezeonu, C. F.; Chidume, G. I.; Udedi, S. C. Insecticidal properties of volatile extracts of orange peels. *Bioresour. Technol.* **2001**, *76*, 273–274.
- (2) Salvatore, A.; Borkosky, S.; Willink, E.; Bardón, A. Toxic effects of lemon peel constituents on *Ceratitis capitata*. *J. Chem. Ecol.* **2004**, *30*, 323–333.
- (3) Stange, R. R.; Midland, S. L.; Sims, J. J. An antifungal compound produced by grapefruit and Valencia orange after wounding of the peel. *J. Nat. Prod.* **1993**, *56*, 1627–1629.
- (4) Jayaprakaha, G. K.; Signh, R. P.; Pereira, J.; Sakariah, K. K. Limonoids from *Citrus reticulata* and their moulting inhibiting activity in mosquito *Culex quinquefasciatus* larvae. *Phytochemistry* **1997**, *40*, 142–146.
- (5) Berdegué, M.; White, K. K.; Trumble, J. T. Feeding deterrence of *Spodoptera exigua* (Lepidoptera: Noctuidae) larvae by low concentrations of linear furanocoumarins. *Environ. Entomol.* **1997**, *26*, 912–919.
- (6) Su, H. C. F.; Horvat, R. Isolation and characterization of four major components from insecticidally active lemon peel extract. *J. Agric. Food Chem.* **1987**, *35*, 509–511.
- (7) Del Río, J. A.; Arcas, M. C.; Botia, J. M.; Ortuño, A. M. Involvement of phenolic compounds in the antifungal defence mechanisms of *Olea europaea* L. and *Citrus* sp. *Recent Res. Dev. Agric. Food Chem.* **2004**, 331–341.
- (8) Del Río, J. A.; Gómez, P.; Arcas, M. C.; Botia, J. M.; Ortuno, A. Changes in the leaves of polymethoxyflavones and flavanones as part of the defense mechanism of *Citrus sinensis* (Cv. Valencia Late) fruits against *Phytophthora citrophthora*. *J. Agric. Food Chem.* **2004**, *52*, 1913–1917.
- (9) Stange, R. R.; Midland, S. L.; Sims, J. J.; Greg McCollum, T. Differential effects of citrus peel extracts on growth of *Penicillium digitatum*, *P. italicum*, and *P. expansum*. *Physiol. Mol. Plant Pathol.* **2002**, *61*, 303–311.
- (10) Siskos, E. P.; Konstantopoulou, M. A.; Mazomenos, B. E.; Jervis, M. Insecticidal activity of *Citrus aurantium*, fruit, leaf and shoot extracts against adults of the olive fruit fly *Bactrocera oleae* Gmel. (Diptera: Tethritidae). *J. Econ. Entomol.* **2007**, *100*, 1215–1220.
- (11) Dreyer, D. L.; Huey, P. F. Coumarins of *Citrus macroptera*. *Phytochemistry* **1973**, *12*, 3011–3013.
- (12) Bellevue, F. H.; Woster, P. M.; Edwards, D. J.; He, K.; Hollenberg, P. F. Synthesis and biological evaluation of 6,7-dihydroxybergamottin (6,7-DHB), a naturally occurring inhibitor of cytochrome P450 3A4. *Bioorg. Med. Chem. Lett.* **1997**, *7*, 2593–2598.
- (13) Hadaček, F.; Müller, C.; Werner, A.; Greger, H.; Proksch, P. Analysis isolation and insecticidal activity of linear furanocoumarins and other derivatives from *Peucedanum* (Apiaceae: Apiaceae). *J. Chem. Ecol.* **1994**, *20*, 2035–2054.
- (14) Zobel, A. M.; Wang, J.; March, R. E.; Brown, S. A. Identification of eight coumarins occurring with psoralen, xanthotoxin, and bergapten on leaf surfaces. *J. Chem. Ecol.* **1991**, *17*, 1859–1870.
- (15) Buiarelli, F.; Cartoni, G. P.; Coccioli, F.; Leone, T. Analysis of bitter essential oils from orange and grapefruit by high-performance liquid chromatography with microbore columns. *J. Chromatogr. A* **1996**, *730*, 9–16.
- (16) Sasaki, H.; Taguchi, H.; Endo, T.; Yosioka, I. The constituents of *Glehnia littoralis* FR SCHMID et MIQ. Structure of a new coumarin glycoside, osthenol-7-O- β -gentiobioside. *Chem. Pharm. Bull.* **1980**, *28*, 1847–1852.
- (17) Velenciennes, E.; Smadja, J.; Conan, J. Y. Screening for biological activity and chemical composition of *Euodia borbonica* var. *borbonica* (Rutaceae) a medicinal plant in Reunion Island. *J. Ethnopharmacol.* **1999**, *58*, 663–668.
- (18) Tukey, J. W. Some selected quick and easy methods of statistical analysis. *Trans. N. Y. Acad. Sci. Ser. 2* **1953**, *16*, 88–97.
- (19) Ohta, T.; Maruyama, T.; Nagahashi, M.; Miyamoto, Y.; Hosoi, S.; Kiuchi, F.; Yamazoe, Y.; Tsukamoto, S. Paradisin C: A new CYP3A4 inhibitor from grapefruit juice. *Tetrahedron* **2002**, *58*, 6631–6635.
- (20) Wangensteen, H.; Molden, E.; Christensen, H.; Malterud, K. E. Identification of epoxybergamottin as a CYP3A4 inhibitor in grapefruit peel. *Eur. J. Clin. Pharmacol.* **2003**, *58*, 663–668.
- (21) Calcagno, M. P.; Coll, J.; Lloria, J.; Faini, F.; Alonso-Amelot, M. E. Evaluation of synergism in the feeding deterrence of some furanocoumarins on *Spodoptera littoralis*. *J. Chem. Ecol.* **2002**, *28*, 175–191.
- (22) Stevenson, P. C.; Simmonds, M. S. J.; Yule, M. A.; Veitch, N. C.; Kite, G. C.; Irwin, D.; Legg, M. Insect antifeedant furanocoumarins from *Tetradium danielli*. *Phytochemistry* **2003**, *63*, 41–46.

- (23) Harborne, J. B.; Baxter, H. *Phytochemical Dictionary: A Handbook of Bioactive Compounds from Plants*; Taylor & Francis: London, 1993.
- (24) Yang, L. L.; Wang, M. C.; Chen, L. G.; Wang, C. C. Cytotoxic activity of coumarins from the fruits of *Cnidium monnieri* on leukaemia cell lines. *Planta Med.* **2003**, *62*, 1091–1095.
- (25) Escoubas, P.; Fukushi, Y.; Lajide, L.; Mizutani, J. A new method for fast isolation of insect antifeedant compounds from complex mixtures. *J. Chem. Ecol.* **1992**, *18*, 1819–32.
- (26) Berenbaum, M. Toxicity of a furanocoumarin to armyworms: A case of biosynthetic escape from insect herbivores. *Science* **1978**, *201*, 532–534.
- (27) Diawara, M. M.; Trumble, J. T.; White, K. K.; Carson, W. G.; Martinez, L. A. Toxicity of linear furanocoumarins to *Spodoptera exigua*: Evidence for antagonistic interactions. *J. Chem. Ecol.* **1993**, *19*, 2473–2484.
- (28) Berenbaum, M. R.; Nitao, J. K.; Zangerl, A. R. Adaptive significance of furanocoumarin diversity in *Pastina sativa* (Apiaceae). *J. Chem. Ecol.* **1991**, *17*, 207–215.
- (29) Feng, R.; Isman, M. B. Selection for resistance to azadirachtin in the green peach aphid *Myzus persicae*. *Experientia* **1995**, *51*, 831–833.

Received for review February 13, 2008. Revised manuscript received May 17, 2008. Accepted May 19, 2008.

JF800446T