

Published on Web 11/06/2007

Photo-Leucine Incorporation Reveals the Target of a Cyclodepsipeptide Inhibitor of Cotranslational Translocation

Andrew L. MacKinnon, Jennifer L. Garrison, Ramanujan S. Hegde, and Jack Taunton* Department of Cellular and Molecular Pharmacology, University of California, San Francisco, California 94158 Received August 19, 2007; E-mail: taunton@cmp.ucsf.edu

Photoaffinity labeling is a powerful tool to identify protein targets of biologically active small molecules and probe the structure of ligand binding sites, especially in the case of integral membrane proteins.¹ A difficulty with photoaffinity labeling is that the small molecule must retain biological activity after derivatization with a photoreactive group such as benzophenone² or 3-trifluoromethyl-3-phenyl diazirine.³ While these substituents are effective photocross-linkers and have been incorporated into small molecules,⁴ peptides,⁵ and proteins,⁶ their large size can potentially interfere with small molecule/protein interactions. The amino acid isosteres, photo-leucine and photo-methionine, have smaller alkyl diazirine side chains and were recently used to probe protein-protein interactions after random biosynthetic incorporation in cells7 or via site-specific native protein ligation.⁸ By contrast, the potential for incorporating these novel photoreactive amino acids into natural product scaffolds has not been explored. In this communication we report (1) an improved synthesis of photo-leucine, (2) incorporation of photo-leucine and an alkyne click chemistry tag into a cyclodepsipeptide inhibitor of protein secretion, and (3) identification of the inhibitor's target by photoaffinity labeling.



HUN-7293 (1) is a fungal cyclodepsipeptide that was first identified as an inhibitor of vascular cell adhesion molecule (VCAM) expression.⁹ Recently, we¹⁰ and others¹¹ discovered that related cyclodepsipeptides potently block the cotranslational translocation of VCAM and a subset of other proteins into the endoplasmic reticulum (ER), an early step in the biogenesis of secretory and membrane proteins. Biochemical experiments revealed that these compounds act at the ER membrane to perturb interactions between nascent ribosome-associated VCAM chains and the translocation channel. A heterotrimeric membrane protein, the Sec61 complex, forms the structural core of this channel¹² and thus emerged as a potential direct target of **1**.

To identify the target, we designed photoaffinity probe 2, in which photo-leucine replaces leucine at position 4 of 1. We substituted *N*-methylphenylalanine at position 5, previously shown to have a negligible effect on potency.¹³ A propargyl substituent was installed at position 1 to enable Cu(I)-catalyzed conjugation of a rhodamine—azide reporter¹⁴ (click chemistry) after photo-cross-linking under native conditions (Figure 1). These conservative



Figure 1. Photoaffinity labeling strategy to identify the target of 2.

Scheme 1. Synthesis of Boc-(S)-Photo-Leucine



modifications were aimed at preserving the biological potency of 1 while providing the necessary functionality for target identification.

Synthesis of **2** required Boc-(*S*)-photo-leucine (**6**), which we prepared via ozonolysis of commercially available **4**,¹⁵ followed by formation of the diazirine by the method of Church and Weiss (Scheme 1).¹⁶ Boc-(*S*)-photo-leucine was efficiently coupled to peptides (EDCI-HOAt) or deprotected (4 N HCl) to give the free amino acid in quantitative yield. This route is a significant improvement over the original six-step synthesis of (*S*)-photo-leucine, which proceeded in low yield and required enzymatic resolution of a racemic intermediate.⁷ Synthesis of **2** followed the solution-phase route developed by Boger and co-workers¹³ with slight modifications. The diazirine side chain was stable to ambient light and the acidic and basic conditions used to prepare **2**. We also synthesized **3** as a photostable control compound. Both **2** and **3** were equipotent to the natural product **1** at inhibiting VCAM expression in transfected cells (EC₅₀ ~ 25 nM, Figure S1).

We incubated a crude ER microsome fraction with 500 nM 2 and irradiated the mixture with \sim 350 nm light for 1 min. Proteins were denatured in 1% sodium dodecylsulfate (SDS) and subjected to standard click chemistry conditions using a rhodamine–azide reporter (see Supporting Information). Following electrophoresis, in-gel fluorescent scanning revealed a major rhodamine-labeled protein with an apparent molecular weight of \sim 50 kDa (Figure 2A, lane 1). Labeling of this protein required both UV light (lane



Figure 2. (A) Photo-cross-linking of 2 in the presence of ER microsomes, followed by click chemistry with a rhodamine-azide reporter. An asterisk marks the major photo-cross-linked protein. (B) Sec61a antibody or rabbit serum (control) was added to photolysis/click reactions, and immunoprecipitates (IP) analyzed by in-gel fluorescent scanning. (C) Proteoliposomes derived from control extracts (lane 3), Sec61-depleted extracts (lane 1), or depleted extracts replenished with purified Sec61 (lane 2) were photolyzed in the presence of 2 and analyzed by click chemistry/in-gel fluorescent scanning. The major cross-linked protein at \sim 50 kDa is shown.

5) and 2 (lane 6) and was competed by excess 3 (lanes 2-4). Weak labeling of two additional proteins (~60 and ~40 kDa) was not competed by excess 3 and is likely nonspecific. Background labeling by the rhodamine-azide, independent of UV light or 2, was also observed (lanes 5 and 6).

The major cross-linked protein migrated on SDS gels with the same relative mobility as Sec61 α , the largest subunit of the Sec61 complex (Figure S2). Consistent with Sec 61α as the primary target of 2, the \sim 50 kDa rhodamine-labeled protein was immunoprecipitated directly from the click reaction mixture with an antibody raised against Sec61 α , but not with a control antibody (Figure 2B).

To independently confirm that Sec61 α is the photo-cross-linked protein, we prepared proteoliposomes reconstituted from either a detergent extract of ER microsomes, a Sec61-depleted extract, or a depleted extract replenished with purified Sec61 complex.¹⁷ Depletion of the intact Sec61 complex from the detergent extract using antibodies against Sec61 β^{18} was ~90% efficient (as judged by immunoblotting for the α and β subunits), whereas other ER proteins were not affected (Figure S3). Compared to proteoliposomes derived from control extracts (Figure 2C, lane 3), photocross-linking of 2 to the \sim 50 kDa protein was significantly diminished in Sec61-depleted proteoliposomes (lane 1) and was restored after adding back purified Sec61 complex (lane 2). We conclude that Sec61 α , thought to form the channel through which all proteins transit as they enter the secretory pathway, is the primary target of 2 in the ER.

Photo-cross-linking of 2 to Sec61 α is remarkably selective given the proteomic complexity of the ER, which contains hundreds of membrane and luminal proteins;¹⁹ Sec61a is a relatively minor constituent (~0.7% of total ER protein).²⁰ We attribute this high selectivity to two factors. First, due to its similar size, shape, and

hydrophobicity, photo-leucine likely forms intimate contacts with Sec61 α in a manner similar to the leucine side chain of 1. Second, the short lifetime (nanoseconds)²¹ of the carbene derived from **2** (Figure 1) ensures that molecules not bound to Sec61 α are rapidly quenched by intramolecular rearrangement, solvent, or membrane lipids. Despite the short lifetime, the photo-cross-linking yield of 2 to Sec61 α was estimated to be ~23% (Figure S4). Thus, depending on the specific application, alkyl diazirines may offer advantages over the widely used benzophenone cross-linker. The modular cyclodepsipeptide scaffold of 1 should facilitate installation of a diazirine at multiple positions. High-resolution mapping of the cross-linking site(s) by mass spectrometry will likely shed light on the mechanism by which these compounds selectively inhibit cotranslational translocation.

Acknowledgment. This work was supported by the NIH (GM81644), the intramural research program of the NICHD, and the Scleroderma Research Foundation.

Supporting Information Available: Detailed experimental section, synthesis, and spectral characterization of new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- (1) For reviews, see: (a) Sadakane, Y.; Hatanaka, Y. Anal. Sci. 2006, 22, 209-218. (b) Kotzyba-Hibert, F.; Kapfer, I.; Goeldner, M. Angew. Chem., Int. Ed. Engl. 1995, 34, 1296-1312
- (3)
- Dormán, G.; Prestwich, G. D. Biochemistry 1994, 33, 5661–5673.
 Brunner, J. Annu. Rev. Biochem. 1993, 62, 483–514.
 For recent examples, see: (a) Kotake, Y.; Sagane, K.; Owa, T.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, Within and M. S. Sagara, K.; Owa, T.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, H.; Uesugi, M.; Ishihama, Y.; Iwata, W.; Mizui, K.; Mimori-Kiyosue, Y.; Shimizu, K.; Mimori-Kiyosue, Y.; Mimori-Kiyos (4)Klyosue, I., Simmizi, H., Ocsugi, M., Ishinana, I., Iwata, W., Mizui, Y. *Nat. Chem. Biol.* 2007, 22, advanced online publication. (b) Fuwa, H.; Takahashi, Y.; Konno, Y.; Watanabe, N.; Miyashita, H.; Sasaki, M.; Natsugari, H.; Kan, T.; Fukuyama, T.; Tomita, T.; Iwatsubo, T. *ACS Chem. Biol.* 2007, 2, 408–418.
 (5) Kauer, J. C.; Erickson-Viitanen, S.; Wolfe, H. R., Jr.; DeGrado, W. F. J.
- Biol. Chem. 1986, 261, 10695-10700.
- (6) (a) Chin, J. W.; Martin, A. B.; King, D. S.; Wang, L.; Shultz, P. G. Proc. Natl. Acad. Sci. U.S.A. 2002, 99, 11020-11024. (b) High, S.; Martoglio, B.; Görlich, D.; Andersen, S. S. L.; Ashford, A. J.; Giner, A.; Hartmann, E.; Prehn, S.; Rapoport, T. A.; Dobberstein, B.; Brunner, J. J. Biol. Chem. **1993**, *268*, 26745–26751.
- (7) Suchanek, M.; Radzikowska, A.; Thiele, C. Nat. Methods 2005, 2, 261-268.
- (8) Vila-Perelló, M.; Pratt, M. R.; Tulin, F.; Muir, T. W. J. Am. Chem. Soc. 2007. 129. 8068-8069.
- (9)(a) Foster, C. A.; Dreyfuss, M.; Mandak, B.; Meingassner, J. G.; Naegeli, H. U.; Nussbaumer, A.; Oberer, L.; Scheel, G.; Swoboda, E. M. J. Dermatol. 1994, 21, 847–854. (b) Boger, D. L.; Keim, H.; Oberhauser, B.; Schreiner, E. P.; Foster, C. A. J. Am. Chem. Soc. 1999, 121, 6197-6205
- (10) Garrison, J. L.; Kunkel, E. J.; Hegde, R. S.; Taunton, J. Nature 2005, 436, 285–289.
- (11) Besemer, J.; Harant, H.; Wang, S.; Oberhauser, B.; Marquardt, K.; Foster, C. A.; Schreiner, E. P.; de Vries, J. E.; Dascher-Nadel, C.; Lindley, I. L. Nature 2005, 436, 290-293.
- (12) Osborne, A. R.; Rapoport, T. A.; van den Berg, B. Annu. Rev. Cell Dev. Biol. 2005, 21, 529–550.
- (13) Chen, Y.; Bilban, M.; Foster, C. A.; Boger, D. L. J. Am. Chem. Soc. 2002, 124, 5431–5440.
- (14)(a) Speers, A. E.; Cravatt, B. F. Chem. Biol. 2004, 11, 535-546. (b) Cohen, M. S.; Hadjivassiliou, H.; Taunton, J. Nat. Chem. Biol. 2007, 3, 156-160.
- (15) Cornish, V. W.; Hahn, K. M.; Shultz, P. G. J. Am. Chem. Soc. 1996, 118, 8150-8151.
- (16) Church, R. F. R.; Weiss, M. J. J. Org. Chem. 1970, 35, 2465-2471.
- Görlich, D.; Rapoport, T. A. Cell 1993, 75, 615-630. (17)
- (18) Fons, R. D.; Bogert, B. A.; Hegde, R. S. J. Cell Biol. 2003, 160, 529-539
- (19) Gilchrist, A.; Au, C. E.; Hiding, J.; Bell, A. W.; Fernandez-Rodriguez, J.; Lesimple, S.; Nagaya, H.; Roy, L.; Gosline, S. J. C.; Hallett, M.; Paiement, J.; Kearney, R. E.; Nilsson, T.; Bergeron, J. J. M. Cell 2006, 127, 1265-1281.
- (20) Guth, S.; Völzing, C.; Müller, A.; Jung, M.; Zimmermann, R. *Eur. J. Biochem.* 2004, *271*, 3200–3207.
 (21) Ford, F.; Yuzawa, T.; Platz, M. S.; Matzinger, S.; Fülscher, M. *J. Am. Chem. Soc.* 1998, *120*, 4430–4438.

JA076250Y