

Communication

Combinatorial evolution of fast-conducting highly selective K⁺-channels via modularly tunable directional assembly of crown ethers

Changliang Ren, Jie Shen, and Huaqiang Zeng

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.7b04335 • Publication Date (Web): 24 Aug 2017

Downloaded from <http://pubs.acs.org> on August 24, 2017

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

Combinatorial evolution of fast-conducting highly selective K^+ -channels via modularly tunable directional assembly of crown ethers

Changliang Ren,[†] Jie Shen,[†] and Huaqiang Zeng^{*,†}

[†] Institute of Bioengineering and Nanotechnology, 31 Biopolis Way, The Nanos, Singapore 138669

ABSTRACT: We describe here a modularly tunable molecular strategy for construction and combinatorial optimization of highly efficient K^+ -selective channels. In our strategy, highly robust supramolecular H-bonded 1D ensemble was used to order the appended crown ethers in such a way that they roughly stack on top of each other to form a channel for facilitated ion transport across the membrane. Among 15 channels that all prefer K^+ over Na^+ ions, channel molecule **5F8** shows the most pronounced optimum for K^+ while disfavoring all other biologically important cations (e.g., Na^+ , Ca^{2+} and Mg^{2+}). With a K^+/Na^+ selectivity of 9.8 and an EC_{50} value of 6.2 μM for K^+ ion, **5F8** is clearly among the best synthetic potassium channels developed over the past decades.

Natural ion channels are supremely elegant in structure and particularly so in function. For instance, natural KcsA K^+ -channel is at least 10,000 times more permeant to K^+ than Na^+ ions while conducting K^+ ions at an exceptional speed of 10^8 ions per second.¹ High selectivity in ion transport however currently present a major challenge to the actively researched field of artificially developed channel systems,² and channels having very moderate K^+/Na^+ selectivity of ≥ 10 still remain difficult to develop.^{3a-c} In constructing synthetic ion channels, crown ethers are one type of privileged ion-transporting units that are featured in many approaches. Examples include the hydraphiles by Gokel,^{2f,2g} as well as crowns conjugated to a peptide scaffold by Voyer,^{2e,3d} ferrocene units by Hall,^{3e} a rigid rod-like molecule by Matile^{3f} and (thio)urea groups^{3b-c} by Barboiu.

We recently devised a mono-peptide-based supramolecular chiral synthon with a built-in H-bonding capacity, able to align appended R_1 - R_3 groups at the same side via an H-bonding-mediated directional self-assembling process (Figure 1a-b).^{4a-c} Supported by the crystal structure of **Fmoc-Phe-C4** (Figure 1b),^{4a} this synthon turns out to be highly robust in guiding molecular packing into a predictable one-dimensionally aligned columnar structure.^{4a-c} Prompted by these findings, we thought that modification of the H-bonded 1D network with ion-transporting units such as 15-crown-5 or 18-crown-6 at the R_1 position (Figures 1c-e, S1 and S2, and Table S1) might lead to a new family of ion channels such as **6I8** (**6-Ile-C8**), likely endowed with ability to discriminate biologically important metal ions such as Na^+ , K^+ , Ca^{2+} and Mg^{2+} . One distinct trait of our system is its intrinsic high modularity in backbone, enabling an evolutionary approach with a parallel synthesis of library members to be




Figure 1. a) A supramolecular chiral synthon for aligning side chains of the same type (e.g., R_1 - R_3) at the same side. b) Crystal structure of **Fmoc-Phe-C4**,^{4a} demonstrating the ability of the synthon to form a H-bonded 1D columnar stack with R_1 - R_3 aligned to the same side. The O and H atoms involved in intermolecular H-bonds are highlighted as small red and gray balls, respectively. c) Combinatorial diversifications at the R_1 and R_2 positions for discovering efficient and selective ion channels. Molecules containing $R_2 = \text{Gly} = G$ were used to elucidate the importance of chiral center in the self-assembling process. d) Structures of control compounds. e) Top view of the computationally optimized most stable structure of H-bonded 1D ensemble formed from **6I8** (Figure S2). f) Gelation of a mixed solvent (hexane:toluene = 4:1, v:v) by **5I8** and **6I8** and the corresponding SEM images. The gels were visualized using Sudan III.

applied for rapid discovery of ion channels with good selectivity.

We firstly began with evaluation of **5I8** (**5-Ile-C8**) and **6I8** having an n-octyl side chain at the R₃ position. Consistent with predictable ability to form a H-bonded 1D structure via intermolecular H-bonds, both **5I8** and **6I8** self-assemble into well-defined fibers as evident from their ability to gel a mixed solvent (hexane:toluene = 4:1, v:v) and from the images of the dried gels obtained using scanning electron microscope (SEM, Figure 1f) and transmission electron microscope (TEM) (Figures S3-S6). The ¹H NMR dilution experiments carried out by diluting a solution containing **5I8** or **6I8** from 100 to 0.8 mM in CDCl₃ reveal concentration-dependent chemical shifts for the two amide protons involved in forming intermolecular H-bonds, additionally supporting the formation of intermolecularly H-bonded 1D structure (Figure S7). Biphasic extraction experiments using equal volumes of H₂O containing 7 metal ions (e.g., Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺, Ca²⁺ and Mg²⁺; each at 0.10 mM) in their nitrate salts and CHCl₃ containing an organic host (e.g., **15C5**, **18C6**, **5I8** or **6I8** at 0.70 mM) were carried out at 25 °C to determine their ion-binding affinity and selectivity using inductively coupled plasma mass spectrometry (ICP-MS). ICP data compiled in Table S2 show that all four ligands could extract Na⁺, K⁺ and Ca²⁺ ions to similar extents, suggesting H-bonding network in **5I8** or **6I8** does not noticeably interfere with the intrinsic ion binding ability of the attached crown ether units. But do note that the observed competitive ion extraction behaviors likely can't be simply extrapolated to what may happen in lipid membrane as ion extraction extent and selectivity are strongly influenced by the extraction solvent and the presence of other metal ions as recently reported by Bartsch.^{5a-b}

Indeed, ion transport activities assessed using the pH-sensitive HPTS assay^{4d} at a final channel concentration of 22 μM over a duration of 5 min (Figures 2 and 3) surprisingly demonstrate very different Na⁺ and K⁺ transports across the membrane by **5I8** (R_{Na}⁺ = 6.1% and R_K⁺ = 17%) and **6I8** (R_{Na}⁺ = 6.4% and R_K⁺ = 60%).⁵ In sharp contrast to **6I8**, both the isolated crown ethers (e.g., **15C5** and **18C6**) and molecules containing a non-chiral center (e.g., **5G8** and **6G8**) exhibit very poor ion transport and low ion selectivity (Figure 2b). These differential channel activities in ion transport and selectivity suggest (1) important roles played by the chiral center and built-in H-bonding motif in organizing crown ethers into a functional assembly and (2) one dimensionally aligned 18-crown-6 units could differ very substantially in function from the ensemble based on 15-crown-5 motifs. As illustrated in Table S1 and Figures S1 and S2, side chains of the amino acids at the R₂ position do modify the relative orientation among the vertically aligned crown units, further highlighting the feasibility of using the mono-peptide-based scaffold for a combinatorial search of more selective and faster ion-conducting channels.

A combinatorial screening was then attempted via synthesis of a small library containing another eight crown ether-containing channel molecules derived from alanine (A), valine (V), leucine (L) and phenylalanine (F) with R₃ = C8. At the same final concentration of 22 μM, a common trend shared by all channel molecules is that they all transport K⁺ preferentially over Na⁺ ions with R_K⁺/R_{Na}⁺ values ranging from 1.4 to 7.1 (Figures 2b and 3).⁵ As similarly observed by others,^{3b-c} K⁺ transport generally appears to be faster with channels containing stacked 15-crown-5 than




Figure 2. a) The HPTS-based assay for proton transport study using the pH-sensitive HPTS (8-Hydroxypyrene-1,3,6-trisulfonic acid) trapped inside larger unilamellar vesicles (LUVs) with a pH gradient of 7 to 8. To determine ion selectivity, the external buffer was changed from 100 mM NaCl to 100 mM MCl or 50 mM MCl₂. b) Selected K⁺ transport curves at a channel concentration of 22 μM, illustrating progressively enhanced transports of K⁺ ions upon fine-tuning R₂ group. After subtracting background intensity at t = 0, the ratiometric values of I₄₆₀/I₄₀₃ were normalized with ratiometric value of I₄₆₀/I₄₀₃ at t = 0 s as 0% and that of I₄₆₀/I₄₀₃ at t = 300 s (obtained after addition of triton) as 100%. For more K⁺ and Na⁺ transports, see Figures S8-S11.

those with 18-crown-6, and transport curves (**5F8**, **5F10**, etc), which exceed 100% after normalization, are indicative of high K/Na selectivity where proton influx is more efficient than Na influx at high concentration. While all four channels derived from alanine and valine (**5A8**, **6A8**, **5V8** and **6V8**) turn out to be poor ion conductors with the highest R_{Na}⁺ and R_K⁺ values being 11% and 34%, respectively, those based on leucine and phenylalanine (**5L8**, **5F8** and **6F8**) are highly active toward K⁺ ions with a R_K⁺ value going beyond 100% (Figure 2b), which corresponds to a point where the internal pH was raised from 7 to larger than the pH of external solution. In particular, **5F8**-mediated K⁺ transport is the fastest among all channels studied and is so fast that the time needed for the internal pH to reach an equilibrium pH of 8 takes merely 25 sec. This extremely fast conduction suggests that a R_K⁺/R_{Na}⁺ value of 2.5 at 22 μM underestimate the ion selectivity of **5F8**. Expectedly, upon lowering the concentration from 22 μM down to 8.8 μM at which a smoothly growing curve was obtained with 97% of K⁺ transfer completed in 5 min, the R_K⁺/R_{Na}⁺ value was increased to 14.7 (Figure 4a).^{5c} Similarly for **5L8**, the R_K⁺/R_{Na}⁺ value was determined to be 8.3 at 19.8 μM (Figure




Figure 3. Ion transport percentages determined over a duration of 5 min at 22 μM for all channel molecules except for **5L8** (19.8 μM), **5F8** (8.8 μM), **5F10** (5.5 μM) and **6F8** (19.8 μM). $R_K^+ = (I_K^+ - I_0)/(I_{\text{Triton}} - I_0)$ wherein I_K^+ and I_0 (background intensity) are the ratiometric values of I_{460}/I_{403} at $t = 300$ s before addition of triton, and I_{Triton} is the ratiometric value of I_{460}/I_{403} at $t = 300$ s right after addition of triton with internal/external buffers containing 100 mM NaCl/100 mM KCl, i.e., $(\text{NaCl})_{\text{in}}/(\text{KCl})_{\text{out}}$. For R_{Na}^+ , the internal/external buffers are $(\text{NaCl})_{\text{in}}/(\text{NaCl})_{\text{out}}$.




Figure 4. Transport activities toward Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺, Ca²⁺ and Mg²⁺ ions at a final concentration of (a) 8.8 μM for **5F8** and (b) 5.5 μM for **5F10**, respectively. c) Changes in fluorescence intensity of Safranin O ($\lambda_{\text{ex}} = 522$ nm, $\lambda_{\text{em}} = 581$ nm; 60 nM) after addition of **5F8** and **5F10** at 5 μM and Valinomycin (VA) from 0.02 to 5 μM . The 10 mM HEPES buffer at pH 7 inside LUVs contains 100 mM KCl while external HEPES buffer at pH 7 contains either 100 mM KCl or 100 mM NaCl. In c), the starting background intensity F_0 at $t = 0$ was set to be 1.

S11c). In terms of both ion transport rate and selectivity, **5F8** is considerably much better than the recently reported crown ether-based K⁺-selective channels^{3b,3c} the best of which requires a higher channel concentration of 25 μM to reach a R_K^+ value of 100% but shows a low R_K^+/R_{Na}^+ value of only 6.7.^{3c}

On the basis of **5F8**, our secondary screening looked into the impact the chain length of R₃ group might have on the channel's performance. Our results show that channels **5F8** and **5F10** containing longer alkyl chains (e.g., **C8** and **C10**, Figure 3) are far more efficient and selective in ion transport than **5F4** and **5F6** with shorter ones. A more careful and systematic comparison between **5F8** and **5F10** reveals (1) these two channels display a similar trend in ion selectivity, i.e., their transport activities decrease in the order of K⁺ > Rb⁺ >> Cs⁺ > Na⁺ > Li⁺ \approx Ca²⁺ \approx Mg²⁺ and (2) the more selective **5F8** ($R_K^+/R_{\text{Na}}^+ = 14.7$)^{5c} transports K⁺ ions less efficiently than the less selective **5F10**

($R_K^+/R_{\text{Na}}^+ = 10.0$). The corresponding EC_{50} values (e.g., the concentration required to attain 50% ion permeation) for K⁺ and Rb⁺ ions are 6.2 and 9.4 μM for **5F8** and 2.4 and 3.9 μM for **5F10** (Figure S12), respectively. These comparisons in EC_{50} value indicate that the longer **C10** chains in **5F10** might enhance the stability and/or increase the open probability of the H-bonded channel structure in the membrane more significantly than the shorter **C8** chains do, particularly at low concentrations of < 6 μM .

With an existence of a difference in ion concentration between the interior and exterior of a LUV, selective ion transport will lead to an accumulation of cations on one side of the membrane, resulting in membrane polarization that can be analyzed using Safranin O, a voltage sensitive dye. Upon changing the ion concentration in the external buffer from 100 mM KCl to 100 mM NaCl, both **5F8** and **5F10** at 5 μM cause large changes in fluorescence intensity by > 100% relative to the signal at $t = 50$ s




Figure 5. Single channel current traces of (a) the dimeric gramicidin A recorded at 200 mV and (b) **5F8** recorded at -200 mV, respectively, in symmetrical solutions (*cis* chamber = *trans* chamber = 1 M KCl).

(Figures 4c and S13a-b). Compared to changes of 164% by Valinomycin at 0.2 μ M (Figures 4c and S13c), < 10% by gramicidin A (Figure S13d) and < 20% by other synthetic highly selective K^+ channels,^{3c} such large degrees of membrane polarization caused by **5F8** and **5F10** are very remarkable, establishing highly selective transport of K^+ ions.

Taken together, the above data demonstrate that **5F8** and **5F10** markedly outperform all other 13 channels studied herein, and **5F8** is the most selective in K^+ transport among all. The ability of **5F8** to function as an ion channel rather than a carrier in mediating ions across the membrane was then unambiguously confirmed by the observation of single channel current traces recorded in both symmetrical (*cis* chamber = *trans* chamber = 1 M KCl) and asymmetric (*cis* chamber = 1M KCl and *trans* chamber = 1 M NaCl) baths (Figures 5b and S14d). On the basis of I-V curves obtained (Figure S14a and c), potassium conduction rate (γ_{K^+}) and K^+/Na^+ selectivity were determined to be 26.4 ± 2.8 pS and 9.8, respectively, demonstrating that **5F8**-mediated transport of K^+ ions occurs in a fast and highly selective fashion.

In conclusion, we have established here a privileged monopeptide-based scaffold toward creation of K^+ -selective ion channels via directional assembly of crown ethers. Characterized by high tunability in structure and reliability in channel formation, our channel system represents an unusual example among the hitherto artificially developed ion channel systems. Accordingly, the most advantageous feature of our approach is its intrinsic modularity in appended side chains, which is mostly independent of and thus exerts a tolerable degree of influences on the main chain's H-bonding functionality and self-assembling propensity, thereby enabling consistent formation of crown ether-containing H-bonded 1D structures. Such easily accessible tunability subsequently allows a combinatorial optimization of channels' ion transport efficiency and selectivity, quickly giving rise to two fast-conducting K^+ -selective channels (e.g., **5F8** and **5F10**) that are comparable to the best K^+ -selective channels described in literature over the years.^{2,3}

■ ASSOCIATED CONTENT

Supporting Information. Synthetic procedures and a full set of characterization data including 1H NMR, ^{13}C NMR and MS for all channel molecules as well as ion transport study, single channel current recording, molecular modeling, SEM/TEM images, 1H NMR dilution experiments and ICP data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

hqzeng@ibn.a-star.edu.sg

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENT

This work was supported by the Institute of Bioengineering and Nanotechnology (Biomedical Research Council, Agency for Science, Technology and Research, Singapore) and the Singapore National Research Foundation under its Environment and Water Research Programme and administered by PUB.

■ REFERENCES

- (1) Doyle, D. A.; Morais Cabral, J.; Pfuetzner, R. A.; Kuo, A.; Gulbis, J. M.; Cohen, S. L.; Chait, B. T.; MacKinnon, R. *Science* **1998**, *280*, 69.
- (2) (a) Vargas Jentzsch, A.; Hennig, A.; Mareda, J.; Matile, S. *Acc. Chem. Res.* **2013**, *46*, 2791; (b) Sakai, N.; Matile, S. *Langmuir* **2013**, *29*, 9031; (c) Montenegro, J.; Ghadiri, M. R.; Granja, J. R. *Acc. Chem. Res.* **2013**, *46*, 2955; (d) Fyles, T. M. *Acc. Chem. Res.* **2013**, *46*, 2847; (e) Otis, F.; Auger, M.; Voyer, N. *Acc. Chem. Res.* **2013**, *46*, 2934; (f) Gokel, G. W.; Negin, S. *Acc. Chem. Res.* **2013**, *46*, 2824; (g) Gokel, G. W.; Mukhopadhyay, A. *Chem. Soc. Rev.* **2001**, *30*, 274; (h) Davis, A. P.; Sheppard, D. N.; Smith, B. D. *Chem. Soc. Rev.* **2007**, *36*, 348; (i) Davis, J. T.; Okunola, O.; Quesada, R. *Chem. Soc. Rev.* **2010**, *39*, 3843; (j) Zhao, Y.; Cho, H.; Widanapathirana, L.; Zhang, S. *Acc. Chem. Res.* **2013**, *46*, 2763; (k) Gong, B.; Shao, Z. *Acc. Chem. Res.* **2013**, *46*, 2856; (l) Si, W.; Xin, P.; Li, Z.-T.; Hou, J.-L. *Acc. Chem. Res.* **2015**, *48*, 1612; (m) Huo, Y. P.; Zeng, H. Q. *Acc. Chem. Res.* **2016**, *49*, 922.
- (3) (a) Wright, A. J.; Matthews, S. E.; Fischer, W. B.; Beer, P. D. *Chem. Eur. J.* **2001**, *7*, 3474; (b) Sun, Z.; Barboiu, M.; Legrand, Y.-M.; Petit, E.; Rotaru, A. *Angew. Chem., Int. Ed.* **2015**, *127*, 14681; (c) Gilles, A.; Barboiu, M. *J. Am. Chem. Soc.* **2016**, *138*, 426; (d) Otis, F.; Racine-Berthiaume, C.; Voyer, N. *J. Am. Chem. Soc.* **2011**, *133*, 6481; (e) D. Hall, C.; J. Kirkovits, G.; C. Hall, A. *Chem. Comm.* **1999**, 1897; (f) Sakai, N.; Gerard, D.; Matile, S. *J. Am. Chem. Soc.* **2001**, *123*, 2517.
- (4) (a) Ren, C. L.; Ng, G. H. B.; Wu, H.; Chan, K.-H.; Shen, J.; Teh, C.; Ying, J. Y.; Zeng, H. Q. *Chem. Mater.* **2016**, *28*, 4001; (b) Ren, C. L.; Chen, F.; Zhou, F.; Shen, J.; Su, H. B.; Zeng, H. Q. *Langmuir* **2016**, *32*, 13510; (c) Ren, C. L.; Shen, J.; Chen, F.; Zeng, H. Q. *Angew. Chem., Int. Ed.* **2017**, *56*, 3847. (d) Zhao, H. Q.; Sheng, S.; Hong, Y. H.; Zeng, H. Q. *J. Am. Chem. Soc.* **2014**, *136*, 14270.
- (5) (a) Bartsch, R. A.; Jeon, E.-G.; Walkowiak, W.; Apostoluk, W. *J. Memb. Sci.* **1999**, *159*, 123; (b) Talanova, G. G.; Elkarim, N. S. A.; Hanes, R. E.; Hwang, H.-S.; Rogers, R. D.; Bartsch, R. A. *Anal. Chem.* **1999**, *71*, 672; (c) Although the values of R_M^+ and ratio **R** (e.g., R_{K^+}/R_{Na^+}) could be used to quickly and reliably gauge the relative ion selectivity across different classes of channel molecules, ratio **R** could only serve as a good approximation of true ion selectivity intrinsic to any channel molecule (See Table S3 and Figure S15 for a detailed discussion involving EC_{50} values). Instead, the best approach to determine the ion selectivity is via the use of single channel current measurement.

Graphic for the Table of Contents

