

CHEMISTRY

A European Journal

A Journal of



Accepted Article

Title: Streamlined synthesis and evaluation of teichoic acid fragments

Authors: Daan van der Es, Francesca Berni, Wouter F. J. Hogendorf, Nico Meeuwenoord, Diana Laverde, Angela van Diepen, Herman S. Overkleeft, Dmitri V. Filippov, Cornelis H. Hokke, Johannes Huebner, Gijsbert A. van der Marel, and Jeroen Codée

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Eur. J.* 10.1002/chem.201800153

Link to VoR: <http://dx.doi.org/10.1002/chem.201800153>

Supported by
ACES

WILEY-VCH

Streamlined synthesis and evaluation of teichoic acid fragments

Daan van der Es,^{at} Francesca Berni,^{at} Wouter F. J. Hogendorf,^a Nico Meeuwenoord,^a Diana Laverde,^b Angela van Diepen,^c Herman S. Overkleeft,^a Dmitri V. Filippov,^a Cornelis H. Hokke,^c Johannes Huebner,^b Gijbert A. van der Marel^a and Jeroen D. C. Codée^{a*}

Abstract: Teichoic acids (TAs) are key components of the Gram-positive bacterial cell wall that are composed of alditol phosphate repeating units, decorated with alanine or carbohydrate appendages. Because of their microheterogeneity, pure well-defined TAs for biological or immunological evaluation cannot be obtained from natural sources. We here present a streamlined automated solid phase synthesis approach for the rapid generation of well-defined glycosylated, glycerol-based TA oligomers. Building on the use of a “universal” linker system and fluororous tag purification strategy, a library of glycerolphosphate pentadecamers, decorated with various carbohydrate appendages, is generated. These are used to create a structurally diverse TA-microarray, which is used to reveal, for the first time, the binding preferences of anti-LTA antibodies at the molecular level.

Teichoic acids (TAs) are a diverse class of biopolymers composed of repeating alditol phosphates and carbohydrates. They are major constituents of the cell wall of most Gram-positive bacteria and can amount up to 70% of the dry weight of the cell wall. Two classes of TAs exist: wall teichoic acids (WTAs) that are covalently attached to the peptidoglycan, and lipoteichoic acids (LTAs) that are anchored in the cell membrane through a glycolipid. TAs are involved in determining cell shape, cell division and nutrient uptake.¹ They are key virulence factors and of importance to biofilm formation and antibiotic resistance.² TAs are targeted by the complement system (lectin ligands)^{3,4} and are recognized by opsonophagocytic antibodies.⁵ The most common structural TA-motif is a polymeric glycerol phosphate backbone decorated with various substituents, including D-alanine esters and carbohydrate appendages. The TAs from the opportunistic pathogens *Enterococcus faecalis*, *Enterococcus faecium* and *Staphylococcus aureus* contain α -kojibiosyl (D-glucose-(α -1,2)-glucosyl)⁶, α -glucosyl⁵ or N-acetyl α -glucosamine residues,⁷ respectively. Positively charged glucosamine groups are also found in certain LTA types.⁸ Insight into how TAs interact with enzymes, receptors and antibodies may guide new antibiotic and vaccine developments and well-defined TA fragments are indispensable tools to fuel these studies.^{9,10} Because of the microheterogeneity of naturally sourced TAs, organic synthesis is the method of choice to access these compounds. We, and others, have previously reported different synthetic strategies to get access to well-defined TA structures,^{11–13} and from our initial solid phase assembly campaign, in which we generated a small set of six

TA-fragments, we identified a glycerol phosphate hexamer bearing a single glucosyl substituent as a lead synthetic antigen.¹⁰ To be able to generate larger TA oligomers with a more diverse substitution pattern, we here report the development of a second generation automated solid phase approach to generate a library of TA oligomers, featuring different carbohydrate appendages at various positions on the TA chain. The method builds on the use of a “universal linker” system and the implementation of a fluororous-tagging strategy.¹⁴ The generated TA fragments have been used to create a TA-microarray to characterize the binding specificity of a monoclonal antibody against *Staphylococcus epidermidis* LTA as well as serum raised against isolated LTA from *E. faecalis* and serum raised against a well-defined TA-BSA conjugate vaccine. These studies reveal, for the first time, the recognition patterns for anti-LTA antibodies at the molecular level, and indicate how Ab-TA binding specificity (or lack thereof) depends on the vaccination strategy used.

Our synthetic strategy, the targeted TA fragments and the used building blocks are depicted in Scheme 1. We aimed at a library, comprising glycerol phosphates with a backbone of 15 monomers in length (Scheme 1A) and containing either an α -glucosyl, an α -glucosaminyl or an α -N-acetyl glucosaminyl at the C-2 of a glycerol moiety at the beginning, middle and/or end of the pentadecamers. TA fragments with a cluster of carbohydrate substituents were generated as well. In all, 16 structures (**1-16**) were pursued as depicted in Scheme 1A. To assemble these TA fragments the required glycerol phosphoramidite building blocks **17-20** (Scheme 1B) were synthesized as described previously.¹¹ In addition we synthesised the fluororous aminospacer phosphoramidite **21**¹⁵ (See SI) to equip all compounds with a purification handle and, after deprotection, a ligation handle. The (commercially available) solid support used for the generation of the TA-library bears a 3-aminopropane-1,2-diol linker that is attached to the resin through a urea linkage (**22**, Scheme 1C).¹⁶ This linker obviates the need for tailor made “pre-loaded” resins, which require the assembly of additional building blocks. The synthesis of non-substituted glycerol phosphate 15-mer **1** commenced with acid-mediated removal of the DMT-group in **22** followed by 5-(benzylthio)-1-H-tetrazole (BTT)-mediated condensation of phosphoramidite **17**. Oxidation of the intermediate phosphite to the phosphotriester and capping of the unreacted alcohols completed the first coupling cycle. Fourteen repetitions of this cycle delivered the resin bound pentadecamer, of which the DMT group was removed by a final acid treatment. After installation of the fluororous linker, treatment with ammonia (2 M in methanol) released the pentadecamer from the resin. Prolonged treatment with aqueous ammonia then removed all remaining cyanoethyl groups from the phosphotriesters. Purification of the crude 15-mer was readily accomplished using conventional reversed phase HPLC. Though we anticipated the need for a fluororous stationary phase to separate the target

^a Leiden Institute of Chemistry, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands

^b Division of Paediatric Infectious Diseases, Dr. von Hauner Children's Hospital, Ludwig-Maximilians-University, Munich, Germany

^c Department of Parasitology, Leiden University Medical Center, Albinusdreef 2, 2333 ZA Leiden, The Netherlands

† D.v.d.E. and F.B. contributed equally to this work.

*Corresponding Author: Dr. J.D.C. Codée: jcodee@chem.leidenuniv.nl


Electronic Supplementary Information (ESI) available: Experimental procedures and analytical data (PDF). See DOI: 10.1039/x0xx00000x

compound from the deletion sequences, the C₈F₁₇-tail proved to be lipophilic enough to provide a similar result with a regular C18 column.

Partially protected pentadecamer **23** was obtained in 17% overall yield, based on resin loading. The synthesis of pentadecamer TAs **24-32**, decorated with a single monosaccharide at three different positions in the glycerol phosphate backbone was undertaken next. After coupling

glycosylated phosphoramidites **18**, **19** or **20** to the resin, fourteen benzyl-substituted glycerol phosphate units were installed and the constructs were capped with fluororous phosphoramidite **21**. The isolated yields for glucose decorated **24** and N-Cbz-glucosamine decorated **25** (17% and 14% respectively) were in line with the yield for undecorated oligomer **23**, the N-acetyl-glucosamine decorated **26** was obtained in 7% overall yield. In parallel, pentadecamers **27-29**, bearing the carbohydrate appendage at the side of the spacer, were synthesized in yields ranging from 8% to 15%. To complete the first series of compounds, the glucose- and glucosamine-decorated pentadecamers, having the substituent in the middle of the glycerol phosphate chain (**30**, **31** and **32**), were assembled (yields ranging from 7-16%). Using the same protocol, fluororous and partially protected GTA-fragments **33-38**, bearing three monosaccharides, clustered or spread over the pentadecamer backbone, were prepared. The installation of the three carbohydrate substituted glycerol phosphate moieties proceeded with comparable efficiency as coupling of the non-substituted glycerol phosphates (as judged from DMT count on the synthesizer) and the TA fragments bearing three carbohydrates were removed by Pd catalyzed hydrogenolysis to provide the pure TA fragments (**1-16**) in 50-90% yield.

Next, with the library of glycosylated TA-pentadecamers in hand we generated TA-microarrays to probe the interaction with anti-LTA antibodies.¹⁷ Besides the TA library, generated here, we printed several TA-fragments that were previously synthesized. These include a series of non-substituted TA-fragments of 6, 10, 14 and 20 glycerol phosphate units in length (**39-42**, Figure 1A),¹⁸ TA-hexamer **43** and kojibiosyl-TA hexamer **44**. We have previously shown that TA-hexamer **43** can be used as a synthetic antigen in a TA-conjugate vaccine, which can offer protection against *E. faecalis* in both active and passive immunization strategies.¹⁰ Notably, while kojibiose has been identified as appendage in *E. faecalis* TA, hexamer **44** proved to be an ineffective synthetic



Scheme 1. A) Target TA fragments. B) Building blocks used. C) Solid phase TA synthesis. Reagents and conditions: a) 3% DCA, toluene; b) **17**, **18**, **19** or **20**, 5-(benzylthio)-1H-tetrazole, acetonitrile; c) I₂, pyridine, H₂O, acetonitrile; d) Ac₂O, N-methylimidazole, 2,6-lutidine, acetonitrile; e) NH₃, MeOH; then NH₄OH, H₂O. **23**: 17%, **24**: 17%, **25**: 14%, **26**: 7%, **27**: 10%, **28**: 15%, **29**: 8%, **30**: 16%, **31**: 16%, **32**: 7%, **33**: 5%, **34**: 6%, **35**: 5%, **36**: 6%, **37**: 8%, **38**: 7%. f) H₂, Pd(0), H₂O. **1**: 66%, **2**: 60%, **3**: 86%, **4**: 41%, **5**: 83%, **6**: 88%, **7**: 36%, **8**: 51%, **9**: 79%, **10**: 39%, **11**: 79%, **12**: 81%, **13**: 84%, **14**: 63%, **15**: 64%, **16**: 84%.




Figure 1. TA-microarrays. A) Previously synthesized TA-fragments. B) Schematic representation of teichoic acid microarray set up and sera evaluation. C) Exemplary fluorescence readout of slides after incubation with serum. D) IgM and IgG levels in the commercially available mouse anti- *S. epidermidis* monoclonal antibody (IBT Bioservices). E) IgM and IgG levels in serum obtained from rabbits immunized with LTA isolated from *E. faecalis* strain 12030.²² F) IgM and IgG levels in rabbit serum raised against a BSA-43 conjugate.¹⁰ Data are represented at three concentrations (30 μM , 10 μM and 3 μM respectively, each spotted in triplicate) and are normalized to the highest intensity.

antigen.

In the array, we also included *E. faecium* WTA fragment **45** that features a di-*N*-acetyl galactosamine-glycerolphosphate backbone,¹⁹ as a negative control. All TA-fragments were printed in varying concentrations (3, 10 and 30 μM) on epoxide functionalized microarray glass slides.²⁰ With the generated microarrays, we probed the binding of a commercially available (IBT Bioservices) mouse anti *S. epidermidis* monoclonal antibody (used at a 1:6000 dilution), serum obtained from rabbits immunized with LTA isolated from *E. faecalis* strain 12030²¹ (used at a 1:500 dilution), and rabbit serum raised against the previously reported BSA-TA **43** conjugate¹⁰ (used at a 1:500 dilution) (Figure 1B). The fluorescence intensities (Figure 1C) were quantified and are graphed in Figures 1D, E and F. Figure 1D depicts the interaction of the monoclonal antibodies against *S. epidermidis* LTA with the immobilized TAs. It becomes immediately apparent that various TA stretches are recognized. There is no selective recognition for either of the carbohydrate appendages and it seems that the monoclonal antibody mainly recognizes the glycerol phosphate backbone. The appendage of multiple carbohydrate substituents (fragments **11-16**) or larger carbohydrates (as in kojibiosyl TA **44**) seems to perturb binding of the antibody. The fact that undecorated or minimally decorated TA stretches are the main interaction partners for the monoclonal antibody, accounts for the fact that this antibody is cross reactive to various Gram-positive bacteria, which share the glycerolphosphate backbone as a common epitope.^{5,9b} We next analyzed the serum, obtained from rabbits immunized with LTA isolated from *E. faecalis* strain 12030. Here, large

differences are observed in the binding of IgM and IgG antibodies. IgM antibodies in the serum bound both the glycosylated and non-glycosylated TAs, again indicating the glycerolphosphate backbone as the prime recognition target. Surprisingly, the IgG antibodies only bound glycosylated fragments. No binding of IgG antibodies to the 'terminally' glycosylated fragments **2-4** and **43** was detected, rather a preference for internally functionalized fragments (as present in TA fragments **8-16**) was observed. This lack of selective recognition of one glycosylated TA over another, can explain the cross-reactivity of serum, raised against *E. faecalis* 12030, with respect to other Gram-positive bacteria such as *S. aureus* and *E. faecium*. Previously, the cross-reactivity of the serum was attributed to recognition of the non-decorated glycerolphosphate backbone by IgG antibodies, which clearly is not observed for the IgG antibodies probed with these arrays.²² Lastly, we investigated the serum raised against a BSA-TA **43** conjugate (Figure 1F). This microarray shows that the serum mainly contains TA-recognizing antibodies of the IgG-type, and only very low levels of IgM antibodies, confirming the effectiveness of the used vaccine modality. Besides conjugate-component TA **43**, also TA **2**, essentially an elongated analogue of TA **43**, and compounds **8** and **11**, harboring the full structure of **43** embedded in or attached to extra glycerol phosphate residues, are recognized, showing the high specificity of the serum for TA stretches with a terminal glucose appendage. The selective recognition observed with this array highlights the power of the analytical potential of the here presented arrays and shows, for the first time, the binding specificity of antibodies at the molecular level. In addition, they indicate that it is possible to

generate antibodies against a well-defined TA epitope, that do not (or only minimally) cross react with other (non-substituted) TA-fragments. Where it could be expected that the glycerolphosphate backbone represents the immunodominant epitope in LTA, it is clear that TA sequence specific antibodies can be raised if the appropriate conjugate is used for vaccination. In summary, an efficient automated solid phase synthesis for the rapid generation of a library of TA oligomers, decorated with a diverse pattern of carbohydrate substituents, is described. The methodology allows for the generation of long glycerol phosphate TA stretches, with carbohydrate appendages positioned at will along on the backbone. The assembled TAs were used to generate TA-microarrays, which have been employed to evaluate the binding preferences of monoclonal antibodies, as well as antibodies in serum, raised against different TA preparations. The arrays have revealed the difference in binding selectivity (or lack thereof) of antibodies, of varying origin. The arrays have shown that very selective antibodies can be generated to recognize specific and well-defined TA epitopes, with minimal binding to the common glycerol phosphate backbone. This in turn implies that it should be possible to generate antibodies that selectively recognize bacterium specific TA stretches for vaccination strategies targeting a specific Gram-positive bacterium. Clearly, the use of TAs isolated from bacterial sources for vaccine purposes can only lead to non-selective serum. The arrays will be valuable tools to probe other biomolecules, such as host immune system lectins or bacteriophage proteins, for binding and they can be used to investigate enzymes involved in the biosynthesis and diversification of TAs.

Acknowledgements

We thank the Netherlands Organization for Scientific Research (NWO, ECHO, grant number 711.011.014) and the EU (Marie-Curie ITN GLYCOVAX, grant number 675671) for financial support.

Keywords Gram-positive bacteria, automated synthesis, microarrays, glycerol phosphate.

Notes and references

- (a) F. C. Neuhaus, J. Baddiley, *Microbiol. Mol. Biol. Rev.* **2003**, *67*, 686–723; (b) S. Brown, J. P. Santa Maria, S. Walker, *Annu. Rev. Microbiol.* **2013**, *67*, 313–336; (c) M. G. Percy, A. Gründling, *Annu. Rev. Microbiol.* **2014**, *68*, 81–100; (d) O. Schneewind, D. Missiakas, *J. Bacteriol.* **2014**, *196*, 1133–1142.
- (a) C. Weidenmaier, A. Peschel, *Nat. Rev. Microbiol.* **2008**, *6*, 276–287; (b) C. Rockel, T. Hartung, *Front. Pharmacol.* **2012**, *3*, 1–19. (c) N. J. Lynch, S. Roscher, T. Hartung, S. Morath, M. Matsushita, D. N. Maennel, M. Kuraya, T. Fujita and W. J. Schwaeble, *J. Immunol.*, **2004**, *172*, 1198–1202.
- K. Kurokawa, K. Takahashi and B. L. Lee, *Immunobiology*, **2016**, *221*, 1091–1101.
- D.-J. Jung, J.-H. An, K. Kurokawa, Y.-C. Jung, M.-J. Kim, Y. Aoyagi, M. Matsushita, S. Takahashi, H.-S. Lee, K. Takahashi and B. L. Lee, *J. Immunol.*, **2012**, *189*, 4951–4959.
- S. Kodali, E. Vinogradov, F. Lin, N. Khoury, L. Hao, V. Pavliak, C. H. Jones, D. Laverde, J. Huebner, K. U. Jansen, A. S. Anderson and R. G. K. Donald. *J. Biol. Chem.*, **2015**, *290*, 19512–19526.
- J. Huebner, Y. Wang, W. A. Krueger, L. C. Madoff, G. Marrisiosian, S. Boisot, D. A. Goldmann, D. L. Kasper, A. O. Tzianabos, G. B. Pier. *Infect Immun.*, **1999**, *67*, 1213–1219.
- A. R. Sanderson, J. L. Strominger, S. G. Nathenson and R. Sanderson, *J. Biol. Chem.*, **1962**, *237*, 3603–3613.
- M. Cot, A. Ray, M. Gilleron, A. Vercellone, G. Larrouy-Maumus, E. Armau, S. Gauthier, G. Tiraby, G. Puzo and J. Nigou, *PLoS One*, **2011**, *6*, e26316.
- (a) C. E. Martin, F. Broecker, S. Eller, M. A. Oberli, C. Anish, C. L. Pereira, P. H. Seeberger, *Chem. Commun.*, **2013**, *49*, 7159–7161; (b) Q. Chen, J. Dintaman, A. Lees, G. Sen, D. Schwartz, M. E. Shirliff, S. Park, J. C. Lee, J. J. Mond, C. M. Snapper, *Infect. Immun.*, **2013**, *81*, 2554–2561; (c) Z. Zhou, W. Ding, C. Li, Z. Wu, *J. Carb. Chem.*, **2017**, *36*, 205–219.
- D. Laverde, D. Wobser, F. Romero-Saavedra, W. Hogendorf, G. van der Marel, M. Berthold, A. Kropec, J. Codee and J. Huebner, *PLoS One*, **2014**, *9*, e110953.
- (a) W. F. J. Hogendorf, A. Kropec, D. V. Filippov, H. S. Overkleeft, J. Huebner, G. A. van der Marel and J. D. C. Codée, *Carbohydr. Res.*, **2012**, *356*, 142–151; (b) W. F. J. Hogendorf, L. N. Lameijer, T. J. M. Beenakker, H. S. Overkleeft, D. V. Filippov, J. D. C. Codée and G. A. Van der Marel, *Org. Lett.*, **2012**, *14*, 848–851.
- (a) W. F. J. Hogendorf, N. Meeuwenoord, H. S. Overkleeft, D. V. Filippov, D. Laverde, A. Kropec, J. Huebner, G. A. Van der Marel and J. D. C. Codée, *Chem. Commun.*, **2011**, *47*, 8961–8963; (b) For the first synthesis of a TA-fragment see: C. A. A. van Boeckel, G. M. Visser, J. P. G. Hermans and J. H. van Boom, *Tetrahedron Lett.*, **1981**, *22*, 4743–4746; (c) For the first automated synthesis of a TA-fragment see: P. Westerduin, G. H. Veeneman, Y. Pennings, G. A. van der Marel, J. H. van Boom, *Tetrahedron Lett.*, **1987**, *28*, 1557–1560; (d) For the first synthesis of a LTA-fragment see: K. Fukase, T. Matsumoto, N. Ito, T. Yoshimura, S. Kotani and S. Kusumoto, *Bull. Chem. Soc. Jpn.*, **1992**, *65*, 2643–2654; (e) For a pivotal example of the synthesis of a carbohydrate-containing TA-fragment see: C. M. Pedersen, I. Figueroa-Perez, B. Lindner, A. J. Ulmer, U. Zähringer and R. R. Schmidt, *Angew. Chemie Int. Ed.*, **2010**, *49*, 2585–2590.
- For a review on all current synthetic methods towards teichoic acids see: D. van der Es, W. F. J. Hogendorf, H. S. Overkleeft, G. A. van der Marel and J. D. C. Codée, *Chem. Soc. Rev.*, **2017**, *46*, 1464–1482.
- W. Zhang, *Chem. Rev.*, **2009**, *109*, 749–795.
- D. V. Filippov, D. J. van Zoelen, S. P. Oldfield, G. A. van der Marel, H. S. Overkleeft, J. W. Drijfhout and J. H. van Boom, *Tetrahedron Lett.*, **2002**, *43*, 7809–7812.
- A. Yagodkin, J. Weisel and A. Azhayev, *Nucleosides, Nucleotides and Nucleic Acids*, **2011**, *30*, 475–489.
- S. Park, J. C. Gildersleeve, O. Blixt and I. Shin, *Chem. Soc. Rev.*, **2013**, *42*, 4310–4326.
- C. Theilacker, A. Kropec, F. Hammer, I. Sava, D. Wobser, T. Sakinc, J. D. C. Codée, W. F. J. Hogendorf, G. A. van der Marel and J. Huebner, *J. Infect. Dis.*, **2012**, *205*, 1076–1085.
- D. van der Es, N. A. Groenina, D. Laverde, H. S. Overkleeft, J. Huebner, G. A. van der Marel and J. D. C. Codée, *Bioorg. Med. Chem.*, **2016**, *24*, 3893–3907.

- 20 For the used protocol see: A. van Diepen, C. H. Smit, L. van Egmond, N. B. Kabatereine, A. Pinot de Moira, D. W. Dunne and C. H. Hokke, *PLoS Negl. Trop. Dis.*, **2012**, *6*, e1922.
- 21 J. Huebner, A. Quaas, W. A. Krueger, D. A. Goldmann and G. B. Pier, *Infect. Immun.*, **2000**, *68*, 4631–4636.
- 22 The non-selective binding of the IgM antibodies may also contribute to the cross reactivity of the serum.

WILEY-VCH


Accepted Manuscript

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Using a streamlined automated solid phase synthesis approach, well-defined glycosylated, glycerol-based teichoic acid oligomers were synthesized. These are used to create a structurally diverse teichoic acid microarray, which is used to reveal - for the first time - the binding preferences of anti-LTA antibodies at the molecular level.



Daan van der Es, Francesca Berni,
Wouter F. J. Hogendorf, Nico
Meeuwenoord, Diana Laverde, Angela
van Diepen, Herman S. Overkleeft,
Dmitri V. Filippov, Cornelis H. Hokke,
Johannes Huebner, Gijsbert A. van der
Marel and Jeroen D. C. Codée* **Page**
No. – Page No.
Streamlined synthesis and evaluation
of teichoic acid fragments