Organic & Biomolecular Chemistry

COMMUNICATION



View Article Online View Journal | View Issue

Cite this: Org. Biomol. Chem., 2014, **12**, 3813 Received 3rd April 2014, Accepted 14th April 2014 DOI: 10.1039/c4ob00705k

www.rsc.org/obc

Practical synthesis of natural plant-growth regulator 2-azahypoxanthine, its derivatives, and biotin-labeled probes[†]

Kazutada Ikeuchi,^a Ryosuke Fujii,^a Shimpei Sugiyama,^a Tomohiro Asakawa,^a Makoto Inai,^a Yoshitaka Hamashima,^a Jae-Hoon Choi,^b Tomohiro Suzuki,^b Hirokazu Kawagishi*^{b,c,d} and Toshiyuki Kan*^a

We describe a practical, large-scale synthesis of the "fairy-ring" plant-growth regulator 2-azahypoxanthine (AHX), and its biologically active hydroxyl metabolite (AOH) and riboside derivative (AHXr). AHXr, a biosynthetic intermediate, was synthesized from inosine *via* a biomimetic route. Biotinylated derivatives of AHX and AHXr were also synthesized as probes for mechanistic studies.

"Fairy rings" arising from fungus-stimulated plant growth occur worldwide, and were first reported in 1675, as reviewed in Nature in 1884.¹ In 2010, we reported that, in the case of the fungus *Lepista sordida*, the "fairy" is a plant growth stimulator, which we identified as 2-azahypoxanthine (AHX: 1).² AHX exhibited growth-regulating activity towards not only turf grass, but also other plants tested, even from different families. Furthermore, this compound increased the seed yields of rice and wheat in pot-growth and field experiments,² suggesting that it might find practical application in agriculture. We also showed that 2-aza-8-oxohypoxanthine (AOH: 3) is a common, biologically active metabolite of AHX (1) in plants (Fig. 1).³

AHX (1) was chemically synthesized from 5-aminoimidazole-4-carboxamide (AICA 2; an intermediate in the purine metabolic pathway in animals, plants, and microorganisms), and converted into AOH (3) by xanthine oxidase-mediated reaction. We hypothesized that plants themselves produce AHX (1) and AOH (3) through a similar pathway, and indeed, we



Fig. 1 Structures of AHX derivatives.

demonstrated that endogenous AHX (1) and AOH (3) are formed *via* the purine pathway in plants.³ We considered that riboside and/or ribotide derivatives of AICA (5, 7) and AHX (4, 6) might also be involved in the biosynthetic pathway. Furthermore, biotinylated derivatives of AHX and AHX riboside (AHXr: 4) are of interest as potential tools for mechanistic studies to identify receptor(s) and related proteins. Herein we report efficient synthetic routes to AHX (2), AOH (3), AHXr (4) and also biotinylated derivatives of AHX and AHXr.

First, we wished to develop a practical synthesis of AHX (1) and AOH (3). Although synthesis of 1 has already been reported,⁴ optimization for large scale-preparation is necessary to obtain sufficient material for field experiments. As shown in Scheme 1, the synthesis was commenced with inexpensive 2·HCl. Upon treatment of 2 with sodium nitrite under acidic conditions, the desired diazonium formation proceeded smoothly to provide diazoimidazole carboxamide (DICA: 8). Although a hundred-gram scale AHX (1) was prepared according to the reported procedure (method A),⁴ we found that the



Scheme 1 Practical synthesis of AHX (1) and AOH (3).

^aSchool of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan. E-mail: kant@u-shizuoka-ken.ac.jp;

Fax: +81-54-264-5745; Tel: +81-54-264-5746

^bResearch Institute of Green Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan. E-mail: achkawa@ipc.shizuoka.ac.jp; Fax: +81-54-238-4885; Tel: +81-54-238-4885

^cGraduate School of Science and Technology, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

^dGraduate School of Agriculture, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

 $[\]dagger$ Electronic supplementary information (ESI) available: Experimental procedures and NMR spectral data and bioassay of 4, 6, 13 and 24. See DOI: 10.1039/ c4ob00705k

Communication

triazine ring of **1** can be constructed by heating **8** in methanol at 60 °C (method B). Furthermore method B was more superior to method A in terms of yield as well as handling. By means of this transformation, together with activated carbon treatment, **1** was obtained on an almost decagram scale. Conversion of **1** to **3** was carried out by enzymatic oxidation with xanthine oxidase.⁵ Purification of **3** was also accomplished by recrystallization, and chromatographic purification was unnecessary (see the ESI† for details).

With the desired biologically active natural products 1 and 3 in hand, we turned our attention to the synthesis of derivatives that would be suitable as probes for mechanistic studies, following on from our previous work.⁶ During studies of polyphenols such as catechins⁷ and flavones,⁸ we had found that a terminal amine or azide group is favorable for facile incorporation of a probe moiety without the need for protection of the phenol group.^{7b,8a} Thus, we decided to incorporate a terminal azide group onto AHX (1). However, direct alkylation of 1 with alkyl halide did not proceed smoothly. In order to enhance the reactivity of the linker, we designed the probe precursor 10, in which the linker group is connected through the acetal group. Alkylation reaction was performed after in situ protection of the nitrogen of imidazole and amide with a TMS group by treatment with N,O-bis(trimethylsilyl)acetamide (BSA), as shown in Scheme 2. Upon treatment of 1 with 1-azido-5-(chloromethyloxy)pentane 9 in the presence of DBU and BSA, the desired reaction proceeded smoothly to provide 10 and 11 as a 1.8:1 mixture. After separation of each regioisomer, the azide 10 was used as the AHX probe precursor. The Hüisgen condensation reaction⁹ of azide 10 and acetylene 12 containing a biotin group in the presence of CuSO₄ and sodium ascorbate proceeded smoothly to afford 13.10 Considering the convenience of the Hüisgen reaction, this synthetic strategy should be applicable to incorporation of a wide range of functional units, such as a fluorescent moiety for imaging or a carrier protein for synthesis of an immunogen.^{7c} We found that the coupling reaction did not require a tedious purification step, although of biotin-containing probes is sometimes purification troublesome.

Based on the proposed biosynthetic pathway of AHXr (4) and AHX-ribotide (AHXR: 6), we next planned to synthesize 4 and 6 from inexpensive inosine (14) as shown in Scheme 3.¹¹ Although conversion of the carbon atom at the C-2 position of



Scheme 2 Hüisgen reaction of azide 10 and alkyne 12.



Scheme 3 Preparation of AHXr (4) and AHXR (6).

14 into the nitrogen atom of 4 seemed to be a challenging task, a similar aminolysis of the pyrimidine ring of inosine (14) has been reported.¹² After protection of the hydroxyl groups of 14 with a TBS group, incorporation of a 2,4-dinitrophenyl group on the amide nitrogen was carried out via nucleophilic aromatic substitution reaction of 1-chloro-2,4dinitrobenzene to give 15. Upon treatment of 15 with ethylenediamine in THF, the desired aminolysis reaction on the imidamide ring proceeded smoothly to give 17 with the release of dihydro-imidazole 16.12 After selective deprotection of the TBS ether on the primary alcohol of 17 by treatment with NH₄F, construction of the triazine ring of 19 was performed by treatment of 18 with sodium nitrite and hydrochloric acid, as employed in the preparation of 1. Removal of the 2,4-dinitrophenyl protecting group of 19 was carried out by treatment with ethylenediamine as a nucleophile. In the original report,¹² treatment of 15 with ethylenediamine in DMF caused simultaneous ring opening reaction and dinitrobenzene cleavage reaction to afford the -CONH₂ derivative. However, the 2,4-dinitrophenyl group played a key role in the efficient preparation of 4, 6 and the probe precursor 23, and a selective aminolysis reaction of 15 was needed. Chemoselective reaction of 15 was accomplished by changing the solvent to THF from DMF. Finally, TBAF-mediated deprotection of the TBS groups of 20 provided AHXr (4).¹⁰ On the other hand, AHXR (6) was synthesized by incorporation of phosphate ester into 20 by the phosphoramidite method,13 followed by the removal of the TBS groups and the benzyl group to give 6.¹⁰

With an efficient synthetic method for AHXr (4) in hand, we turned our attention to the synthesis of a biotin-labeled AHXr probe compound, starting from the synthetic intermediate **19**. Since AHXr (4) is a biosynthetic precursor of AHX (2) in plants, the biotinylated AHXr probe should enable the identification of proteins involved in the biosynthesis of AHX. Since the



Scheme 4 Synthesis of biotin-labelled AHXr probe 24.

ribose unit of the AHXr probe (24) can serve as a hydrophilic linker unit of the AHX probe, an efficient and flexible synthesis of 24 would be useful. As shown in Scheme 4, the alkylation reaction of 19 with 9 proceeded smoothly to provide 21. Utilizing a similar method to that used for preparation of 4, removal of the 2,4-dinitrophenyl group and the TBS group of 21 provided a reactive probe precursor 23. Next, coupling with biotin was examined. Upon treatment of azide 23 and acetylene 12 under the conditions employed for preparation of 13, the desired Hüisgen reaction proceeded smoothly to provide 24¹⁰ without any need for purification. This coupling reaction is compatible with the reactive hydroxyl group and the amide group. Synthesis of other kinds of probe molecules from 10 and 23 is in progress in our laboratories, and the details will be reported in due course.

Conclusion

We have developed a practical, large-scale synthesis of the plant-growth regulator AHX (1), as well as an efficient synthetic route for AOH (3), AHXr (4) and AHXR (6), respectively. Biotinlabeled derivatives of AHX (13) and AHXr (24) were also synthesized. The probe precursors 10 and 23 should also be suitable for preparation of other probe molecules.

This work was partially supported by the Programme for Promotion of Basic and Applied Researches for Innovations in Bio-oriented Industry, and by a Grant-in-Aid for Scientific Research on Innovative Areas of "Chemical Biology of Natural Products" from MEXT.

Notes and references

- 1 H. Evershed, Nature, 1884, 29, 384.
- 2 J.-H. Choi, K. Fushimi, N. Abe, H. Tanaka, S. Maeda, A. Morita, M. Hara, R. Motohashi, J. Matsunaga, Y. Eguchi,

N. Ishigaki, D. Hashizume, H. Koshino and H. Kawagishi, *ChemBioChem*, 2010, **11**, 1373.

- 3 (a) J.-H. Choi, T. Ohnishi, Y. Yamakawa, S. Takeda, S. Sekiguchi, W. Maruyama, K. Yamashita, T. Suzuki, A. Morita, T. Ikka, R. Motohashi, Y. Kiriiwa, H. Tobina, T. Asai, S. Tokuyama, H. Hirai, N. Yasuda, K. Noguchi, T. Asakawa, S. Sugiyama, T. Kan and H. Kawagishi, *Angew. Chem., Int. Ed.*, 2014, 53, 1552; (b) Our report was covered in Nature and C&E News; A. Mitchinson, *Nature*, 2014, 505, 298.A. Mitchinson, *C&E News*, 2014, 92, 25.
- 4 Y. Wang and M. F. G. Stevens, *Bioorg. Med. Chem. Lett.*, 1996, 6, 185.
- 5 Xanthine oxidase from buttermilk (XOD; $0.28 \text{ U} \text{ mg}^{-1}$ powder) was purchased from Oriental Yeast Co. (Tokyo, Japan).
- 6 (a) S. Sasaki, H. Suzuki, H. Ouchi, T. Asakawa, M. Inai, R. Sakai, K. Shimamoto, Y. Hamashima and T. Kan, Org. Lett., 2014, 16, 564; (b) H. Fuwa, Y. Takahashi, Y. Konno, N. Watanabe, H. Miyashita, M. Sasaki, H. Natsugari, T. Kan, T. Fukuyama, T. Tomita and T. Iwatsubo, ACS Chem. Biol., 2007, 2, 408; (c) T. Kan, Y. Kita, Y. Morohashi, Y. Tominari, S. Hosoda, T. Tomita, H. Natsugari, T. Iwatsubo and T. Fukuyama, Org. Lett., 2007, 9, 2055.
- 7 (a) K. Shimizu, T. Asakawa, N. Harada, D. Fukumoto, H. Tsukada, T. Asai, S. Yamada, T. Kan and N. Oku, *PLoS One*, 2014, 9, e85520; (b) A. Yoshida, Y. Hirooka, Y. Sugata, M. Nitta, T. Manabe, S. Ido, K. Murakami, R. K. Saha, T. Suzuki, M. Ohshima, A. Yoshida, K. Itoh, K. Shimizu, N. Oku, T. Furuta, T. Asakawa, T. Wakimoto and T. Kan, *Chem. Commun.*, 2011, 47, 1794; (c) Y. Hirooka, N. Nitta, T. Furuta and T. Kan, *Synlett*, 2008, 3234.
- 8 (a) A. Hiza, Y. Tsukaguchi, T. Ogawa, M. Inai, T. Asakawa, Y. Hamashima and T. Kan, *Heterocycles*, 2014, 88, 1371;
 (b) T. Asakawa, A. Hiza, M. Nakayama, M. Inai, D. Oyama, H. Koide, K. Shimizu, T. Wakimoto, N. Harada, H. Tsukada, N. Oku and T. Kan, *Chem. Commun.*, 2011, 47, 2868; (c) T. Furuta, M. Nakayama, H. Suzuki, H. Tajimi, M. Inai, H. Nukaya, T. Wakimoto and T. Kan, *Org. Lett.*, 2009, 11, 2233.
- 9 For a recent review of Hüisgen reaction, see: (a) S. Dedola, S. A. Nepogodiev and R. A. Field, *Org. Biomol. Chem.*, 2007, 5, 1006; (b) P. Wu and V. V. Fokin, *Aldrichimica Acta*, 2007, 40, 7; (c) H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem.*, *Int. Ed.*, 2001, 40, 2004.
- 10 Interestingly, biotin probes 13 and 24, as well as AHXr (4) and AHXR (6), possessed comparable biological activity to natural AHX (1). For details, see the ESI.[†]
- 11 Direct condensation reaction of AHX (1) and tetraacetyl ribose was unsatisfactory; for details, see the ESI.†
- 12 L. D. Napoli, A. Messere, D. Montesarchio and G. Piccialli, J. Chem. Soc., Perkin Trans. 1, 1997, 2079.
- 13 For a review of the phosphoramidite method, see: R. P. Iyer, *Tetrahedron*, 1992, **48**, 2223.