

Article

Subscriber access provided by TULANE UNIVERSITY

Palladium-catalyzed Aerobic anti-Markovnikov Oxidation of Aliphatic Alkenes to Terminal Acetals

Saki Komori, Yoshiko Yamaguchi, Yasutaka Kataoka, and Yasuyuki Ura

J. Org. Chem., Just Accepted Manuscript • DOI: 10.1021/acs.joc.8b02919 • Publication Date (Web): 21 Jan 2019

Downloaded from http://pubs.acs.org on January 21, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

7

8 9 10

11 12

13

14

15

16

17

18

19 20 21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

Palladium-catalyzed Aerobic anti-Markovnikov Oxidation of Aliphatic Alkenes to Terminal Acetals

Saki Komori, Yoshiko Yamaguchi, Yasutaka Kataoka, and Yasuyuki Ura*

Department of Chemistry, Biology, and Environmental Science, Faculty of Science, Nara Women's University, Kitauoyanishi-machi, Nara 630-8506, Japan

ABSTRACT: Terminal acetals were selectively synthesized from various unbiased aliphatic terminal alkenes and 1,2-, 1,3-, or 1,4diols using a $PdCl_2(MeCN)_2/CuCl$ catalyst system in the presence of *p*-toluquinone under 1 atm of O₂ and mild reaction conditions. The slow addition of terminal alkenes suppressed the isomerization to internal alkenes successfully. Electron-deficient cyclic alkenes, such as *p*-toluquinone, were key additives to enhance the catalytic activity and the anti-Markovnikov selectivity. The halogen groups in the alkenes were found to operate as directing groups, suppressing isomerization and increasing the selectivity efficiently.

INTRODUCTION

Realization of anti-Markovnikov selectivity in the oxidation of terminal alkenes is challenging.¹⁻³ Although ketones are typically obtained in palladium-catalyzed Wacker-type oxidation (Scheme 1a), aldehydes can be formed preferentially from terminal alkenes with oxygen or nitrogen directing groups^{1,4-8} and vinylarenes.^{1-3,9-12} In the case of vinylarenes, iron-^{13,14} and ruthenium^{15,16}-catalyzed reactions affording aldehydes have been reported; these reactions proceed via a tandem epoxidationisomerization pathway. On the other hand, unbiased aliphatic terminal alkenes (other than ethylene) are known to be difficult to convert into aldehydes selectively.¹⁷⁻¹⁹ Grubbs et al. reported anti-Markovnikov Wacker-type oxidation of the unbiased aliphatic alkenes using nitrite salts as co-catalysts under aerobic conditions.²⁰ Che et al. achieved iron porphyrin-catalyzed anti-Markovnikov oxidation in the presence of iodosylbenzene as an oxidant.14

The situation is similar in the formation of acetals, which are protected compounds of aldehydes or ketones, from terminal alkenes. Alkenes with electron-deficient groups, 21-25 alkenes with oxygen, nitrogen, or sulfur directing groups,²⁶⁻²⁸ 1,5-dienes,²⁸ and vinylarenes^{21,28-32} are known to afford terminal acetals preferentially by reactions with alcohols or diols, depending on the reaction conditions; however, unbiased aliphatic alkenes normally afford Markovnikov products (internal acetals³³⁻³⁶ or ketones^{19,21,30,34,37}) and remain challenging substrates for anti-Markovnikov oxidation (Scheme 1a). Here we report the palladiumcatalyzed aerobic anti-Markovnikov oxidation of aliphatic alkenes to terminal acetals under mild reaction conditions (Scheme 1b). The key features of this reaction include: 1) the slow addition of alkenes, which can suppress the isomerization of terminal alkenes into internal alkenes, 2) addition of electrondeficient cyclic alkenes as auxiliary ligands, which enhance the catalytic activity and anti-Markovnikov selectivity,^{9,29} and 3) halogen groups in alkenes, which act as directing groups despite their weakly coordinating ability, affording high anti-Markovnikov selectivity.

RESULTS AND DISCUSSION

1-Octene (1a) and pinacol (2a, 3.0 equiv) were treated with

Scheme 1. Markovnikov and anti-Markovnikov Oxidation of Aliphatic Terminal Alkenes

a) conventional oxidation of aliphatic alkenes to aldehydes, ketones, or acetals



b) anti-Markovnikov oxidation of aliphatic alkenes to terminal acetals (this work)



catalytic amounts of PdCl₂(MeCN)₂ (10 mol%) and CuCl (20 mol%) in t-BuOH at 40 °C under 1 atm of O₂ (Table 1). Under these conditions, terminal acetal 3aa was formed in only 8% yield along with the formation of 2- and 3-octanones (entry 1). Isomerization of **1a** to 2-octenes was also observed during the reaction (39%). The formation of 3-octanone would occur by the Wacker-type oxidation of 2-octenes. Because the isomerization would be catalyzed by in situ formed Pd-H species, we expected that, by maintaining the concentration of **1a** at a low level, the isomerization may be suppressed efficiently. When slow addition of **1a** to the reaction mixture by a syringe pump (7 h) was performed, the formation of 2-octenes was suppressed significantly (19%), and the yield of 3aa and anti-Markovnikov selectivity were both increased (entry 2). The effect of electrondeficient cyclic alkenes as additives was then examined (entries 3-15). We previously found that these alkenes can operate as ligands to enhance the catalytic activity of the

Table 1. Optimization of Reaction Conditions^a

Table 2. Scope of Aliphatic Alkenes^a

n-Hex + HO + H									
1a	2a	atm)) 3aa	<i>n</i> -∺ex 4a					
Entry	Additive	Yield of 3aa (%) ^b	Yield of 4a (%) ^b	Selectiv- ity (%) ^c					
1 ^{<i>d</i>}	none	8	25 ^e	24					
2	none	34	28	55					
3	maleic anhydride	50	17	75					
4	maleimide	44	30	59					
5	N-methylmaleimide	49	33	60					
6	N-phenylmaleimide	49	18	73					
7	BQ ^f	21	70	23					
8	CIBQ	37	53	41					
9	2,6-Cl₂BQ	25	60	29					
10	Cl ₄ BQ	38	33	54					
11	MeOBQ	32	32 16						
12	PhBQ	51	51 35						
13	MeBQ	53	34	61					
14	$MeBQ^{g}$	34	15	69					
15 ^{<i>h</i>}	MeBQ	25	53	32					
16 ^{<i>i</i>}	MeBQ	47	13	78					
17 ^{i,j}	MeBQ	5	13	28					

^a Reaction conditions: 1a (0.50 mmol), 2a (1.50 mmol), PdCl₂(MeCN)₂ (0.050 mmol), CuCl (0.10 mmol), additive (0.50 mmol), *t*-BuOH (2.0 mL), 40 °C, O₂ (1 atm). 1a was added over 7 h by a syringe pump, and the reaction mixture was stirred for an additional 1 h (8 h in total). ^b Determined by ¹H NMR. ^c Selectivity = 3aa/(3aa+4a). ^d 1a was added at a time (without a syringe pump). ^e Total yield of 4a (11%) and 3-octanone (14%).
^f BQ = *p*-benzoquinone. ^g 10 mol% of MeBQ (0.050 mmol) was used. ^h *t*-AmylOH was used as a solvent. ⁱ Under Ar. ^j Without CuCl.

anti-Markovnikov oxidation of vinylarenes to aldehydes9 and terminal acetals²⁹ under aerobic conditions. Maleic anhydride and maleimides were found to increase both the yield of 3aa and the selectivity (entries 3-6). p-Benzoquinone (BQ), which is often used as an oxidant in Pd-catalyzed reactions, gave low selectivity for the product (entry 7). Halogen substituted BQs and MeOBO were also inefficient (entries 8-11). On the other hand, PhBQ and MeBQ gave higher yields of 3aa along with moderate selectivity (entries 12 and 13). Reducing the loading of MeBQ to 10 mol% was ineffective (entry 14). The use of tamylOH instead of t-BuOH as a solvent lowered the yield of 3aa and the selectivity (entry 15). Although the reaction proceeded even under an argon atmosphere (entry 16), the absence of both O₂ and CuCl resulted in low yield of **3aa** (entry 17). Thus, MeBQ also functioned as an oxidant, especially in the presence of CuCl. Indeed, in entries 13 and 16, 30% and 58% of MeBQ were converted into methyl-p-hydroquinone (MeHQ), respectively, according to ¹H NMR analysis.

The optimized reaction conditions using MeBQ as an additive were then applied to various aliphatic terminal alkenes (Table 2). By increasing the steric hindrance at the substituents, the anti-Markovnikov selectivity increased (entries 2 and 3). 4-

~		PdCl ₂ (MeCN) ₂ (10 mol%) CuCl (20 mol%) MeBQ (1.0 equiv)		6) R			
1	* HO' X 2a	<i>t-</i> BuOH, 40 ^c O ₂ (1 atr	°C, 8 h n)	3	*	R 4	
Entry	Proc	duct		Yield of 3 (%) ^{b,c}	Yield of 4 (%) ^b	Selectivity (%) ^d	
1	~~~~	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3aa	53(37)	32	62	
2	\downarrow	J°×	3ba	55(47)	18	75	
3 ^e	X	°∕∕	3ca	73(46)	12	86	
4	\bigcirc	°××	3da	74(61)	25	75	
5 ^f	ci	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3ea	77(60)	8	91	
6 ^{<i>g</i>}	Br	↓°×	3fa	84(76)	3	97	
7 ^f	Br	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3ga	86(75)	7	92	
8	Br	°××	3ha	77(63)	21	79	
9 ^g	ı~~~	J°×	3ia	40(33)	4	91	
10 ^f	BnO	°×	3ja	60(49)	21	74	
11 ^e	BnO	°× °×	3ka	56(52)	27	67	
12 ^{e,h}	но	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3la	66(60)	18	79	
13 ^{e,h}	BnO	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	3ma	68(51)	12	85	

^{*a*} Reaction conditions: **1** (2.0 mmol), **2a** (6.0 mmol), PdCl₂(MeCN)₂ (0.20 mmol), CuCl (0.40 mmol), MeBQ (2.0 mmol), *t*-BuOH (8.0 mL), 40 °C, O₂ (1 atm). **1** was added over 7 h by a syringe pump, and the reaction mixture was stirred for an additional 1 h (8 h in total). ^{*b*} Determined by ¹H NMR. ^{*c*} The values in parentheses are isolated yields. ^{*d*} Selectivity = **3**/(**3**+**4**). ^{*e*} For 12 h (slow addition 7 h + additional 5 h). ^{*f*} No slow addition, for 3 h. ^{*s*} No slow addition, for 1 h. ^{*h*} Maleic anhydride was used instead of MeBO.

Phenyl-1-butene also gave a good product yield with good selectivity (entry 4). We found that linear alkenes with a chloro or

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

bromo group at the terminal carbon afforded high yields of terminal acetals with high selectivity (entries 5-8). The carbonhalogen bonds remained intact in these products. 4-Iodo-1-butene also gave the terminal acetal in high selectivity, albeit the yield was low (entry 9). The low conversion of 4-iodo-1-butene (47%) indicates that a side reaction such as carbon-iodine bond cleavage occurred, deactivating the catalyst. In entries 5-7 and 9, the slow addition of alkenes was not necessary because the isomerization of alkenes did not compete. As a control experiment, 1-bromobutane (1.0 equiv) was used as another additive for the reaction of 1-octene with pinacol to examine whether the bromo group in haloalkane is also effective. However, almost no improvement in the product yields (3aa: 53%, 4a: 29%) and the selectivity (65%) was observed compared to those shown in entry 1, Table 2. These results indicate that the halogen substituents in the alkenes act as directing groups, even though they are regarded as weakly coordinating ligands in general. We also attempted terminal acetal formation from 4-bromo-1-butene and pinacol (10 equiv) using a PdCl₂(MeCN)₂ (10 mol%)/BQ (2.0 equiv, as an oxidant in this case)/DMF system (60 °C, 6 h), which has been used for vinylarenes, allyl ethers, and 1,5dienes²⁸; however, only 8% of **3fa** and 2% of **4f** were formed (89% conversion). Thus, because there are only weakly coordinating compounds in the present system (t-BuOH and pinacol would not be so favorable as ligands due to steric repulsion against other ligands on palladium), even halogen substituents, which are small and have weak coordination ability, may be able to coordinate to palladium. Related to the above results for haloalkenes, allylic fluorides are known to afford β-fluorinated aldehydes in a PdCl₂(PhCN)₂/CuCl₂/AgNO₂ catalyst system, although mechanistic investigations suggest that the anti-Markovnikov selectivity is attributed to the inductive effect of

Table 3. Scope of Diols^a



^{*a*} Reaction conditions: **1f** (2.0 mmol), **2** (6.0 mmol), PdCl₂(MeCN)₂ (0.20 mmol), CuCl (0.40 mmol), MeBQ (2.0 mmol), *t*-BuOH (8.0 mL), 40 °C, O₂ (1 atm), 8 h. ^{*b*} Determined by ¹H NMR. ^{*c*} The values in parentheses are isolated yields. ^{*d*} Selectivity = **3**/(**3**+**5**+**4f**). ^{*e*} Isolated as a mixture of **3fb** and **5fb**. ^{*f*} The value was calculated from the NMR yield of **3fb** and the ratio of **3fb** to **5fb** in the isolated mixture. ^{*g*} For 28 h. ^{*h*} For 3 h.

fluorine substituents, unlike the present reaction.³⁸ Other alkenes with oxygen functional groups such as ethers (entries 10 and 11), carboxy groups (entry 12), and ester groups (entry 13) also afforded moderate yields of the corresponding terminal acetals with good to high selectivity.

By using 4-bromo-1-butene as a terminal alkene, the scope of diols was then examined (Table 3). In the case of the primary diol 2,2-dimethyl-1,3-propanediol, internal acetal **5fb** was also formed as a by-product in addition to terminal acetal **3fb** and **4f**, resulting in decreased selectivity (entry 1). The relatively less bulky primary diol can also operate as a nucleophile, resulting in internal nucleophilic attack on the coordinated alkene (*vide infra*). On the other hand, the tertiary diols afforded high product yields with high selectivity (entries 2 and 3).

The reaction of 4-bromo-1-butene with pinacol was monitored by ¹H NMR spectroscopy (Figure 1). CDCl₃ was used as a co-solvent (*t*-BuOH/CDCl₃ = 1:1). We found that 3-bromobutanal, **6f**, an anti-Markovnikov Wacker-type oxidation product, was formed as the main oxidized compound in the early stages, and **6f** appeared to be gradually converted into **3fa** by reaction with pinacol catalyzed by an in situ generated acid. Because no induction period was observed in the formation of **3fa**, an alternative direct conversion from **1f** and pinacol to **3fa** (which does not go through **6f**) seems to be a competing reaction.

Based on these experimental results, a proposed mechanism for the formation of terminal acetals and ketones is shown in Scheme 2. To the LPdCl₂ species, terminal alkene 1 and MeBQ coordinate to afford 7. Then, t-BuOH attacks the coordinated alkene preferentially at the terminal carbon to afford alkyl palladate 8. Although electronic factors favor internal nucleophilic attack, the steric repulsion between R and t-Bu groups facilitates terminal nucleophilic attack. Subsequent elimination of HCl and β-H elimination result in the formation of alkenyl *t*butyl ether 9, which is converted into terminal acetal 3 via aldehyde 6 or directly, catalyzed by an acid. If *t*-BuOH attacks the internal carbon of the coordinated alkene in 7, ketone 4 is formed via similar steps to those for the formation of 6. The formed Pd-H species isomerizes the terminal alkene 1 into internal alkenes. The reduction of Pd(II) to Pd(0) with release of HCl and the reoxidation of the Pd(0) species by CuCl₂ reproduces the PdCl₂ species, and the reduced CuCl is reoxidized to CuCl₂ by O₂ and HCl. The Pd(0) species can be also reoxidized to Pd(II) by MeBQ and HCl, especially in the presence of CuCl, as observed in entry 16, Table 1. In the case of 4-bromo-1-butene, chelate coordination occurs and, thus, terminal nucleophilic attack is more favorable than internal nucleophilic attack because of the formation of a stable five-membered ring after the attack. The suppression of isomerization from the terminal alkene into internal alkenes by Pd-H species can be also rationalized by chelate coordination (Scheme 3). Other haloalkenes would also have similar chelate coordination effects. We propose that the electron-deficient cyclic alkenes such as MeBQ coordinate to Pd to facilitate the nucleophilic attack and the reduction of Pd(II) to Pd(0), as well as the stabilization of Pd(0)species, suppressing its deactivation to Pd black.^{9,29} As another possibility, the catalytically active species may be Pd-Cu heterobimetallic complexes^{17,39-42} and the electron-deficient cyclic alkenes may control the reactivity of the bimetallic complexes through the coordination to Cu.



Figure 1. Time course of the conversion of 1f into 3fa, 6f, and 4f.

Scheme 2. Proposed Mechanism

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60



Scheme 3. Suppression of the Isomerization by Chelate Coordination



CONCLUSION

We have developed a Pd-catalyzed aerobic anti-Markovnikov oxidation of aliphatic alkenes to terminal acetals. The use of a bulky nucleophile, *t*-BuOH, resulted in the anti-Markovnikov selectivity. Slow addition of terminal alkenes efficiently suppressed the isomerization into internal alkenes. The addition of electron-deficient cyclic alkenes such as MeBQ enhanced the catalytic activity and the selectivity. Halogen groups at appropriate positions in the alkenes were found to operate as directing groups to afford the corresponding terminal acetals in high yield and selectivity. We believe that these findings would be helpful for the further development of anti-Markovnikov Wacker-type oxidation and related reactions of aliphatic terminal alkenes.

EXPERIMENTAL SECTION

General Information. Unless otherwise indicated, all reactions were performed under an oxygen atmosphere (1 atm). PdCl₂(MeCN)₂,⁴³ 4-iodo-1-butene,⁴⁴ 5-hexenyl benzyl ether,⁴⁵ 9-decenyl benzyl ether⁴⁵ and 5-hexenoic acid benzyl ester⁴⁶ were prepared as described in the literature. t-BuOH was purchased from Nacalai tesque Co. Ltd. and was purified by distillation. Other chemicals were also commercially available and were used without further purification. Flash column chromatography was performed using silica gel SILICYCLE SiliaFlash F60 (40-63 µm, 230-400 mesh). NMR spectra were recorded on a Bruker AV-300N (300 MHz (1H), 75 MHz (13C)) spectrometer or a JEOL JNM AL-400 (400 MHz (1H), 100 MHz (^{13}C)) spectrometer. Chemical shift values (σ) were expressed relative to SiMe₄. Infrared spectra were measured on a JASCO FT/IR-6100 spectrophotometer. High-resolution mass spectra were recorded on a JEOL JMS-T100LC spectrometer (ESI-TOF MS) with positive ionization mode.

Synthesis of Terminal Acetals 3.

General Procedure for the Synthesis of 3. Method A: To a reaction vessel, CuCl (39.6 mg, 0.40 mmol), PdCl₂(MeCN)₂ (51.9 mg, 0.20 mmol), MeBQ (244 mg, 2.0 mmol), and pinacol (709 mg, 6.0 mmol) were added, and O₂ was purged. To the mixture, t-BuOH (8 mL) was added and the reaction mixture was stirred at 40 °C. Immediately, a terminal alkene (2.0 mmol) was added slowly over 7 h by a syringe pump, and the reaction mixture was stirred for an additional 1-5 h (8-12 h in total). The solvent and volatile materials were evaporated under vacuum. To the residue, dichloromethane (10 mL) and silica gel (2.5 g) were added, and the solvent was evaporated to adsorb the residue on the silica gel. Purification by silica gel column chromatography (hexane/ethyl acetate = 20/1 with 1% triethylamine) afforded the product as a pale yellow oil. Method B: This method is similar to Method A. After the reaction completed, the solvent and volatile materials were evaporated under vacuum. To the residue, a saturated aqueous NaHCO3 solution and diethyl ether were added to extract. The aqueous layer was further extracted with diethyl ether (twice). The combined organic layer was washed with water and brine, and was dried over anhydrous sodium sulfate. After filtration, the solvent was evaporated under vacuum. To the residue, dichloromethane (10 mL) and silica gel (2.5 g) were added, and the solvent was evaporated to adsorb the residue on the silica gel. Purification by silica gel column chromatography (hexane/ethyl acetate = 20/1with 1% triethylamine) afforded the product as a pale yellow oil. Method C: To a reaction vessel, CuCl (39.6 mg, 0.40 mmol), PdCl₂(MeCN)₂ (51.9 mg, 0.20 mmol), MeBQ (244 mg, 2.0 mmol), and pinacol (709 mg, 6.0 mmol) were added, and O₂ was purged. To the mixture, t-BuOH (8 mL) and a terminal alkene (2.0 mmol) were added and the reaction mixture was stirred at 40 °C for 1–28 h. The solvent and volatile materials were evaporated under vacuum. To the residue, dichloromethane (10 mL) and silica gel (2.5 g) were added, and the sol-

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

vent was evaporated to adsorb the residue on the silica gel. Purification by silica gel column chromatography (hexane/ethyl acetate = 20/1 with 1% triethylamine) afforded the product as a pale yellow oil. Method D: This method is similar to Method C. After the reaction completed, the solvent and volatile materials were evaporated under vacuum. To the residue, a saturated aqueous NaHCO₃ solution and diethyl ether were added to extract. The aqueous layer was further extracted with diethyl ether (twice). The combined organic layer was washed with water and brine, and was dried over anhydrous sodium sulfate. After filtration, the solvent was evaporated under vacuum. To the residue, dichloromethane (10 mL) and silica gel (2.5 g) were added, and the solvent was evaporated to adsorb the residue on the silica gel. Purification by silica gel column chromatography (hexane/ethyl acetate = 20/1 with 1% triethylamine) afforded the product as a pale yellow oil.

2-Heptyl-4,4,5,5-tetramethyl-1,3-dioxolane (3aa). Method A was applied (0.17 g, 0.74 mmol, 37% yield). ¹H NMR (300 MHz, CDCl₃) δ 5.02 (t, J = 5.4 Hz, 1H), 1.61–1.54 (m, 2H), 1.40–1.27 (m, 10H), 1.19 (s, 12H), 0.87 (t, J = 6.9 Hz, 3H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 101.0, 81.5, 36.5, 31.8, 29.6, 29.2, 24.5, 24.2, 22.6, 22.1, 14.1. HRMS (ESI): m/z calcd for C₁₄H₂₇O₂ [M-H]⁺ 227.2011, found 227.2019.

2-(3-Methylbutyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(**3ba**). Method A was applied (0.19 g, 0.95 mmol, 47% yield). ¹H NMR (300 MHz, CDCl₃) δ 5.01 (t, J = 5.1 Hz, 1H), 1.61– 1.51 (m, 3H), 1.31–1.23 (m, 2H), 1.19 (s, 12H), 0.88 (d, J = 6.6Hz, 6H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 101.2, 81.5, 34.3, 33.4, 28.0, 24.2, 22.5, 22.1. HRMS (ESI): m/z calcd for C₁₂H₂₃O₂ [M-H]⁺ 199.1698, found 199.1691.

2-(3,3-Dimethylbutyl)-4,4,5,5-tetramethyl-1,3-dioxolane (**3ca**). Method A was applied (196 mg, 0.91 mmol, 46% yield). ¹H NMR (300 MHz, CDCl₃) δ 4.99 (t, J = 5.1 Hz, 1H), 1.60– 1.52 (m, 2H), 1.31–1.20 (m, 2H), 1.19 (s, 12H), 0.87 (s, 9H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 101.7, 81.6, 38.3, 31.7, 30.0, 29.2, 24.2, 22.1. HRMS (ESI): m/z calcd for C₁₃H₂₅O₂ [M-H]⁺ 213.1855, found 213.1860.

2-(3-Phenylpropyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(3da). Method A was applied (0.30 g, 1.21 mmol, 61% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.23–7.11 (m, 5H), 5.02 (t, J = 5.1 Hz, 1H), 2.62 (t, J = 7.5 Hz, 2H), 1.76–1.66 (m, 2H), 1.63– 1.57 (m, 2H), 1.16 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 142.2, 128.4, 128.2, 125.6, 100.7, 81.6, 35.8, 35.7, 26.1, 24.2, 22.1. HRMS (ESI): m/z calcd for C₁₆H₂₃O₂ [M-H]⁺ 247.1698, found 247.1707.

2-(5-Chloropentyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(3ea). Method C was applied (284 mg, 1.20 mmol, 60% yield). ¹H NMR (300MHz, CDCl₃) δ 5.02 (t, J = 5.4 Hz, 1H), 3.52 (t, J = 6.9 Hz, 2H), 1.82–1.73 (m, 2H), 1.64–1.56 (m, 2H), 1.52– 1.37 (m, 4H), 1.19 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 100.7, 81.6, 45.0, 36.2, 32.5, 26.9, 24.2, 23.8, 22.0. HRMS (ESI): m/z calcd for C₁₂H₂₂ClO₂ [M-H]⁺ 233.1308, found 233.1306.

2-(3-Bromopropyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(**3fa**). Method C was applied (384 mg, 1.53 mmol, 76% yield). The spectral data for **3fa** were in accordance with those reported in the literature.⁴⁷

2-(4-Bromobutyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(**3ga**). Method C was applied (395 mg, 1.49 mmol, 75% yield). ¹H NMR (300MHz , CDCl₃) δ 5.03 (t, *J* = 4.8 Hz, 1H), 3.40 (t, *J* = 6.9 Hz, 2H), 1.95–1.85 (m, 2H), 1.64–1.48 (m, 4H), 1.19 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 100.5, 81.7, 35.4, 33.6, 32.7, 24.2, 23.2, 22.0. HRMS (ESI): *m*/*z* calcd for C₁₁H₂₀BrO₂ [M-H]⁺ 263.0647, found 263.0637.

2-(6-Bromohexyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(**3ha**). Method A was applied (366 mg, 1.25 mmol, 63% yield). ¹H NMR (300MHz, CDCl₃) δ 5.02 (t, J = 5.1 Hz, 1H), 3.39 (t, J = 6.9 Hz, 2H), 1.90–1.80 (m, 2H), 1.62–1.55 (m, 2H), 1.46– 1.35 (m, 6H), 1.19 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 100.8, 81.6, 36.3, 33.9, 32.7, 28.7, 28.0, 24.2 (2C), 22.0. HRMS (ESI): m/z calcd for C₁₃H₂₄BrO₂ [M-H]⁺ 291.0960, found 291.0970.

2-(3-Iodopropyl)-4,4,5,5-tetramethyl-1,3-dioxolane (3ia). Method D was applied (196 mg, 0.66 mmol, 33% yield). The spectral data for **3ia** were in accordance with those reported in the literature.⁴⁷

2-(5-Benzyloxypentyl)-4,4,5,5-tetramethyl-1,3-dioxolane (**3ja).** Method D was applied (299 mg, 0.97 mmol, 49% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.30–7.17 (m, 5H), 4.96 (t, *J* = 5.1 Hz, 1H), 4.43 (s, 2H), 3.39 (t, *J* = 6.6 Hz, 2H), 1.59–1.50 (m, 4H), 1.37–1.32 (m, 4H), 1.13 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 138.6, 128.3, 127.6, 127.4, 100.8, 81.5, 72.8, 70.3, 36.4, 29.7, 26.2, 24.3, 24.2, 22.0. HRMS (ESI): *m/z* calcd for C₁₉H₃₀O₃Na [M+Na]⁺ 329.2093, found 329.2076.

2-(9-Benzyloxynonyl)-4,4,5,5-tetramethyl-1,3-dioxolane (**3ka**). Method B was applied (374 mg, 1.03 mmol, 52% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.27–7.16 (m, 5H), 4.97 (t, *J* = 5.1 Hz, 1H), 4.43 (s, 2H), 3.39 (t, *J* = 6.6 Hz, 2H), 1.59–1.49 (m, 4H), 1.34–1.22 (m, 12H), 1.13 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 138.5, 128.2, 127.4, 127.3, 100.8, 81.3, 72.7, 70.3, 36.3, 29.6, 29.5, 29.3 (3C), 26.0, 24.3, 24.1, 21.9. HRMS (ESI): *m/z* calcd for C₂₃H₃₈O₃Na [M+Na]⁺ 385.2719, found 385.2703.

2-(4-Carboxybutyl)-4,4,5,5-tetramethyl-1,3-dioxolane

(31a). Method B was applied. Maleic anhydride was used instead of MeBQ as an additive. Water was used instead of a saturated aqueous NaHCO₃ solution for extraction. After filtration of the drying agent, the solvent was evaporated and volatile materials were further removed under vacuum at 40 °C overnight to afford **31a** as a pale yellow oil (276 mg, 1.20 mmol, 60% yield). ¹H NMR (400 MHz, CDCl₃) δ 11.42 (br, 1H), 5.00 (t, *J* = 5.2 Hz, 1H), 2.32 (t, *J* = 7.6 Hz, 2H), 1.68–1.55 (m, 4H), 1.47–1.38 (m, 2H), 1.16 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 179.7, 100.5, 81.7, 35.9, 33.9, 24.6, 24.1, 23.9, 21.9. IR (neat) v 1710 cm⁻¹ (C=O). HRMS (ESI): *m/z* calcd for C₁₂H₂₁O₄ [M-H]⁺ 229.1440, found 229.1431.

2-(4-Benzyloxycarbonylbutyl)-4,4,5,5-tetramethyl-1,3-dioxolane (3ma). Method B was applied. Maleic anhydride was used instead of MeBQ as an additive (326 mg, 1.02 mmol, 51% yield). ¹H NMR (300 MHz, CDCl₃) δ 7.37–7.31 (m, 5H), 5.11 (s, 2H), 5.01 (t, *J* = 5.1 Hz, 1H), 2.37 (t, *J* = 7.5 Hz, 2H), 1.74–1.56 (m, 4H), 1.48–1.38 (m, 2H), 1.18 (s, 12H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 173.4, 136.1, 128.5, 128.1, 100.6, 81.6, 66.1, 36.0, 34.2, 24.9, 24.2, 24.0, 22.0. IR (neat) v 1738 cm⁻¹ (C=O). HRMS (ESI): *m*/*z* calcd for C₁₉H₂₇O₄ [M-H]⁺ 319.1909, found 319.1922.

2-(3-Bromopropyl)-5,5-dimethyl-1,3-dioxane (3fb) and 2-(**2-bromoethyl)-2,5,5-trimethyl-1,3-dioxane (5fb).** Method D was applied. 2,2-Dimethyl-1,3-propanediol was used instead of pinacol. Purification by silica gel column chromatography afforded a mixture of **3fb** and **5fb** as a pale yellow oil (323 mg, 1.36 mmol, 68% total yield, **3fb:5fb** = 2.1:1). ¹H NMR for **3fb** (300MHz, CDCl₃) δ 4.46 (t, J = 4.8 Hz, 1H), 3.61–3.38 (m, 6H), 2.06–1.97 (m, 2H), 1.81–1.75 (m, 2H), 1.18 (s, 3H), 0.72 (s, 3H). ¹H NMR for **5fb** (300MHz, CDCl₃) δ 3.61–3.38 (m, 6H), 2.28 (t, J = 8.4 Hz, 2H), 1.38 (s, 3H), 1.03 (s, 3H), 0.85 (s, 3H). ¹³C{¹H} NMR for **3fb** (75 MHz, CDCl₃) δ 101.1, 77.2, 33.7, 33.3, 30.1, 27.2, 22.9, 21.8. ¹³C{¹H} NMR for **5fb** (75 MHz, CDCl₃) δ 98.3, 70.4, 42.9, 29.8, 27.2, 22.8, 22.3, 19.9. HRMS (ESI): m/z calcd for C₉H₁₇BrKO₂ [M+K]⁺ 275.0049, found 275.0045.

2-(3-Bromopropyl)-4,4,6,6-tetramethyl-1,3-dioxane (3fc). Method D was applied. 2,4-Dimethyl-2,4-pentanediol was used instead of pinacol (347 mg, 1.31 mmol, 66% yield). ¹H NMR (300MHz, CDCl₃) δ 4.91 (t, J = 4.8 Hz, 1H), 3.43 (t, J = 6.9 Hz, 2H), 2.03–1.94 (m, 2H), 1.73–1.66 (m, 2H), 1.58 (d, J = 13.8 Hz, 1H), 1.45 (d, J = 13.8 Hz, 1H), 1.32 (s, 6H), 1.22 (s, 6H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 89.2, 71.0, 45.5, 34.1, 33.8, 33.3, 27.7, 25.1. HRMS (ESI): *m*/*z* calcd for C₁₁H₂₁O₂ [M-Br]⁺ 185.1542, found 185.1550.

2-(3-Bromopropyl)-4,4,7,7-tetramethyl-1,3-dioxepane

(3fd). Method D was applied. 2,5-Dimethyl-2,5-hexanediol was used instead of pinacol (360 mg, 1.29 mmol, 65% yield). ¹H NMR (300MHz, CDCl₃) δ 4.73 (t, J = 5.7 Hz, 1H), 3.40 (t, J = 6.9 Hz, 2H), 1.94–1.84 (m, 2H), 1.79–1.60 (m, 6H), 1.22 (s, 6H), 1.18 (s, 6H). ¹³C{¹H} NMR (75 MHz, CDCl₃) δ 90.9, 74.7, 35.5, 35.3, 33.7, 30.0, 28.7, 25.8. HRMS (ESI): m/z calcd for C₁₂H₂₃O₂ [M-Br]⁺ 199.1698, found 199.1693.

ASSOCIATED CONTENT

The Supporting Information is available free of charge on the ACS Publications website at DOI: xxx.

¹H and ¹³C NMR spectra for **3** (PDF)

AUTHOR INFORMATION

Corresponding Author

*E-mail for Y.U.: ura@cc.nara-wu.ac.jp.

ORCID

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

58 59

60

Yasuyuki Ura: 0000-0003-0484-1299

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This study was supported by JSPS KAKENHI Grant Numbers JP16H01028 in Precisely Designed Catalysts with Customized Scaffolding, JP25410116, JP18K05122, and JP18H03914.

REFERENCES

(1) Muzart, J. Aldehydes from Pd-catalysed oxidation of terminal olefins. *Tetrahedron* **2007**, *63*, 7505-7521.

47 (2) Guo, J.; Teo, P. Anti-Markovnikov oxidation and
48 hydration of terminal olefins. *Dalton Trans.* 2014, 43, 695249 6964.

(3) Dong, J. J.; Browne, W. R.; Feringa, B. L. PalladiumCatalyzed anti-Markovnikov Oxidation of Terminal Alkenes. *Angew. Chem., Int. Ed.* 2015, *54*, 734-744.

- (4) Takacs, J. M.; Jiang, X.-t. The Wacker Reaction and Related Alkene Oxidation Reactions. *Curr. Org. Chem.* 2003, 7, 369-396.
- 56 (5) Dong, J. J.; Fañanás-Mastral, M.; Alsters, P. L.; Browne,
 57 W. R.; Feringa, B. L. Palladium-Catalyzed Selective Anti-

Markovnikov Oxidation of Allylic Esters. *Angew. Chem., Int. Ed.* **2013**, *52*, 5561-5565.

(6) Michel, B. W.; Steffens, L. D.; Sigman, M. S. In *Organic Reactions*; John Wiley & Sons: Hoboken, 2014; Vol. 84, p 75-413.

(7) Jira, R. In *Applied Homogeneous Catalysis with Organometallic Compounds*; 2nd ed.; Cornils, B., Herrmann, W. A., Eds.; Wiley-VCH: Weinheim, 2002; Vol. 1, p 386-405.

(8) Mann, S. E.; Benhamou, L.; Sheppard, T. D. Palladium(II)-Catalysed Oxidation of Alkenes. *Synthesis* **2015**, *47*, 3079-3117.

(9) Nakaoka, S.; Murakami, Y.; Kataoka, Y.; Ura, Y. Maleimide-assisted anti-Markovnikov Wacker-type oxidation of vinylarenes using molecular oxygen as a terminal oxidant. *Chem. Commun.* **2016**, *52*, 335-338.

(10) Teo, P.; Wickens, Z. K.; Dong, G.; Grubbs, R. H. Efficient and Highly Aldehyde Selective Wacker Oxidation. *Org. Lett.* **2012**, *14*, 3237-3239.

(11) Wright, J. A.; Gaunt, M. J.; Spencer, J. B. Novel anti-Markovnikov regioselectivity in the Wacker reaction of styrenes. *Chem. Eur. J.* **2006**, *12*, 949-955.

(12) Dong, G.; Teo, P.; Wickens, Z. K.; Grubbs, R. H. Primary Alcohols from Terminal Olefins: Formal Anti-Markovnikov Hydration via Triple Relay Catalysis. *Science* **2011**, *333*, 1609-1612.

(13) Chowdhury, A. D.; Ray, R.; Lahiri, G. K. An iron catalyzed regioselective oxidation of terminal alkenes to aldehydes. *Chem. Commun.* **2012**, *48*, 5497-5499.

(14) Chen, G.-Q.; Xu, Z.-J.; Zhou, C.-Y.; Che, C.-M. Selective oxidation of terminal aryl and aliphatic alkenes to aldehydes catalyzed by iron(iii) porphyrins with triflate as a counter anion. *Chem. Commun.* **2011**, *47*, 10963-10965.

(15) Jiang, G.; Chen, J.; Thu, H.-Y.; Huang, J.-S.; Zhu, N.; Che, C.-M. Ruthenium Porphyrin-Catalyzed Aerobic Oxidation of Terminal Aryl Alkenes to Aldehydes by a Tandem Epoxidation–Isomerization Pathway. *Angew. Chem.*, *Int. Ed.* **2008**, *47*, 6638-6642.

(16) Chen, J.; Che, C.-M. A Practical and Mild Method for the Highly Selective Conversion of Terminal Alkenes into Aldehydes through Epoxidation–Isomerization with Ruthenium(IV)–Porphyrin Catalysts. *Angew. Chem., Int. Ed.* **2004**, *43*, 4950-4954.

(17) Feringa, B. L. Catalytic oxidation of alk-1-enes to aldehydes. J. Chem. Soc., Chem. Commun. **1986**, 909-910.

(18) Wenzel, T. T. Oxidation of olefins to aldehydes using a palladium-copper catalyst. *J. Chem. Soc., Chem. Commun.* **1993**, 862-864.

(19) Ogura, T.; Kamimura, R.; Shiga, A.; Hosokawa, T. Reversal of Regioselectivity in Wacker-Type Oxidation of Simple Terminal Alkenes and Its Paired Interacting Orbitals (PIO) Analysis. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1555-1557. (20) Wickens, Z. K.; Morandi, B.; Grubbs, R. H. Aldehyde-Selective Wacker-Type Oxidation of Unbiased Alkenes Enabled by a Nitrite Co-Catalyst. *Angew. Chem., Int. Ed.* **2013**, *52*, 11257-11260.

(21) Hosokawa, T.; Ohta, T.; Kanayama, S.; Murahashi, S. Palladium(II)-catalyzed acetalization of terminal olefins bearing electron-withdrawing substituents with optically active diols. *J. Org. Chem.* **1987**, *52*, 1758-1764.

2

60

(22) Hosokawa, T.; Yamanaka, T.; Murahashi, S.-I.
Palladium(II)-catalysed asymmetric acetalization of alkenes. *J. Chem. Soc., Chem. Commun.* **1993**, 117-119.

- 3 (23) Hosokawa, T.; Yamanaka, T.; Itotani, M.; Murahashi,
 4 S.-I. Palladium(II)-catalyzed asymmetric acetalization of
- 5 alkenes. J. Org. Chem. **1995**, 60, 6159-6167.
- 6 (24) Jia, L.; Jiang, H.; Li, J. Palladium(II)-catalyzed
 7 oxidation of acrylate esters to acetals in supercritical carbon
 8 dioxide. *Chem. Commun.* 1999, 985-986.
- 9 (25) Kishi, A.; Sakaguchi, S.; Ishii, Y. Acetalization of
- 10 Alkenes Catalyzed by Pd(OAc)₂/NPMoV Supported on 11 Activated Carbon under a Dioxygen Atmosphere. *Org. Lett.*
- 11 Activated Carbon under a Dioxygen Atmosphere. *Org. Lett.* 12 **2000**, *2*, 523-525.
- (26) Lai, J.; Shi, X.; Dai, L. Reversal of regiochemistry of
 Wacker-type reactions oriented by heteroatoms. J. Org. *Chem.* 1992, 57, 3485-3487.
- (27) Hosokawa, T.; Aoki, S.; Murahashi, S.-I. Palladium(II)Catalyzed Acetalization of Allylic Acetates and Its
 Utilization for the Synthesis of 2-Cyanovinyl Ketones.
 Synthesis 1992, 558-561.
- 19 (28) Yamamoto, M.; Nakaoka, S.; Ura, Y.; Kataoka, Y.
 20 (28) Yamamoto, M.; Nakaoka, S.; Ura, Y.; Kataoka, Y.
 21 Palladium-catalyzed synthesis of terminal acetals via highly
 22 selective anti-Markovnikov nucleophilic attack of pinacol
 23 on vinylarenes, allyl ethers, and 1,5-dienes. *Chem. Commun.*23 2012, 48, 1165-1167.
- (29) Matsumura, S.; Sato, R.; Nakaoka, S.; Yokotani, W.;
 Murakami, Y.; Kataoka, Y.; Ura, Y. Palladium-Catalyzed
 Aerobic Synthesis of Terminal Acetals from Vinylarenes
 Assisted by π-Acceptor Ligands. *ChemCatChem* 2017, 9,
 751-757.
- (30) Hosokawa, T.; Ataka, Y.; Murahashi, S.-I. Catalysis of
 Pd(II)-Catalyzed Acetalization of Alkenes with Diols. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 166-169.
- 32 (31) Chowdhury, A. D.; Lahiri, G. K. A generalized
 33 approach for iron catalyzed chemo- and regioselective
 34 formation of anti-Markovnikov acetals from styrene
 35 derivatives. *Chem. Commun.* 2012, 48, 3448-3450.
- (32) Kumar, M. A.; Swamy, P.; Naresh, M.; Reddy, M. M.;
 Rohitha, C. N.; Prabhakar, S.; Sarma, A. V. S.; Kumar, J. R.
 P.; Narender, N. Iodine-catalyzed tandem synthesis of terminal acetals and glycol mono esters from olefins. *Chem.*
- *Commun.* **2013**, *49*, 1711-1713.
- 40 (33) Hosokawa, T.; Makabe, Y.; Shinohara, T.; Murahashi,
- 41 S.-I. SYNTHESIS OF NATURAL AND UNNATURAL 42 FRONTALIN. *Chem. Lett.* **1985**, *14*, 1529-1530.
- (34) Lloyd, W. G.; Luberoff, B. J. Oxidations of olefins with alcoholic palladium(II) salts. *J. Org. Chem.* **1969**, *34*, 3949-3952.
- 46 (35) Kongkathip, B.; Sookkho, R.; Kongkathip, N.
 47 CONVENIENT SYNTHETIC ROUTE TO (+-)48 FRONTALIN. *Chem. Lett.* 1985, 14, 1849-1850.
 49

(36) Muzart, J. Palladium-catalysed reactions of alcohols. Part C: Formation of ether linkages. *Tetrahedron* **2005**, *61*, 5955-6008.

(37) Speziali, M. G.; Costa, V. c. V.; Robles-Dutenhefner, P. c. A.; Gusevskaya, E. V. Aerobic Palladium(II)/Copper(II)-Catalyzed Oxidation of Olefins under Chloride-Free Nonacidic Conditions. *Organometallics* **2009**, *28*, 3186-3192.

(38) Chu, C. K.; Ziegler, D. T.; Carr, B.; Wickens, Z. K.; Grubbs, R. H. Direct Access to β -Fluorinated Aldehydes by Nitrite-Modified Wacker Oxidation. *Angew. Chem., Int. Ed.* **2016**, *55*, 8435-8439.

(39) Jiang, Y.-Y.; Zhang, Q.; Yu, H.-Z.; Fu, Y. Mechanism of Aldehyde-Selective Wacker-Type Oxidation of Unbiased Alkenes with a Nitrite Co-Catalyst. *ACS Catal.* **2015**, *5*, 1414-1423.

(40) Keith, J. A.; Nielsen, R. J.; Oxgaard, J.; Goddard, W. A. Unraveling the Wacker Oxidation Mechanisms. *J. Am. Chem. Soc.* **2007**, *129*, 12342-12343.

(41) Hosokawa, T.; Nomura, T.; Murahashi, S.-I. Palladium– copper–DMF complexes involved in the oxidation of alkenes. *J. Organomet. Chem.* **1998**, *551*, 387-389.

(42) Hosokawa, T.; Aoki, S.; Takano, M.; Nakahira, T.; Yoshida, Y.; Murahashi, S.-I. Palladium(II)-catalysed oxidation of carbon–carbon double bonds of allylic compounds with molecular oxygen; regioselective formation of aldehydes. *J. Chem. Soc., Chem. Commun.* **1991**, 1559-1560.

(43) Michelin, R. A.; Facchin, G.; Uguagliati, P. Functionalized isocyanides as ligands. Synthesis of 2-(chloromethyl)- and 2-(iodomethyl)phenyl isocyanides and their transition-metal complexes. *Inorg. Chem.* **1984**, *23*, 961-969.

(44) Hodgson, D. M.; Kloesges, J.; Evans, B. Asymmetric Synthesis of Terminal *N*-tert-Butylsulfinyl Aziridines from Organoceriums and an α -Chloroimine. *Org. Lett.* **2008**, *10*, 2781-2783.

(45) Itoh, T.; Matsueda, T.; Shimizu, Y.; Kanai, M. Copper-Catalyzed Oxyboration of Unactivated Alkenes. *Chem. Eur. J.* **2015**, *21*, 15955-15959.

(46) Nookaraju, U.; Kumar, P. Total synthesis of (+)petromyroxol via tandem α -aminoxylation–allylation and asymmetric dihydroxylation–S_N2 cyclization approach. *RSC Adv.* **2015**, *5*, 63311-63317.

(47) Stowell, J. C.; Polito, M. A. A facile procedure for producing γ -halo butyraldehyde acetals. *J. Org. Chem.* **1992**, *57*, 2195-2196.

ACS Paragon Plus Environment

A graphic entry for the Table of Contents (TOC)