


RSC Advances



This article can be cited before page numbers have been issued, to do this please use: L. Zhang, Y. Li, L. Y. Jin and F. Liang, *RSC Adv.*, 2015, DOI: 10.1039/C5RA13708J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

COMMUNICATION

Metal-free C—N cross-coupling of electrophilic compounds and N-haloimides†

Cite this: DOI: 10.1039/x0xx00000x

Luyan Zhang,^a Yanru Li,^a Long-Yi Jin^c and Fushun Liang^{*,a,b}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x


www.rsc.org/

When DBU is added, the cross-coupling reaction between alkyl halides (halogen = Cl, Br and I) and N-haloimides (halogen = Cl, Br) occurs, resulting in the formation of aminated products. A halogen bond activated nucleophilic substitution mechanism was proposed. The methodology represents an elegant example of applying halogen bond activation strategy in organic transformation.

Amine is important raw material and intermediate in medicine, pesticide, as well as liquid crystal material. Phthalimides are versatile synthetic tools that are primarily employed in the preparation of amines via the Gabriel reaction.¹ Gabriel reaction has proven to be a very effective means of forming *N*-alkylphthalimides. Starting from *N*-substituted phthalimides, for example, the use of the Nefkens' reagent, *N*-alkylphthalimides can also be prepared.² Adimurthy et al. found an L-proline catalyzed transamidation method of phthalimide with amines, to achieve new amines.³ A. K. Yadav and L. D. S. Yadav used *N*-tosylhydrazone and phthalimides to prepare amine.⁴ Additionally, transition metal-catalyzed C—N coupling has been developed as a powerful method.⁵ Herein, we would like to report a new type of cross-coupling reaction⁶ between alkyl halides⁷ and *N*-haloimides, which provides an efficient and straightforward method for the synthesis of amines and functionalized amines.

In our recent research, we have developed *N*-haloimides to be a versatile amination reagent used in organic transformation.⁸ In further work, we found that, with organobases like DBU as the activator,⁹ benzyl chloride can react with *N*-chlorophthalimide (NCP, 1.2 equiv), giving the corresponding imidation product in 91% yield. It was not surprising that no reaction can occur in the absence of DBU (eq 1). The possible mechanism for the imidation is depicted in Scheme 1.¹⁰ The interaction between NCP and DBU gives halogen bond adduct I,¹¹ which delivers an ion pair intermediate II containing highly electrophilic chlorine cation and potentially nucleophilic phthalimide anion.¹² The nucleophilic substitution of

alkyl halides by phthalimide anion affords the imidation product.¹³ We think that the halogen bond between 1-chloro-2,3,4,6,7,8,9,10-octahydropyrimido[1,2-*a*]azepinium (III) and alkyl halide may facilitate the nucleophilic substitution reaction.⁹ The mutual activation induced by halogen bonding makes otherwise inert reaction to take place.¹⁴




Scheme 1 Proposed mechanism

Our initial optimization of the model reaction commences from benzyl chloride (1a). The impact of various parameters on the efficiency of the reaction of 1a with NCP (2a) is summarized in Table 1. In the presence of DBU (1.2 equiv), the reaction mixture conducted in DCE at room temperature, afforded product 3a in 31%

COMMUNICATION

isolated yield (entry 1). Other solvents screened include MeCN, DMF and DMSO (entries 2–4). DMSO proved to be the most efficient (85% yield) (entry 4).¹⁵ The reaction mixture performed at 60 °C gave product **3a** in improved yield (91%) and shortened reaction time (5 min) (entry 5). Besides DBU, NaOH, *t*-BuONa, PPh₃, DBN, MTBD, and DABCO were selected as the activators. MTBD gave similar reactivity and DBN exhibits relatively lower efficiency (entries 6 and 7). However, NaOH, *t*-BuONa, PPh₃ and DABCO appear to be inert (entries 8–11). Catalytic amount of DBU, e.g., 0.2 equiv, could not drive the reaction to completion (entry 12), which may support the presence of the halogen bonding interaction between 1-chloro-2,3,4,6,7,8,9,10-octahydropyrimido[1,2-*a*]azepinium and alkyl halide, i.e., adduct **III** in Scheme 1.¹⁶


Table 1 Optimization of the reaction conditions.^a

Entry	Temp (°C)	Activator	Solvent ^b	Time(min)	Yield (%) ^c
1	rt	DBU	DCE	180	31
2	rt	DBU	MeCN	120	56
3	rt	DBU	DMF	120	57
4	rt	DBU	DMSO	30	85
5	60	DBU	DMSO	5	91
6	60	DBN	DMSO	180	61
7	60	MTBD	DMSO	6	88
8	60	NaOH	DMSO	180	n.r
9	60	<i>t</i> -BuONa	DMSO	180	n.r
10	60	Ph ₃ P	DMSO	180	n.r
11	60	DABCO	DMSO	180	n.r
12	60	DBU (0.2 equiv)	DMSO	180	16

^a Reactions were carried out with **1a** (1.0 mmol), **2a** (1.2 equiv) and activator (1.2 equiv) in solvent (2.0 mL). ^b Dry solvents were used in all cases to prevent the hydrolysis of benzyl chloride. ^c Isolated yield.

With optimized reaction conditions in hand (Table 1, entry 5), we investigated this methodology by reacting various alkyl halides with NCP (Table 2).¹⁷ The scope of this new coupling reaction is fairly broad. Alkyl chlorides examined includes benzyl chloride, *n*-butyl chloride, α -chlorocyclopentanone and α -chloro *N,N*-dimethylacetamine. Benzyl chloride is highly efficient, while *n*-butyl chloride exhibits relatively lower efficiency. α -Chlorocyclopentanone and α -chloro *N,N*-dimethylacetamine gave satisfactory yields. Primary alkyl bromides containing various substituents like benzyl, allyl, ethyl, homobenzyl, isobutyl are suitable substrates. A lot of functional groups are compatible with the reaction conditions, such as methoxy, dimethoxyacetal, carbonyl and ester. Secondary alkyl


bromides like isopropyl bromide and diphenylmethyl bromide can also afford the corresponding aminated product, albeit in low yields. However, cyclic alkyl bromide, i.e., cyclohexyl bromide, was inefficient. Alkyl iodide like methyl iodide worked very well. In one word, alkyl halides RX (X = Cl, Br and I) exhibit increased reactivity in the order of RCl, RBr and RI. The gram-scale synthesis of **3a** (1.51 g) was also achieved in 92% yield by DBU-mediated reaction of benzyl bromide and NCP.

Table 2 Cross-coupling reaction of alkyl halides with NCP^{a,b}

^a Reaction conditions: **1** (1.0 mmol), **2a** (1.2 equiv) and DBU (1.2 equiv) in DMSO (2.0 mL). ^b Isolated yields.

Interestingly, we found that dihalides are also suitable for the cross-coupling reaction (Scheme 2). In the reaction of 1,2-dibromoethane with NCP (1.2 equiv), two types of products could be prepared, depending the feed ratio of DBU employed. When 1.2 equiv of DBU was used, the mixture afforded monobromo


substituted amine **3q** in 56% yield. In the case of 2.2 equiv of DBU, further elimination of HBr occurs, giving rise to enamine **4** in 53% yield. However, no elimination of HBr or HCl was observed for 1,4-dibromobutane, 1,6-dibromooctane and 1,2-dichloroethane substrates, even though excess amount of DBU was used. The corresponding imidation products **3r-t** with a pendant bromine or chlorine atom were attained in low to moderate yields, which may be potentially useful in further transformation.¹⁹ Mixed dihalides were also subjected to the reaction sequence. For example, 1-bromo-4-chlorobutane gave chlorine atom-tolerant imidation product **3u** as the only product, due to the different reactivity between bromine and chlorine atoms.



Scheme 2 Cross-coupling reaction of dihalides with NCP.

We investigated the efficacy of *N*-haloimides by reacting them with benzyl bromide (Scheme 3). Different *N*-haloimide exhibits different reactivity. NBP is less efficient than NCP (65%). Similar trend was observed for *N*-halosuccinimides. NCS gave product **7** in 51% yield while NBS and NIS could not react. From Scheme 1 one can see there are two types of halogen bonding interaction, *N*-haloimide and DBU (adduct I), and the resulting chlorine cation in 1-chloro-2,3,4,6,7,8,9,10-octahydropyrimido[1,2-*a*]azepinium with alkyl halide RX (adduct III). For the latter, chlorine atom gives stronger halogen bonding ability than bromine atom, due to the larger electron negativity, thus leading to R—X bond cleavage easier (nucleophilic substitution is the rate-determining step). *N*-Bromosaccharin was a competent reagent, affording 2-benzylbenzo[*d*]isothiazol-3(2*H*)-one 1,1-dioxide **8** in 89% yield. Other *N*-haloimides like 1,3-dibromo-5,5-dimethyl hydantoin and *N*-bromoacetamide proved to be inefficient. All the above reactions

indicated relatively high efficiency and broad scope for both alkyl halides and *N*-haloimides.



Scheme 3 Cross-coupling reaction of benzyl bromide with *N*-haloimides.

In summary, we have developed a highly efficient and metal-free cross-coupling reaction of alkyl halides and *N*-haloimides. With this strategy, various alkyl- and functionalized alkylamines were synthesized. The scope of the reaction is broad in terms of both alkyl halides and *N*-haloimides. A halogen bond activated nucleophilic substitution mechanism was proposed. This work demonstrated the feasibility and potential of halogen bonding activation in organic transformation.

Financial support from the National Natural Science Foundation of China (21172034 and 21372039), and the 111 Project is gratefully acknowledged.

Notes and references

- ^a Department of Chemistry, Northeast Normal University, Changchun 130024, China. Email: liangfs112@nenu.edu.cn; Fax: (+86) 431-85099759
^b Key Laboratory for UV-Emitting Materials and Technology of Ministry of Education, Northeast Normal University, Changchun 130024, China
^c Faculty of Chemistry, Yanbian University

- (a) M. S. Gibson and R. W. Bradshaw, *Angew. Chem. Int. Ed.*, 1968, **7**, 919; (b) J. C. Sheehan and William A. Bolhofer, *J. Am. Chem. Soc.*, 1950, **72**, 2786.
- (a) G. H. L. Nefkens, *Nature*, 1960, **185**, 309; (b) G. H. L. Nefkens and G. I. Tesser, *J. Am. Chem. Soc.*, 1961, **83**, 1263; (c) P. M. Worster, C. C. Leznoff and C. R. McArthur, *J. Org. Chem.*, 1980, **45**, 174; (d) E. G. Bowen and D. J. Wardrop, *J. Am. Chem. Soc.*, 2009, **131**, 6062.
- S. N. Rao, D. C. Mohan and S. Adimurthy, *Org. Lett.*, 2013, **15**, 1496.
- A. K. Yadav and L. D. S. Yadav, *RSC Adv.*, 2014, **4**, 34764.
- Selected reviews: (a) G. Evano, N. Blanchard and M. Toumi, *Chem. Rev.*, 2008, **108**, 3054; (b) J. Bariwal and E. Van der Eycken, *Chem. Soc. Rev.*, 2013, **42**, 9283; (c) J. F. Hartwig, *Acc. Chem. Res.*, 2008, **41**, 1534; (d) J. P. Wolfe, S. Wagaw, J.-F. Marcoux and S.

- Buchwald, *Acc. Chem. Res.*, 1998, **31**, 805; (e) D. S. Surry and S. L. Buchwald, *Angew. Chem. Int. Ed.*, 2008, **47**, 6338.
- 6 Cross-coupling of halides: (a) J. Zhou and G. C. Fu, *J. Am. Chem. Soc.*, 2004, **126**, 1340; (b) X. Wu, J. W. T. See, K. Xu, H. Hirao, J. Roger, J. -C. Hierso and J. Zhou, *Angew. Chem. Int. Ed.*, 2014, **53**, 13573; (c) O. Gutierrez, J. C. Tellis, D. N. Primer, G. A. Molander and M. C. Kozlowski, *J. Am. Chem. Soc.*, 2015, **137**, 4896; (d) N. A. Owston and G. C. Fu, *J. Am. Chem. Soc.*, 2010, **132**, 11908; (e) Z. Lu and G. C. Fu, *Angew. Chem. Int. Ed.*, 2010, **49**, 6676; (f) H. Xu, C. Zhao, Q. Qian, W. Deng and H. Gong, *Chem. Sci.*, 2013, **4**, 4022; (h) C. Zhao, X. Jia, X. Wang and H. Gong, *J. Am. Chem. Soc.*, 2014, **136** (50), 17645; (i) C.-T. Yang, Z.-Q. Zhang, J. Liang, J.-H. Liu, X.-Y. Lu, H.-H. Chen and L. Liu, *J. Am. Chem. Soc.*, 2012, **134**, 11124.
- 7 Alkyl halides are ubiquitous building blocks throughout organic chemistry owing to their participation in radical reactions, nucleophilic substitution reactions and organometallic cross-coupling reactions. (a) H. Togo, *Advanced Free Radical Reactions for Organic Synthesis* (Elsevier, 2004); (b) F. A. Carey and R. J. Sundberg, *Advanced Organic Chemistry* 5th edn (Springer, 2007); (c) A. Rudolph and M. Lautens, *Angew. Chem. Int. Ed.*, 2009, **48**, 2656; (d) J. Terao and N. Kambe, *Acc. Chem. Res.*, 2008, **41**, 1545; (e) N. A. Owston and G. C. Fu, *J. Am. Chem. Soc.*, 2010, **132**, 11908.
- 8 (a) H. Tan, M. Li and F. Liang, *RSC Adv.*, 2014, **4**, 33765; (b) Y. Wei, S. Lin and F. Liang, *Org. Lett.*, 2012, **14**, 4202; (c) Y. Wei, S. Lin, F. Liang and J. Zhang, *Org. Lett.*, 2013, **15**, 852; (d) Y. Wei, F. Liang and X. Zhang, *Org. Lett.*, 2013, **15**, 5186; (e) Y. Li, L. Zhang, H. Yuan, F. Liang and J. Zhang, *Synlett*, 2015, **26**, 116.
- 9 For a review of NBS activation by Lewis base, see: (a) S. E. Denmark, W. E. Kuester and M. T. Burk, *Angew. Chem. Int. Ed.*, 2012, **51**, 10938; Selected papers: (b) A. Sakakura, A. Ukai and K. Ishihara, *Nature*, 2007, **445**, 900; (c) S. A. Snyder and D. S. Treitler, *Angew. Chem. Int. Ed.*, 2009, **48**, 7899; (d) X. L. Cui and R. S. Brown, *J. Org. Chem.*, 2000, **65**, 5653; (e) X. Huang, M. Seid and J. W. Keillor, *J. Org. Chem.*, 1997, **62**, 7495.
- 10 A possible radical pathway was excluded via a control experiment with TEMPO as the radical trapping reagent. Also, the model reaction worked well under dark.
- 11 For reviews on halogen bonding, see: (a) P. Metrangolo, H. Neukirch, T. Pilati and G. Resnati, *Acc. Chem. Res.*, 2005, **38**, 386; (b) P. Auffinger, F. A. Hays, E. Westhof and P. S. Ho, *Proc. Natl. Acad. Sci. USA*, 2004, **101**, 16789; (c) P. Metrangolo and G. Resnati, *Chem. Eur. J.*, 2001, **7**, 2511; Selected papers on halogen bonding: (d) F. Sladojevich, E. McNeill, J. Börgel, S.-L. Zheng and T. Ritter, *Angew. Chem. Int. Ed.*, 2015, **54**, 3712; (e) I. Castellote, M. Moron, C. Burgos, J. Alvarez-Builla, A. Martin, P. Gomez-Sal and J. J. Vaquero, *Chem. Commun.*, 2007, **43**, 1281; (f) K. Raatikainen and K. Rissanen, *Chem. Sci.*, 2012, **3**, 1235; (g) Y.-M. Wang, J. Wu, C. Hoong, V. Rauniyar and F. D. Toste, *J. Am. Chem. Soc.*, 2012, **134**, 12928; (h) P. Metrangolo, F. Meyer, T. Pilati, G. Resnati and G. Terraneo, *Angew. Chem. Int. Ed.*, 2008, **47**, 6114; (i) C. K. Tan, L. Zhou and Y.-Y. Yeung, *Synlett*, 2011, **10**, 1335; (j) F. Chen, C. K. Tan and Y.-Y. Yeung, *J. Am. Chem. Soc.*, 2013, **135**, 1232.
- 12 For an excellent review on halogen-bonded adducts are pre-reactive complexes, see: (a) A. C. Legon, *Angew. Chem. Int. Ed.*, 1999, **38**, 2686. Selected papers: (b) S. M. Walter, F. Kniep, E. Herdtweck and S. M. Huber, *Angew. Chem. Int. Ed.*, 2011, **50**, 7187; (c) R. Castelli, S. Schindler, S. M. Walter, F. Kniep, H. S. Overkleeft, G. A. Van der Marel, S. M. Huber and J. D. C. Codée, *Chem. Asian J.*, 2014, **9**, 2095; (d) F. Kniep, L. Rout, S. M. Walter, H. K. V. Bensch, S. H. Jungbauer, E. Herdtweck and S. M. Huber, *Chem. Commun.*, 2012, **48**, 9299.
- 13 Both S_N1 and S_N2 pathway are possible, depending on different structure of alkyl halides employed.
- 14 Halogen bonding has received relatively little attention from synthetic organic chemists. Early report: (a) A. Bruckmann, M. A. Pena and C. Bolm, *Synlett*, 2008, **06**, 900; (b) O. Coulembier, F. Meyer and P. Dubois, *Polym. Chem.*, 2010, **1**, 434; Huber's work: (c) F. Kniep, S. M. Walter, E. Herdtweck and S. M. Huber, *Chem. – Eur. J.*, 2012, **18**, 1306; (d) F. Kniep, S. H. Jungbauer, Q. Zhang, S. M. Walter, S. Schindler, I. Schnapperelle, E. Herdtweck and S. M. Huber, *Angew. Chem., Int. Ed.*, 2013, **52**, 7028; (e) S. H. Jungbauer, S. M. Walter, S. Schindler, L. Rout, F. Kniep and S. M. Huber, *Chem. Commun.*, 2014, **50**, 6281; (f) S. H. Jungbauer, S. Schindler, F. Kniep, S. M. Walter, L. Rout and S. M. Huber, *Synlett*, 2013, **24**, 2624.
- 15 For halogen cation stabilized by DMSO, see: (a) Y. Ashikari, A. Shimizu, T. Nokami and J. Yoshida, *J. Am. Chem. Soc.*, 2013, **135**, 16070; (b) A. Shimizu, R. Hayashi, Y. Ashikari, T. Nokami and J. Yoshida, *Beilstein J. Org. Chem.*, 2015, **11**, 242; (c) S. Song, X. Sun, X. Li, Y. Yuan and N. Jiao, *Org. Lett.*, 2015, **17**, 2886.
- 16 That DBU binding to the halogen in any step will help us to understand why stoichiometric amount of DBU is required in the reaction.
- 17 When KI-starch test paper was dipped into the reaction system, it turns blue after water treatment. This indicates the formation of dihalogen.
- 18 The C—C double bond is tolerant under such conditions. One possible reason is no reaction occurs on the C—C double bond due to the low concentration of the dihalogen. The other reason is halogenation takes place, followed by a rapid elimination. Thus, the net result is the C—C double bond is intact.
- 19 No second coupling (imidation) takes place.