

# A Journal of the Gesellschaft Deutscher Chemiker A Deutscher Chemiker GDCh International Edition www.angewandte.org

### **Accepted Article**

Title: Ligand-controlled Palladium-catalyzed Carbonylation of Alkynols: Highly Selective Synthesis of  $\alpha$ -Methylene- $\beta$ -lactones

Authors: Yao Ge, Fei Ye, Jiawang Liu, Ji Yang, Anke Spannenberg, Haijun Jiao, Ralf Jackstell, and Matthias Beller

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.202006550

Link to VoR: https://doi.org/10.1002/anie.202006550

# WILEY-VCH

COMMUNICATION

#### WILEY-VCH

## Ligand-controlled Palladium-catalyzed Carbonylation of Alkynols: Highly Selective Synthesis of $\alpha$ -Methylene- $\beta$ -lactones

Yao Ge<sup>+</sup>, Fei Ye<sup>+</sup>, Jiawang Liu, Ji Yang, Anke Spannenberg, Haijun Jiao, Ralf Jackstell, and Matthias Beller\*

[a] Dr. Y. Ge, Dr. F. Ye, Dr. J. Liu, J. Yang, Dr. A. Spannenberg, Dr. H. Jiao, Dr. R. Jackstell, Prof. Dr. M. Beller Leibniz-Institute for Catalysis, Albert-Einstein-Straße 29a, 18059 Rostock, Germany E-mail: <u>matthias.beller@catalysis.de</u>

[b] Dr. F. Ye

Key Laboratory of Organosilicon Chemistry and Material Technology of Ministry of Education, Key Laboratory of Organosilicon Chemistry and Material Technology of Zhejiang Province, Hangzhou Normal University, No. 2318, Yuhangtang Road, 311121, Hangzhou, PR China <sup>†</sup> These authors contributed equally to this work.

Supporting information for this article is given via a link at the end of the document.

**Abstract:** The first general and regioselective Pd-catalyzed cyclocarbonylation to give  $\alpha$ -methylene- $\beta$ -lactones is reported. Key to the success for this process is the use of a specific sterically demanding phosphine ligand based on *N*-arylated imidazole (L11) in the presence of Pd(MeCN)<sub>2</sub>Cl<sub>2</sub> as pre-catalyst. A variety of easily available alkynols provide under additive-free conditions the corresponding  $\alpha$ -methylene- $\beta$ -lactones in moderate to good yields with excellent regio- and diastereoselectivity. The applicability of this novel methodology is showcased by the direct carbonylation of biologically active molecules including natural products.

 $\alpha$ -Alkylidene- $\beta$ -lactones have emerged as important synthetic targets due to their occurrence in a variety of natural compounds and biologically active molecules (Scheme 1).<sup>[1]</sup> For example, lactones  $A^{[1a-c]}$  and  $B^{[1d]}$  were separated from Grazielia species and Disynaphia multicrenulata. Moreover, C exhibits promising inhibitory activities against certain fungal pathogens.<sup>[1e]</sup> In addition,  $\alpha$ -alkylidene- $\beta$ -lactones **D** and **E** were studied as potent and selective inhibitors for serine hydrolase ABHD16A.[1f-g] Following this work, F also served as a sensitive probe for detecting ABHD16A activity in mouse brain membrane lysates.<sup>[1g]</sup> Apart from these medicinal applications, specifically a-methylene- $\beta$ -lactones attracted interest in material sciences; e.g. welldefined copolymers with controllable molecular weight and polydispersity were prepared by narrow ring-opening polymerization. Here, the vinylidene groups of the lactones could be further functionalized, producing well-defined blocks with designable segments.<sup>[2]</sup> In organic chemistry, the high degree of functional groups in a compact manner allows diverse synthetic utilizations and makes this class of compounds interesting building blocks.<sup>[3]</sup> More specifically, owing to their inherent strain in the four-membered ring, they readily undergo ring opening reactions with a wide range of nucleophiles by either acyl C-O or alkyl C-O bond cleavage.[3f,3g] Besides the usual electrophilic sites of the carbonyl and oxetane carbon atoms contained in  $\beta$ lactones, Michael-type additions of nucleophiles and radicals at the methylene carbon atom are feasible<sup>[3h]</sup> and offer attractive possibilities for preparative purposes. Notably, the carboxyactivated exo-methylene group serves as reactive dienophile, while the  $\alpha,\beta$ -unsaturated carbonyl moiety promises potential as heterodiene for [4+2]-cycloadditions.<sup>[3e]</sup> Moreover, α-methylene- $\beta$ -lactones constitute convenient allene equivalents as

demonstrated in the decarboxylation to allenes on thermolysis.<sup>[3b,3e]</sup>



Scheme 1. Natural compounds and biologically active molecules with  $\alpha\text{-}$  methylene- $\beta\text{-}lactone$  skeleton

Considering the value of  $\alpha$ -alkylidene- $\beta$ -lactones in organic synthesis, significant interest in their preparation exists.<sup>[4]</sup> Because of the dense functionalization, mainly special synthetic methods have been developed for this class of compounds including [2+2]-cycloaddition of ketenes,<sup>[3n,4a-c]</sup> lactonization of  $\beta$ -hydroxycarboxylic acids or derivatives,<sup>[3d,4d-g]</sup> elimination of selenoxide from  $\alpha$ -methyl-substituted lactones,<sup>[3b]</sup> and deoxygenation of  $\beta$ -peroxylactones.<sup>[3f,4g-h]</sup>

More recently, also rhodium- or palladium-catalyzed carbonylation of alkynols have been disclosed.<sup>[5]</sup> Thus, *a*-(triorganosilyl)-methylene- $\beta$ -lactones (Scheme 2a),<sup>[5a-b]</sup> *a*-(alkoxycarbonyl)-methylene- $\beta$ -lactones (Scheme 2b),<sup>[5c-d]</sup> and (*Z*)- $\alpha$ -chloro/bromo-alkylidene- $\beta$ -lactones (Scheme 2b)<sup>[5e-f]</sup> were obtained. Notably, the synthesis of parent  $\alpha$ -methylene- $\beta$ -lactones gained less success due to the highly reactive *exo*-methylene double bond. In fact, to the best of our knowledge, there is only one reported example for the carbonylation of 1-methyl-2-butyn-1-ol described, leading to 4,4-dimethyl-3-methyleneoxetan-2-one in 5% yield with probably polymeric esters as other products through a palladium-catalyzed process (Scheme 2c).<sup>[5g]</sup>

Despite these problems, we thought the cyclocarbonylation of propargylic alcohols in the presence of an improved catalyst would offer a most straightforward and atom-efficient access to these products. Thus, we became attracted by this challenge.

### COMMUNICATION

Based on our interest in the development of carbonylation reactions,<sup>[6]</sup> herein, we report the first general and highly selective Pd-catalyzed carbonylation of propargylic alcohols to provide a family of new  $\alpha$ -methylene- $\beta$ -lactones (Scheme 2d).



Scheme 2. Rhodium- or palladium-catalyzed cyclocarbonylation of alkynols

At the beginning of our studies, 1-ethynyl-1-cyclohexanol 1a was chosen as model substrate. To identify a suitable catalyst system, a variety of ligands (in the case of diphosphine ligands 2 mol%, in the case of monophosphine ligands 4 mol%) were tested in the presence of  $[Pd(\pi-cinnamyl)Cl]_2$  (Figure 1). Initially, the reactivity of bidentate phosphines L1-L5 with different backbones and chelating units was evaluated. When L1 (Xantphos), L2 (BINAP), L4 (d<sup>t</sup>bpx) and L5 were applied, almost equal amounts of the desired  $\beta$ -lactone 2a and butenolide 3a were obtained. L3 (DPPF) proved to be not suitable at all, leading to 3a in 55% with 3% yield for 2a. No progress was achieved when monodentate ligand L6 (BuPAd<sub>2</sub>) was used in this reaction. However, in the presence of L7 (P<sup>4</sup>Bu<sub>3</sub>) a slightly improved regioselectivity (71/29) was obtained. Based on this result, we assumed that tert-butyl groups may have a positive influence on the desired branched selectivity. Thus, other monodentate ligands L8-L11 were tried with different backbones bearing tert-butyl substituents on the phosphorus atom. Indeed, more-sterically hindered L8 (JohnPhos) gave 80/20 selectivity and 57% yield of 2a. When the N-phenylpyrrole-based ligand L9 was applied, a similar selectivity (77/23) and a higher yield (67%) were observed. To further increase the steric bulk of the ligand, we introduced substituents on the ortho position of the phenyl group resulting in the new ligand L10, which was prepared in good yield in two reaction steps (see SI Scheme S1 for detail). With this ligand in hand, the regioselectivity could be improved to 97/3, albeit the reactivity was affected negatively (30% yield of 2a). Finally, to our delight, the 1-(2,6diisopropylphenyl)-1H-imidazole-based phosphine L11,[7] gave an acceptable yield (56%) and excellent 98/2 regioselectivity.



**Figure 1.** Pd-catalyzed cyclocarbonylation of 1-ethynyl-1-cyclohexanol **1a**: Influence of phosphine ligands. Reaction conditions: **1a** (0.5 mmol), [Pd( $\pi$ -cinnamyl)Cl<sub>2</sub>] (1.0 mol%), bidentate ligand (2.0 mol%) or monodentate ligand (4.0 mol%), and CO (40 bar) in THF (2.0 mL) at 100 °C for 20 h. The ratio of **2a/3a** and the yield were determined by GC using mesitylene as the internal standard.

Next, we tried to improve the reactivity further in the presence of **L11** as the ligand of choice (see SI; Table S2-S6). Variation of critical parameters (solvent, pre-catalyst, temperature, pressure) revealed the following optimal conditions: 1 mol% Pd(MeCN)<sub>2</sub>Cl<sub>2</sub> as catalyst precursor, metal/ligand ratio of 1: 6, methyl *tert*-butyl ether (MTBE) as solvent, 100 °C, 40 bar CO. Hence, the desired  $\beta$ -lactone **2a** was obtained in excellent yield (98%) and regioselectivity (98/2).

With optimized reaction conditions established, a range of easily available and structurally diverse propargylic alcohols were examined. It is worthy to note that all of the desired  $\alpha$ -methylene- $\beta$ -lactones were obtained in isolated yield with excellent regioselectivity and diastereoselectivity. Notably, the latter is likely to be controlled by the substrates. The alkynols with different substituents (dimethyl, phenyl, ketal) on the 3- or 4-position of cyclohexyl group were transformed into the corresponding products 2a-2e in yields of 60-97% and excellent selectivity. This protocol can be readily scaled-up to carbonylation of 1.0 gram of 1a. This reaction proceeded smoothly, providing 2a in 92% yield. Substrates 1f-1i containing heteroatoms (oxygen, sulfur, nitrogen) proved to be viable too and gave the corresponding  $\beta$ lactones 2f-2i in 81-98% yields with >20/1 selectivity. Fivemembered ring substrates such as 1j can be also applied successfully in this carbonylation reaction (83% yield of 2j). In

### COMMUNICATION

case of the carbonylation of the 1-ethynylcyclododecan-1-ol 1k, the use of L10 instead of L11 led to a higher yield of 2k (88%). Noncyclic alkynols 11-1r bearing different alkyl and benzyl groups underwent lactonization smoothly and gave the desired products 2I-2r in 38-90% yields. By increasing the catalyst loading (5 mol% Pd(MeCN)<sub>2</sub>Cl<sub>2</sub>, 30 mol% L11), the corresponding products 2m and 2q were isolated in 90% and 49% yield, respectively, with >20/1 regioselectivity. When  $\alpha$ -monoalkyl-substituted propynyl alcohol 1s was subjected to the optimized conditions, 2s was obtained, albeit in a lower isolated yield. On the other hand, starting from  $\alpha$ -monoaryl-substituted alkynol **1t**, carbonylation proceeded at increased catalyst loading to give 2t in 58% yield. Interestingly, dicarbonylated product 2u was obtained directly in 56% isolated yield by carbonylation of dialkynol 1u. It should be noted that the synthesis of such multiply  $\beta$ -lactone is not an easy task. In fact, to our knowledge no such transformation has been described yet.





[a] Unless otherwise noted, all reactions were performed in MTBE (2.0 mL) at 100 °C for 20 h in the presence of 1 (0.5 mmol), Pd(MeCN)<sub>2</sub>Cl<sub>2</sub> (1.3 mg, 0.005 mmol), L11 (11.2 mg, 0.03 mmol), and CO (40 bar). Isolated yields were given before the parentheses. The NMR yields (values within the parentheses), regioselectivity of 2/3 and diastereoselectivity of 2 were determined by crude <sup>1</sup>H NMR analyses using dibromomethane as the internal standard. [b] 1a (1 g, 8 mmol), Pd(MeCN)<sub>2</sub>Cl<sub>2</sub> (21 mg, 0.08 mmol), L11 (178 mg, 0.48 mmol), MTBE (30 mL). [c] 0.1 mmol of 1 was used.

The importance of this novel methodology is showcased by the late-stage modification of biologically active and natural products,<sup>[8]</sup> which provides easy access to diverse  $\alpha$ -methylene- $\beta$ -lactones, highlighting the substrate scope of this protocol and its potential utility in organic synthesis. Due to the poor solubility of some of the complex substrates, typically 5 mol% of palladium catalyst was applied. Under otherwise similar conditions, in all cases the reactions proceeded well with excellent regio- and diastereoselectivities. More specifically, tropinone-derived propargylic alcohol 1v delivered the desired product 2v, with good efficiency (81% yield). Pentoxifyllin, a drug with anti-inflammatory properties, was transformed to the corresponding product 2w smoothly (85% yield). Recently, much attention has been paid to steroidal containing spiro-heterocycles for their characteristic physiological activities.<sup>[9]</sup> Thus, we investigated reactions of pharmaceutically relevant steroidal alkynols: ethynyl estradiol, ethisterone, levonorgestrel, and lynestrenol, which are used for contraception and gynecological disorders. All these compounds participated efficiently in this transformation to provide the carbonylative products 2x-2z, 2aa in high yields (85-93%). Notably, the molecular structure of ethynyl estradiol derivative 2x was unambiguously confirmed by X-ray structure analysis.<sup>[10]</sup> Similarly,  $\alpha$ -methylene- $\beta$ -lactones **2ab-2af** derived from other steroid hormones such as dihydrocholesterol, stanolone and epiandrosterone, were obtained in 41-87% yields with excellent selectivity. Moreover, homopropargylic alcohols also proved to be suitable substrates and afforded the corresponding 5-membered products with excellent regioselectivity (see SI, Scheme S2).



[a] Unless otherwise noted, all reactions were performed in MTBE (2.0 mL) at 100 °C for 20 h in the presence of 1 (0.1 mmol), Pd(MeCN)<sub>2</sub>Cl<sub>2</sub> (1.3 mg, 0.005 mmol), L11 (11.2 mg, 0.03 mmol), and CO (40 bar). Isolated yields were given before the parentheses. The NMR yields (values within the parentheses), regioselectivity of 2/3 and diastereoselectivity of 2 were determined by crude <sup>1</sup>H NMR analyses using dibromomethane as the internal standard. [b] 0.5 mmol of 1 was used.

It should be noted that more than 80% of the here described  $\alpha$ -methylene- $\beta$ -lactones are prepared for the first time. This clearly demonstrates the synthetic value of this novel

### COMMUNICATION

methodology. We assumed that our new products can be conveniently used as interesting basic building blocks.<sup>[3]</sup> Thus, to showcase their utility, selected follow-up transformations were conducted by using 2b as the starting material (Scheme 3). To illustrate the possibility to prepare functionalized acrylic acid derivatives,  $\alpha$ -methylene- $\beta$ -lactone **2b** readily underwent ring opening with benzylamine in the presence of Pd(OAc)<sub>2</sub>, affording  $\beta$ -hydroxy amide **4** in 61% yield. Furthermore,  $\alpha$ -methylene- $\beta$ lactones provide an easy and efficient entry into  $\alpha$ -alkylidene- $\beta$ lactones applying cross metathesis in the presence of Grubbs II catalyst. Indeed, a good yield of 5 was obtained with high Zselectivity (stereochemistry determined by NOESY study, see SI for detail). Particularly, this route allows for the efficient preparation of focused libraries of  $\beta$ -lactones, which have found use as biological research probes and therapeutic agents.<sup>[11]</sup> Addition of carbon- or hetero-nucleophiles gives access to  $\alpha$ alkylated *β*-lactones. Exemplarily, the Rh-catalyzed conjugate addition of phenylboronic acid to 2b provided 6 in 40% yield and the treatment of **2b** with thiophenol and triethylamine provided  $\alpha$ -(thiomethyl)- $\beta$ -lactone 7 via a nucleophilic conjugate addition. Finally, four-membered thiolactones can be made in a facile manner by employing Lawesson's reagent. The synthesis of  $\alpha$ methylene- $\beta$ -S-thiolactone 8 illustrated the diverse possibilities for the construction of novel sulfur heterocycles.



**Scheme 3.** Further synthetic transformations of **2b**. Reagents and conditions: (i) Pd(OAc)<sub>2</sub>, PPh<sub>3</sub>, BnNH<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt for 16 h. (ii) Grubbs' 2nd generation, 1octene, CH<sub>2</sub>Cl<sub>2</sub>, reflux for 10 h. (iii) [Rh(cod)Cl]<sub>2</sub>, KOH/H<sub>2</sub>O, PhB(OH)<sub>2</sub>, dioxane, 60 °C for 1h. (iv) PhSH, Et<sub>3</sub>N, MeOH, rt, 24 h. (v) Lawesson's Reagent, PhCH<sub>3</sub>, reflux for 6 h.

Regarding the mechanism of this novel carbonylation reaction, in principle two main pathways are possible (Scheme 4, a): 1) Initially, the active palladium hydride species I could be generated in situ by the combination of palladium precursor with phosphine ligands.<sup>[12]</sup> in which an excess of phosphine ligand (L11) is needed to reduce the initial Pd(II) precursor. After coordination of the alkyne to this complex followed by migratory insertion into the Pd-H bond, the corresponding alkenyl-Pd complex II should be obtained, which is transformed into the corresponding acyl complex III via CO coordination and insertion. Finally, intramolecular nucleophilic attack of hydroxyl on the acyl carbonyl leads to the formation of the desired lactone and regeneration of the [Pd-H]<sup>+</sup> species. Alternatively, the Pd(II) precursor is reduced in situ to a Pd(0) species (probably by an excess amount of phosphine ligands). The Pd(0) species undergoes insertion into the oxygen-hydrogen bond of alkynol affording the corresponding alkoxypalladium complex. Then, insertion of CO into palladium-oxygen bond would give the Pd acyl species. Intramolecular addition of the palladium hydride to the triple bond would form metallacycle complex, which leads to the formation of the desired lactone and regenerates the catalyst (see SI, Scheme S4). In order to differentiate between these two possibilities, control experiments were performed. As shown in Scheme 4b (entry 1), the carbonylation of propargylic alcohol 1a was also carried out with a Pd(0) pre-catalyst. However, in the presence of Pd(dba)<sub>2</sub> under the standard reaction conditions, no conversion was observed. In contrast, using Pd(dba)<sub>2</sub> in the presence of 2 mol% of hydrochloric acid gave the desired product 2a in 98% yield (Scheme 4b, entry 2). These experiments provided clear evidence for a mechanism involving catalytically active palladium hydride species. Although the detailed reaction mechanism of the cyclocarbonylation of propargylic alcohols remains to be further elucidated, based on our previous studies on alkoxycarbonylations<sup>[6b,13]</sup> as well as mechanistic studies by Cole-Hamilton, Drent and Sparkes<sup>[14]</sup>, it is most likely that this reaction goes through the Pd hydride mechanism shown in Scheme 4 (a).[15]



Scheme 4. (a) Proposed mechanism. (b) Control experiments.

HCI (2 mol%)

98

2

In summary, we developed the first catalyst system for a general and selective cyclocarbonylation of alkynols to produce synthetically useful  $\alpha$ -methylene- $\beta$ -lactones. By applying a distinctive ligand, a wide range of propargylic alcohols was efficiently transformed into the corresponding  $\alpha$ -methylene- $\beta$ lactones in good yields (up to 98%) with high regio- and diastereoselectivity (>20/1). The applicability of this methodology is specifically highlighted by the functionalization of biologically active and natural molecules. Combining this novel procedure with established functionalizations allows for an efficient preparation of privileged  $\beta$ -lactone scaffolds. This efficient procedure features the following advantages: high atom economy, additive free reaction conditions, availability of substrates and obtained excellent selectivities. It complements the current methodologies for carbonylations in organic synthesis as shown by the synthesis of 30 products; the vast majority of them are new.

### COMMUNICATION

#### Acknowledgements

This work is supported by the State of Mecklenburg-Vorpommern and the BMBF, Germany. We thank the analytical team of LIKAT.

# **Keywords**: alkynol • palladium • phosphine • carbonylation • catalysis • $\alpha$ -methylene- $\beta$ -lactone

- a) F. Bohlmann, C. Zdero, R. M. King, H. Robinson, *Phytochemistry* 1981, 20, 1069-1075; b) F. Bohlmann, C. Zdero, R. M. King, H. Robinson, *Phytochemistry* 1983, 22, 2860-2862; c) F. Bohlmann, A. H. K. Paul, *Tetrahedron Lett.* 1984, 25, 1697-1700; d) A.N. de Gutierrez, A. Bardón, C. A. N. Catalán, T. B. Gedris, W. Herz, *Biochem. Syst. Ecol.* 2001, 29, 633-647; e) M. J. Dai, M. Seleem, X. Yin, US 10087190 B1, 2018; f) S. Kamat, K. Camara, W. H. Parsons, D.-H. Chen, M. M. Dix, T. D. Bird, A. R. Howell, B. F. Cravatt, *Nat. Chem. Biol.* 2015, *11*, 164-171; g) W. H. Parsons, M. J. Kolar, S. S. Kamat, A. B Cognetta III, J. J. Hulce, E. Saez, B. B. Kahn, A. Saghatelian, B. F. Cravatt, *Nat. Chem. Biol.* 2016. *12*, 367-372; h) B. F. Cravatt, S. S. Kamat, W. H. Parsons, A. R. Howell, K. Camara, A. Anderson, WO 2016/069542 A2, 2016.
- a) Y.-C. Xu, H. Zhou, X.-Y. Sun, W.-M. Ren, X.-B. Lu, *Macromolecules* 2016, *49*, 5782-5787; b) Y.-C. Xu, W.-M. Ren, H. Zhou, G.-G. Gu, X.-B. Lu, *Macromolecules* 2017, *50*, 3131-3142.
- a) W. Adam, R. Albert, L. Hasemann, V. O. Nava Salgado, B. Nestler, [3] E.-M. Peters, K. Peters, F. Prechtl, H. G. von Schnering, J. Org. Chem. 1991. 56. 5782-5785; b) R. L. Danheiser, Y. M. Choi, M. Menichincheri, E. J. Stoner, J. Org. Chem. 1993, 58, 322-327; c) I. Martinez, A. E. Andrews, J. D. Emch, A. J. Ndakala, J. Wang, A. R. Howell, Org. Lett. 2003, 5, 399-402; d) R. Raju, A. R. Howell, Org. Lett. 2006, 8, 2139-2141; e) W. Adam, L. Hasemann, Tetrahedron Lett. 1991, 32, 7033-7036; f) W. Adam, L. Hasemann, Chem. Ber. 1990, 123, 1449-1451; f) C. A. Malapit, D. R. Caldwell, N. Sassu, S. Milbin, A. R. Howell, Org. Lett. 2017, 19, 1966-1969; g) J. Xia, L. Kong, X. Zhou, G. Zheng, X. Li, Org. Lett. 2017, 19, 5792-15795; h) C. A. Malapit, I. K. Luvaga, D. R. Caldwell, N. K. Schipper, A. R. Howell, Org. Lett. 2017, 19, 4460-4463; i) W. Adam, V. O. Nava Salgado, J. Org. Chem. 1995, 60, 578-584; j) G. Zhu, W. Shi, H. Gao, Z. Zhou, H. Song, W. Yi, Org. Lett. 2019, 21, 4143-4147; k) Y. Xu, L. Zhang, M. Liu, X. Zhang, X. Zhang, X. Fan, Org. Biomol. Chem. 2019, 17, 8706-8710; I) M. Bian, K. Ma, H. Mawjuda, X. Yu, X. Li, H. Gao, Z. Zhou, W. Yi, Org. Biomol. Chem. 2019, 17, 6114-6118; m) W. Adam, V. O. Nava Salgado, E.-M. Peters, K. Peters, H. G. von Schnering, Chem. Ber. 1993, 126, 1481-1486; n) A. Pommier, J.-M. Pons, Synthesis 1993, 441-459; o) Y. Wang, R. L. Tennyson, D. Romo, Heterocycles 2004, 64, 605-658.
- [4] a) G. B. Payne, J. Org. Chem. 1966, 31, 718-721; b) G. J. Baxter, R. F. C. Brown, F. W. Eastwood, B. M. Gatehouse, M. C. Nesbit, Aust. J. Chem. 1978, 31, 1757-1767; c) A. P. Masters, T. S. Sorensen, Tetrahedron Lett. 1989, 30, 5869-5872; d) C. Zhang, X. Lu, Synthesis 1996, 586-588; e) A. Bartels, J. Liebscher, Synthesis 1998, 1645-1654; e) G. Roso-Levi, I. Amer, J. Mol. Catal. A: Chem. 1996, 106, 51-56; f) E. M. Campi, K. Dyall, G. Fallon, W. R. Jackson, P. Perlmutter, A. J. Smallridge, Synthesis 1990, 855-856; g) W. Adam, R. Albert, N. Dachs Grau, L. Hasemann, B. Nestler, E.-M. Peters, K. Peters, F. Prechtl, H. G. von Schnering, J. Org. Chem. 1991, 56, 5778-5781; h) W. Adam, L. Hasemann, F. Prechtl, Angew. Chem. Int. Ed. 1988, 27, 1536-1537.
- [5] a) I. Matsuda, A. Ogiso, S. Sato, J. Am. Chem. Soc. 1990, 112, 6120-6121; b) L. A. Aronica, C. Mazzoni, A. M. Caporusso, Tetrahedron 2010, 66, 265-273; c) B. Gabriele, M. Costa, G. Salerno, G. P. Chiusoli, J. Chem. Soc., Chem. Commun. 1994, 1429-1430; d) B. Gabriele, G. Salerno, F. D. Pascali, M. Costa, G. P. Chiusoli, J. Chem. Soc., Perkin Trans 1 1997, 147-154; e) S. Ma, B. Wu, S. Zhao, Org. Lett. 2003, 5, 4429-4432; f) S. Ma, B. Wu, X. Jiang, S. Zhao, J. Org. Chem. 2005, 70, 2568-2575; g) C. S. Consorti, G. Ebeling, J. Dupont, Tetrahedron Lett. 2002, 43, 753-755.
- [6] a) K. Dong, R. Sang, R. Franke, A. Spannenberg, H. Neumann, R. Jackstell, M. Beller, *Angew. Chem. Int. Ed.* 2017, 56, 5267-5271; b) J. Liu, K. Dong, R. Franke, H. Neumann, R. Jackstell, M. Beller, *J. Am.*

Chem. Soc. 2018. 140. 10282-10288; c) J. Liu. H. Li. R. Dühren, J. Liu. A. Spannenberg, R. Franke, R. Jackstell, M. Beller Angew. Chem. Int. Ed. 2017, 56, 11976-11980; d) L. Wu, I. Fleischer, R. Jackstell, I. Profir, R. Franke, M. Beller, J. Am. Chem. Soc. 2013, 135, 14306-14312; e) J. Yang, J. Liu, H. Neumann, R. Franke, R. Jackstell, M. Beller, Science 2019, 366, 1514-1517; f) K. Dong, X. Fang, S. Gülak, R. Franke, A. Spannenberg, H. Neumann, R. Jackstell, M. Beller, Nat. Commun. 2017, 8, 14117; g) Z. Ren, Y. Lyu, X. Song, Y. Ding, Appl. Catal. A: Gen. 2020, 595, 117488; h) K. Nogi, H. Yorimitsu, Chem. Asian J. 2020, 15, 441-449; i) D. J. Jones, M. Lautens, G. P. McGlacken, Nat. Catal. 2019, 2, 843-851; j) S. Zhao, N. P. Mankad, Catal. Sci. Technol. 2019, 9, 3603-3613; k) R. Mancuso, N. Della Cà, L. Veltri, I. Ziccarelli, B. Gabiele, Catalysts 2019, 9, 610; I) J.-B. Peng, H.-Q. Geng, X.-F. Wu, Chem 2019, 5, 526-552; m) K. Ma, B. S. Martin, X. Yin, M. Dai, Nat. Prod. Rep. 2019, 36, 174-219; n) H. Matsubara, T. Kawamoto, T. Fukuyama, I. Ryu, Acc. Chem. Res. 2019, 51, 2023-2035.

- a) T. Schareina, R. Jackstell, T. Schulz, A. Zapf, A. Cott, M. Gotta, M. Beller, Adv. Synth. Catal. 2009, 351, 643–648; b) T. Schulz, C. Torborg, S. Enthaler, B. Schäffner, A. Dumrath, A. Spannenberg, H. Neumann, A. Börner, M. Beller, Chem. Eur. J. 2009, 15, 4528-4533; c) T. Schulz, C. Torborg, B. Schffner, J. Huang, A. Zapf, R. Kadyrov, A. Börner, M. Beller, Angew. Chem. Int. Ed. 2009, 48, 918–921; d) C. Torborg, J. Huang, T. Schulz, B. Schffner, A. Zapf, A. Spannenberg, A. Börner, M. Beller, Chem. Eur. J. 2009, 15, 1329–1336.
- [8] a) M. Moir, J. J. Danon, T. A. Reekie, M. Kassiou, *Exp. Opin. Drug Discov.* 2019, 14, 1137-1150; b) J. Boström, D. G. Brown, R.J. Young, G. M. Keserü, *Nat. Rev. Drug Discovery* 2018, *17*, 709-727.
- a) K. P. Olive, M. A. Jacobetz, C. J. Davidson, A. Gopinathan, D. McIntyre, D. Honess, B. Madhu, M. A. Goldgraben, M. E. Caldwell, D. Allard, K. K. Frese, G. DeNicola, C. Feig, C. Combs, S. P. Winter, H. Ireland-Zecchini, S. Reichelt, W. J. Howat, A. Chang, M. Dhara, L. Wang, F. Rckert, R. Grtzmann, C. Pilarsky, K. Izeradjene, S. R. Hingorani, P. Huang, S. E. Davies, W. Plunkett, M. Egorin, R. H. Hruban, N. Whitebread, K. McGovern, J. Adams, C. Iacobuzio-Donahue, J. Griffiths, D. A. Tuveson, *Science* 2009, *324*, 1457-1461; b) T. Sakata, J. K. Chen, *Chem. Soc. Rev.* 2011, *40*, 4318-4331; c) S. Peukert, K. Miller-Moslin, *ChemMedChem* 2010, *5*, 500-512; d) Z. Zhang, V. Baubet, C. Ventocilla, C. Xiang, N. Dahmane, J. D. Winkler, *Org. Lett.* 2011, *13*, 4786-4789; e) P. Heretsch, A. Bttner, L. Tzagkaroulaki, S. Zahn, B. Kirchner, A. Giannis, *Chem. Commun.* 2011, *47*, 7362-7364.
- [10] CCDC 1994467 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service www.ccdc.cam.ac.uk/structures.
- [11] a) R. Flavin, S. Peluso, P. L. Nguyen, M. Loda, *Future Oncol.* 2010, 6, 551-562; b) A. F. Kluge, R. C. Petter, *Curr. Opin. Chem. Biol.* 2010, 14, 421-427; c) R. D. Richardson, G. Ma, Y. Oyola, M. Zancanella, L. M. Knowles, P. Cieplak, D. Romo, J. W. Smith, *J. Med. Chem.* 2008, 51, 5285-5296; d) T. Böttcher, S. A. Sieber, *Angew. Chem. Int. Ed.* 2008, 47, 4600-4603.
- [12] H. Liu, N. Yan and P. J. Dyson, Chem. Commun. 2014, 50, 7848–7851.
- [13] a) J. Liu, J. Yang, C. Schneider, R. Franke, R. Jackstell, M. Beller, Angew. Chem. Int. Ed. DOI: 10.1002/anie.201915386; b) J. Liu, Q. Liu, R. Franke, R. Jackstell, M. Beller, J. Am. Chem. Soc. 2015, 137, 8556–8563; c) H. Li, X. Fang, R. Jackstell, H. Neumann, M. Beller, Chem. Commun. 2016, 52, 7142-7145; d) H. Li, K. Dong, H. Jiao, H. Neumann, R. Jackstell, M. Beller, Nat. Chem. 2016, 8, 1159-1166; e) X. Fang, H. Li, R. Jackstell, M. Beller, J. Am. Chem. Soc. 2014, 136, 16039-16043; f) H. Li, K. Dong, H. Neumann, M. Beller, Angew. Chem. Int. Ed. 2015, 54, 10239-10243; g) Q. Liu, L. Wu, H. Jiao, X. Fang, R. Jackstell, M. Beller, Angew. Chem. Int. Ed. 2013, 52, 8064-8068.
- [14] a) L. Crawford, D. J. Cole-Hamilton, E. Drent, M. Bühl, *Chem. Eur. J.* **2014**, *20*, 13923-13926; b) L. Crawford, D. J. Cole-Hamilton, M. Bühl, *Organometallics* **2015**, *34*, 438-449; c) A. Shuttleworth, A. M. MilesHobbs, P. G. Pringle, H. A. Sparkes, *Dalton Trans.* **2017**, *46*, 125-137.
- [15] a) M. Beller, A. Krotz, W. Baumann, Adv. Synth. Catal. 2002, 344, 517-524; b) V. V. Grushin, Chem. Rev. 1996, 96, 2011-2034; c) C. Blanco, C. Godard, E. Zangrando, A. Ruiz, C. Claver, Dalton Trans. 2012, 41, 6980-

### COMMUNICATION

6991; d) P. Roesle, L. Caporaso, M. Schnitte, V. Goldbach, L. Cavallo, S. Mecking, *J. Am. Chem. Soc.* 2014, *136*, 16871-16881; e) P. Roesle, C. J. Dürr, H. M. Möller, L. Cavallo, L. Caporaso, S. Mecking, *J. Am. Chem. Soc.* 2012, 134, 17696-17703; f) I. del Río, C. Claver, P. W. N. M. van Leeuwen, *Eur. J. Inorg. Chem.* 2001, 2719-2738.



### COMMUNICATION

#### COMMUNICATION



A special catalyst for tensed problems: The advanced Pd/L11 system enables highly regioselective carbonylation of alkynols for synthesis of a wide range of 4-membered  $\alpha$ -methylene- $\beta$ -lactones.

Yao Ge, Fei Ye, Jiawang Liu, Ji Yang, Anke Spannenberg, Haijun Jiao, Ralf Jackstell, and Matthias Beller\*

Page No. – Page No.

Ligand-controlled Palladiumcatalyzed Carbonylation of Alkynols: Highly Selective Synthesis of  $\alpha$ -Methylene- $\beta$ -lactones