

A novel synthetic approach towards *N*-phenylsuccinimides from γ -lactam-2-carboxylic acid derivatives by reaction with CAN–NaBrO₃

Gopa Barman, Mahuya Roy, Jayanta. K. Ray*

Department of Chemistry, Indian Institute of Technology, Kharagpur 721302, India

Received 4 October 2007; revised 4 December 2007; accepted 12 December 2007

Available online 14 January 2008

Abstract

N-Arylsuccinimides have been synthesized by decarboxylative oxidation of *N*-aryl- γ -lactam-2-carboxylic acids with the dual oxidant CAN/NaBrO₃ in refluxing acetonitrile–water.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: Dual oxidant; Decarboxylative oxidation; *N*-Arylsuccinimides; Radical cation


1. Introduction

Imide derivatives have numerous applications in biology, synthetic and polymer chemistry.^{1,2} Cyclic *N*-arylsuccinimides have high antifungal activity, especially against *Sclerotinia sclerotiorum* and *Botrytis cinerea*.³ However, the availability of simple routes for their synthesis is limited.

We herein report the synthesis of *N*-aryl-substituted succinimides by decarboxylative oxidation of γ -lactam carboxylic acids. Many of the oxidants used for the decarboxylation of carboxylic acids are metal derivatives, such as lead(IV), cobalt(III), silver(II), manganese(III) and thallium(III) salts; chromic acid and non-metallic oxidants have also been reported.⁴

Cerium(IV) compounds, particularly ceric ammonium nitrate [(NH₄)₂Ce(NO₃)₆, CAN], are notable oxidants. Cerium(IV)-promoted oxidations were originally carried out under strongly acidic conditions but recent studies have focused on the development of much milder and more convenient procedures with wider applications.⁵ Ho have found that the dual oxidant system Ce⁴⁺/BrO₃[−] allows the use of cerium(IV) as a catalyst as it is continuously

replenished by the action of the bromate. Using this system, hydroquinones are oxidized to quinones, sulfides to sulfoxides and arylmethanols are transformed into the corresponding carbonyl compounds in good yields.^{6,7} Loudon et al. reported that oxidative decarboxylation of amino acids with [bis(trifluoroacetoxy)iodo]benzene in the presence of pyridine gives the corresponding aldehyde in about 45% yield.⁸ In 1988, Ochiai and Moriaty et al. found that oxidation of L-proline with iodosobenzene (ISB) gave 2-pyrrolidinone under neutral conditions.⁹ The reaction of the γ -lactam-carboxylic acids **1** with CAN–NaBrO₃ in acetonitrile–water (1:1; v/v) at 80 °C furnished exclusively the 1,3-diaryl succinimides **2** in high yields (Scheme 1, Table 1). The starting materials for this study, γ -lactam-2-carboxylic acids **1**, were synthesized by a general method.^{10–13}



Scheme 1. Decarboxylative oxidation of γ -lactam-2-carboxylic acids.

* Corresponding author. Tel.: +91 3222 283326; fax: +91 3222 282252.
E-mail address: jkray@chem.iitkgp.ernet.in (J. K. Ray).


Table 1
Synthesis of *N*-arylsuccinimides **2** from γ -lactam-2-carboxylic acids **1**^a

Substrate <i>N</i> -aryl- γ -lactam-2-carboxylic acid	Product <i>N</i> -arylsuccinimide	Yield (%)
1a R ¹ = R ³ = H, R ² = Cl	2a	87
1b R ¹ = R ³ = H, R ² = F	2b	76
1c R ¹ = R ³ = H, R ² = Br	2c	70
1d R ¹ = R ³ = H, R ² = CH ₃	2d	80
1e R ¹ = R ³ = R ² = H	2e	90
1f R ¹ = Cl, R ² = F, R ³ = H	2f	76

^a Reagents and conditions: All the reactions were carried out with 1 equiv of CAN, and 2 equiv of NaBrO₃ in CH₃CN–H₂O (1:1, v/v) at 80 °C for 5–8 h.

Except for oxalic, malonic and α -hydroxycarboxylic acids, carboxylic acids are resistant to oxidation with cerium(IV). In our case CAN–NaBrO₃ acts as a dual oxidant although the cerium(IV) ion is the actual reagent. Cerium(IV) has a high oxidation potential of +1.61 V compared to the NHE (normal hydrogen electrode),¹⁴ and acts as a single electron oxidant. The oxidation by Ce(IV)/BrO₃[−] occurred via a radical cation in the above reaction. In 1974, Trahanovsky et al. proposed a mechanism for the CAN mediated decarboxylative oxidation of substituted phenyl acetic acids.⁴ However, the mechanism of the reaction is uncertain, and we were unable to isolate any intermediates, although a plausible mechanism may be written as depicted in Scheme 2.¹⁵

The standard electrode potential, $E^0(\text{Ce}^{\text{IV}}/\text{Ce}^{\text{III}})$ is unknown. The formal potential for equal concentrations of cerium(IV) and cerium(III) varies considerably with the nature and concentration of the acidic medium. Cerium(III) is converted to cerium(IV) by the BrO₃[−] ion in the reaction medium, also the BrO₃[−] ion can oxidize the alcohol formed during the course of the reaction.¹⁵ The electronic configurations of Ce^{III} and Ce^{IV} are [Xe]4f¹ and [Xe]4f⁰ where Xe represents the xenon configuration. The stability of the vacant f shell accounts for the ability of cerium to exist in the Ce^{IV} oxidation state.¹⁴



Scheme 2. Mechanistic pathway.

In conclusion, we have developed a novel and simple method for the decarboxylative oxidation of γ -lactam-carboxylic acids to *N*-arylsuccinimides in one step and in good yields. This procedure demonstrates the potential of the CAN–NaBrO₃ reagent system as a decarboxylative oxidizing agent in refluxing acetonitrile–water as solvent. The synthesis allows oxidation under mild conditions using low cost reagents. The experimental simplicity of the reaction opens new opportunities for the use of this reaction in synthetic and industrial processes.

2. Typical experimental procedure

2.1. General procedure for the synthesis of 1,3-diaryl-succinimides **2** from *N*-aryl γ -lactam-2-carboxylic acid **1**

To a flask containing the γ -lactam-2-carboxylic acid **1** (1 mmol) in acetonitrile (10 mL) was added a mixture of ceric ammonium nitrate (1 mmol) and NaBrO₃ (2 mmol) in water (15 mL) and the mixture was stirred for 15 min at room temperature. The reaction mixture was refluxed for 6–8 h (monitored by TLC) and then cooled to room temperature. The solvent was evaporated under reduced pressure and the residue was extracted with CH₂Cl₂. The combined organic layer was washed successively with H₂O, 10% NaHCO₃ solution and then again with more H₂O. After drying the organic layer with Na₂SO₄, the solvent was evaporated under reduced pressure. The product thus obtained was crystallized from an ethyl acetate–petroleum ether mixture.

3. Physical properties and spectral data of representative compounds

3.1. Compound **2a**, 1-(4-chlorophenyl)-3-phenylpyrrolidine-2,5-dione

Light yellow solid; mp 158–162 °C, ¹H NMR (CDCl₃, 200 MHz) δ 2.96 (dd, J = 4.9, 18.5 Hz, 1H), 3.32 (dd, J = 4.5, 8.8 Hz, 1H), 4.16 (dd, J = 4.9, 9.6 Hz, 1H), 7.13–7.54 (m, 9H). ¹³C NMR (CDCl₃, 50 MHz) δ : 36.96, 45.76, 127.24, 127.56, 127.98, 129.15, 129.21, 130.24, 134.26, 134.76, 174.75, 176.33, IR (CHCl₃): ν_{max} 1718, 1685, 1560 cm^{−1}, ESI-MS for C₁₆H₁₂NO₂Cl [M], [M+H⁺] = 286.071 (100%). HRMS (ESI) calcd for C₁₆H₁₂NO₂Cl [M+H⁺]: 286.063, found: 286.060.

3.2. Compound **2b**, 1-(4-fluorophenyl)-3-phenylpyrrolidine-2,5-dione

Light yellow solid, mp 136–140 °C, ¹H NMR (CDCl₃, 200 MHz) δ 3.00 (dd, J = 4.9, 18.6 Hz, 1H), 3.38 (dd, J = 9.5 Hz, 18.7 Hz, 1H), 4.19 (dd, J = 4.9, 9.5 Hz, 1H), 7.12–7.51 (m, 9H). ¹³C NMR (CDCl₃, 50 MHz) δ 37.12, 45.91, 115.98, 116.44, 127.33, 128.14, 128.20, 129.15, 129.31, 132.43, 136.93, 175.04, 176.5, IR (CHCl₃): ν_{max} 1717, 1655, 1509 cm^{−1}, ESI-MS for C₁₆H₁₂NO₂F [M],

$[M+H^+] = 270.088$ (100%). HRMS (ESI) calcd for $C_{16}H_{12}NO_2F$ $[M+H^+]$: 270.085, found: 270.089.

3.3. Compound **2c**, 1-(4-bromophenyl)-3-phenylpyrrolidine-2,5-dione

Light yellow solid, mp 155–158 °C, 1H NMR ($CDCl_3$, 200 MHz) δ 3.01 (dd, $J = 4.9$, 18.6 Hz, 1H), 3.39 (dd, $J = 9.6$, 18.6 Hz, 1H), 4.20 (dd, $J = 4.9$, 9.6 Hz, 1H), 7.22–7.45 (m, 7H), 7.59–7.65 (m, 2H), ^{13}C NMR ($CDCl_3$, 50 MHz) δ 37.07, 45.87, 122.45, 127.30, 127.87, 128.10, 129.79, 130.81, 133.75, 136.78, 174.69, 176.25, IR ($CHCl_3$): ν_{max} 1718, 1655, 1508 cm^{-1} , ESI-MS for $C_{16}H_{12}NO_2Br$ $[M]$, $[M+H^+] = 330.020$ (^{79}Br), 332.020 (^{81}Br).

3.4. Compound **2d**, 1-(p-tolyl)-3-phenylpyrrolidine-2,5-dione

Light yellow solid, mp 145–147 °C (lit. mp 142 °C).¹⁶

3.5. Compound **2e**, 1,3-diphenylpyrrolidine-2,5-dione

Light yellow solid, mp 135–138 °C (lit. mp 140 °C).¹⁶

3.6. Compound **2f**, 1-(3-chloro-4-fluorophenyl)-3-phenylpyrrolidine-2,5-dione

Yellow viscous dense liquid 1H NMR ($CDCl_3$, 200 MHz) δ : 3.00 (dd, $J = 4.9$, 18.7 Hz, 1H), 3.35 (dd, $J = 9.6$, 18.6 Hz, 1H), 4.20 (dd, $J = 4.8$, 9.5 Hz, 1H), 7.20–7.47 (m, 8H), ^{13}C NMR ($CDCl_3$, 50 MHz) δ : 37.02, 45.87, 116.75, 117.20, 121.53, 121.90, 126.29, 126.43,

127.30, 127.74, 128.23, 128.82, 129.04, 129.92, 136.62, 155.26, 160.26, 174.64, 176.20, IR ($CHCl_3$): ν_{max} 1718, 1654, 1500 cm^{-1} , ESI-MS for $C_{16}H_{11}NO_2FCl$ $[M]$, $[M+H^+] = 304.055$.

Acknowledgement

Financial support from DST and CSIR (New Delhi) is gratefully acknowledged.

References and notes

- Reddy, P. Y.; Kondo, S.; Ueno, Y. *J. Org. Chem.* **1997**, 62, 2652.
- Alcaide, B.; Almendros, P.; Cabrero, G.; Ruiz, M. P. *Org. Lett.* **2005**, 7, 3981.
- Takayama, C.; Fujinami, A. *Pestic. Biochem. Physiol.* **1979**, 12, 163.
- Trahanovsky, W. S.; Cramer, J.; Brixius, D. W. *J. Am. Chem. Soc.* **1974**, 96, 1077.
- Molander, A. Gary *Chem. Rev.* **1992**, 92, 29.
- Ho, T. L. *Synth. Commun.* **1979**, 9, 237.
- Ho, T. L. *Synthesis* **1978**, 936.
- Loudon, G. M.; Radhakrishna, A. S.; Almond, M. R.; Blodgett, J. K.; Boutin, R. H. *J. Org. Chem.* **1984**, 49, 4272.
- Ochiai, M.; Inenaga, M.; Nagao, Y.; Moriarty, M. R.; Vaid, K. R.; Duncan, P. M. *Tetrahedron Lett.* **1988**, 29, 6917.
- Kar, G. K.; Ramesh, D.; Chatterjee, B. G.; Ray, J. K. *J. Chem. Res. (M)* **1997**, 568; *J. Chem. Res. (S)* **1997**, 80.
- Ray, J. K.; Kar, G. K.; Roy, B. C.; Brahma, N. K. *Bioorg. Med. Chem.* **1994**, 2, 1417.
- Ray, J. K.; Kar, G. K.; Adhikari, S. D.; Ray, J. K.; Brahma, N. K. *Bioorg. Med. Chem.* **1998**, 6, 2397–2403.
- Haldar, P.; Barman, G.; Ray, J. K. *Tetrahedron* **2007**, 63, 3049.
- Nair, V.; Deepthi, A. *Chem. Rev.* **2007**, 107, 1862.
- Ganin, E.; Amer, I. *Synth. Commun.* **1995**, 25, 3149.
- Mustafa, A.; Asker, W.; Khattab, S.; Zayed, S. M. A. D. *J. Org. Chem.* **1961**, 26, 787.