metal-organic compounds

Acta Crystallographica Section C Crystal Structure Communications

ISSN 0108-2701

Bis[N,N'-(2-chlorobenzylidene)ethylenediamine- $\kappa^2 N,N'$]copper(I) dichloridocuprate(I) acetonitrile solvate

Mohammad Hossein Habibi,^a* Morteza Montazerozohori,^b Kazem Barati,^a Ross W. Harrington^c and William Clegg^c

^aDepartment of Chemistry, University of Isfahan, Isfahan 81746-73441, Iran, ^bChemistry Department, Yasouj University, Yasouj 75914-353, Iran, and ^cSchool of Natural Science (Chemistry), Newcastle University, Newcastle Upon Tyne NE1 7RU, England

Correspondence e-mail: habibi@chem.ui.ac.ir

Received 15 October 2007 Accepted 31 October 2007 Online 24 November 2007

The 1:1 adduct of N,N'-bis(2-chlorobenzylidene)ethylenediamine (cb₂en) with copper(I) chloride proves to be an ionic compound with Cu^I-centred cations and anions, [Cu-(C₁₆H₁₄Cl₂N₂)₂][CuCl₂]·CH₃CN. In the cation, the Cu^I atom has a flattened tetrahedral coordination geometry, with a small bite angle for the chelating ligands, which form a doublehelical arrangement around the metal centre. The anion is almost linear, as expected. The packing of the cations involves intermolecular π - π interactions, which lead to columns of translationally related cations along the shortest unit-cell axis, with anions and solvent molecules in channels between them.

Comment

The coordination chemistry of copper(I) complexes has received increased attention over the last few decades. This is mainly due to the potential application of these complexes in catalytic processes (Fife et al., 1985; Bowker et al., 1988; Bowmarker et al., 2000), photosensitization reactions (Horváth, 1994; Kutal, 1990), light-harvesting studies (Rosi et al., 1999; Dietrich-Buchecker & Sauvage, 1987) and the design of supramolecular arrays (Meghdadi et al., 2002; Foster et al., 2000). Reaction of copper(I) halides, CuX, with nitrogenbased ligands (L) yields $CuXL_n$ adducts. The steric, electronic and conformational effects imparted by the coordinated ligands play an important role in influencing the properties of the resulting metal complexes. A thorough understanding of these effects can serve as the basis for the rational design of complexes with specific and predictable properties. The number of ligands bound to the monovalent Cu^I ion is greatly influenced by both the chemical nature and the geometry of the ligand L, and the particular choice of halogen X (Lange et al., 2000). Although structural reports on $[Cu^{I}L_{4}]^{+}$ complex cations (where L is an N-donor ligand) are numerous (Alcon et al., 2000; Miller & Karpishin, 1999; Panja et al., 2002), there are a limited number of studies of copper(I) complexes with isolated linear dihalidocuprate(I) anions (Amirnasr et al., 2005; Mirkhani et al., 2004). The structures of halidocuprate(I) compounds are exceptionally varied, and the nature of the cation plays an important role in determining the coordination number of Cu^I in halidocuprates and the tendency of the anions to form extended structures (Andersson & Jagner, 1989; Hasselgren et al., 1999). Thus, bulky cations with well screened charges tend to favour the formation of discrete anions in which Cu^I has a low coordination number. It has been recognized that cation properties, such as size, shape and the distribution of positive charge, are of importance for the anion configurations adopted by halidocuprates(I).



In this context, we decided to examine the nature of the complex formed with an unconjugated diimine ligand. The title complex, (I), was prepared by reacting the bidentate ligand N,N'-bis(2-chlorobenzylidene)ethylenediamine (cb₂en) with CuCl. The structure determination of (I) is consistent



Figure 1

The asymmetric unit of (I), showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms have been omitted for clarity, and only one component is shown for each disordered group.

with the stoichiometry of a 1:1 copper(I) chloride–ligand adduct, [CuCl(L)]. The structural connectivity, however, is that of an ionic complex, the asymmetric unit of the structure comprising a bis(ligand)copper(I) cation and a dichlorido-cuprate(I) anion, together with a molecule of acetonitrile solvent, [L_2 Cu]⁺[CuCl_2]⁻·MeCN (Fig. 1).

The anion of (I) has no crystallographic symmetry, and one Cl atom is disordered over two sites. Neither of these gives a completely linear coordination (Table 1). The geometry is in good agreement with that of several other previously reported examples (Kaiser *et al.*, 1974; Engelhardt *et al.*, 1984).

The cation of (I) has a markedly flattened tetrahedral geometry and also lies in a general position with no crystallographic symmetry. The angles at the Cu^I atom in the cation



Figure 2

The stacking of cations of (I) via π - π interactions.



Figure 3

The crystal packing in (I), viewed along the *a* axis, showing the columns of cations, with anions and solvent molecules in channels.

are similar to the corresponding angles for ethylenediamine complexes (Engelhardt et al., 1984), with a small bite angle for the chelating ligand. The two ligands form a double-helical arrangement around the Cu^I atom. Within the cation, the two benzene rings of each ligand are almost parallel to those of the other, with a dihedral angle of 2.73 (16)° for rings C1-C6 and C27-C32, and 0.38 (17)° for rings C11-C16 and C17-C22, and with interplanar spacings of about 3.6 Å in each case. This does not give rise to intramolecular $\pi - \pi$ stacking, since the pairs of rings are substantially displaced laterally, the centroidto-centroid distances being 4.877 (4) and 4.846 (4) Å, respectively. However, the same pairs of rings are similarly close to parallel in adjacent molecules along the short a axis (dihedral angles are the same by symmetry), and here there is intermolecular π - π stacking (Fig. 2), the centroid-to-centroid distances being 3.645 (3) and 3.536 (3) Å, respectively, and the interplanar spacings being 3.532 and 3.523 Å, respectively. This stacking produces columns of cations along the *a* axis, with anions and solvent molecules occupying the channels between them (Fig. 3).

Experimental

The title compound was prepared by the reaction of CuCl and cb_2en (molar ratio 1:1) in an acetonitrile solution at room temperature. The solution was then concentrated under vacuum. Red crystals of (I) were formed by vapour diffusion of diethyl ether into the concentrated solution.

 $V = 3548 (4) \text{ Å}^3$

 $\lambda = 0.8462 \text{ Å}$ $\mu = 1.68 \text{ mm}^{-1}$

T = 120 (2) K

 $R_{\rm int} = 0.000$

Synchrotron radiation

 $0.10 \times 0.10 \times 0.10$ mm

30485 measured reflections

30485 independent reflections

15283 reflections with $I > 2\sigma(I)$

Z = 4

Crystal data

 $[Cu(C_{16}H_{14}Cl_2N_2)_2][CuCl_2] \cdot C_2H_3N$ $M_r = 849.42$ Monoclinic, P_{2_1}/n a = 7.845 (5) Å b = 16.965 (10) Å c = 26.915 (15) Å $\beta = 97.896$ (9)°

Data collection

Bruker APEXII CCD diffractometer Absorption correction: multi-scan (*TWINABS*; Bruker, 2004) $T_{\rm min} = 0.730, T_{\rm max} = 0.850$

Refinement

 $\begin{array}{ll} R[F^2 > 2\sigma(F^2)] = 0.056 & 24 \mbox{ restraints} \\ wR(F^2) = 0.127 & H-atom \mbox{ parameters constrained} \\ S = 0.79 & \Delta\rho_{max} = 0.71 \mbox{ e \AA^{-3}} \\ 30485 \mbox{ reflections} & \Delta\rho_{min} = -0.84 \mbox{ e \AA^{-3}} \\ 456 \mbox{ parameters} \end{array}$

Table 1

Selected geometric parameters (Å, °).

Cu1-N1 Cu1-N2 Cu1-N3 Cu1-N4 Cu2-Cl5	2.024 (3) 2.024 (3) 2.039 (3) 2.014 (3) 2.0862 (14)	Cu2-Cl6 N1-C7 N1-C8 N1-C8 <i>A</i> N2-C9	2.171 (5) 1.277 (4) 1.535 (8) 1.502 (8) 1.534 (8)
N1-Cu1-N2 N1-Cu1-N4 N2-Cu1-N3 N3-Cu1-N4	85.09 (12) 115.87 (13) 115.33 (12) 85.18 (12)	Cl5-Cu2-Cl6 Cu1-N2-C9 Cu1-N2-C10	170.5 (4) 108.7 (4) 133.5 (3)

The small size and weak diffraction of the crystals necessitated the use of synchrotron radiation. The crystal was found to be a nonmerohedral rotation twin about (100), with symmetry-inequivalent reflections overlapping in the diffraction pattern, making it impossible to merge equivalent reflections before the refinement with *SHELXL* HKLF5 style data (*SHELXTL*; Bruker, 2005). The twin law is (1 0 0/ 0 -1 0/-0.8867 0 -1). A number of reflections were rejected in the data processing because of problems with spot shape or masking by the beam-stop. The deposited structure factors are in the format generated by the undocumented LIST 7 instruction of *SHELXTL* (and *SHELXL97*), in which the contributions of the twin components are identified following refinement of the twin fraction; this was 0.5415 (5):0.4585 (5).

All H atoms were placed in idealized positions and constrained to ride on their parent atoms, with C–H distances in the range 0.95– 0.99 Å, with $U_{iso}(H) = 1.2U_{eq}(C)$ [1.5 $U_{eq}(C)$ for methyl H atoms]. One Cl atom of the anion was modelled as disordered over two positions, with occupancy factors 0.55 (2):0.45 (2). Twofold disorder was also found for the CH₂CH₂ bridge of one of the ligands in the cation, with occupancy factors 0.515 (15):0.485 (15); restraints were applied to the displacement parameters of these disordered atoms.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Bruker, 2005); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and local programs.

The authors thank Isfahan University Center of Excellence (Catalyst and Fuel Cells) for financial support of this research, the EPSRC for funding of the National Crystallographic Service, and STFC for access to synchrotron facilities. Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK3174). Services for accessing these data are described at the back of the journal.

References

- Alcon, M. J., Puebla, E. G., Iglesias, M., Monge, M. A. & Sanchez, F. (2000). *Inorg. Chim. Acta*, **306**, 117–123.
- Amirnasr, M., Khalaji, A. D. & Falvello, L. R. (2005). Inorg. Chim. Acta, 359, 713–716.
- Andersson, S. & Jagner, S. (1989). Acta Chem. Scand. Ser. A, 43, 39-41.
- Bowker, M., Hadden, R. A., Houghton, H., Hyland, J. N. K. & Waugh, K. C. (1988). *J. Catal.* **109**, 263–273.
- Bowmarker, G. A., Hanna, J. V., Reid, P. C., Rickard, C. E. F. & White, A. H. (2000). J. Chem. Soc. Dalton Trans. pp. 753–757.
- Bruker (2004). APEX2, SAINT and TWINABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). SHELXTL. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dietrich-Buchecker, C. O. & Sauvage, J.-P. (1987). Chem. Rev. 87, 795-814.
- Engelhardt, L. M., Papasergio, R. I. & White, A. H. (1984). Aust. J. Chem. 37, 2207–2213.
- Fife, D. J., Moore, W. M. & Morse, K. W. (1985). J. Am. Chem. Soc. 107, 7077–7081.
- Foster, C. L., Kilner, C. A., Thornton-Pett, M. & Halcrow, M. A. (2000). Acta Cryst. C56, 319–320.
- Hasselgren, C., Stenhagen, G., Ohrstrom, L. & Jagner, S. (1999). *Inorg. Chim.* Acta, 292, 266–271.
- Horváth, O. (1994). Coord. Chem. Rev. 135, 303-324.
- Kaiser, J., Brewer, G., Schroder, F. A., Taylor, I. F. & Rasmussen, S. E. (1974). J. Chem. Soc. Dalton Trans. pp. 1490–1495.
- Kutal, C. (1990). Coord. Chem. Rev. 99, 213-252.
- Lange, J., Elias, H. & Paulus, H. (2000). Inorg. Chem. 39, 3342-3349.
- Meghdadi, S., Amirnasr, M., Schenk, K. J. & Dehghanpour, S. (2002). *Helv. Chim. Acta*, **85**, 807–812.
- Miller, M. T. & Karpishin, T. B. (1999). Inorg. Chem. 38, 5246-5255.
- Mirkhani, V., Harkema, S. & Kia, R. (2004). Acta Cryst. C60, m343-m344.
- Panja, S., Chowdhury, S., Drew, M. G. B. & Datta, D. (2002). Inorg. Chem. Commun. 5, 304–307.
- Rosi, M., Sgamellotti, A., Franceschi, F. & Floriani, C. (1999). *Inorg. Chem.* 38, 1520–1525.