### Accepted Manuscript

2,4,5-triaryl imidazole probes for the selective chromo-fluorogenic detection of Cu(II). Prospective use of the Cu(II) complexes for the optical recognition of biothiols

Hazem Essam Okda, Sameh El Sayed, Ismael Otri, R. Cristina M. Ferreira, Susana P.G. Costa, M. Manuela M. Raposo, Ramón Martínez-Má ñez, Félix Sancenón

 PII:
 S0277-5387(19)30393-6

 DOI:
 https://doi.org/10.1016/j.poly.2019.05.055

 Reference:
 POLY 13980

To appear in: Polyhedron

Received Date:16 April 2019Revised Date:28 May 2019Accepted Date:28 May 2019



Please cite this article as: H. Essam Okda, S. El Sayed, I. Otri, R.C.M. Ferreira, S.P.G. Costa, M.M.M. Raposo, R. Martínez-Má ñez, F. Sancenón, 2,4,5-triaryl imidazole probes for the selective chromo-fluorogenic detection of Cu(II). Prospective use of the Cu(II) complexes for the optical recognition of biothiols, *Polyhedron* (2019), doi: https://doi.org/10.1016/j.poly.2019.05.055

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# 2,4,5-triaryl imidazole probes for the selective chromofluorogenic detection of Cu(II). Prospective use of the Cu(II) complexes for the optical recognition of biothiols

Hazem Essam Okda,<sup>a,b,c</sup> Sameh El Sayed,<sup>a,b,c</sup> Ismael Otri,<sup>a,b,c</sup> R. Cristina M. Ferreira,<sup>d</sup> Susana P. G. Costa,<sup>d</sup> M. Manuela M. Raposo,<sup>\*d</sup> Ramón Martínez-Máñez,<sup>\*a,b,c</sup> and Félix Sancenón<sup>a,b,c</sup>

<sup>a</sup> Instituto Interuniversitario de Investigación de Reconocimiento Molecular y Desarrollo Tecnológico (IDM), Universitat Politècnica de València, Universitat de València. Spain. E-mail: rmaez@qim.upv.es

<sup>b</sup> Departamento de Química, Universitat Politècnica de València. Camino de Vera s/n, 46022, València, Spain.

<sup>c</sup> CIBER de Bioingeniería, Biomateriales y Nanomedicina (CIBER-BBN).

<sup>d</sup> Centro de Química, Universidade do Minho, Campus de Gualtar, 4710-057, Braga, Portugal. E-mail: mfox@quimica.uminho.pt

### Abstract

The sensing behaviour toward metal cations and biothiols of two 2,4,5-triarylimidazole probes (**3a** and **3b**) is tested in acetonitrile and in acetonitrile-water. In acetonitrile the two probes present charge-transfer absorption bands in the 320-350 nm interval. Among all cations tested only Cu(II) is able to induce bathochromic shifts of the absorption band in the two probes, which is reflected in marked colour changes. Colour modulations are ascribed to the formation of 1:1 Cu(II)-probe complexes in which the cation interacts with the imidazole acceptor heterocycle. Besides, the two probes present intense emission bands (at 404 and 437 nm for **3a** and **3b** respectively) in acetonitrile that are quenched selectively by Cu(II). Probe **3a** is soluble in acetonitrile-water 1:1 (v/v) and Cu(II) also induces bathochromic shifts of the absorption bands. Moreover, the emission bands of probe **3a** in this mixed aqueous solutions is quenched in the

presence of Cu(II). The potential use of the 1:1 complex formed between **3a** and Cu(II) for the chromo-fluorogenic detection of biothiols (GSH, Cys and Hcy) in aqueous environments is also tested. At this respect, addition of GSH, Cys and Hcy to acetonitrile-water 1:1 v/v solutions of **3a**-Cu(II) complex induces a hypsochromic shift of the visible band (reflected in a bleaching of the solutions) with a marked emission increase at 470 nm.

#### Keywords

Imidazole-based probes; Cu(II) detection; biothiols recognition; Cu(II) imaging GSH imaging

#### 1. Introduction

In the last years, the development of chromo-fluorogenic chemosensors for transition metal cations has attracted the attention of researchers around the world.[1] These chromo-fluorogenic chemosensors are generally formed by two subunits, namely the binding site and the signalling group, that can be covalently linked or forming a supramolecular assembly.[2] In the first case, interaction of transition metal cations with the binding sites induce rearrangements in the  $\pi$ -conjugated system of the signalling unit which are reflected in colour and/or emission changes.[3] In the second approach, interaction of the transition metal cation with the binding site induced the displacement of the signalling unit from the initial complex to the solution.[4]

Among transition metal cations, Cu(II) is one of the most abundant essential element in the human body and plays vital roles in several physiological processes. For instance, it has been reported that Cu(II) stimulates the proliferation of endothelial cells and is necessary for the secretion of several angiogenic factors by tumour cells.[5,6] Aside from its biological and environmental importance, copper is widely used in metallurgical, pharmaceutical and agrochemical industries.[7] As a result of the extensive applications of Cu(II) in life science and industry, it has become one of the first hazard environmental pollutants.[8] Despite the important role played by Cu(II) in several biological processes, abnormal levels of this cation can cause serious health problems on humans due to its ability to displace other vital metal ions in some enzyme-catalysed reactions.[9] In addition, high concentrations of Cu(II) in cells was documented to cause toxicity and different neurodegenerative diseases such as Menkes, Wilson's and Alzheimer.[10] Therefore, simple and rapid sensing tools to monitor Cu(II) levels in

biological and environmental media are of importance. Besides, the World Health Organization (WHO) has recommended the maximum allowable level of Cu(II) in drinking water at 2.0 ppm (~30 µM).[11,12]

Currently, several complex analytical techniques are used to detect metal ions such as electrochemical measurements, atomic absorption spectrometry and inductively coupled plasma mass spectroscopy.[13,14] However, these techniques are time-consuming, needs sample pre-treatment, used expensive equipment's, required trained personnel and cannot be used *in situ*.

On the other hand, biothiols (such as the amino acids cysteine (Cys) and homocysteine (Hcy) and the tripeptide glutathione (GSH)) play vital roles in cellular processes related with the damage of cellular components by reactive oxygen species (ROS).[15,16] Levels of biothiols in cells are controlled by the equilibrium between thiols and disulfides. Besides, the relative levels of biothiols can be used as biomarkers for aging, neurodegenerative pathologies and many other diseases such as cancer, AIDS and cystic fibrosis among others.[17,18] Thus, taking into account the above mentioned facts, determination of biothiols especially in biological samples is an important issue. Actually, different analytical methods are available to determination of biothiol levels including ca. HPLC, capillary electrophoresis, and mass spectrometry.[19-21] As an alternative to these classical techniques, in the past two decades, a number of chromofluorogenic probes for biothiol detection have been reported.[22-24] Most of the described examples are based on the chemodosimeter approach and are synthesized taking into account the nucleophilic character of thiol moieties in these biomolecules.[25-31] Also, the displacement approach has been extensively used to design biothiol selective chromo-fluorogenic probes. For this purpose, the emission quenching features of transition metal ions such as Cu(II) has been widely used. The basis of these probes is the generation of non-emissive Cu(II) complexes with fluorophores equipped with coordinating subunits. In the presence of biothiols, there is a demetallation process, restoring the full emission of the fluorophore.[32-37] However, many of the described biothiols chemosensors did not work in pure water or in mixed aqueous environments. Besides, the selectivity achieved in most cases is low because probes are unable to distinguish between Cys, Hcy and GSH. In fact, the synthesis and characterization of probes for the selective recognition of individual biothiol in aqueous environments is of interest.



3

Scheme 1. One pot synthesis of imidazole probes 3a and 3b.

Taking into account our interest in the development of chromo-fluorogenic probes for biomolecules[38-45] we report herein the synthesis, characterization and binding studies toward Cu(II) and biothiols (GSH, Hcy and Cys), of two probes containing 2,4,5-trisubstitued imidazole moieties (**3a** and **3b** in Scheme 1). The prepared probes contain electron donor (furan) and electron acceptor (phenanthroline) rings of different strength covalently linked with an imidazole heterocycle. 2,4,5-triaryl(heteroaryl)-imidazole based chromophores have received increasing attention due to their distinctive optical properties, applications in medicinal chemistry and materials sciences, and as nonlinear optical materials (SHG chromophores, two-photon absorbing molecules) and thermally stable luminescent materials for several applications (such as OLEDs and fluorescent probes).

#### 2. Experimental section

**Materials and methods.** All melting points were measured on a Stuart SMP3 melting point apparatus. TLC analyses were carried out on 0.25 mm thick pre-coated silica plates (Merck Fertigplatten Kieselgel  $60F_{254}$ ) and spots were visualised under UV light. Chromatography on silica gel was carried out on Merck Kieselgel (230-240 mesh). IR spectra were determined on a BOMEM MB 104 spectrophotometer using KBr discs. NMR spectra were obtained on a Bruker Avance III 400 at an operating frequency of 400 MHz for <sup>1</sup>H and 100.6 MHz for <sup>13a</sup> using the solvent peak as internal reference at 25 °C. All chemical shifts are given in ppm using <sup>5</sup>H Me<sub>4</sub>Si = 0 ppm as reference. Assignments were supported by spin decoupling-double resonance and bidimensional heteronuclear correlation techniques. UV/Vis titration profiles were carried out with JASCO V-650 spectrophotometer (Easton, MD, USA). Fluorescence measurements were recorded with

a JASCO FP-8500 spectrophotometer. Commercially available reagents 4-(dimethylamino) benzaldehyde (**1**), 1,2-di(furan-2-yl)ethane-1,2-dione (**2a**), 1,10phenanthroline-5,6-dione (**2b**) and ammonium acetate were purchased from Sigma-Aldrich and Acros and used as received. TLC analyses were carried out on 0.25 mm thick pre-coated silica plates (Merck Fertigplatten Kieselgel  $60F_{254}$ ) and spots were visualized under UV light. Chromatography on silica gel was carried out on Merck Kieselgel (230-240 mesh).

Synthesis of 4-(4,5-di(furan-2-yl)-1H-imidazol-2-yl)-*N*,*N*-dimethylbenzenamine (3a). 4-(dimethylamino) benzaldehyde (1, 0.15 g, 1 mmol), 1,2-di(furan-2-yl)ethane-1,2-dione (2a, 0.19 g, 1 mmol) and NH<sub>4</sub>OAc (1.54 g, 20 mmol) were dissolved in glacial acetic acid (5 mL), followed by stirring and heating at reflux for 8 h. The reaction mixture was then cooled to room temperature, ethyl acetate was added (15 mL) and the mixture was washed with water (3 x 10 mL). After drying the organic phase with anhydrous MgSO<sub>4</sub>, the solution was filtered and the solvent was evaporated to dryness. The resulting crude product was purified by column chromatography (silica gel, DCM/MeOH 100:1 v/v); yield (64 mg, 60%) as Grey solid. Mp = 240.2-240.9 °C.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  = 2.95 (s, 6H), 6.55 (s, 1H), 6.63 (br s, 1H), 6.71 (d, *J* = 2.0 Hz, 1H), 6.77 (dd, *J* = 8.8 and 2.0 Hz, 2H), 6.89 (d, *J* = 2.8 Hz, 1H), 7.68 (br s, 1H), 7.78 (br s, 1H), 7.90 (dd, *J* = 8.8 and 2.0 Hz, 2H), 12.47 (s, 1H, NH) ppm. <sup>13</sup>C NMR (100.6 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  = 39.89, 106.56, 107.81, 111.33, 111.80, 117.50, 118.27, 126.68, 129.15, 141.61, 142.22, 144.79, 147.23, 149.59, 150.53 ppm. IR (Nujol): v = 3570, 1665, 1612, 1530, 1494, 1222, 1202, 1169, 1122, 1074, 1014, 986, 943, 911, 885, 815 cm<sup>-1</sup>. HRMS-EI m/z: calcd for C<sub>19</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>+H<sup>+</sup>: 320.1399; measured: 320.1394.

**Synthesis of 4-(1H-imidazo[4,5-f][1,10]phenanthrolin-2-yl)-***N*,*N*-dimethylbenzenamine (3b). 4-(dimethylamino) benzaldehyde (1, 0.15 g, 1 mmol), 1,10-phenanthroline-5,6-dione (2b, 0.2 g, 1 mmol) and NH<sub>4</sub>OAc (1.54 g, 20 mmol) were dissolved in glacial acetic acid (5 mL), followed by stirring and heating at reflux for 8 h. The reaction mixture was then cooled to room temperature, ethyl acetate was added (15 mL) and the mixture was washed with water (3 x 10 mL). After drying the organic phase with anhydrous MgSO<sub>4</sub>, the solution was filtered and the solvent was

evaporated to dryness. The resulting crude product was purified by column chromatography (silica gel, DCM/MeOH 98:2 v/v); yield (8 mg, 7%) as yellow oil.

<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  = 3.02 (s, 6H), 6.90 (d, *J* = 9.2 Hz, 2H), 7.81-7.84 (m, 2H,), 8.11 (dd, *J* = 8.8 and 2.0 Hz, 2H), 8.92 (dd, *J* = 8.0 and 1.6 Hz, 2H), 9.00 (dd, *J* = 4.0 and 2.0 Hz, 2H), 13.40 (s, 1H, NH) ppm. <sup>13</sup>C NMR (100.6 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  = 40.13, 111.97, 117.29, 123.33, 127.45, 127.62, 129.77, 142.94, 147.34, 147.40, 151.21, 151.88 ppm. IR (liquid film): *v* = 3414, 2925, 1657, 1611, 1502, 1438, 1372, 1351, 1195, 1168, 1125, 1067, 1024, 946, 814, 739 cm<sup>-1</sup>. HRMS-EI m/z: calcd for C<sub>21</sub>H<sub>17</sub>N<sub>5</sub>+H<sup>+</sup>: 340.1562; measured: 340.1560.

**UV-visible and emission measurements.** Stock solutions of the cations (i.e., Cu(II), Pb(II), Mg(II), Ge(II), Ca(II), Zn(II), Co(II), Ni(II), Ba(II), Cd(II), Hg(II), Fe(III), In(III), As(III), Al(III), Cr(III), Ga(III), K(I), Li(I) and Na(I) as perchlorate salts) were prepared at  $(1.0 \times 10^{-3} \text{ mol } \text{L}^{-1})$  in acetonitrile. The concentrations of probes used in spectroscopy measurements were ca.  $(5.0 \times 10^{-5} \text{ mol } \text{L}^{-1})$  and  $(1.0 \times 10^{-5} \text{ mol } \text{L}^{-1})$ . We took care that the maximum addition of cations solutions did not exceed 10% of the volume of the receptor to avoid significant changes in the total solution concentration. In the experiments that required the addition of excess of ions (20 equiv.), corrections of the volume and concentration were made. The UV/Vis and fluorometric titrations were carried out at room temperature (25 °C).

### 3. Results and discussion

**Synthesis and characterization of the probes.** Heteroaromatic commercially available diones **2a** and **2b** containing furyl and phenanthroline rings (see Scheme 1) were used as precursors for the synthesis of **3a** and **3b**. The synthesis of probes **3a** and **3b** was previously described.[45,46] The final probes were prepared through a condensation reaction between **2a** and **2b** diones and 4-(*N*,*N*-dimethylamino) benzaldehyde (**1**) in presence of ammonium acetate and glacial acetic acid at reflux under Debus-Radziszewski imidazole synthesis conditions (see also Scheme 1).[47] Probes were synthesized in fair to moderate yields (Table 1) as a consequence of several factors such as the difficulty in the purification step due to the polarity of the compounds as well as their decomposition during the long reaction times under hard conditions (acetic acid reflux). The final probes **3a** and **3b** were characterized by <sup>1</sup>H and <sup>13</sup>C-

NMR, and IR. The data obtained are in full agreement with the proposed formulations (see Experimental section).

Probe	Yield (%)	δH NH (ppm)		
3a	60	12.45		
3b	7	13.40		

Table 1. Yields and <sup>1</sup>H NMR data of imidazole probes 3a and 3b at 400 MHz in DMSO-D6.

One of the most characteristic <sup>1</sup>H-NMR signals of probes **3a** and **3b** is that corresponding to the N-H proton in the imidazole ring (see Table 1). As could be seen, a clear correlation between the electronic nature of the rings (furan and phenanthroline) directly linked with imidazole and the shift of the N-H proton of this heterocycle was observed. At this respect, probe **3b** bearing the electron deficient phenanthroline system exhibited the highest chemical shift for the nitrogen proton of the imidazole ring (13.40 ppm). On the other hand, probe **3a** functionalized with electron rich furan heterocycle exhibit lower chemical shifts for the nitrogen proton of the imidazole ring (12.45 ppm) when compared to **3b**.

**UV/Vis and emission spectroscopic behaviour of probes in the presence of selected metal cations in acetonitrile.** In a first step, UV-visible changes of both probes in acetonitrile were studied. Acetonitrile solutions of **3a** and **3b** (1.0 x 10<sup>-5</sup> mol L<sup>-1</sup>) showed intense absorption bands in the UV zone (321 nm for **3a** and 341 nm for **3b**). The observed absorption bands are ascribed to charge-transfer transitions between the *N*,*N*-dimethylamino donor moiety and imidazole acceptor heterocycle. Besides, the wavelength of the charge-transfer band was affected by the aromatic units directly connected with the central imidazole heterocycle. At this respect, the presence of electron deficient phenanthroline unit in probe **3b** induced the larger charge-transfer character and the associated absorption appeared at the higher wavelength (341 nm). On the other hand, in probe **3a**, the imidazole core was linked to an electron donating heterocycle (furan) and, as a consequence, absorption bands are shifted to lower wavelengths (321 nm).

Besides, both probes were highly emissive in acetonitrile upon excitation at the maximum of the absorbance of the visible band. At this respect, excitation at 321 nm of acetonitrile solutions of probe **3a** ( $1.0 \times 10^{-5}$  mol L<sup>-1</sup>) induced the appearance of a marked

emission band centred at 404 nm. Nearly the same results were obtained for probe **3b** with an intense emission centred at 437 nm. Besides, the quantum yield of the two probes (determined using pyrene as standard, see Supporting Information for details) was 0.41 and 0.31 for **3a** and **3b** respectively.



**Figure 1.** UV/Vis titration profile of probe **3a** in acetonitrile (1.0 x 10<sup>-5</sup> mol L<sup>-1</sup>) upon addition of increasing quantities of Cu(II) cation (0-10 eq.). Inset: Absorbance at 563 nm vs equivalents of Cu(II) added.

Then, UV-visible changes in acetonitrile solutions of both probes (1.0 x 10<sup>-5</sup> mol L<sup>-1</sup>) in the presence of increasing amounts (from 0.1 to 10 eq.) of selected metal cations (Cu(II), Pb(II), Mg(II), Ge(II), Ca(II), Zn(II), Co(II), Ni(II), Ba(II), Cd(II), Hg(II), Fe(III), In(III), As(III), Al(III), Cr(III), Ga(III), K(I), Li(I) and Na(I)) were tested. The two probes showed nearly the same behaviour and in all cases only Cu(II) induced the progressive appearance of a new redshifted absorption band. Figure 1 shows the set of spectra obtained upon addition of increasing quantities of Cu(II) cation to an acetonitrile solution of probe **3a**. As could be seen in Figure 1, addition of Cu(II) induced a progressive bathochromic shift together with a marked decrease of the band centred at 321 nm with the concomitant appearance of a redshifted absorption at 563 nm. Besides, a marked colour change from colourless to violet was observed.

Nearly the same behaviour was observed with probe **3b** in the presence of Cu(II) cation. Addition of Cu(II) to an acetonitrile solution of **3b** induced the appearance of a new absorbance at 466 nm with a marked colour change from colourless to yellow. Changes upon addition of Cu(II) to acetonitrile solutions of **3a** and **3b** was ascribed to the formation of complexes in which the cation interacts with the acceptor part of the probes (i.e. the imidazole heterocycle).

In a next step, the emission of probes **3a** and **3b** in acetonitrile (1.0 x 10<sup>-5</sup> mol L<sup>-1</sup>) was also studied upon addition of increasing amounts of selected cations. A marked selective response was obtained for Cu(II) that was the unique cation able to induce a remarkable emission quenching for both probes. Figure 2 shows the set of emission spectra obtained upon addition of increasing amounts of Cu(II) to probe **3a**. Addition of Cu(II) induced a progressive quenching of the emission band of **3a** at 404 nm (excitation at ca. 380 nm) together with a hypsochromic shift of ca. 10 nm. A similar quenching and hypsochromic shifts of the emission bands was observed upon addition of Cu(II) to acetonitrile solutions of probe **3b** (see Supporting Information). The obtained results are in accordance with the marked and well-known quenching behaviour of Cu(II) cation.



**Figure 2.** Fluorescence titration profile of probe **3a** in acetonitrile (1.0 x 10<sup>-5</sup> mol L<sup>-1</sup>) upon addition of increasing quantities of Cu(II) cation (0-10 equiv.).

From the UV-visible and emission titration profiles obtained for both probes, the limits of detection (LOD) for Cu(II) in acetonitrile were determined (see Table 2). LODs were assessed using the  $3.3^*(\sigma/s)$  equation, where  $\sigma$  is the standard error of the predicted Y-value vs. each X (i.e. Cu(II) concentration), and *s* is the slope for the linear relationship. Both probes presented similar Cu(II) LODs in the 0.9-2.0  $\mu$ M range.

Table 2. Limits of detection of Cu(II) measured for 3a and 3b probes in acetonitrile.

Probe	UV/Vis [µM]	Emission [µM]	
3a	0.9	1.0	

#### **3b** 1.2 2.0

**Determination of binding stoichiometries and stability constants of probe 3a and 3b with Cu(II) in acetonitrile.** In order to complete the characterization of the complex formed between probe **3a** and **3b** and Cu(II) cation we carried out UV-visible and emission studies to obtain Job's plots to assess the binding stoichiometry and to determine the stability constants.



**Figure 3.** Job's plot for the binding of **3a** with Cu(II). Absorbance at 555 nm was plotted as a function of the molar ratio [Cu(II)]/([3a] + [Cu(II)]). The total concentration of Cu(II) and probe **3a** was  $(2.0 \times 10^{-5} \text{ mol } L^{-1})$ .

Job's plots, constructed using UV-visible measurements in acetonitrile, showed that **3a** and **3b** formed 1:1 complexes with Cu(II). As an example, Figure 3 showed the Job's plot obtained for probe **3a**. Besides, from the UV-visible and fluorescence titration profiles obtained for probes **3a** and **3b** in the presence of Cu(II) cation, the stability constant for the formation of the corresponding 1:1 complexes were determined. The obtained values are shown in Table 3.

Table 3. Logarithms of binding constants measured for the interaction of probes 3a and 3b with Cu(II) in acetonitrile.

Probe ·	Log K		
	UV/Vis	Emission	
3a	5.1 ± 0.1	6.4 ± 0.3	
3b	$5.8 \pm 0.9$	5.8 ± 0.1	

UV/Vis and emission spectroscopic behaviour of probes 3a and 3b in the presence of selected metal cations in aqueous environments. In order to assess the possible use of both probes for Cu(II) detection in real samples, the UV-visible and emission response toward Cu(II) was tested in aqueous environments. Probes 3a and 3b were fully solubilized in water (pH 7.4)-acetonitrile 1:1 v/v. In this medium, both probes showed charge-transfer absorption bands centred at 321 and 339 nm for 3a and 3b respectively. The maxima of the charge-transfer band in water-acetonitrile 1:1 v/v were very similar to those measured in pure acetonitrile, indicating that conformation and hydration of the probes did not change, to any remarkable extent, the energy of their electronic levels in the basal state.

Next, the UV-visible behaviour of aqueous solutions of both probes (5.0×10<sup>-5</sup> mol L<sup>-1</sup>) in the presence of selected metal cations was tested. None of the metal cations tested induced changes in the absorption bands of probe **3b**. In contrast, for probe **3a**, only Cu(II) induced the appearance of a new redshifted absorption at 563 nm as could be seen in Figure 4 (similar to that found in acetonitrile) together with marked colour changes (Figure 5).



**Figure 4.** Absorption of probe **3a** at 563 nm (water (pH 7.4)-acetonitrile 1:1 v/v,  $5.0 \times 10^{-5}$  mol L<sup>-1</sup>) in the presence of 10 eq. of selected metal cations.

More in detail, addition of increasing quantities of Cu(II) cation to water (pH 7.4)acetonitrile 1:1 v/v ( $5.0 \times 10^{-5}$  mol L<sup>-1</sup>) solutions of probe **3a** induced a moderate decrease of the absorption at 321 nm and the simultaneous appearance of a new band centred at 563 nm (see Figure 5). These spectral changes were reflected in a clear colour change from colourless to

violet (see inset in Figure 5). Besides, from the titration profile at 489 nm, a limit of detection for Cu(II) of 2.60  $\mu$ M was determined.



**Figure 5.** UV-Vis titration profile of probe **3a** in water (pH 7.4)-acetonitrile 1:1 v/v (5.0×10<sup>-5</sup> mol L<sup>-1</sup>) upon addition of increasing quantities of Cu(II) cation (0-10 eq.). Inset: colour change when Cu(II) cation was added to water (pH 7.4)-acetonitrile 1:1 v/v solutions of probe **3a**.



**Figure 6.** Fluorescence titration profile of probe **3a** in water (pH 7.4)-acetonitrile 1:1 v/v ( $5.0 \times 10^{-5} \text{ mol L}^{-1}$ ) upon addition of increasing amounts of Cu(II) cation (from 0 to 10 eq.). Inset: Emission intensity at 420 vs equivalents of Cu(II) added.

On the other hand, **3a** was highly emissive in water (pH 7.4)-acetonitrile 1:1 v/v solutions showing a remarkable emission band at 420 nm (excitation at 321 nm). Moreover, fluorescence changes in the presence of selected metal cations was tested. For this purpose, solutions of probe **3a** were excited at the isosbestic point observed in the UV-visible titration profile (ca. 375 nm) and the emission recorded after the addition of different amounts of selected cations. From all cations tested, only Cu(II) was able to induce a response which

consisted of a marked emission quenching of the emission band 420 nm (see Figure 6). The obtained results clearly resemble those obtained for probe **3a** in acetonitrile in the presence of Cu(II) cation. From the emission titration profile (see also Figure 6), a LOD for Cu(II) of 3.4  $\mu$ M for **3a** was determined. This LOD is higher than that found in acetonitrile, most likely as consequence of a stronger solvation of Cu(II) cation in water.

**Spectroscopic detection of biothiols.** In this section we studied the potential use of the **3a**·Cu complex for the recognition of biothiols.[48] Studies were carried out in the presence of selected amino acids (Ala, Arg, Asp, Cys, Gln, Glu, Gly, Hcy, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val) and small peptides (GSH).



**Figure 7.** UV-visible changes of  $3a \cdot Cu$  (6.2 x 10<sup>-6</sup> mol L<sup>-1</sup>) in water (pH 7.4)-acetonitrile 1:1 v/v in the presence of selected amino acids (0.2 eq.) and biothiols (0.2 eq.).



**Figure 8.** Changes in the emission band of  $3a \cdot Cu$  complex (6.2 x 10<sup>-6</sup> mol L<sup>-1</sup>) in water (pH 7.4)-acetonitrile 1:1 v/v upon addition of biothiols (0.2 eq.) and selected amino acids (0.2 eq.).

Water (pH 7.4)-acetonitrile 1:1 v/v solution of complex  $3a \cdot Cu$  (6.2 x 10<sup>-6</sup> mol L<sup>-1</sup>, formed *in situ* by the addition of equimolar quantities of probe and Cu(II) cation) shows an intense absorption band at 563 nm which was the responsible of the strong violet colour observed. From all amino acids (0.2 eq.) and peptides (0.2 eq.) tested, only in the presence of thiol-containing molecules (Cys, Hcy and GSH) a bleaching of the solution was observed (see Figure 7). Besides, water (pH 7.4)-acetonitrile 1:1 v/v solutions of the  $3a \cdot Cu$  complex (6.2 x 10<sup>-6</sup> mol L<sup>-1</sup>) were weakly emissive and only addition of Cys, Hcy and GSH induced a marked emission enhancement at 420 nm (see Figure 8). The chromo-fluorogenic changes observed upon addition of biothiols to the aqueous solutions of  $3a \cdot Cu$  complex are ascribed to a demetallation of the complex, that released the free probe 3a. Finally, from UV-visible and fluorescence titration profiles obtained for  $3a \cdot Cu$ , the LODs for biothiols were determined (Table 4).

Table 4. Limits of detection of biothiols measured for 3a Cu in aqueous environments.

	UV-visible (µM)		Fluorescence (µM)		(μM)	
Complex	GSH	Cys	Нсу	GSH	Cys	Нсу
<b>3a</b> ∙Cu	0.6	1.3	0.8	0.6	0.9	0.7

#### 4. Conclusions

In summary, we show herein the synthesis and chromo-fluorogenic behaviour toward metal cations and biothiols of easy to prepare imidazole-containing probes **3a** and **3b**. Acetonitrile solutions of both probes were characterized by the presence of a charge-transfer absorption band in the 320-350 range. Both probes were moderately emissive in acetonitrile. From all cations tested only Cu(II) induced the growth of new redshifted bands in the 450-570 nm interval. Besides, the emission bands of both receptors were quenched in the presence of Cu(II) cation. These spectral changes were ascribed to a preferential coordination of Cu(II) cation with the electron acceptor imidazole heterocycle. Only probe **3a** changed its UV-visible and emission spectra in the presence of Cu(II) in water-acetonitrile mixtures. On the other hand, weakly emissive complex **3a**·Cu was used to detect biothiols (GSH, Cys and Hcy) in water-acetonitrile 1:1 v/v solution. In the presence of biothiols, the visible bands of the complex disappeared and the emission intensity of the free probe was fully restored. These changes were ascribed to a demetallation process.

#### Acknowledgments

We thank the Spanish Government (MAT2015-64139-C4-1-R) and Generalitat Valenciana (PROMETEO 2018/024). H. E. O. thanks Generalitat Valenciana for his Grisolia fellowship. Thanks are also due to Fundação para a Ciência e Tecnologia (Portugal) for financial support to the Portuguese NMR network (PTNMR, Bruker Avance III 400-Univ. Minho), FCT and FEDER (European Fund for Regional Development)-COMPETEQREN-EU for financial support to the research centre CQ/UM [PEst-C/QUI/UI0686/2013 (FCOMP-01-0124-FEDER-037302], and a post-doctoral grant to R.M.F. Batista (SFRH/BPD/79333/2011).

#### References

- [1] Udhayakumari D, Naha S, Velmathi S, Anal. Methods 2017; 9: 552-578.
- [2] Wu J, Kwon B, Liu W, Anslyn EV, Wang P, Kim JS, Chem. Rev. 2015; 115: 7893-7943.
- [3] Zhang J, Cheng F, Li J, Zhu JJ, Lu Y, Nano Today 2016; 11: 309-329.
- [4] Cheng J, Zhou X, Xiang H, Analyst 2015; 140: 7082-7115.
- [5] Swaminathan S, Gangadaran P, Venkatesh T, Ghosh M, J. Pharm. Biomed. Sci. 2011; 9: 1-15.

[6] Kamble S, Utage B, Mogle P, Kamble R, Hese S, Dawane B, Gacche R, AAPS Pharm. Sci. Tech. 2016; 17: 1030-1041.

[7] Jain AK, Singh RK, Jain S, Raisoni J, Transition Met. Chem. 2008; 33: 243-249.

[8] Goswami S, Sen D, Das NK, Hazra G, Tetrahedron Lett. 2010; 51: 5563-5566.

[9] Xie Q, Zhou T, Yen L, Shariff M, Nguyen T, Kami K, Gu P, Liang L, Rao J, Shi R, Nutr. Diet. Suppl. 2013; 5: 1-6.

[10] Choo XY, Alukaidey L, White AR, Grubman A, Int. J. Alzheimer's Dis. 2013; 2013: 145345.

- [11] Sarkar B, Chem. Rev. 1999; 99: 2535-2544.
- [12] Barnham KJ, Masters CL, Bush AI, Nat. Rev. Drug Discov. 2004; 3: 205-214.
- [13] Ackerman CM, Lee S, Chang CJ, Anal. Chem. 2017; 89: 22-41.
- [14] Hotta H, Tsunoda K, Anal. Sci. 2015; 31: 7-14.
- [15] Townsend DM, Tew KD, Tapiero H, Biomed. Pharmacother. 2003; 57; 145-155.
- [16] Miller JW, Beresford SA, Neuhouser ML, Cheng TY, Song X, Brown EC, Zheng Y,
- Rodriguez B, Green R, Ulrich CM, Am. J. Clin. Nutr. 2013; 97: 827-834.

[17] Dorszewska J, Prendecki M, Oczkowska A, Dezor M, Kozubski W, Curr. Alzheimer Res. 2016; 13: 952-963.

[18] Rozycka A, Jagodzinski PP, Kozubski W, Lianeri M, Dorszewska J, Curr. Genomics 2013; 14: 534-542.

[19] Santa T, Drug Discov. Ther. 2013; 7: 172-177.

[20] Carlucci F, Tabucchi A, J. Chromatogr. B Analyt. Technol. Biomed. Life Sci. 2009; 877: 3347-3357.

[21] Blair IA, Biomed. Chromatogr. 2010; 24: 29-38.

[22] Isik M, Ozdemir T, Turan IS, Kolemen S, Akkaya EU, Org. Lett. 2013; 15: 216-219.

[23] Liao YC, Venkatesan P, Wei LF, Wu SP, Sens. Actuators B Chem. 2016; 232: 732-737.

[24] Lee KS, Park J, Park HJ, Chung YK, Park SB, Kim HJ, Shin IS, Hong JI, Sens. Actuators B Chem. 2016; 237: 256-261.

[25] Kong F, Liu R, Chu R, Wang X, Xu K, Tang B, Chem. Commun. 2013; 49: 9176-9178.

[26] Zhang J, Jiang XD, Shao X, Zhao J, Su Y, Xi D, Yu H, Yue S, Xiao LJ, Zhao W, RSC Adv.2014; 4: 54080-54083.

[27] Wang YW, Liu SB, Ling WJ, Peng Y, Chem. Commun. 2016; 52: 827-830.

[28] Liu J, Sun YQ, Zhang H, Huo Y, Shi Y, Guo W, Chem. Sci. 2014; 5: 3183-3188.

[29] Guan YS, Niu LY, Chen YZ, Wu LZ, Tung CH, Yang QZ, RSC Adv. 2014; 4: 8360-8364.

[30] Fu ZH, Han X, Shao Y, Fang J, Zhang ZH, Wang YW, Peng Y, Anal. Chem. 2017; 89: 1937-1944.

[31] Yang YL, Zhang FM, Wang YW, Zhang BX, Fang R, Fang JG, Peng Y, Chem. As. J. 2015; 10: 422-426.

[32] Fu ZH, Yan LB, Zhang X, Zhu FF, Han XL, Fang J, Wang YW, Peng Y, Org. Biomol. Chem. 2017; 15: 4115-4121.

[33] Singh Y, Arun S, Singh BK, Dutta PK, Ghosh T, RSC Adv. 2016; 6: 80268-80274.

[34] Lee SH, Lee JJ, Shin JW, Min KS, Kim C, Dyes Pigm. 2015; 116: 131-138.

[35] Maheshwaran D, Nagendraraj T, Manimaran P, Ashokkumar B, Kumar M, Mayilmurugan R,

Eur. J. Inorg. Chem. 2017; 1007-1016.

[36] Kim YS, Park GJ, Lee SA, Kim C, RSC Adv. 2015; 5: 31179-31188.

[37] You GR, Lee JJ, Choi YW, Lee SY, Kim C, Tetrahedron 2016; 72: 875-881.

[38] Santos-Figueroa LE, Llopis-Lorente A, Royo S, Sancenón F, Martínez-Máñez R, Costero AM, Gil S, Parra M, ChemPlusChem 2015; 80: 800-804.

[39] Marín-Hernández C, Santos-Figueroa LE, El Sayed S, Pardo T, Raposo MMM, Batista RMF, Costa SPG, Sancenón F, Martínez-Máñez R, Dyes Pigm. 2015; 122: 50-58.

[40] Lo Presti M, El Sayed S, Martínez-Máñez R, Costero AM, Gil S, Parra M, Sancenón F, New J. Chem. 2016; 40: 9042-9045.

[41] El Sayed S, de la Torre C, Santos-Figueroa LE, Pérez-Paya E, Martínez-Máñez R, Sancenón F, Costero AM, Parra M, Gil S, RSC Adv. 2013; 3: 25690-25693.

[42] El Sayed S, de la Torre C, Santos-Figueroa LE, Martínez-Máñez R, Sancenón F, Supramol. Chem. 2015; 4: 244-254.

[43] Marín-Hernández C, Santos-Figueroa LE, Moragues ME, Raposo MMM, Batista RMF,
Costa SPG, Pardo T, Martínez-Máñez R, Sancenón F, J. Org. Chem. 2014; 79: 1075210761.

[44] Okda HE, El Sayed S, Otri I, Ferreira RCM, Costa SPG, Raposo MMM, Martínez-Máñez R, Sancenón F, Dyes Pigm. 2019; 162: 303-308.

[45] Nakashima K, Fukuzaki Y, Nomura R, Shimoda R, Nakamura Y, Kuroda N, Akiyama S, Irgum K, Dyes Pigm. 1998; 38: 127-136.

[46] Bian, ZQ, Wang KZ, Jin LP, Polyhedron 2002; 21: 313-319.

[47] Okda HE, El Sayed S, Ferreira RCM, Costa SPG, Raposo MMM, Martínez-Máñez R, Sancenón F, Dyes Pigm. 2018; 159: 45-48.

[48] Mandal S, Das G, Singh R, Shukla R, Bharadwaj PK, Coord. Chem. Rev. 1997; 160: 191-235.

Graphical abstract synopsis:

Chromo-fluorogenic recogntion of Cu(II) cation and bithiols was achieved using two imidazolebased probes.

Graphical abstract pictogram:

