This article was downloaded by: [University of California Santa Cruz] On: 23 October 2014, At: 15:04 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

A New and Efficient Method for the Protection of Alcohols and Phenols by Using Hexamethyldisilazane in the Presence of Anhydrous Ferric Chloride under Mild Reaction Conditions

Batool Akhlaghinia^a

^a Department of Chemistry , Damghan University of Basic Sciences , Damghan, Iran Published online: 30 Jan 2007.

To cite this article: Batool Akhlaghinia (2007) A New and Efficient Method for the Protection of Alcohols and Phenols by Using Hexamethyldisilazane in the Presence of Anhydrous Ferric Chloride under Mild Reaction Conditions, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:3, 687-694, DOI: <u>10.1080/10426500601047487</u>

To link to this article: <u>http://dx.doi.org/10.1080/10426500601047487</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the

Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



A New and Efficient Method for the Protection of Alcohols and Phenols by Using Hexamethyldisilazane in the Presence of Anhydrous Ferric Chloride under Mild Reaction Conditions

Batool Akhlaghinia

Department of Chemistry, Damghan University of Basic Sciences, Damghan, Iran

Alcohols and phenols are protected with hexamethyldisilazane in the presence of anhydrous ferric chloride in good to excellent yields in acetonitrile. This method is highly selective for the conversion of primary alcohols in the presence of secondary and tertiary alcohols and also for conversion phenols.

Keywords Alcohol; ferric chloride; hexamethyldisilazane HMDS; phenol; trimethylsilylether

INTRODUCTION

The importance of hydroxyl-group functionality in organic chemistry is reflected by the continuous appearance in the literature of methods for developing complex structures based on the chemical interconversion of this functional group. A limiting factor in some cases is the presence of more than one oxygen function of comparable reactivity in the molecule to be transformed. This difficulty is circumvented by the use of protective groups, which is a practice that in theory allows free manipulation of the desired functionality. However, it is only in a few cases that this practice reaches an acceptable practical level, mostly due to the difficulties involved in the selectivity of protecting and of regenerating specific hydroxyl groups. The role of silyl groups has already been recognized as an important part of organic chemistry from both analytical and synthetic points of view, especially as a protecting group in many syntheses of reasonable complexity.¹ Generally the formation of

Received July 29, 2005; accepted August 21, 2006.

The authors gratefully acknowledge the partial support of this study by the Damghan University Research Council.

Address correspondence to Batool Akhlaghinia, Department of Chemistry, Damghan University of Basic Sciences, Damghan, Iran. E-mail: b_akhlaghinia@dubs.ac.ir

silyl ethers is carried out by treatment of alcohols with silyl chlorides or silvl triflates in the presence of a base,² Li₂S,³ and sometimes a nonionic super base catalyst.⁴ However, some of these methods frequently suffer from drawbacks, such as a lack of reactivity or difficulty in the removal of amine salts derived from the reaction of byproduced acids and cobases during the silvlation reaction. 1,1,1,3,3,3-Hexamethyldisilazane (HMDS) is a stable, commercially available, and cheap reagent for trimethylsilylation of hydrogen-labile substrates,^{1a} giving ammonia as the only byproduct. On the other hand, silvlation using this silazantype reagent is nearly neutral and does not need precautions. However, the low silvlating power of HMDS is the main drawback for its application, which needs forceful conditions and long reaction times in many cases. A variety of catalysts, such as (CH₃)₃SiCl,⁵ sulfonic acids⁶ ZnCl₂,⁷ complexes of metal chlorideswithnitrogen ligands,⁸ zirconium sulfophenyl phosphonate,⁹ K-10 montmorilonite,¹⁰ and a special type of catalyst having the general formula X¹-NH-X² in which at least one of the groups X^1, X^2 , is electron-withdrawing, such as an ester or amide group, has also been reported for the silvlation of a wide range of functional groups using HMDS.¹¹

As part of our ongoing research to develop new synthetic methodologies involving various new reagents,¹² we perceived that anhydrous ferric chloride, which is readily available, might be a useful catalyst for trimethylsilylation because of its mild Lewis acidity. In this article we report that anhydrous ferric chloride can be used as an efficient catalyst for the trimethylsilylation of alcohols and phenols.

RESULTS AND DISCUSSION

In the present study the trimethylsilylation of alcohols and phenols in the presence of a catalytic amount of $FeCl_3$ was investigated. The protection of benzyl alcohol under the catalytic effect of $FeCl_3$ was studied. Benzyl trimethylsilylether was produced immediately in a quantitative yield. We applied these reaction conditions for the protection of structurally different alcohols and phenols (Scheme 1).

$$2 \text{ ROH } + H_3C - \underbrace{\underset{\substack{I \\ CH_3}}{\text{Si}-\text{N}}}_{\text{R: primary, secondary, tertiary alkyl and phenyl}}^{\text{CH}_3 \text{ H} \quad \underset{\substack{I \\ CH_3}}{\text{CH}_3} \underbrace{\underset{\substack{I \\ CH_3}}{\text{FeCl}_3}}_{\text{FeCl}_3} \underbrace{\underset{\substack{I \\ CH_3}}{\text{FeCl}_3}}^{\text{CH}_3} 2 \begin{array}{c} \text{RO-Si} - \text{CH}_3 + \text{NH}_3 \\ \\ \text{R: primary, secondary, tertiary alkyl and phenyl}} \\ \end{array}$$

SCHEME 1





In order to optimize reaction conditions, we first examined the effect of different molar ratios of the alcohol/HMDS/catalyst. The best results were obtained using the ratio 1:0.7:0.07. Trimethylsilylethers were produced in quantitative yields (Table I).

Compounds that were trimethylsilylated in this way are primary, secondary, and tertiary alcohols and phenols. Generally, the reactions were completed at r.t. and were accompanied by the evolution of NH_3 gas from the reaction mixture.

The mechanism of these transformations is unclear. One hypothesis may be that $FeCl_3$ acts as a Lewis acid, which may polarize the Si-N bond in HMDS to produce the effective silvlating agent 1 (Scheme 2). A rapid reaction of 1 with alcohol yields 2 with a concomitant release of the corresponding silvlether. The reaction of 2 with a second molecule of alcohol produces the corresponding silvlether and the Fe(III) ammine complex 3. The cleavage of 3 leads to the fast evolution of NH₃ and release of the catalyst, which re-enters the catalytic cycle (Scheme 2).

This method was found to be highly selective. In a binary mixture of benzyl alcohol and 3-phenyl-1-propanol, benzylic alcohol was completely converted to the corresponding silylether, while only a 5% conversion was observed for the propanol derivative (Table II). Excellent

ROH	Reaction time	Conversion $(\%)^a$	Isolated yield (%)
ОН	Immediately	100	92
П	$2.5 \mathrm{h}2$	100	90
OH			
	Immediately	100	93
ОН	10 min	100	96
ОН	Immediately	100	98
	5 h	100	97
	3 h	100	96
ОН	3 h	100	98
ОН	3.5 h	100	97
O ₂ N OH	20 min	100	91
СІ	10 min	100	92
ОН	Immediately	100	98
ОН			
	Immediately	100	95

TABLE I Trimethyl
silylation of Alcohols and Phenols Catalyzed by $\ensuremath{\text{FeCl}}_3$ in Dry Acetonitrile

690

ROH	Reaction time	Conversion $(\%)^a$	Isolated yield (%)
ОН	Immediately	100	95
ОН	Immediately	100	99
OH	Immediately	100	90
——————————————————————————————————————	46 h	100	94
ОН	32 h	80	60
ОН	10 min	100	95
OH	30 min	100	97
OH	70 min	100	90
ОН	25 min	100	94
CI	2 h	100	95
OH	75 min	90	85
ОН	1.5 h	65	58

TABLE I Trimethyl
silylation of Alcohols and Phenols Catalyzed by $FeCl_3$ in Dry Acetonitrile
 (Continued)

^{*a*}Yields were determined by GC using an internal standard.

Binary mixture	Products	Reaction time	Yield (%)
ОН	OSiMe ₃	Immediately	100
ОН	ОН		95
ОН	OSiMe ₃	Immediately	100
ОН	ОН		100
ОН	OSiMe ₃	Immediately	100
ОН	ОН		96
ОН	OSiMe ₃	4.5 h	100
ОН	ОН		100
ОН	OSiMe ₃	Immediately	100
ОН	ОН		100
ОН	OSiMe ₃	Immediately	100
ОН	ОН		95
ОН	OSiMe ₃	Immediately	100
ОН	ОН		97

TABLE II The Selective Reaction of Different Binary Mixtures with HMDS/FeCl_3 $% \mathcal{A}_{3}$

Binary mixture	Products	Reaction time	Yield (%) ^a
ОН	OSiMe3	6.5 h	100
ОН	ОН		97
ОН	OSiMe ₃	Immediately	100
OH OH	OH		97

TABLE II The Selective Reaction of Different Binary Mixtures with HMDS/FeCl $_3$

^aYields were determined by an GC using internal standard.

selectivity was also observed for the conversion of primary alcohols in the presence of a secondary alcohol as well as of secondary alcohols in the presence of tertiary ones (Table II). Selectivity was observed in the conversion of benzyl alcohol in the presence of phenol (Table II).

In conclusion, the methodology described in this article shows that $FeCl_3$ is an effective catalyst for the trimethylsilylation of alcohols and phenols. Advantages of our protocol are simplicity, short reaction times, mild reaction conditions, as well as good yields. We expect this method to be a useful addition to modern synthetic methodologies.

EXPERIMENTAL

Chemicals were obtained from Merck (Darmstadt, Germany) or Fluka (Buchs, Switzerland). FTIR spectra were recorded on a Perkin Elmer (Norwalk, Connecticut, USA) RXI spectrometer.NMR spectra were recorded on a Bruker Avance DPX 250 MHz instrument. Products were purified by column chromatography, and purity determination of the products was accomplished by GLC on a Shimadzu model GC 10-A instrument using *n*-octane as an internal standard or by TLC on a silica-gel polygram on SIL G/UV 254 plates.

Typical Procedure for the Protection of Benzyl Alcohol

To a mixture of benzyl alcohol (0.108 g, 1 mmol) and HMDS (0.112 g, 0.7 mmol) in dry CH_3CN (3–5 mL), $FeCl_3$ (0.011 g, 0.07 mmol) was added, and the reaction mixture was stirred at room temperature. GC

analysis showed that the reaction was completed immediately. The reaction mixture was filtered. Column chromatography of the filtrate on silica gel using *n*-hexane as an eluent gave benzyl trimethylsilylether in a 98% yield.

REFERENCES

- (a) M. Lalonde and T. H. Chan, Synthesis, 817 (1985); (b) T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, 2nd ed. (Wiley, New York, 1991); (c) P. J. Kocienski, In Protective Groups, R. Enders, and R. Noyori, eds., (Thieme, Stuttgart, 1994), 28–42.
- [2] S. K. Ghaudhary, and O. Hernandez, Tetrahedron Lett., 20, 99 (1979).
- [3] G. A. Olah, B. G. B. Gupta, S. C. Narang, and R. Malhotra, J. Org. Chem., 44, 4272 (1979).
- [4] (a) B. A. D. Sa, D. Mc Leod, and J. G. Verkade, J. Org. Chem., 62, 5057 (1997); (b) B.
 A. D. Sa and J. G. Verkade, J. Am. Chem. Soc., 118, 2832 (1996).
- [5] (a) S. H. Langer, S. Connell, and J. Wender, J. Org. Chem., 23, 50 (1958); (b) P. Gauttret, S. El-Ghamarti, A. Legrand, D. Coutrier, and B. Rigo, Synth. Commun., 26, 707 (1996).
- [6] A. G. T. Goldschmidt, German Patent 2758884, Chem. Abstr., 90, 65309 (1979).
- [7] H. Firouzabadi and B. Karimi, Synth. Commun., 23, 1633 (1993).
- [8] (a) H. Firouzabadi, A. R. Sardarian, Z. Khayat, and B. Karimi, Synth. Commun., 27, 2709 (1997); (b) A. Bongini, G. Cardillo, M. Orena, and S. Sandri, Synthesis, 618 (1979).
- [9] M. Curini, F. Epifano, M. C. Marcotullio, O. Rosati, and U. Constantino, Synth. Commun., 29, 541 (1999).
- [10] Z. H. Zhang, T. S. Li, F. Yang, and C. G. Fu, Synth. Commun., 28, 3105 (1998).
- [11] C. A. Bruynes and T. K. Jurriens, J. Org. Chem., 47, 3966 (1982).
- [12] (a) B. Akhlaghinia, M. Asadi, E. Safaei, and M. Heydarpoor, *Phosphorus, Sulfur, and Silicon*, **179**, 2099 (2004); (b) B. Akhlaghinia, M. Asadi, E. Safaei, and M. Heydarpoor, *J. Porphyrins and Phthalocyanies*, **8**, 1285 (2004); (c) B. Akhlaghinia and A. R. Pourali, *Synthesis*, 1747 (2004); (d) B. Akhlaghinia, *Phosphorus, Sulfur, and Silicon*, **179**, 1783 (2004); (e) B. Akhlaghinia and S. Tavakoli, *Synthesis*, 1775 (2005); (f) B. Akhlaghinia, S. Tavakoli, M. Asadi, and E. Safaei, *J. Porphyrins and Phthalocyanies*, **10**, 167 (2006).