IP Organocatalysis Very Important Paper

International Edition: DOI: 10.1002/anie.201606480 German Edition: DOI: 10.1002/ange.201606480

An Organocatalytic Cope Rearrangement

Dainis Kaldre and James L. Gleason*

Abstract: The first example of an organocatalytic Cope rearrangement is reported. Acyclic and cyclic acyl hydrazides catalyze the rearrangement of 1,5-hexadiene-2-carboxaldehydes via iminium ion formation. A correlation between ring size and catalyst activity was observed for the cyclic hydrazides, with seven- and eight-membered-ring catalysts being the most active. Diazepane carboxylate 5c (10 mol%) catalyzed the rearrangement of a range of dienes at room temperature in acetonitrile using triflic acid as a co-catalyst. Preliminary proof of principle for asymmetric catalysis was provided by rearrangement of 3,3-dimethyl-7-phenyl-1,5-heptadiene-2-carboxaldehyde in the presence of a novel 7-substituted diazepane carboxylate.

 ${oldsymbol{P}}$ ericyclic reactions play a central role in the synthesis of complex natural products.^[1] Whereas significant progress has been made in the catalysis of cycloadditions, most notably the Diels-Alder reaction,^[2] catalytic asymmetric methods of sigmatropic and electrocyclic reactions are less well developed.^[3] In both the Cope and Claisen rearrangements, transition-metal coordination to the alkenes has played a prominent role, and complexes of palladium, platinum, and nickel have all been reported to facilitate [3,3]-sigmatropic rearrangements.^[4] These methods have been extended to several catalytic asymmetric Claisen rearrangements^[5] and a single example of an enantioselective Cope rearrangement, the gold-catalyzed rearrangement of 1-cyclopropylidiene-1,5hexadienes.^[6] In the Claisen rearrangement, Lewis acid coordination of the oxygen atom may also accelerate the [3,3]-rearrangement by weakening the C-O bond and/or stabilizing charge on the oxygen atom in the transition state.^[7] This has enabled several catalytic asymmetric Claisen rearrangements that are catalyzed by chiral Lewis acid complexes^[8] and chiral protic organocatalysts.^[9,10] Finally, a few sporadic reports have suggested that stoichiometric acid can promote the Cope rearrangement of acyl-1,5-dienes,^[11] and the use of catalytic amounts of acid has been shown to promote the electrocyclic closure of acyl-1,3,5-hexatrienes.[3a,d,12] Herein, we expand upon the possibility of acid catalysis to report the first organocatalytic Cope rearrangement and identify a new seven-membered-ring hydrazide

 [*] D. Kaldre, Prof. J. L. Gleason Department of Chemistry McGill University
 801 Sherbrooke W., Montreal, QC, H3A 0B8 (Canada)
 E-mail: jim.gleason@mcgill.ca
 Homepage: http://gleason-group.mcgill.ca

 Supporting information and the ORCID identification number(s) for
 the author(s) of this article can be found under: http://dx.doi.org/10.1002/anie.201606480.

Angew. Chem. Int. Ed. 2016, 55, 1-6

catalyst capable of iminium ion formation from hindered enals.

The concept of LUMO-lowering catalysis via iminium ion formation is well established for Diels–Alder cycloadditions.^[13] We envisioned extending LUMO-lowering catalysis to other pericyclic reactions such as the Cope and Claisen rearrangements. We began by examining the accelerating effect on the Cope rearrangement of 1,5-hexadiene by an iminium ion substituent at the 2-position (see Figure 1). This



Figure 1. Activation modes for Cope and Claisen rearrangements.

position was chosen as it would be impervious to effects of conjugation/deconjugation and would lend itself more easily to eventual asymmetric catalysis. An inherent limitation is the significant A-1,3 strain that arises in the formation of an iminium ion from an α -substituted enal/enone with a secondary amine. Indeed, there are only a few such examples of iminium catalysis with secondary amines,^[14] although primary amine catalysis is often a viable solution,^[15] and thus this is an area of opportunity for new catalyst development.

4-Phenyl-1,5-hexadiene-2-carboxaldehyde (1a)was chosen as the model substrate as the Cope equilibrium was expected to favor the conjugated product 2a. Indeed, thermolysis of 1a in toluene for 24 h at 110°C proceeded to completion and resulted in the isolation of 2a in 94% yield. Potential catalysts were initially screened in $[D_4]$ methanol at 60°C, conditions under which the uncatalyzed rearrangement of 1a was observed to proceed in only 21% yield over 24 h (Table 1). The addition of proline (50 mol%) slightly increased the conversion (29%) whereas other simple secondary amines, such as pyrrolidine, proline methyl ester, and MacMillan's second-generation catalyst (3),^[16] each as the hydrochloride salt, did not afford any observable catalysis. As noted above, the negligible levels of catalysis were presumed to result from significant A-1,3 strain in the iminium ion; indeed, we did not observe any iminium ion formation by ¹H NMR spectroscopy with simple secondary amine catalysts. We reasoned that a more nucleophilic amine might increase the rate of iminium ion formation. N-Acyl hydrazides have been found to form iminium ions at high rates in the Diels-Alder reaction, with the higher reactivity

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Wiley Online Library

Table 1: Catalyst screen.

	CHO 50 mol% catalyst		СНО	
	Ph 1a	50 mol% HCl	Ph	2a
Entry	Catalyst	Solvent	T [°C]	Yield [%] ^[a]
1	_[b]	CD₃OD	60	21
2	proline ^[b]	CD ₃ OD	60	29
3	pyrrolidine	CD₃OD	60	21
4	3	CD ₃ OD	60	16
5	4a	CD₃OD	60	34
6	4 b	CD₃OD	60	45
7	5 a	CD₃OD	60	18
8	5 b	CD₃OD	60	35 ^[c]
9	4 b	CD₃CN	23	19
10	5 a	CD₃CN	23	6
11	5 b	CD₃CN	23	28
12	5 c	CD₃CN	23	83
13	5 d	CD₃CN	23	89

[a] Determined by ¹H NMR spectroscopy with mesitylene as the internal standard. All reactions were run for 24 h unless otherwise noted.
[b] Without an acid co-catalyst. [c] Acetal and oxa-Michael products observed.



initially ascribed to the α -effect.^[14a,17] We were pleased to find that several acyl hydrazides afforded a measurable improvement in catalytic activity. For instance, cyclic hydrazides **4a** and **4b** afforded modest yields, with conversions up to 45% (Table 1, entries 5 and 6).

Examination of cyclic hydrazides showed that whereas a five-membered-ring hydrazide (5a) was inferior to acyclic hydrazides, the corresponding six-membered-ring catalyst 5b was significantly more active, with the starting material being fully consumed within 24 h (Table 1, entries 7 and 8). However, the mass balance of the reaction was negatively affected by the formation of a mixture of acetal and oxa-Michael addition products that result from addition of the solvent, [D₄]methanol. Switching to [D₃]acetonitrile alleviated this problem, and at room temperature, conditions under which the uncatalyzed reaction does not proceed to any measureable extent, six-membered-ring catalyst 5b afforded the product in 28%, a definite advantage over the smaller catalyst 5a, which was largely ineffective under these conditions (Table 1, entries 10 and 11). Given the higher reactivity of the larger-ring catalyst, we extended our study to the seven-membered diazepane carboxylate 5c and were pleased to observe a significant improvement in reactivity along with a yield of 83% (Table 2, entry 12). Given the continued trend towards higher reactivity with increasing ring size, we also prepared eight-membered-ring catalyst 5d. Whereas a slight increase in reactivity was noted (89% yield), its inherently more difficult synthesis discouraged further development. It is important to note that primary amines, such as aniline, benzylamine, 1,2-cyclohexanediaTable 2: Optimization of the acid co-catalyst.

Entry	Catalyst	Co-catalyst (mol%)	Yield [%] ^[b]
1	5 b	TsOH (50)	25
2	5 b	HCI (50)	28
3	5 b	HBr (50)	38
4	5 b	HClO₄ (50)	94
5	5 b	TfOH (50)	96
6	5 b	TfOH (20)	75
7	5 b	TfOH (10)	43
8	5 c	TfOH (10)	95

[a] General reaction conditions: **1** a, catalyst as noted in equivalent mol% to acid, CD₃CN, 23 °C. [b] Determined by ¹H NMR spectroscopy with mesitylene as the internal standard. Tf=trifluoromethanesulfonyl, Ts = *para*-toluenesulfonyl.

mine, *O*-methyl hydroxylamine, alanine, and glycine methyl ester, all failed to afford any rearrangement product at 23 °C, highlighting the unique reactivity of the cyclic hydrazide catalysts.

The acid co-catalyst was optimized using the less reactive six-membered-ring catalyst **5b**. We found that stronger acids afforded significantly improved reaction rates (Table 2). Using either triflic or perchloric acid with **5b**, each at 50 mol% loading, the reaction proceeded to completion within 24 h. Although reducing the catalyst loading with less reactive catalyst **5b** resulted in a significant diminution in yield, applying these conditions with 10 mol% of the more active diazepane carboxylate **5c** allowed the rearrangement to proceed to completion within 24 h and afforded the Cope product in 95% yield (isolated in 90% yield; Table 3).

Diazepane catalyst 5c is highly effective with a range of 1,5-diene-2-carboxaldehydes (Table 3). Good yield was observed for *n*-pentyl-substituted substrate 1b, indicating that alkene conjugation with a phenyl group, as in product 2a, is not required for reactivity. Substituents were also tolerated at the 3- and 6-positions (entries 3 and 4). E Substitution was also tolerated at the 1-position (entry 5), but the reaction was slower, which is presumably due to the presence of an axial methyl group in a probable chair-like transition state.^[18] Intriguingly, substitution at the 5-position was not tolerated (1 f, entry 6) under the standard reaction conditions, and decomposition was observed with higher loadings. In contrast, 1f does undergo rearrangement under standard thermal conditions (toluene, 110°C, 24 h) to afford 2f in 64% yield. Finally, we found that geminal dimethyl substitution was surprisingly well tolerated at the 3-position (entries 7 and 8), again demonstrating the unique reactivity of the diazepane carboxylate in forming highly hindered α -substituted iminium ions.

We initially envisioned a simple catalytic cycle in which iminium ion formation would facilitate the Cope rearrangement through simple LUMO lowering (dashed arrow, Scheme 1). DFT calculations $(B3LYP/6-31G^*)^{[19]}$ clearly indicated an accelerating effect of electron-withdrawing substituents on the Cope rearrangement. The rearrangement of 1,5-hexadiene was predicted to have an activation energy of 35.8 kcal mol⁻¹. Incorporating an aldehyde at the 2-position lowered the predicted barrier to 30.5 kcal mol⁻¹, and formation of a simple iminium (R = H in Scheme 1) reduced this

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim





[a] Yield of isolated product. [b] Substrate **1d** was used as a 72:28 *E/Z* mixture. [c] At 40 °C for 96 h. [d] Substrate **1 f** was used as a 69:31 *E/Z* mixture. [e] Use of 50 mol% **5 c**/25 mol% TfOH led to significant decomposition. [f] Run for 48 h.

barrier even further to 10.1 kcalmol⁻¹. However, while the predicted transition states for 1,5-hexadiene and 1,5-hexadiene-2-carboxaldehyde correspond to an archetypical concerted sigmatropic rearrangement, the iminium ion is predicted to proceed through a shallow-energy intermediate $(\Delta G = 9.7 \text{ kcal mol}^{-1})$ that bears a positive charge on the 5-position. Thus the reaction might be viewed as a conjugate addition of the alkene to the α,β -unsaturated iminium ion, followed by fragmentation of the cation to complete the rearrangement; in many ways, the transformation would be similar to the palladium-catalyzed Cope rearrangements originally described by Overman and co-workers.^[4a,b] Thus far, the only evidence for a stepwise reaction is computational, and it is noteworthy that the rearrangement of 1 f, which bears a methyl group at the 5-position and would thus be expected to stabilize the intermediate cation, does not afford any rearranged product. Although stabilizing the cation might allow it to follow an alternative pathway, we



Scheme 1. A) Proposed catalytic cycle. Energies computed at the B3LYP/6-31G* level of theory for R = H. ΔG values given in kcal mol⁻¹. B) Structures of the computed transition state and the intermediate cation.

did not identify any side products with this particular substrate.

Whereas detailed mechanistic work remains to be done, we have not been able to observe the iminium ion intermediates directly, suggesting that iminium ion formation is rate-limiting. Importantly, we have found that the N-methyl analogue of 5c does not catalyze the Cope rearrangement, which is consistent with iminium ion formation and suggests that simple proton catalysis of the Cope rearrangement is not operative under our conditions.^[20] It was presumed that the high catalytic activity of 5c/5d arises from faster iminium ion formation relative to their smaller-ring congeners. Owing to the increased A-1,3 strain resulting from the larger bond angles in the seven-membered ring, the stability of the iminium intermediates is predicted to decrease as the ring size increases,^[21] which is opposite to the observed reactivity trend. As suggested by Tomkinson and co-workers, the higher reactivity of 5c may result from the reduced proton affinity of the amine, facilitating easier protonation of the oxygen atom during iminium ion formation.^[17f,22]

The catalyst system reported here will open up new avenues for reaction discovery. In particular, the high reactivity of the diazepane carboxylates with α -substituted enals has a wide range of possible applications in catalysis and

www.angewandte.org

makes them an intriguing platform for chiral catalyst development. To this end, we assessed the potential for asymmetric catalysis using substrate **1g**. This is a challenging substrate as the new stereocenter is remote to the positioning of the catalyst (see the TS structure in Scheme 1). We examined two catalyst architectures, one with a stereocenter adjacent to the nucleophilic nitrogen atom (**5e/5f**; Scheme 2) and one with the stereocenter adjacent to the carbamate (**5g/5h**; Scheme 2). To our surprise, catalysts bearing the substituent



Scheme 2. Asymmetric variant of the Cope rearrangement.

adjacent to the nucleophilic nitrogen atom gave minimal enantioinduction. In contrast, **5g** and **5h**, with substitution adjacent to the carbamate nitrogen atom, afforded the desired product in 28 and 47 % *ee*, respectively.^[23,24] The observation that substitution further from the reactive site was more effective was unexpected and may result from gearing of the carbamate by the adjacent stereocenter. This is an intriguing control element for asymmetric catalysis that should enable the rational development of catalysts for reactions that involve α -substituted iminium ions.

In conclusion, we have developed the first organocatalytic Cope rearrangement and identified diazepane carboxylates as unique catalysts for the activation of α -substituted enals. The method is viable with a range of 1,5-hexadienyl-2-carboxyladehydes. Computational evidence suggests a possible stepwise mechanism, and preliminary examination of chiral catalysts revealed that remote stereocontrol elements are effective through gearing of the hydrazide carboxylate. The diazepane catalysts represent an intriguing platform for future development in iminium catalysis of α -substituted enals.

Experimental Section

To a solution of ethyl 1,2-diazepane-1-carboxylate (**5c**; 1.3 mg, 75 µmol, 10 mol%) in 275 µL MeCN were added 26.5 µM of a stock solution of TfOH (0.283 M, 7.5 µmol, 10 mol%) and mesitylene (internal standard). After 5 min, the mixture was added to neat 2-methylene-4-phenylhex-5-enal (**1a**; 14.0 mg, 0.075 mmol, 1 equiv). The reaction was mixed for 2 min and monitored by ¹H NMR spectroscopy for 24 h. The reaction was diluted with EtOAc (2 mL), washed with saturated NaHCO₃ (2 mL), extracted with EtOAc (2 × 5 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo. The product was purified by flash chromatography on silica gel (hexanes/ EtOAc 95:5), providing **2a** as an oil (12.6 mg, 0.065 mmol, 90%). IR (film): $\tilde{\nu}$ =3028, 2925, 1689, 964 cm⁻¹; ¹H NMR (400 MHz, CDCl₃):

$$\begin{split} &\delta = 9.56 \text{ (s, 1 H), 7.34-7.26 (m, 4 H), 7.21 (t, J = 6.9 Hz, 1 H), 6.40 (d, J = 15.9 Hz, 1 H), 6.30 (s, 1 H), 6.18 (dt, J = 15.6, 6.4 Hz, 1 H), 6.04 (s, 1 H), 2.46-2.36 ppm (m, 4 H); ^{13}\text{C NMR} (100 MHz, CDCl_3): \\ &\delta = 194.62, 149.28, 137.48, 134.70, 130.74, 129.25, 128.50, 127.04, 125.98, 31.10, 27.65 ppm. HRMS:$$
m/*z*calcd for C₁₃H₁₄ONa⁺: 209.0937 [*M*+Na]⁺; found: 209.0929.

Acknowledgements

We thank the Natural Science and Engineering Research Council of Canada for funding. D.K. thanks the Canadian Institutes of Health Research for a fellowship.

Keywords: allylic strain · Cope rearrangements · hydrazides · iminium ions · organocatalysis

- a) K. C. Nicolaou, S. A. Snyder, T. Montagnon, G. Vassilikogiannakis, *Angew. Chem. Int. Ed.* **2002**, *41*, 1668–1698; *Angew. Chem.* **2002**, *114*, 1742–1773; b) G. Desimoni, G. Tacconi, A. Barco, G. P. Pollini, *ACS Monogr.* **1983**, *180*, 443; c) J. Poulin, C. M. Grise-Bard, L. Barriault, *Chem. Soc. Rev.* **2009**, *38*, 3092– 3101.
- [2] a) E. J. Corey, Angew. Chem. Int. Ed. 2002, 41, 1650-1667;
 Angew. Chem. 2002, 114, 1724-1741; b) P. Merino, E. Marqués-López, T. Tejero, R. Herrera, Synthesis 2009, 1-26.
- [3] a) L. M. Bishop, J. E. Barbarow, R. G. Bergman, D. Trauner, Angew. Chem. Int. Ed. 2008, 47, 8100-8103; Angew. Chem.
 2008, 120, 8220-8223; b) E. E. Maciver, S. Thompson, M. D. Smith, Angew. Chem. Int. Ed. 2009, 48, 9979-9982; Angew. Chem. 2009, 121, 10164-10167; c) S. Müller, B. List, Angew. Chem. Int. Ed. 2009, 48, 9975-9978; Angew. Chem. 2009, 121, 10160-10163; d) L. M. Bishop, R. E. Roberson, R. G. Bergman, D. Trauner, Synthesis 2010, 2233; e) E. E. Maciver, P. C. Knipe, A. P. Cridland, A. L. Thompson, M. D. Smith, Chem. Sci. 2012, 3, 537.
- [4] a) L. E. Overman, F. M. Knoll, J. Am. Chem. Soc. 1980, 102, 865–867; b) L. E. Overman, E. J. Jacobsen, J. Am. Chem. Soc. 1982, 104, 7225–7231; c) J. L. van der Baan, F. Bickelhaupt, Tetrahedron Lett. 1986, 27, 6267–6270; d) L. E. Overman, A. F. Renaldo, J. Am. Chem. Soc. 1990, 112, 3945–3949; e) N. J. Kerrigan, C. J. Bungard, S. G. Nelson, Tetrahedron 2008, 64, 6863–6869.
- [5] a) K. Akiyama, K. Mikami, *Tetrahedron Lett.* 2004, 45, 7217–7220; b) E. C. Linton, M. C. Kozlowski, J. Am. Chem. Soc. 2008, 130, 16162–16163; c) M. E. Geherty, R. D. Dura, S. G. Nelson, J. Am. Chem. Soc. 2010, 132, 11875–11877; d) T. Cao, E. C. Linton, J. Deitch, S. Berritt, M. C. Kozlowski, J. Org. Chem. 2012, 77, 11034–11055.
- [6] R. J. Felix, D. Weber, O. Gutierrez, D. J. Tantillo, M. R. Gagné, *Nat. Chem.* 2012, 4, 405–409.
- [7] a) K. Takai, I. Mori, K. Oshima, H. Nozaki, *Tetrahedron Lett.* 1981, 22, 3985–3988; b) B. M. Trost, G. M. Schroeder, *J. Am. Chem. Soc.* 2000, 122, 3785–3786; c) M. Hiersemann, L. Abraham, Org. Lett. 2001, 3, 49–52; d) G. Koch, P. Janser, G. Kottirsch, E. Romero-Giron, *Tetrahedron Lett.* 2002, 43, 4837–4840; e) J. Rehbein, S. Leick, M. Hiersemann, *J. Org. Chem.* 2009, 74, 1531–1540.
- [8] a) K. Maruoka, S. Saito, H. Yamamoto, J. Am. Chem. Soc. 1995, 117, 1165–1166; b) L. Abraham, R. Czerwonka, M. Hiersemann, Angew. Chem. Int. Ed. 2001, 40, 4700–4703; Angew. Chem. 2001, 113, 4835–4837; c) L. Abraham, M. Körner, M. Hiersemann, Tetrahedron Lett. 2004, 45, 3647–3650; d) J.-C.

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Marié, Y. Xiong, G. K. Min, A. R. Yeager, T. Taniguchi, N. Berova, S. E. Schaus, J. A. Porco, Jr., *J. Org. Chem.* **2010**, *75*, 4584–4590; e) J. Tan, C.-H. Cheon, H. Yamamoto, *Angew. Chem. Int. Ed.* **2012**, *51*, 8264–8267; *Angew. Chem.* **2012**, *124*, 8389–8392.

- [9] a) C. Uyeda, E. N. Jacobsen, J. Am. Chem. Soc. 2008, 130, 9228 –
 9229; b) C. Uyeda, E. N. Jacobsen, J. Am. Chem. Soc. 2011, 133, 5062 5075.
- [10] For alternative catalytic approaches to the Claisen rearrangement, see: a) T. P. Yoon, D. W. C. MacMillan, *J. Am. Chem. Soc.* 2001, *123*, 2911–2912; b) L. Candish, D. W. Lupton, *Chem. Sci.* 2012, *3*, 380–383.
- [11] W. G. Dauben, A. Chollet, *Tetrahedron Lett.* 1981, 22, 1583– 1586.
- [12] P. P. Painter, B. M. Wong, D. J. Tantillo, Org. Lett. 2014, 16, 4818-4821.
- [13] K. A. Ahrendt, C. J. Borths, D. W. C. MacMillan, J. Am. Chem. Soc. 2000, 122, 4243–4244.
- [14] a) J. L. Cavill, J.-U. Peters, N. C. O. Tomkinson, *Chem. Commun.* 2003, 728–729; b) V. Terrasson, A. van der Lee, R. Marcia de Figueiredo, J. M. Campagne, *Chem. Eur. J.* 2010, *16*, 7875–7880; c) B. P. Bondzic, T. Urushima, H. Ishikawa, Y. Hayashi, *Org. Lett.* 2010, *12*, 5434–5437; d) A. Quintard, A. Lefranc, A. Alexakis, *Org. Lett.* 2011, *13*, 1540–1543; e) E. K. Kemppainen, G. Sahoo, A. Valkonen, P. M. Pihko, *Org. Lett.* 2012, *14*, 1086–1089; f) Y. Wei, N. Yoshikai, *J. Am. Chem. Soc.* 2013, *135*, 3756–3759; g) E. K. Kemppainen, G. Sahoo, A. Piisola, A. Hamza, B. Kótai, I. Pápai, P. M. Pihko, *Chem. Eur. J.* 2014, *20*, 5983–5993.
- [15] a) T. Kano, Y. Tanaka, K. Osawa, T. Yurino, K. Maruoka, *Chem. Commun.* 2009, 1956–1958; b) K. Ishihara, K. Nakano, *J. Am. Chem. Soc.* 2005, *127*, 10504–10505; c) A. Erkkilä, P. M. Pihko, M.-R. Clarke, *Adv. Synth. Catal.* 2007, *349*, 802–806; d) O. Lifchits, C. M. Reisinger, B. List, *J. Am. Chem. Soc.* 2010, *132*, 10227–10229; e) O. Lifchits, M. Mahlau, C. M. Reisinger, A. Lee, C. Farès, I. Polyak, G. Gopakumar, W. Thiel, B. List, *J. Am. Chem. Soc.* 2013, *135*, 6677–6693.

- [16] J. F. Austin, D. W. C. MacMillan, J. Am. Chem. Soc. 2002, 124, 1172–1173.
- [17] a) M. Lemay, W. W. Ogilvie, Org. Lett. 2005, 7, 4141-4144;
 b) M. Lemay, W. W. Ogilvie, J. Org. Chem. 2006, 71, 4663-4666;
 c) J. L. Cavill, R. L. Elliott, G. Evans, I. L. Jones, J. A. Platts, A. M. Ruda, N. C. O. Tomkinson, Tetrahedron 2006, 62, 410-421;
 d) M. Lemay, J. Trant, W. W. Ogilvie, Tetrahedron 2007, 63, 11644;
 e) M. Lemay, L. Aumand, W. W. Ogilvie, Adv. Synth. Catal. 2007, 349, 441-447;
 f) J. B. Brazier, J. L. Cavill, R. L. Elliott, G. Evans, T. J. K. Gibbs, I. L. Jones, J. A. Platts, N. C. O. Tomkinson, Tetrahedron 2009, 65, 9961-9966.
- [18] The corresponding Z-substituted alkene isomerized to the more stable E isomer prior to rearrangement.
- [19] a) K. A. Black, S. Wilsey, K. N. Houk, J. Am. Chem. Soc. 1998, 120, 5622-5627; b) D. A. Hrovat, J. Chen, K. N. Houk, W. T. Borden, J. Org. Chem. 2000, 65, 7456-7460; c) Y. Xia, F. Zhou, Y. Li, W. Li, J. Mol. Struct. THEOCHEM 2009, 904, 69-73.
- [20] Use of triflic acid alone led to decomposition, and a combination of triethylamine and triflic acid did not result in any catalysis.
- [21] DFT calculations at the B3LYP/6-31G* level of theory suggested that the six- and seven-membered-ring iminium ions are about 2.5 and 3.7 kcalmol⁻¹ less stable, respectively, than the fivemembered ring.
- [22] DFT-computed proton affinities: $5a: 220.3 \text{ kcal mol}^{-1}$; $5b: 219.5 \text{ kcal mol}^{-1}$; $5c: 218.9 \text{ kcal mol}^{-1}$; $5d: 218.4 \text{ kcal mol}^{-1}$.
- [23] Note that catalysts 5g and 5h afforded the opposite major enantiomer relative to catalysts 5e and 5f. The absolute stereochemistry of 2g was not assigned.
- [24] We found that camphor-based sulfonyl hydrazide, CaSH (see H. He, B. J. Pei, H.-H. Chou, T. Tian, W.-H. Chan, A. W. M. Lee, *Org. Lett.* 2008, 10, 2421–2424), which is effective in Diels– Alder reactions of α-unsubstituted enals, was inactive in the Cope rearrangement.

Received: July 4, 2016 Published online: ■■ ■■, ■■■

www.angewandte.org



Communications



Communications



An Organocatalytic Cope Rearrangement



Seven-membered catalysts: Diazepane carboxylates catalyze the Cope rear-rangement of 1,5-hexadiene-2-carboxal-dehydes under mild conditions. This process constitutes the first example of an organocatalytic Cope rearrangement and highlights the efficient formation of α -substituted iminium ions with diazepane carboxylates.

6 www.angewandte.org

C 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2016, 55, 1-6

These are not the final page numbers!