ORIGINAL RESEARCH



Synthesis of thiocarbohydrazide and carbohydrazide derivatives as possible biologically active agents

Kiran Gangarapu · Sarangapani Manda · Anvesh Jallapally · Sreekanth Thota · Subhas S. Karki · Jan Balzarini · Erik De Clercq · Harukuni Tokuda

Received: 23 March 2013/Accepted: 25 July 2013 © Springer Science+Business Media New York 2013

Abstract A series of new β -isatin aldehyde-*N*,*N'*-thiocarbohydrazone, bis- β -isatin thiocarbohydrazones, bis- β isatin carbohydrazones was synthesized by condensation of 5-substituted isatin with thiocarbohydrazide or carbohydrazide. The chemical structures of the newly synthesized compounds were confirmed by FT-IR, ¹H NMR, and mass spectral analysis. The synthesized compounds were evaluated for in vitro antiviral activity against various strains of DNA and RNA viruses, but exhibited moderate antiviral activity compared with the reference compounds. Among all the compounds **6c** exhibited the highest chemoprevention activity in a two-stage mouse-skin carcinogenesis test.

Keywords Isatin · Thiocarbohydrazide · Carbohydrazide · Antiviral · Anti-influenza · Carcinogenesis

K. Gangarapu (🖂)

Department of Pharmaceutical Chemistry, Kakatiya Institute of Pharmaceutical Sciences, Pembarthy (V), Hasanparthy (M), Warangal 506 371, AP, India e-mail: gangakiran1905@gmail.com

K. Gangarapu

Center for Pharmaceutical Sciences, IST, JNTU, Kukatpally, Hyderabad 500085, India

S. Manda

Department of Pharmaceutical Chemistry, University College of Pharmaceutical Sciences, Kakatiya University, Warangal 506 009, AP, India

A. Jallapally

Department of Pharmaceutical Chemistry, Acharya and BM Reddy College of Pharmacy, Soldevanahalli, Bangalore, KN, India

Introduction

Isatin is a versatile lead molecule for designing potential antiviral agents. Isatin derivatives display broad spectrum pharmacological properties, including cytotoxic (Reddy et al., 2012; Alcaide et al., 2012), antibacterial (Akhaja and Raval, 2011), antifungal (Reddy et al., 2011), antiviral (Kang et al., 2011; Bacani et al., 2011; Sin et al., 2009), anti-HIV (Bal et al., 2005), and antimalarial (Hans et al., 2011) activities. The first clinically approved antiviral agent, *N*-methylisatin- β -thiosemicarbazone (methisazone) was active against smallpox and vaccinia viruses (Bauer, 1985). N,N'-disubstituted thiosemicarbazone derivatives of isatin were inhibitory to HIV-1 replication (Teitz and Ronen, 1984). N-methylisatin-β-thiosemicarbazone is an inhibitor of Japanese encephalitis virus infection (Sebastain et al., 2008). The general configuration of the isatin molecule resembles that of the purines and also tryptophan and

S. Thota

Department of Chemistry, Colorado State University, Fort Collins 80523, CO, USA

S. S. Karki Department of Pharmaceutical Chemistry, KLE University's College of Pharmacy, Rajajinagar, Bangalore, KN, India

J. Balzarini · E. De Clercq Rega Institute for Medical Research, KU Leuven, Leuven, Belgium

H. Tokuda Department of Complementary and Alternative Medicine, Clinical R&D, Graduate School of Medical Science Kanazawa University, Kanazawa, Japan 2007). Viral pathogens are responsible for many human diseases and examples of diseases caused by viruses include the common cold, smallpox, AIDS, cold sores, human herpes virus, papilloma virus, etc. (Prado-Prado et al., 2009). The worldwide market for antiviral drugs is estimated to be approximately 20 billion US dollars and is expected to grow rapidly as new therapies become available. The need for effective antiviral drugs is further emphasized by the lack of vaccines for most respiratory tract virus infections [adenovirus, rhinovirus, parainfluenza virus and respiratory syncytial virus (RSV), the widely occurring human papilloma viruses, and herpes simplex virus (HSV), varicella-zoster virus (VZV), Epstein-Barr virus, cytomegalovirus (CMV), and the vast array of hemorrhagic fever viruses (De Clercq, 2002)]. Millions of people worldwide are affected by infectious diseases caused by viruses. Further, widespread viral resistance has renewed the interest in the quest for new antiviral agents. Earlier work could clearly define the structural

caspase, MAO and COX-2 enzymes (Medvedev et al.,

requirements necessary for the retention of antiviral activity by heterocyclic thiosemicarbazone in neurovaccinia infections in mice. Viral infections primarily affect the respiratory and gastrointestinal system; epithelium, mucous membranes and endothelium of skin, mouth and genitals; lymphoid tissue, liver and other organs; and the central nervous system (Bruce *et al.*, 2007).

Recently, we have reported that β -isatin-N,N'-thiocarbohydrazone derivatives could have antimicrobial and antioxidant activity (Kiran et al., 2012a). Thiosemicarbazones were the first antiviral compounds recognized to possess broad-spectrum antiviral activity against a range of DNA and RNA viruses [Castro et al., 2011]. Earlier reports have described antiviral activity of isatin-β-thiosemicarbazone and isatin-B-semicarbohydrazone against different viral strains [Quenelle et al., 2006; Pandeya et al., 2005]. However, the biological activities of isatin-β-thiocarbohydrazone and isatin-\beta-carbohydrazone have not been evaluated against DNA and RNA viruses. In connection with our ongoing work on isatin (Kiran et al., 2012b), we are now reporting a simple and efficient method for the synthesis of some β -isatin-N.N'-thiocarbohydrazone (4ag), bis- β -isatin thiocarbohydrazone, and bis- β -isatin thiocarbohydrazone derivatives (6a-e) as depicted in Schemes 1 and 2. These compounds were screened for in vitro







Scheme 2 Schematic steps of Bis-isatin derivatives synthesis

antiviral activity and also in a two stage mouse-skin carcinogenesis test.

Experimental

Materials and methods

Melting points (m.p.) were determined in open capillaries, using the Toshniwal melting point apparatus, expressed in °C and were uncorrected. FT-IR spectra (KBr) were recorded on a thermo Nicolet Nexus 670S series. NMR spectra were recorded on an Avance-300 MHz in DMSO d_6 solvent using TMS as an internal standard (chemical shifts in δ , ppm). Mass spectra were obtained on a LC-MSD-Trap-SL. Reactions were monitored using thin layer chromatography (TLC) on aluminum-backed precoated silica gel 60 F₂₅₄ plates (E Merck). The purity of the compounds was checked on silica gel-coated aluminum sheets by TLC. Column chromatography was performed by using Qualigen's silica gel (60-120 mesh). All the solvents were AR grade and distilled before use. 5-Substituted isatins were purchased from Himedia (Mumbai, India). Indole-3-carboxaldehyde was obtained from E. Merck (India). All substituted aldehydes were purchased from Sd. Fine Chemicals.

General procedure

Synthesis of isatin derivatives

The synthesis of isatin derivatives was carried out using a modified Sandmeyer methodology (Henry and Blatt, 1964). *N*-benzyl isatin is prepared by a microwave method as reported in the literature (Shmidt *et al.*, 2008). All the isatin derivatives exhibit same physical properties as reported in the literature.

Synthesis of thiocarbohydrazide (1)

The synthesis of thiocarbohydrazide was carried out by the Taguchi method. The physical data of the synthesized

6a: R= H; **6b:**R=Cl; **6c:** R= CH₃; **6d:**R=NO₂; **6e**=H, X=S

compound are comparable and show m.p 170–172 °C similar to that reported in the literature (Zhou *et al.*, 2010).

Synthesis of monothiocarbohydrazones (2a-g)

An equimolar mixture of thiocarbohydrazide (0.01 mol), substituted aldehyde (0.01 mol), and a few drops of glacial acetic acid were taken in 30 ml of ethanol in 250 ml conical flask. The resulting mixture was heated under reflux for 1 h. Then the reaction mixture was cooled overnight and the precipitate was collected by filtration. Further it was purified by recrystallization with ethanol. The synthesized compounds were comparable with the reported literature (Mohamed *et al.*, 1991).

Synthesis of β -isatin aldehyde-N,N'-thiocarbohydrazones (4a-g)

To a solution of monothiocarbohydrazone (2a-g, 0.01 mol) and substituted isatin (3, 0.01 mol) in ethanol, a few drops of glacial acetic acid were added to initiate the reaction. The progress of the reaction was monitored by TLC. The residue was cooled, kept overnight and the precipitate was collected by filtration. The solid was purified by column chromatography, and subsequently by recrystallization with ethanol.

Synthesis of bis-isatin derivatives (6a-e)

A mixture of thiocarbohydrazide or carbohydrazide (0.01 mol) was mixed with appropriate substituted isatin derivatives (0.02 mol) in ethanol–acetic acid (1:1). The reaction mixture was refluxed for 2 h. Then the reaction mixture was cooled, kept overnight and the precipitate was collected by filtration. The resulting solid was purified by column chromatography and subsequently by recrystallization with ethanol.

Spectral data of the synthesized compounds

3-[(1H-indol-3-ylmethylidene)amino]-1-{[(3Z)-2-oxo-2, 3-dihydro-1H-indol-3-ylidene]amino}thiourea (4a) IR (v cm⁻¹): 3226 (br, N–H, lactam), 3138, 3070 (w, C–H, Ar), 1705 (s, C=O, lactam), 1337 (s, C=S), 1538 (s, C=N), 1463 (s, C=C). ¹H-NMR (DMSO- d_6) δ ppm: 14.1 (s, 1H, NH), 12.2 (s, 1H, NH), 11.7 (s, 1H, NH), 11.3 (s, 1H, NH), 8.67 (1H, =CH–), 8.47 (1H, N=CH), 7.87 (dd, J = 7.4, 1.6 Hz, 1H), 7.64–7.52 (m, 3H), 7.49–7.40 (m, 2H), 7.34 (td, J = 7.4, 1.5 Hz, 1H), 6.59 (d, J = 7.5 Hz, 1H). ¹³C NMR (DMSO- d_6) δ ppm: 173.9 (C=S), 162 (C=O), 143 (C=N), 110–137 (Ar–C); mass (ESI–MS): m/z 363 (M+1, 100 %), 385 (M+Na, 80 %). Elemental analysis for C₁₈H₁₄N₆OS: calculated (%) C: 59.65; H: 3.89; N: 23.19; found: C: 59.29; H: 3.24; N: 23.45.

3-{[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino}-1-[thiophen-2-ylmethylidene)amino]thiourea (4b) IR (v cm⁻¹): 3316 (N–H), 3026 (w, C–H, Ar), 1708 (s, C=O, lactam), 1368 (s, C=S), 1395 (s, C=N), 1528 (s, CH=N). ¹H-NMR (DMSO- d_6) δ ppm: 14.7 (s, 1H, NH), 13.3 (s, 1H, –NH), 11.9 (s, 1H, –NH), 8.90 (s, 1H), 7.65–7.59 (m, 2H), 7.57–7.45 (m, 2H), 7.31 (dd, J = 7.5, 2.8 Hz, 1H), 7.27–7.20 (m, 1H), 6.96 (dd, J = 7.5, 1.5 Hz, 1H); mass (ESI–MS): *m*/z 330 (M+1, 100 %). Elemental analysis for C₁₄H₁₁N₅OS₂: calculated (%) C: 51.05; H: 3.37; N: 21.26; found: C: 51.02; H: 3.34; N: 21.18.

l-(*{*[4-(dimethylamino)phenyl]methylidene]amino)-3-{[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino} thiourea (4c) IR (v cm⁻¹): 3218 (−NH), 3016 (w, C−H, Ar), 2700 (C−H, str), 1702 (s, C=O, lactam), 1358 (s, C=S), 1528 (s, C=N), 1528 (s, CH=N). ¹H-NMR (DMSO- d_6) δ ppm: 14.9 (s, 1H, −NH), 14.5 (s, 1H, −NH), 13.0 (s, 1H, −NH), 8.06 (s, 1H), 7.76 (dd, J = 7.5, 1.5 Hz, 1H), 7.61 (dd, J = 7.5, 1.7 Hz, 1H), 7.53–7.46 (m, 3H), 7.28–7.20 (m, 1H), 6.85–6.79 (m, 2H), 3.02 (s, 6H); mass (ESI–MS): m/z 367 (M+1, 100 %), 389 (M+Na, 40 %). Elemental analysis for C₁₈H₁₈N₆OS: calculated (%) C: 59.00; H: 4.95; N: 22.93; found: C: 59.02; H: 4.86; N: 22.78.

I-{[(4-Nitrophenyl)methylidene]amino}-3-{[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino}thiourea (4d) IR (ν cm⁻¹): 3120 (–NH), 3006 (w, C–H, Ar), 1690 (s, C=O, lactam), 1358 (s, C=S), 1528 (s, C=N), 1528 (s, CH=N). ¹H-NMR (DMSO-d₆) δ ppm: 14.2 (s, 1H,–NH), 13.6 (s, 1H, –NH), 12.0 (s, 1H, –NH), 8.06 (s, 1H), 7.64 (dd, J = 7.5, 1.5 Hz, 1H), 7.64 (dd, J = 7.5, 1.7 Hz, 1H), 7.42–7.26 (m, 3H), 7.28–7.20 (m, 1H), 6.85–6.79 (m, 2H); mass (ESI-MS): m/z 369 (M+1, 100 %), 391 (M+Na, 30 %). Elemental analysis for C₁₆H₁₂N₆O₃S: calculated (%) C: 52.17; H: 3.28; N: 22.81 found: C: 52.12; H: 3.24; N: 22.76.

1-{[(4-Chlorophenyl)methylidene]amino}-3-{[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino}thiourea (4e) IR (v cm⁻¹): 3345 (N–H), 3050 (C–H, Ar), 1630 (C=O, lactam), 900–690 (Ar–H), 785 (C–Cl); ¹H-NMR (DMSO- d_6) δ ppm; 14.7 (s, 1H), 13.6 (s, 1H), δ 11.2 (s, 1H), 7.93 (s, 1H), 7.85–7.79 (m, 2H), 7.68 (dd, J = 7.5, 1.6 Hz, 1H), 7.60 (dd, J = 7.4, 1.6 Hz, 1H), 7.48 (td, J = 7.4, 1.5 Hz, 3H), 7.27–7.20 (m, 1H); mass (ESI–MS): m/z 357 (M+, 100 %), 359 (M+2, 38 %). Elemental analysis for C₁₆H₁₂N₆O₃S: calculated (%) C: 53.71; H: 3.38; N: 19.57; found: C: 53.68; H: 3.42; N: 19.76.

I-{[(3,4-Dimethoxyphenyl)methylidene]amino}-3-{[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino}thiourea (*4f*) IR (ν cm⁻¹): 3348 (N–H), 3020 (C–H, Ar), 1648 (C=O), 1313 (C=S); ¹H-NMR (DMSO-*d*₆) δ ppm; 14.7 (s, 1H), 13.1 (s, 1H), δ 10.9 (s, 1H), 7.93 (s, 1H), 7.71–7.57 (m, 4H), 7.48 (td, J = 7.5, 1.5 Hz, 1H), 7.14–7.08 (m, 2H), 3.79 (s, 6H); mass (ESI–MS): *m*/*z* 384 (M+1, 100 %). Elemental analysis for C₂₄H₂₁N₅OS: calculated (%) C: 56.38; H, 4.47; N, 18.27; found: C: 56.34; H: 4.46; N: 18.23.

3-{[(3Z)-1-benzyl-2-oxo-2,3-dihydro-1H-indol-3-ylidene] amino}-1-[(phenylmethylidene)amino]thiourea (**4g**) IR (ν cm⁻¹): 3360 (N–H), 3050 (C–H, Ar), 1630 (C=O, lactam), 900–690 (Ar, oop); ¹H-NMR (DMSO-d₆) δ ppm; 14.7 (s, 1H), 13.6 (s, 1H), δ 11.2 (s, 1H), 7.93 (s, 1H), 6.8–7.5 (Ar–H), 3.2 (s, –CH₂, 2H); mass (ESI–MS): *m*/ z 428 (M+, 100 %), 450 (M+Na, 38 %). Elemental analysis for C₁₈H₁₇N₅O₃S: calculated (%) C: 66.81; H: 4.63; N: 16.94; found: C: 66.78; H: 4.58; N: 16.78.

1,3-Bis({[(3Z)-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino})urea (**6a**) IR (ν cm⁻¹): 3350–3050 (NH, NH), 3070 (C–H, Ar), 1742 (s, C=O, lactam), 1608, 1472 (C=C, Ar), 1650 (C=O), 1610 (C=N). ¹H-NMR (DMSO-*d*₆) ppm; δ 13.2 (s, 2H), 7.93 (s, 2H), 7.15–7.79 (m, 6H); mass (ESI-MS): *m*/*z* 349 (M+1, 100 %), 371 (M+Na, 52 %). Elemental analysis for C₁₇H₁₂N₆O₃: calculated (%) C:58.62; H:3.47; N:24.13; found: C: 58.58; H: 3.46; N: 24.20.

1,3-Bis(*{*[(*3Z*)-5-*chloro-2-oxo-2,3-dihydro-1H-indol-3-ylidene]amino})urea (6b) IR (ν cm⁻¹): 3250–3050 (NH, NH), 3010 (C–H, Ar), 1642 (s, C=O, lactam), 1608, 1472 (C=C, Ar), 1650 (C=O), 1610 (C=N). ¹H-NMR (DMSO-<i>d*₆) ppm; δ 13.2 (s, 2H), 7.93 (s, 2H), 7.15–7.79 (m, 6H); mass (ESI–MS): m/z 417 (M+1, 100 %), 439 (M+Na, 46 %).Elemental analysis for C₁₇H₁₀Cl₂N₆O₃: calculated (%) C:48.94; H:2.42;N: 20.14; found: C: 48.89; H: 2.46; N: 20.20.

 $\begin{array}{l} 1,3-Bis(\{[(3Z)-5-methyl-2-oxo-2,3-dihydro-1H-indol-3-yli-dene]amino\})urea(6c) & IR (v \ cm^{-1}) \ 3198 \ (NH), \ 3010 \ (C-H, \ Ar), \ 2980 \ (-CH \ str), \ 1730 \ (C=O), \ 1680 \ (s, \ C=O, \ lactam), \ 1608, \ 1472 \ (C=C, \ Ar), \ 1620 \ (C=N). \ ^1H-NMR \end{array}$

(DMSO- d_6) ppm; δ 11.2 (s, 2H), 10.2 (s, 2H), 7.15–7.79 (m, 6H) 2.2 (s, -CH₃); mass (ESI–MS): m/z 377 (M+1, 100 %), 386 (M+Na, 51 %). Elemental analysis for C₁₉H₁₆ N₆O₃: calculated (%) C: 60.63; H: 4.28; N: 22.33; found: C: 60.69; H: 4.26; N: 22.32.

1,3-Bis(*{*[(*3Z*)-5-*nitro-2-oxo-2,3-dihydro-1H-indol-3-ylidene] amino}*)*urea* (*6d*) IR (ν cm⁻¹): 3097 (NH), 3050 (C–H, Ar), 1702 (C=O), 1618 (s, C=O, lactam), 1608, 1472 (C=C, Ar), 1626 (C=N) 1355 (–NO₂). ¹H-NMR (DMSO-*d*₆) ppm; δ 13.5 (s, 2H), 11.5 (s, 2H), 7.15–7.79 (m, 6H); mass (ESI– MS): *m/z* 439 (M+1, 100 %). Elemental analysis for C₁₇H₁₀ N₈O₇: calculated (%) C: 46.58; H: 2.30; N: 25.56; found: C: 46.53; H: 2.26; N: 25.54.

1,3-Bis(*{*[(*3Z*)-2-*oxo*-2,3-*dihydro*-1*H*-*indo*]-3-*y*lidene]amino})) thiourea (*6e*) IR (ν cm⁻¹): 3150, 3010 (NH, NH) 1650 (C=O), 1610 (C=N), 1350 (NCSN). ¹H-NMR (DMSO-*d*₆) δ ppm; δ 13.5 (s, 2H), 7.93 (s, 2H), 7.15–7.79 (m, 6H); mass (ESI–MS): *m*/*z* 365 (M+1, 100 %). Elemental analysis for C₁₇H₁₂ N₆O₂S: calculated (%) C: 56.04; H: 3.32; N: 23.06; found: C: 56.03; H: 3.36; N: 23.04.

Biological activities

Antiviral activity assays

The synthesized compounds were evaluated for antiviral activity according to well-established procedures by De

Fig. 1 Inhibitory effects of compounds 6b and 6c on mouse-skin carcinogenesis induced by DMBA and TPA.
a Positive control [DMBA (390 nmol) + TPA (1.7 nmol)];
b TPA + 85 nmol of 6b;
c TPA + 85 nmol of 6c

Clercq (1981, 1986). The compounds were evaluated for antiviral activity by using the following viruses: human cytomegalovirus (HCMV) strains AD-169 and Davis, herpes simplex virus type 1 (HSV-1) strain KOS, thymidine kinasedeficient (TK⁻), HSV-1 KOS strain resistant to ACV (ACV^r), herpes simplex virus type 2 (HSV-2) strain G, vaccinia virus, vesicular stomatitis virus (VSV), VZV strain Oka, TK⁻ VZV strain 07-1, RSV strain Long, VSV, coxsackie B4, parainfluenza 3, Reovirus-1, Sindbis, Punta Toro, feline coronavirus (FIPV: feline infectious peritonitis virus), influenza A virus subtypes H1N1 and H3N2, and influenza B virus. Antiviral assays were carried out in Crandell-Rees feline kidney cells (feline corona virus and feline herpes virus), HeLa cell cultures (vesicular stomatitis virus, coxsackie virus B4, and RSV), human embryonic lung (HEL) cell cultures [HSV-1 (KOS), HSV-2 (G), vaccinia virus, vesicular stomatitis virus and HSV-1 TK⁻ KOS ACV^r], Madin Darby canine kidney (MDCK) cells (influenza A H1N1 subtype, influenza A H3N2 subtype, and influenza B) and Vero cell cultures (parainfluenza-3 virus, reovirus-1, sindbis virus, coxsackie virus B4, and Punta Toro virus). The HEL, Vero, and HeLa cell lines used in this study were monitored for mycoplasma contamination and were found to be mycoplasma-free.

Screening for inhibition of virus-induced cytopathic effect in vitro

Confluent cell cultures in microtiter 96-well plates were inoculated with 100 CCID₅₀ of virus (1 CCID₅₀ being the





virus dose to infect 50 % of the cell cultures) or with 20 plaque forming units (PFU) (for VZV) and the cell cultures were incubated in the presence of varying concentrations of the test compounds. Viral cytopathicity or plaque formation (VZV) was recorded as soon as it reached completion in the control virus-infected cell cultures that were not treated with the test compounds. Antiviral activity was expressed as the EC_{50} or compound concentration required to reduce virus-induced cytopathicity or viral plaque formation by 50 %.

Cytotoxicity assays

Cytostatic activity were based on the inhibition of HEL cell growth. HEL cells were seeded at 5×10^3 cells/well into 96-well microtiter plates and allowed to proliferate for 24 h. Then, medium containing different concentrations of the test compounds was added. After 3 days of incubation at 37 °C, the cell number was determined with a Coulter

counter. The cytostatic concentration was calculated as CC_{50} (the compound concentration required to reduce cell growth by 50 % relative to the number of cells in the untreated controls). CC_{50} values were estimated from graphic plots of the number of cells (percentage of control) as a function of the concentration of the test compounds. Cytotoxicity was expressed as minimum cytotoxic concentration (MCC) or the compound concentration that causes a microscopically detectable alteration of cell morphology.

Anti-influenza virus activity

The synthesized compounds were evaluated for their antiviral activity against three influenza virus subtypes [A/ Puerto Rico/8/34 (H1N1); A/Hong Kong/7/87 (H3N2), and B/Hong Kong/5/72]. Antiviral activity was estimated from the inhibitory effect on virus-induced cytopathic effect, as determined by microscopical examination and/or the

Table 1 Inhibitory activity of synthesized isatin derivatives against DNA viruses

S. no.	Compounds	Human embryonic cell culture (HEL) EC_{50}^{a} (μM)							Crandell–Rees feline kidney cell culture (CRFK)		Human embryonic cell culture (HEL)	
		Cytomegalovirus		Herpes simplex	Herpes simplex	Vaccinia virus	Vesicular stomatitis	Herpes simplex	Feline corona	Feline herpes	Varicella-zoster virus (VZV)	
		AD-169 strain	Davis strain	virus-1 (KOS)	virus-2 (G)		virus	virus-ITK- KOS ACV ^r	virus	virus	TK + VZV strain	TK- VZV strain
1	4a	12 ± 3.5	15 ± 0.4	>20	>20	>20	>20	>20	>20	>20	>4	7.4
2	4b	16 ± 3.7	11.32 ± 0.08	>20	>20	>20	>20	>20	>20	>20	>20	>20
3	4c	>20	>20	>100	>100	>100	>100	>100	>100	>100	>100	>100
4	4d	>20	>20	>100	>100	>100	>100	>100	>100	>100	>20	>20
5	4e	>20	>20	>100	>100	>100	>100	>100	>100	>100	>20	>20
6	4f	>20	>20	>20	>20	>20	>20	>20	>100	>100	>20	>20
7	4g	>20	>100	>100	>100	>100	>100	>100	>100	>100	>100	>20
8	6a	>20	>100	>100	>100	>100	>100	>100	>100	>100	>20	>100
9	6b	>20	>20	>100	>100	>100	>100	>100	>100	19.7	>20	>20
10	6c	>100	>100	>100	>100	>100	>100	>100	>100	55.1	>100	>100
11	6d	>100	>100	>100	>100	>100	>100	>100	>100	>100	>20	>20
12	6e	>20	>20	>100	>100	>100	>100	>100	>100	>100	>20	>20
13	Ganciclovir	8.1 ± 0.2	5.5 ± 2.36	0.02	0.01	>100	>100	4	>100	1.4	-	-
14	Cidofovir	1.0 ± 0.05	0.78 ± 0.49	2	1	22	>250	0.9	_	_	-	-
15	Brivudin	-	_	0.04	112	4	>250	50	-	-	2.1	143
16	Acyclovir	-	_	0.4	0.2	>250	>250	22	_	_	0.019	104.5
17	$\begin{array}{c} HHA \\ (\mu g \ ml^{-1}) \end{array}$	-	-	-	-	-	-	-	4.2	11.0	-	-
18	$\begin{array}{c} UDA \\ (\mu g \ ml^{-1}) \end{array}$	-	_	-	-	-	-	-	1.6	1.2	-	-

"-" not performed

 $^{\rm a}$ Required to reduce virus-induced cytopathogenicity by 50 %

formazan-based MTS cell viability test. The EC₅₀ (50 % Effective concentration), or concentration producing 50 % inhibition of virus-induced cytopathic effect, was determined by visual scoring of the CPE, or by measuring the cell viability with the colorimetric formazan-based MTS assay. Cytotoxicity of the test compounds was expressed as the compound concentration causing minimal changes in cell morphology (MCC), or the concentration causing 50 % cytotoxicity (CC₅₀), as determined by the MTS assay.

Two-stage mouse-skin carcinogenesis test

Specific pathogen-free female ICR mice (6 weeks old, body weight approx. 30 g) were obtained from Japan SLC Inc., Shizuoka, Japan, and the animals were housed, five per polycarbonate cage, in a temperature-controlled room at 24 ± 2 °C and given food and water ad libitum throughout the experiment. Animals were divided into three experimental groups containing 15 mice each. The back ($2 \times 8 \text{ cm}^2$) of each mouse was shaved with surgical clippers, and the mice were topically treated with 7,12dimethyl benz[a]anthracene (DMBA) (100 µg, 390 nmol) in acetone (0.1 ml) an initiating treatment. One week after the initiation, papilloma formation was promoted twice weekly by the application of 12-*O*-tetradecanoylphorbol-13-acetate (TPA) (1 µg, 1.7 nmol) in acetone (0.1 ml) to the skin (Fig. 1). One hour before each treatment with TPA, the mice were treated with the samples (85 nmol) in acetone (0.1 ml). The incidence of papillomas was examined weekly over a period of 20 weeks (Tanaka *et al.*, 2001), the percentages of mice bearing papillomas (Fig. 2a) and the average number of papillomas per mouse (Fig. 2b) were recorded.

Results and discussion

A series of new β -isatin aldehyde-N.N'-thiocarbohydrazone, bis-β-isatin thiocarbohydrazones, and bis-β-isatin carbohydrazones derivatives was synthesized by microwave-oriented reaction by two-step procedure as depicted in Schemes 1 and 2. The synthesized compounds were purified by column chromatography. All of the derivatives were characterized by FT-IR, ¹H NMR, and mass spectral data. The FT-IR spectra of the compounds revealed absorption bands in the region $1,509-1,610 \text{ cm}^{-1}$ corresponding to C=N stretching bands. Absence of carbonyl (C=O) peak around 1.715 cm^{-1} revealed the formation of Schiff bases. The ¹H NMR spectrum of the synthesized compounds showed only one set of signals for the aromatic protons around 6.73-7.80 ppm, while the -NH of the isatin and -NH of hydrazones gave rise to two different signals in the 11-14 ppm range.

 Table 2
 Inhibitory activity of synthesized isatin derivatives against RNA Viruses

S.	Compound	Antiviral activity EC_{50}^{*} (μM)									
no.		HeLa cell culture	es	Vero cell cultures							
		Vesicular stomatitis virus	Coxsackie virus B4	Respiratory syncytial virus	Para-influenza- 3 virus	Reovirus- 1	Sindbis virus	Coxsackie virus B4	Punta Toro virus		
1	4 a	>20	>20	>20	>20	>20	>20	>20	>20		
2	4b	>4	>4	>4	>20	>20	>20	>20	>20		
3	4c	>100	>100	>100	>100	>100	>100	>100	>100		
4	4d	>100	>100	>100	>20	>20	>20	>20	>20		
5	4e	>100	>100	>100	>100	>100	>100	>100	>100		
6	4f	>100	>100	>100	>20	>20	>20	>20	>20		
7	4g	>20	>20	>20	>100	>100	>100	>100	>100		
8	6a	>100	>100	>100	>20	>20	>20	>20	>20		
9	6b	>100	>100	>100	>20	>20	>20	>20	>20		
10	6c	>100	>100	>100	>20	>20	>20	>20	>20		
11	6d	>100	>100	>100	>100	>100	>100	>100	>100		
12	6e	>100	>100	>100	>20	>20	>20	>20	>20		
13	$\begin{array}{c} \text{DS5,000} \\ (\mu g \ ml^{-1}) \end{array}$	1	>100	0.8	>100	>100	100	>100	>100		
14	(S)-DHPA	>250	>250	>250	>250	>250	>250	>250	>250		
15	Ribavirin	4	112	4	85	146	>250	250	25		

^a Required to reduce virus-induced cytopathogenicity by 50 %

Antiviral activity

The novel β -isatin aldehyde-N,N'-thiocarbohydrazones (4a-g) and bis-isatin (6a-e) were evaluated against various strains of DNA viruses (i.e., human cytomegalovirus (HCMV), vaccinia virus (VACV), HSV type 1 (HSV-1) and type 2 (HSV-2), acyclovir-resistant, ACV^r), VZV (Table 1). The test compounds were also evaluated against several RNA viruses (i.e., VSV, coxsackie B4 virus, RSV, parainfluenza virus type 3, reovirus-1, sindbis virus and Punta Toro virus (Table 2). The EC_{50} was calculated as the effective concentration of compound that decreased the percentage of formazan production in compound treated, virus-infected cells to 50 % of that produced by compound-free uninfected cells. The 50 % cytostatic concentration (CC_{50}) was calculated as the concentration required to reduce cell growth by 50 %. MCC is the minimum cytotoxic concentration that causes a microscopically detectable alteration of cell morphology. The HEL, Vero, and HeLa cell lines used in this study were regularly monitored for mycoplasma contamination and found to be mycoplasma-free. The antiviral activity was measured for twelve isatin derivatives using a cell-protection assay. The data are recorded in Tables 1, 2, 3, and 4.

There was only moderate activity of the **4a** and **4b** compounds against both strains of HCMV. Although at concentrations close to their toxicity levels, no specific antiviral effects were noted with the evaluated compounds against HSV-1, (HSV-1 strain KOS), HSV-2 (HSV-2 strain G), HSV-1 (ACV resistant KOS strain), vaccinia virus, vesicular stomatitis virus, coxsackie virus B4, respiratory syncytial virus, parainfluenza-3 virus, reovirus-1, sindbis virus, and Punta Toro virus. None of the compounds exhibited specific antiviral effects (i.e., minimal antiviral effective concentration \geq fivefold lower than minimal cytotoxic concentration) against any of the viruses evaluated.

Anti-influenza virus activity

The anti-influenza virus activities of a series of isatin derivatives were determined by measuring their inhibitory effects on virus replication in MDCK cells using three

Table 3 Anti-influenza virus activity and cytotoxicity of compounds in MDCK cell cultures

Compound	Cytotoxicity		Antiviral EC ^c ₅₀ (μM)						
	$\overline{CC_{50}^{a}(\mu M)}$	Minimum cytotoxic concentration ^b	Influenza A I subtype	Influenza A H1N1 subtype		Influenza A H3N2 subtype		Influenza B	
			Visual CPE score	MTS	Visual CPE score	MTS	Visual CPE score	MTS	
4a	51	20	>4	>4	>4	>4	>4	>4	
4b	46	20	>4	>4	>4	>4	>4	>4	
4c	>100	>100	>100	>100	>100	>100	>100	>100	
4d	>100	>100	>100	>100	>100	>100	>100	>100	
4e	>100	>100	>100	>100	>100	>100	>100	>100	
4f	>100	>100	39 ± 5.5	37 ± 25	27 ± 7.0	27 ± 7.3	45 ± 0.0	37 ± 8.75	
4g	>100	>100	>100	>100	>100	>100	>100	>100	
6a	>100	>100	>100	>100	>100	>100	>100	>100	
6b	>100	>100	>100	>100	>100	>100	>100	>100	
6c	46	≥20	>20	>20	>20	>20	>20	>20	
6d	>100	>100	47 ± 2.5	55 ± 26	72 ± 27	33 ± 1.7	47 ± 2.5	38 ± 1.3	
6e	>100	>100	72 ± 27	61 ± 32	72 ± 27	34 ± 9.5	39 ± 5.5	48 ± 31	
Oseltamivir carboxylate	>100	>100	0.4 ± 0.0	0.35 ± 0.2	10.25 ± 9.7	5.45 ± 4.05	6.7 ± 5.3	2.5 ± 0.7	
Ribavirin	>100	>100	2.5 ± 0.5	2.8 ± 0.25	3 ± 1.0	2 ± 0.2	2.4 ± 1.6	2.45 ± 0.25	
Amantadine	>200	>200	5	3.2	0.9	0.8	>200	>200	
Rimantadine	>200	>200	>200	>200	0.4	0.6	>200	>200	

MDCK cells Madin Darby canine kidney cells

^a 50 % Cytotoxic concentration, as determined by measuring the cell viability with the colorimetric formazan-based MTS assay

^b Minimum compound concentration that causes a microscopically detectable alteration of normal cell morphology

^c 50 % Effective concentration, or concentration producing 50 % inhibition of virus-induced cytopathic effect, as determined by visual scoring of the CPE, or by measuring the cell viability with the colorimetric formazan-based MTS assay

	, , ,					
S.no	Compound	Human embryonic lung cell cultures	Crandell-Rees feline kidney cells	HeLa cell cultures	Vero cell cultures	
		Minimum cytotoxic concentration ^a (µM)				
1	4a	60	86	≥20	100	
2	4b	100	63	20	100	
3	4c	≥100	>100	>100	>100	
4	4d	100	>100	>100	100	
5	4e	100	>100	>100	>100	
6	4f	100	100	>100	≥20	
7	4g	≥100	≥100	100	>100	
8	6a	≥100	>100	>100	100	
9	6b	100	>100	>100	100	
10	6с	>100	>100	>100	100	
11	6d	≥100	>100	>100	>100	
12	6e	100	>100	>100	100	
13	Acyclovir	>400	-	-	_	
14	Brivudin	>300	-	-	_	
15	Ganciclovir	>350	>100	_	-	
16	Cidofovir	>300	-	-	_	
17	DS-5000 (µg ml)	-	-	>100	>100	
18	(S)-DHPA	-	-	>250	>250	
19	Ribavirin	-	-	>250	>250	

Table 4 Cytotoxicity for synthesized isatin derivatives in different cell cultures

^a Minimum cytotoxic concentration that causes a microscopically detectable alteration of normal cell morphology



Fig. 2 Inhibition effects of compounds **6b** and **6c** on DMBA–TPAinduced carcinogenesis. Tumor formation in all mice was initiated with DMBA (390 nmol) and promoted with TPA (1.7 nmol) twice weekly, 1 week after initiation. **a** Percentage of mice with papillomas.

strains of the influenza subtype. The antiviral data obtained by microscopic evaluation of the virus-induced CPE were confirmed by the MTS cell viability assay and were expressed as the EC₅₀. The cytotoxicities of the compounds were expressed as the MCC (the compound concentration causing minimal alterations in cell morphology as estimated by microscopy) or the CC₅₀ (the compound



b Average number of papillomas/mouse. *Circle* control [DMBA (390 nmol) + TPA (1.7 nmol)]; *filled square* TPA + 85 nmol of **6b**; *triangle* TPA + 85 nmol of **6c**

concentration causing 50 % reduction in cell viability based on the MTS assay).

None of the twelve analogs were able to inhibit the cytopathic effects of three influenza virus subtypes i.e., H1N1, H3N2, and influenza B (Table 3), except **4f**, **6d**, and **6e** that showed some activity below the non-toxic concentration of 100 μ M.

Two-stage mouse-skin carcinogenesis test

In the present study, we selected two compounds 6b and 6c among the ten compounds to examine their effects on in vivo two-stage carcinogenesis using mouse-skin papillomas induced by DMBA as an initiator and TPA as a promoter. During the test, no significant toxic effects, such as inflammation and lesional damage (edema, erosion, and ulcer), were observed on the areas of mouse skin topically treated with test compounds. In our experiment, the body weight of the animals was not affected by the treatment with test compounds. As shown in Fig. 2a, the positive control (DMBA and TPA) group in papilloma-bearing mice appears rapidly from week 6, and reached 100 % within 10 weeks of promotion. On the other hand, treatment with compound 6c (85 nmol) along with DMBA/TPA reduced the percentage of papilloma-bearing mice to 40-70 % during weeks 10-15, and thereafter 90 % during week 18 and reached 100 % at week 20. As shown in Fig. 2b, in the positive control group, the number of papillomas per mouse formed increased rapidly after week 6 and reached 7.9 papillomas/mouse at week 20. On the other hand, mice treated with compound 6c bore only 5.1 papillomas/mouse over the period of week 20. A moderate protection is observed for the compound 6b. The percentage of papillomas increased rapidly from week 7 and reached 100 % after week 17, and in this group only 1.2, 3.9, and 6.1 papillomas were formed per mouse at 10, 15, and 20 weeks of promotion. From these results, the inhibitory effects of compound 6c on mouse two-stage carcinogenesis induced by DMBA and TPA were apparently more potent than those of compound 6b. Thus, as demonstrated above, the in vivo anti-tumor promoting activity of compound 6c showed stronger chemoprevention activity than that of compound **6b**. These results suggest that isatin lead molecules may be useful as cancer chemopreventive agents.

Conclusion

We have synthesized twelve compounds by reacting substituted/unsubstituted 2,3-dihydro-2,3-dioxindole and thiocarbohydrazide/carbohydrazide in alcohol in the presence of glacial acetic acid. All synthesized compounds were characterized by FT-IR, NMR, and mass spectra. All the compounds were screened for antiviral activity against a various strains of DNA and RNA viruses and some of them, exhibited moderate antiviral activity compared with the reference compounds. Further, the anti-carcinogenic activity of compounds **6b** and **6c** was examined by a two stage-carcinogenesis test using DMBA as initiator and TPA as a promoter. The data suggested that compound **6c** exhibited stronger chemoprevention activity than compound **6b**. Thus these results suggest that isatin lead molecules could also be useful as cancer chemopreventive agents.

Acknowledgments The authors are very much thankful to M. Sadalaxmi, President, Kakatiya Institute of Pharmaceutical Sciences, for providing the laboratory facilities. The antiviral research was performed by grants of the KU Leuven (GOA 10/14). The authors thank Leentje Persoons, Frieda De Meyer, Anita Camps, Lies Van den Heurck and Steven Carmans for excellent technical assistance.

References

- Akhaja TN, Raval JP (2011) 1,3-dihydro-2H-indol-2-ones derivatives: design, synthesis, in vitro antibacterial, antifungal and antitubercular study. Eur J Med Chem 46:5573–5579
- Alcaide B, Almendros P, Aragoncillo C, Campillos GG, Arnó M, Domingo LR (2012) Scandium-catalyzed preparation of cytotoxic 3-functionalized quinolin-2-ones: regioselective ring enlargement of isatins or imino isatins. ChemPlusChem 77:563– 569
- Bacani PMC, Reis MB, Serpeloni JM, Calvo TR, Vilegas W, Varanda EA, Cólus IMS (2011) Mutagenicity and genotoxicity of isatin in mammalian cells in vivo. Mutat Res 719:47–51
- Bal TR, Anand B, Yogeeswari P, Sriram D (2005) Synthesis and evaluation of anti-HIV activity of isatin beta-thiosemicarbazone derivatives. Bioorg Med Chem Lett 15:4451–4455
- Bauer DJ (1985) A history of the discovery and clinical application of antiviral drugs. Br Med Bull 41:309–314
- Bauer DJ, Sadler PW (1960) New antiviral chemotherapeutic agent active against smallpox infection. Lancet 21:1110–1101
- Bruce F, Richard HP, Pamela CC (2007) Lippincott's illustrated reviews: microbiology. Lippincott Williams & Wilkins, Hagertown, pp 47–51
- Castro EF, Fabian LE, Caputto ME, Gagey D, Finkielsztein LM, Moltrasio GY, Moglioni AG, Campos RH, Cavallaro LV (2011) Inhibition of bovine viral diarrhea virus RNA synthesis by thiosemicarbazone derived from 5,6-dimethoxy-1-indanone. J Virol 85:5436–5445
- De Clercq E (2002) Strategies in the design of antiviral drugs. Nat Rev Drug Discov 1:13–25
- DeClercq E, Balzarini J, Torrence PF, Mertes MP, Schmidt CL, Shugar D, Barr PJ, Jones AS, Verhelst G, Walker RT (1981) Thymidylate synthetase as target enzyme for the inhibitory activity of 5-substituted 2'-deoxyuridines on mouse leukemia L1210 cell growth. Mol Pharmacol 19:321
- DeClercq E, Holý A, Rosenberg I, Sakuma T, Balzarini J, Maudgal PC (1986) A novel selective broad-spectrum anti-DNA virus agent. Nature 323:464–467
- Hans RH, Wiid IJF, Helden PDV, Wan B, Frazblau SG, Gut J, Rosenthal PJ, Chilbale K (2011) Novel thiolactone–isatin hybrids as potential antimalarial and antitubercular agents. Bioorg Med Chem Lett 21(7):2055–2058
- Henry G, Blatt AH (1964) Organic synthesis collective, 2nd edn. Wiley, New York, pp 327–334
- Kang IJ, Wang LW, Hsu TA, Yueh A, Lee CC, Lee YC, Lee CY, Chao YS, Shih SR, Chern JH (2011) Isatin-β-thiosemicarbazones as potent herpes simplex virus inhibitors. Bioorg Med Chem Lett 21:1948–1952
- Kiran G, Maneshwar T, Rajeshwar Y, Sarangapani M (2012a) Microwave-assisted synthesis, characterization, antimicrobial and antioxidant activity of some new isatin derivatives. J Chem 2013:1–7. doi:10.1155/2013/192039

- Kiran G, Sarangapani M, Rajeshwar Y, Karki SS, Balzarini J, De Clercq E, Tokuda H (2012b) Microwave assisted synthesis, characterization of some new isatin and thiophene derivatives as cytotoxic and chemopreventive agents. Lett Drug Des Discov 9:934–941
- Medvedev AE, Buneeva O, Glover V (2007) Biological targets for isatin and its analogues: implications for therapy. Biologics 1:151–162
- Mohamed A, Badawy A, Sayed A, Hady A (1991) Reaction of isatin with thiocarbohydrazide: a correction. Arch Pharm (Weinheim) 324:349–351
- Pandeya SN, Smitha S, Jyoti M, Sridhar SK (2005) Biological activities of isatin and its derivatives. Acta Pharm 55:27–46
- Prado-Prado FJ, de la Vega OM, Uriarte E, Ubeira FM, Chou KC, González-Díaz H (2009) Unified QSAR approach to antimicrobials. 4. Multi-target QSAR modeling and comparative multidistance study of the giant components of antiviral drug–drug complex networks. Bioorg Med Chem 17:569–575
- Quenelle DC, Keith KA, Kern ER (2006) In vitro and in vivo evaluation of isatin-beta-thiosemicarbazone and marboran against vaccinia and cowpox virus infections. Antiviral Res 71:24–30
- Reddy BVS, Rajeswari N, Sarangapani M, Reddy GR, Madan Ch, Kumar KP, Rao MS (2011) Iodine-catalyzed conjugate addition of indoles onto en-1,4-dione: a novel synthesis of 3-(1-(1Hindol-3-yl)-2-oxo-2-phenylethyl)indolin-2-ones as antibacterial and antifungal agents. Bioorg Med Chem Lett 21(21):6510–6514
- Reddy BVS, Rajeswari N, Sarangapani M, Prashanthi Y, Jones GR, Anthony A (2012) Iodine-catalyzed condensation of isatin with

indoles: a facile synthesis of di(indolyl)indolin-2-ones and evaluation of their cytotoxicity. Bioorg Med Chem 22:2460–2463

- Sebastain L, Desai A, Shampur MN, Permul Y, Vasanthapuram R (2008) *N*-methylisatin-beta-thiosemicarbazone derivative (SCH 16) is an inhibitor of Japanese encephalitis virus infection in vitro and in vivo. Virol J 22(5):64
- Shmidt MS, Reverdito AM, Kremenchuzky L, Perillo IA, Blanco MM (2008) Simple and efficient microwave assisted N-alkylation of isatin. Molecules 13:831–840
- Sin N, Venables SNBL, Combrink KD, Gulqeze HB, Yu KL, Civiello RL, Thuring J, Wang XA, Yang Z, Zadjura L, Marino A, Kadow KF, Cianci CW, Clarke J, Genovesi ESV, Medina I, Lamb L, Krystal M, Meanwell NA (2009) Respiratory syncytial virus fusion inhibitors. Part 7: structure–activity relationships associated with a series of isatin oximes that demonstrate antiviral activity in vivo. Bioorg Med Chem Lett 19:4857–4862
- Tanaka R, Minami T, Tsujimoto K, Matsunga S, Tokuda H, Nishino H, Terada Y, Yoshitake A (2001) Cancer chemopreventive agents, serratane-type triterpenoids from Picea jezoensis. Cancer Lett 172:119–126
- Teitz Y, Ronen D (1984) Inhibition of the synthesis of Moloney leukemia virus structural proteins by N-methylisatin-beta-4', 4'-diethylthiosemicarbazone. Antimicrob Agents Chemother 26: 913–916
- Zhou J, Wu D, Gou J (2010) Optimization of the production of thiocarbohydrazide using the Taguchi method. Chem Technol Biotechnol 85:1402–1406