This article was downloaded by: [University Library Utrecht] On: 12 November 2012, At: 23:47 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

### α-

Naphthyltriphenylphosphonium Peroxodisulfate: A Selective and Regenerable Reagent for Oxidation of Organic Compounds

Moslem Mansour Lakouraj<sup>a</sup>, Mahmood Tajbakhsh<sup>a</sup> & Farhad Ramzanian-Lehmali<sup>a</sup>

<sup>a</sup> Department of Chemistry, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran Version of record first published: 04 Jun 2008.

To cite this article: Moslem Mansour Lakouraj, Mahmood Tajbakhsh & Farhad Ramzanian-Lehmali (2008):  $\alpha$ -Naphthyltriphenylphosphonium Peroxodisulfate: A Selective and Regenerable Reagent for Oxidation of Organic Compounds, Phosphorus, Sulfur, and Silicon and the Related Elements, 183:6, 1388-1395

To link to this article: http://dx.doi.org/10.1080/10426500701648051

### PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <u>http://www.tandfonline.com/page/terms-and-conditions</u>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



# $\alpha\text{-Naphthyltriphenylphosphonium Peroxodisulfate: A Selective and Regenerable Reagent for Oxidation of Organic Compounds$

### Moslem Mansour Lakouraj, Mahmood Tajbakhsh, and Farhad Ramzanian-Lehmali

Department of Chemistry, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran

Alcohols, hydroquinones, tetrahydropyranyl ethers, trimethylsilyl ethers, oximes, semicarbazones, and phenylhydrazones were efficiently converted to the corresponding carbonyl compounds using  $\alpha$ -naphthyltriphenylphosphonium peroxodisulfate (NTPPPODS) in acetonitrile at reflux conditions. This reagent could be regenerated and reused, which represents significant advantages over the known reagents.

Keywords  $\alpha$ -Naphthyltriphenylphosphonium peroxodisulfate; oximes; phenylhydrazones; semicarbazones; tetrahydropyranyl ethers; trimethylsilyl ethers

### INTRODUCTION

The oxidation of organic compounds under mild conditions has found valuable applications in modern organic synthesis. So far, significant improvments have been achieved by the use of new oxidizing agents.<sup>1–8</sup> Unfortunately, many of the reagents mentioned in literature suffer from at least one of the following disadvantages: cost of preparation, hygroscopicity, photosensitivity, instability, lack of selectivity, and tedious work-up procedure with accompanying reagent decomposition.

The peroxodisulfate ion is one of the strongest oxidizing agents and its standard redox potential in aqueous solution is about 2.01V, but there are only a few reports concerning the capability of peroxodisulfate for oxidative transfomations.<sup>9–16</sup>

In connection with our ongoing work on the application of peroxodisulfate based oxidants to the selective oxidation of organic

Received 9 January 2007; accepted 20 August 2007.

We are thankful to the Mazandaran University Research Council for partial support of this work.

Address correspondence to Moslem Mansour Lakouraj, Department of Chemistry, Faculty of Sciences, University of Mazandaran, P.O. Box 47416, Babolsar, Iran. E-mail: lakouraj@umz.ac.ir



(NTPPPODS)

### FIGURE 1

compounds,<sup>9,17</sup> we report here the oxidation of structurally different alcohols, hydroquinones, tetrahydropyranyl ethers, trimethylsilyl ethers, oximes, semicarbazones, and phenylhydrazones using  $\alpha$ -naphthyltriphenylphosphonium peroxodisulfate in acetonitrile at reflux conditions.  $\alpha$ -Naphthyltriphenylphosphonium peroxodisulfate (NTPPPODS, Figure 1) as an oxidant can overcome most of the abovementioned disadvantages for the oxidation of organic compounds.

### **RESULTS AND DISCUSSION**

 $\alpha$ -Naphthyltriphenylphosphonium peroxodisulfate was readily prepared from an aqueous solution of  $\alpha$ -naphthyltriphenylphosphonium chloride and potassium peroxodisulfate at room temperature and characterized spectroscopically, as well as by elemental analysis. The content of active oxidizing agent (S<sub>2</sub>O<sub>8</sub><sup>2-</sup>) was determined by titrimetric method.<sup>18</sup> This reagent is a stable white powder, which can be stored for months without losing its activity.

The effect of solvent on the oxidation reaction was evaluated by carrying out the oxidation in different solvents. Oxidation of benzyl alcohol with  $\alpha$ -naphthyltriphenylphosphonium peroxodisulfate was carried out in dichloromethane, chloroform, tetrahydrofuran, *n*-hexane, acetonitrile, and in a mixture of water and acetonitrile (70:30) at reflux temperature. The results are presented in Table I.

As indicated in Table I, a 1:0.5 molar ratio of benzyl alcohol and  $\alpha$ naphthyltriphenylphosphonium peroxodisulfate in acetonitrile or in a mixture of water and acetonitrile gave excellent yield of benzaldehyde within five minutes.

In addition, the effect of temperature on this reaction was studied by carrying out the oxidation of benzyl alcohol at lower temperatures; longer reaction time was necessary to achieve complete conversion to benzaldehyde, however.

In order to evaluate the general applicability of this method we have treated benzylic, allylic and aliphatic alcohols,

Solvent	Reaction time (min)	Yield (%) <sup>b</sup>	
CH <sub>3</sub> CN	2	98	
H <sub>2</sub> O/CH <sub>3</sub> CN (70/30)	5	96	
CHCl <sub>3</sub>	60	4	
CH <sub>2</sub> Cl <sub>2</sub>	60	5	
$n-C_6H_{12}$	60	0	
THF	60	0	

TABLE I Oxidation of Benzylal cohol in Different Solvents Using  $\rm NTPPPODS^a$ 

 $^a$ Reactions were carried out at reflux conditions using substrate/oxidant (2:1).

<sup>b</sup>Yields refer to isolated products or GC.

hydroquinones, tetrahydropyranyl and trimethylsilyl ethers with  $\alpha$ -naphthyltriphenylphosphonium peroxodisulfate in refluxing acetonitrile. The results are shown in Table II.

It is evident from Table II, that aromatic substrates, including bezylic alcohols as well as tetrahydropyranyl and trimethylsilyl ethers, were rapidly oxidized to the corresponding carbonyl compounds in excellent yields under the reaction conditions, whereas aliphatic compounds needed longer reaction time for oxidation.

It is noteworthy that, unlike other hydrolytic oxidative methods, the major drawback of overoxidation of the resulting aldehydes was not observed under this reaction conditions.  $\alpha$ , $\beta$ -Unsaturated derivatives are oxidized very efficiently without affecting the olefinic bond and the reaction is essentially chemoselective (Table II).

It is interesting to mention that NTPPPODS can selectively oxidize hydroquinones in the presence of alcohols in refluxing chloroform. When an equimolar mixture of hydroquinone and a benzylic or an aliphatic alcohol was treated with NTPPPODS, only the hydroquinone was oxidized, while the alcohol remained unaffected (Scheme 1).



Substrate	Product	Reagent /substrate	Time (min)	Yield $(\%)^{b,c}$
$C_6H_5CH_2OH$	$C_6H_5CHO$	0.5	2	98
$p-MeOC_6H_4CH_2OH$	$p-MeOC_6H_4CHO$	0.5	$^{2}$	95
o-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	o-ClC <sub>6</sub> H <sub>4</sub> CHO	0.5	3	97
p-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	$p-ClC_6H_4CHO$	0.5	$^{2}$	99
p-BrC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	p-BrC <sub>6</sub> H <sub>4</sub> CHO	0.5	4	97
$p-NO_2C_6H_4CH_2OH$	$p-NO_2C_6H_4CHO$	1	15	65
C <sub>6</sub> H <sub>5</sub> CH(OH)CH <sub>3</sub>	$C_6H_5COCH_3$	0.5	3	95
C <sub>6</sub> H <sub>5</sub> CH(OH)CH <sub>2</sub> CH <sub>3</sub>	$C_6H_5COCH_2CH_3$	0.5	4	97
$(C_6H_5)_2$ CHOH	$(C_6H_5)_2CO$	0.5	$^{2}$	95
$C_6H_5CH_2CH_2OH$	$C_6H_5CH_2CHO$	1	20	93
$CH_3(CH_2)_5CH_2OH$	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CHO	1	20	84
$CH_3(CH_2)_6CH_2OH$	$CH_3(CH_2)_6CHO$	1	20	87
C <sub>6</sub> H <sub>5</sub> CH=CHCH <sub>2</sub> OH	C <sub>6</sub> H <sub>5</sub> CH=CHCHO	1	10	95
Cyclohexanol	Cyclohexanone	1	15	89
Hydroquinone	Quinone	0.5	<b>2</b>	99
tert-Butylhydroquinone	tert-Butylquinone	0.5	$^{2}$	95
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	C <sub>6</sub> H <sub>5</sub> CHO	0.5	3	98
(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> CHOSiMe <sub>3</sub>	$(C_6H_5)_2CO$	0.5	5	96
p-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	p-ClC <sub>6</sub> H <sub>4</sub> CHO	0.5	4	99
p-BrC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	p-BrC <sub>6</sub> H <sub>4</sub> CHO	0.5	3	97
p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CHO	1	40	74
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CHO	1	30	92
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CHO	1	40	81
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>6</sub> CH <sub>2</sub> OSiMe <sub>3</sub>	$CH_3(CH_2)_6CHO$	1	45	84
C <sub>6</sub> H <sub>5</sub> CH=CHCH <sub>2</sub> OSiMe <sub>3</sub>	C <sub>6</sub> H <sub>5</sub> CH=CHCHO	0.5	<b>5</b>	86
Hydroquinone trimethylsilylether	Quinone	0.5	3	94
tert-Butylhydroquinone	tert-Butylquinone	0.5	<b>2</b>	98
trimethylsilylether				
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> OTHP	$C_6H_5CHO$	0.5	10	96
$(\tilde{C}_6H_5)_2$ CHOTHP	$(C_6H_5)_2CO$	0.5	12	95
o-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OTHP	o-ClC <sub>6</sub> H <sub>4</sub> CHO	0.5	8	95
$p-\text{Me}\tilde{C}_6H_4CH_2OTHP$	$p-MeC_6H_4CHO$	0.5	10	94
p-BrC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OTHP	p-BrC <sub>6</sub> H <sub>4</sub> CHO	0.5	8	93
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CH <sub>2</sub> OTHP	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>5</sub> CHO	1	30	87
CH <sub>3</sub> CH=CHCH <sub>2</sub> OTHP	CH <sub>3</sub> CH=CHCHO	0.5	17	91
Cyclohexanol tetrahydropyranyl ether	Cyclohexanone	1	25	90

## TABLE II Oxidation of Alcohols and Hydroquinones and OxidativeDeprotection of Trimethylsilyl and Tetrahydropyranyl Ethers withNTPPPODS $^a$

<sup>a</sup>Reactions were carried out in water and acetonitrile (70/30) at reflux temperature.

<sup>b</sup>All products were characterized spectroscopically (<sup>1</sup>H NMR, IR) and showed physical and spectral data in accordance with their structures and also in accord with those of authentic samples.

<sup>c</sup>Yields refer to pure isolated products; the eluent in column chromarography was EtOAc/*n*-hexane (1:5) for alcohols, and EtOAc/*n*-hexane (1:10) for TMS and THP ethers.

As summarized in Table III, our experiments show, that oximes, phenylhydrazones and semicarbazones can be deprotected to the corresponding aldehydes and ketones in water and acetonitrile (70/30) at reflux temperature.

TABLE III Deprotection of Phenylhydrazones, Oximes and
Semicarbazones to the Corresponding Carbonyl Compounds Using
NTPPPODS <sup>a</sup>

Substrate	Product	Reagent /substrate	Time (min)	Yield $(\%)^{b,c}$
$4C_6H_5CHNNHC_6H_5$	$C_6H_5CHO$	1.5	15	96
$o-ClC_6H_4CHNNHC_6H_5$	$o-ClC_6H_4CHO$	1.5	15	95
$p-\mathrm{ClC}_6\mathrm{H}_4\mathrm{CHNNHC}_6\mathrm{H}_5$	p-ClC <sub>6</sub> H <sub>4</sub> CHO	1.5	12	94
p-MeOC <sub>6</sub> H <sub>4</sub> CHNNHC <sub>6</sub> H <sub>5</sub>	$p-MeOC_6H_4CHO$	1.5	12	96
$p-{ m MeC_6H_4CHNNHC_6H_5}$	$p-MeC_6H_4CHO$	1.5	12	94
$p-NO_2C_6H_4CHNNHC_6H_5$	$p-NO_2C_6H_4CHO$	1.5	45	78
$(C_6H_5)_2CNNHC_6H_5$	$(C_6H_5)_2CO$	1.5	18	95
$C_6H_5CH_2CH_2CHNNHC_6H_5$	$C_6H_5CH_2CH_2CHO$	1.5	40	83
Cyclohexanone Phenylhydrazone	Cyclohexanone	1.5	30	85
C <sub>6</sub> H <sub>5</sub> CH=CHCHNNHC <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> CH=CHCHO	1.5	25	91
C <sub>6</sub> H <sub>5</sub> CHNOH	$C_6H_5CHO$	2	20	96
$p-MeC_6H_4CHNOH$	$p-MeC_6H_4CHO$	2	18	93
p-MeOC <sub>6</sub> H <sub>4</sub> CHNOH	$p-MeOC_6H_4CHO$	2	15	97
m-ClC <sub>6</sub> H <sub>4</sub> C(CH <sub>3</sub> )NOH	m-ClC <sub>6</sub> H <sub>4</sub> COCH <sub>3</sub>	2	20	93
o-ClC <sub>6</sub> H <sub>4</sub> C(CH <sub>3</sub> )NOH	$o-ClC_6H_4COCH_3$	2	18	94
p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> C(CH <sub>3</sub> )NOH	p-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> COCH <sub>3</sub>	2	60	81
$(C_6H_5)_2CNOH$	$(C_6H_5)_2CO$	2	20	97
C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH <sub>2</sub> CHNOH	$C_6H_5CH_2CH_2CHO$	2	40	87
Cyclohexanone Oxime	Cyclohexanone	2	45	85
C <sub>6</sub> H <sub>5</sub> CH=CHCHNOH	C <sub>6</sub> H <sub>5</sub> CH=CHCHO	2	20	91
$C_6H_5CHNNHCONH_2$	$C_6H_5CHO$	1.5	20	96
p-ClC <sub>6</sub> H <sub>4</sub> CHNNHCONH <sub>2</sub>	p-ClC <sub>6</sub> H <sub>4</sub> CHO	1.5	20	91
$o-ClC_6H_4CHNNHCONH_2$	$o-ClC_6H_4CHO$	1.5	20	95
p-BrC <sub>6</sub> H <sub>4</sub> CHNNHCONH <sub>2</sub>	$p ext{-}BrC_6H_4CHO$	1.5	20	93
m-MeOC <sub>6</sub> H <sub>4</sub> CHNNHCONH <sub>2</sub>	m-MeOC <sub>6</sub> H <sub>4</sub> CHO	1.5	15	95
$p-MeOC_6H_4CHNNHCONH_2$	$p-MeOC_6H_4CHO$	1.5	18	92
o-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CHNNHCONH <sub>2</sub>	o-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CHO	1.5	55	<b>78</b>
(C <sub>6</sub> H <sub>5</sub> ) <sub>2</sub> CNNHCONH <sub>2</sub>	$(C_6H_5)_2CO$	1.5	20	96
$C_6H_5CH_2CH_2CHNNHCONH_2$	$C_6H_5CH_2CH_2CHO$	1.5	30	92
$C_6H_5CH=CHCHNNHCONH_2$	$C_6H_5CH=CHCHO$	1.5	25	92

<sup>a</sup>Reactions were carried out in water and acetonitril (70/30) at reflux temperature.

<sup>b</sup>All products were characterized spectroscopically (<sup>1</sup>H NMR, IR) and showed physical and spectral data in accordance with their structures and also in accord with those of authentic samples.

<sup>c</sup>Yields refer to pure isolated products; the eluent in column chromarography was EtOAc/*n*-hexane (1:4).

Similarly, this reagent can selectively deprotect oximes, semicarbazones, and phenylhydrazones in the presence of 2,4dinitrophenylhydrazones to the coreesponding carbonyl compounds in refluxing acetonitrile. When an equimolar mixture of an oxime, a semicarbazone, or a phenylhydrazone and 2,4-dinitrophenylhydrazone was treated with NTPPPODS, the oxime, semicarbazone or phenylhydrazone were deprotected but 2,4-dinitrophenylhydrazone remained unchanged (Scheme2).



**SCHEME 2** 

Another interesting feature of this method is, that the  $\alpha$ -naphthyltriphenylphosphonium cation could be recycled and reused for regeneration of the reagent. On completion of the reaction, the residual solid materials were collected, dissolved in DMSO, and treated with freshly prepared aqueous K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> to give  $\alpha$ -naphthyltriphenylphosphonium peroxodisulfate quantitatively (Scheme 3).



#### **SCHEME 3**

In conclusion, we have developed a convenient and efficient method for oxidation of structurally different alcohols, hydroquinones, tetrahydropyranyl ethers and trimethylsilyl ethers to the corresponding carbonyl compounds and the regeneration of carbonyl compounds from oximes, phenylhydrazones and semicarbazones. Our new method has significant advantages over the existing methods such as selective and recyclable procedure, high yields and mild reaction conditions.

### **EXPERIMENTAL**

All the alcohols and hydroquinones were purchased from Fluka and Merck. The reactions were monitored by TLC using silica gel plates. The products were purified by flash column chromatography on silica gel (Merck; 230–400 mesh) and were identified by comparison of their IR and <sup>1</sup>H NMR spectra and physical data with those of authentic samples. <sup>1</sup>H NMR spectra were measured at 90 MHz with a JEOL JNM-EX 90 spectrometer with tetramethylsilane as an internal reference and DMSO-d<sub>6</sub> as solvent. IR spectra were recorded with a Pye-unicam SP 1100 spectrophotometer. Elemental analyses were performed on a LECO 250 instrument.

## Preparation of $\alpha$ -Naphthyltriphenylphosphonium Peroxodisulfate (NTPPPODS)

To an aqueous solution of  $\alpha$ -naphthyltriphenylphosphonium chloride (0.878 g, 2 mmol, 10 mL) was added an aqueous solution of potassium peroxodisulfate (0.32 g, 1.2 mmol). The mixture was stirred at room temperature for 20 min. The resulting white solid product was filtered, washed with cooled distilled water (10 mL) and dried in vacuum desiccators. Yield 0.969 g, 97%; m.p. = 221–223°C.<sup>1</sup>H NMR (DMSO):  $\delta$  = 5.61 (d, J = 16.3 Hz, 4H, CH<sub>2</sub>), 7.02–8.27 (m, 44H, arom. H). IR (KBr)  $\nu$ (cm<sup>-1</sup>): 1290, 1260, 1092, 1058, 700, 685, 590, 554; Elemental analysis: Calcd for C<sub>58</sub>H<sub>48</sub>P<sub>2</sub>S<sub>2</sub>O<sub>8</sub>: C, 69.72%; H, 4.84%; S, 6.41%, Found: C, 69.75%; H, 4.79%; S, 6.38%.

### General Procedure for Oxidation of Alcohols, Hydroquinones Tetrahydropyranyl and Trimethylsilyl Ethers, and Deprotection of Oximes, Phenylhydrazones and Semicarbazones

A mixture of the substrate (1 mmol), acetonitrile (10 mL) and NTPP-PODS (0.5-2 mmol), were placed in a round bottomed flask and stirred for the specified time (Tables II and III) at reflux conditions. After completion of the reaction (monitored by TLC), the reaction mixture was cooled to room temperature and filtered. The solid material was washed with acetonitrile (10 mL). From the combined filtrates the solvent was evaporated under reduced pressure and the resulting crude material was purified on a silica-gel column with an appropriate eluent. The pure carbonyl compounds were obtained in 65–99% yield.

## Regeneration of $\alpha$ -Naphthyltriphenylphosphonium Peroxodisulfate

The used reagent was washed with diethyl ether (2  $\times$  10 mL). The filter cake was dissolved in DMSO (5 mL) and added dropwise to a freshly

prepared aqueous solution of potassium peroxodisulfate. The mixture was stirred at room temperature for 20 min. Then, the solid material was separated by filtration, repeatedly washed with water to remove the free peroxodisulfate, and dried in a vacuum desiccator over calcium chloride to obtain the regenerated reagent.

### REFERENCES

- [1] G. A. Rajkumar and V. Murugesan, Synth. Commun., 29, 2105 (1999).
- [2] N. Alessandro, L. Liberatore, L. Tonucci, A. Morvillo, and M. Bressan, J. Mol. Catal. A, 175, 83 (2001).
- [3] F. Shirini, I. Mohammadpoor-Baltork, Z. Hejazi, and P. Heravi, Bull. Korean Chem. Soc., 24, 517 (2003).
- [4] A. Shaabani, S. Naderi, A. Rahmati, Z. Badri, M. Darvishi, and D. G. Lee, Synthesis, 3023 (2005).
- [5] M. Hunsen, Tetrahedron Lett., 46, 1651 (2005).
- [6] P. A. Anthony and K. Yoonhyun, Tetrahedron Lett., 46, 5417 (2005).
- [7] J. Y. Huang, S. J. Li, and Y. G. Wang, Tetrahedron Lett., 47, 5637 (2006).
- [8] M. Lei, R. J. Hu, and Y. G. Wang, Tetrahedron, 62, 8928 (2006).
- [9] (a) M. M. Lakouraj and K. Bahrami, J. Chem. Res (S)., 222 (2000); (b) M. Tajbakhsh, I. Mohamadpoor-Baltork, and F. Ramzanian-Lehmali, J. Chem. Res (S), 185 (2001);
  (c) M. Tajbakhsh, I. Mohamadpoor-Baltork, and F. Ramzanian-Lehmali, Phosphorus, Sulfur, and Silicon, 178, 2621 (2003); (d) M. Tajbakhsh, M. M. Lakouraj, and F. Ramzanian-Lehmali, Synlett, 11, 1724 (2006).
- [10] H. Firozabadi, P. Salehi, and I. Mohamadpour-Baltork, Bull. Chem. Soc. Jpn., 65, 2878 (1992).
- [11] J. C. Jung, H. C. Choi, and Y. H. Kim, Tetrahedron Lett., 7, 3581 (1993).
- [12] F. Chen, J. Wang, C. Guan, J. Yang, and H. Zhang, Synth. Commun., 26, 2, 253 (1996).
- [13] I. Mohammadpoor, A. R. Hajipour, and H. Mohammadi, Bull. Chem. Soc. Jpn., 71, 1649 (1998).
- [14] A. R. Hajipour, I. Mohammadpour-Baltork, and G. Kianfar, Bull. Chem. Soc. Jpn., 71, 2655 (1998).
- [15] I. Mohammadpoor-Baltork, A. R. Hajipour, and R. Haddadi, J. Chem. Res (S), 102 (1999).
- [16] G. H. Imanzadeh, M. R. Zamanloo, H. Eskandari, and A. R. Banae, *Bull. Chem. Soc. Jpn.*, **27**, 119 (2006).
- [17] (a) M. M. Lakouraj and A. Keyvan, J. Chem. Res (S), 206 (1999); (b) B. Movassagh, M. M. Lakouraj, and K. Ghodrati, Synth. Commun., 30, 3597 (2000); (c) B. Movassagh, M. M. Lakouraj, and K. Ghodrati, Synth. Commun., 32, 847 (2002); (d) M. Tajbakhsh, I. Mohamadpoor-Baltork, and F. Ramzanian-Lehmali, J. Chem. Res (S), 710 (2003); (e) M. Tajbakhsh, I. Mohamadpoor-Baltork, and F. Ramzanian-Lehmali, Phosphorus, Sulfur, and Silicon, 178, 2617 (2003); (f) M. Tajbakhsh, M. M. Lakouraj, M. H. Gholami, and F. Ramzanian-Lehmali, Phosphorus, Sulfur, and Silicon, 179, 1731 (2004); (g) M. Tajbakhsh, M. M. Lakouraj, and A. Fadavi, Synth. Commun., 34, 1173 (2004); (h) M. Tajbakhsh, M. M. Lakouraj, and K. Yadoolahzadeh, Phosphorus, Sulfur, and Silicon, 180, 2431 (2005); (i) Lakouraj, M. Tajbakhsh, F. Shirini, and M. V. Asadi Tamami, Synth. Commun., 35, 775 (2005).
- [18] W. Minghu, Y. Guichun, and C. Zuxing, React. Funct. Polym., 44, 97 (2000).