

ORIGINAL ARTICLE

Cyclic lipodepsipeptides verlamelin A and B, isolated from entomopathogenic fungus *Lecanicillium* sp.

Kei-ichi Ishidoh¹, Hiroshi Kinoshita¹, Yasuhiro Igarashi², Fumio Ihara³ and Takuya Nihira^{1,4}

Verlamelin and its new derivative (verlamelin B) were isolated from fermentation broth of entomopathogenic fungus *Lecanicillium* sp. HF627. As the structural elucidation of verlamelin so far was only preliminary, we studied and determined the absolute structure of these two compounds to be cyclo(5S-hydroxytetradecanoic acid-D-alloThr/Ser-D-Ala-L-Pro-L-Gln-D-Tyr-L-Val). This is the first study that precisely analyzed the structure of verlamelin.

The Journal of Antibiotics (2014) 67, 459–463; doi:10.1038/ja.2014.22; published online 2 April 2014

Keywords: cyclic lipodepsipeptide; entomopathogenic fungi; *Lecanicillium*; *Verticillium lecanii*

INTRODUCTION

Entomopathogenic fungi are fungal parasites of insects. During their infection to insects, they are known to evade insect immunities and kill the host insects by secreting several bioactive substances.^{1–3} As the high productivity has been well established by continuous discovery of novel bioactive compounds,^{4–6} entomopathogenic fungi have been regarded as a source of bioactive compounds. Especially, those belonging to the genus *Lecanicillium* (formerly *Verticillium lecanii*) can be good source of novel compounds, as exemplified by the isolation of indolosesquiterpenes (lecanindoles A–D⁷), phenopicolinic acid derivatives (vertilecanins⁸) and pregnanes.^{9,10}

Here we would like to report isolation and structure determination of two compounds, verlamelin (verlamelin A, **1**) and its new derivative (verlamelin B, **2**) from entomopathogenic fungus *Lecanicillium* sp. Although the planar structure of verlamelin has already been reported, its structure determination was only preliminary, because these reports lacked detailed information on the signal assignments of ¹H/¹³C, the signal connectivity in HSQC/HMBC and the D/L configuration of the component amino acids.^{11–13} Furthermore, the configuration of a hydroxyl group on the 5-hydroxytetradecanoic acid moiety has not yet been determined. Therefore, in order to completely establish the structure of verlamelin, we conducted a detailed structure determination of **1** and **2**.

RESULTS AND DISCUSSIONS

Lecanicillium sp.

HF627 was cultivated under static conditions in liquid medium containing sucrose 8%, yeast extract 4%, CaCO₃ 0.5% and HP20 1%. The whole broth was extracted with *n*-butanol. Compounds **1** and **2** were purified by a series of steps with silica gel column

chromatography, C18 column chromatography and preparative reverse phase HPLCs.

Both major compound **1** and minor compound **2** were obtained as colorless solid. The molecular formula of **1** was identified as C₄₅H₇₁N₇O₁₁ based on high-resolution fast atom bombardment MS (HRFABMS) (observed *m/z* 886.5289 [M + H]⁺, calcd 886.5290 for C₄₅H₇₂N₇O₁₁). The molecular formula of **2** was C₄₄H₆₉N₇O₁₁ (HRFABMS, observed *m/z* 872.5140 [M + H]⁺, calcd 872.5134 for C₄₄H₇₀N₇O₁₁), suggesting that **2** is a one-methylene-shorter derivative of **1**, considering the almost identical UV/Vis spectrum (λ_{\max} 224, 278 nm).

All the ¹³C signals and all the ¹H signals were assigned using DEPT135, HSQC, COSY and HMBC (Table 1). ¹³C NMR of the two compounds detected eight carbonyl carbons in the range of δ_C 170–174, and ¹H NMR in DMSO-*d*₆ revealed eight broad resonances, which were not detectable in the measurement using methanol-*d*₄, suggesting that certain numbers of amide linkages were present in these compounds. Overlapping methylene signals in the high magnetic field suggested that both compounds have an alkyl chain. Based on these data, **1** and **2** were assumed to bear typical lipopeptide substructures. By COSY and HMBC correlations, the component amino acids were estimated to be valine (Val), alanine (Ala), tyrosine (Tyr), threonine (Thr) (**1**) or serine (Ser) (**2**), glutamine (Gln) and proline (Pro). The remaining substructure was deduced to be a 5-hydroxytetradecanoic acid moiety by comparison of the δ_C values of C-2 to C-6 with those on cordycommunin, which has the 5-hydroxytetradecanoic acid substructure.¹⁴ As δ_H of the oxymethine H-5 was detected relatively downfield (4.79 (**1**), 4.77 (**2**)), the C-5 was judged to be involved in ester bonding. The connecting sequence of components was finally determined as

¹International Center for Biotechnology, Osaka University, Osaka, Japan; ²Biotechnology Research Center, Toyama Prefectural University, Toyama, Japan; ³National Institute of Fruit Tree Science, Ibaraki, Japan and ⁴Mahidol University–Osaka University Collaborative Research Center for Bioscience and Biotechnology, Faculty of Science, Mahidol University, Bangkok, Thailand

Correspondence: Professor T Nihira, Mahidol University–Osaka University Collaborative Research Center for Bioscience and Biotechnology, Faculty of Science, Mahidol University, 272 Rama VI Road, Ratchathewi, Bangkok 10400, Thailand.

E-mail: nihira@icb.osaka-u.ac.jp

Received 19 November 2013; revised 26 February 2014; accepted 5 March 2014; published online 2 April 2014

Table 1 NMR spectroscopic data (400 MHz, DMSO-d₆) for verlamelin A (1) and verlamelin B (2)

Verlamelin A (1)			Verlamelin B (2)		
Position	δC , mult.	δH , mult. (J in Hz)	Position	δC , mult.	δH , mult. (J in Hz)
L-Val			L-Val		
1	170.8, qC	—	1	170.8, qC	—
2	59.4, CH	3.84, t (7.3)	2	59.3, CH	3.86, t (7.3)
3	29.2, CH	1.91–2.04, ^b m	3	29.2, CH	1.97–2.02, ⁱ m
4	18.5, CH ₃	0.82, d (6.4)	4	18.5, CH ₃	0.81, d (6.4)
4'	19, CH ₃	0.84, d (6.4)	5	19, CH ₃	0.84, d (6.4)
NH	—	8.46, bd (6.9)	NH	—	8.41, bd (6.9)
D-Tyr			D-Tyr		
1	172.7, qC	—	1	172.5, qC	—
2	54.2, CH	4.61, m	2	54.0, CH	4.64, m
3	38.0, CH ₂	2.75, m	3	38.1, CH ₂	2.73, m
4	127.5, qC	—	4	127.5, qC	—
5, 9	130.3, CH	7.04, d (8.7)	5, 9	130.3, CH	7.02, d (8.2)
6, 8	114.8, CH	6.60, d (8.7)	6, 8	114.8, CH	6.59, d(8.2)
7	155.8, qC	—	7	155.8, qC	—
7-OH	—	9.09, bs	7-OH	—	9.11, bs
NH	—	7.50, bd (9.6)	NH	—	7.55, bd (8.1)
L-Gln			L-Gln		
1	171.0, qC	—	1	170.9, qC	—
2	52.7, CH	4.04, m	2	52.6, CH	4.09, m
3	26.8, CH ₂	1.71, m; 1.91–2.04, ^b m	3	27.1, CH ₂	1.71, m; 1.89, m
4	31.8, CH ₂	1.91–2.04, ^b m	4	31.8, CH ₂	1.97–2.02, ⁱ m
CONH ₂	173.7, qC	6.74, bs; 7.21, bs	CONH ₂	173.8, qC	6.74, bs; 7.21, bs
NH	—	7.78, bd (8.2)	NH	—	7.75, bd (8.2)
L-Pro			L-Pro		
1	170.9, qC	—	1	171.0, qC	—
2	60.2, CH	4.33, m	2	60.1, CH	4.33, m
3	29.2, CH ₂	1.83–1.84, ^d m; 2.11–2.16, ^c m	3	29.2, CH ₂	1.81, ^j m; 2.09, m
4	24.2, CH ₂	1.83–1.84, ^d m	4	24.2, CH ₂	1.81, ^j m
5	46.8, CH ₂	3.48, m; 3.62, m	5	46.7, CH ₂	3.45, m; 3.54, m
D-Ala			D-Ala		
1	171.0, qC	—	1	170.6, qC	—
2	46.9, CH	4.48, m	2	47.0, CH	4.48, m
3	16.3, CH ₃	1.18, d (6.4)	3	16.2, CH ₃	1.15, d (6.4)
NH	—	7.80, bd (6.4)	NH	—	7.76, bd (6.4)
D-allo-Thr			D-Ser		
1	170.5, qC	—	1	170.6, qC	—
2	58.3, CH	4.20, dd (7.1, 8.7)	2	55.3, CH	4.27, m
3	66.4, CH	3.99, m	3	61.3, CH ₂	3.64, m
4	20.0, CH ₃	1.08, d (6.4)	3-OH	—	5.27, bt (4.6)
3-OH	—	5.27, bs	NH	—	7.63, bd (7.7)
NH	—	7.48, bd (9.2)			
5(S)-hydroxytetradecanoic acid			5(S)-hydroxytetradecanoic acid		
1	172.2, qC	—	1	172.6, qC	—
2	34.7, CH ₂	1.91–2.04, ^b m; 2.11–2.16, ^c m	2	34.3, CH ₂	1.97–2.02, ⁱ m; 2.21, m
3	19.8, CH ₂	1.33–1.35, ^e m	3	19.8, CH ₂	1.33–1.37, ^k m
4	32.7, CH ₂	1.33–1.35, ^e m; 1.42–1.46, ^f m	4	32.6, CH ₂	1.33–1.37, ^k m; 1.42–1.44, ^l m
5	73.2, CH	4.79, m	5	73.2, CH	4.77, m
6	33.5, CH ₂	1.42–1.46, ^f m	6	33.6, CH ₂	1.42–1.44, ^l m
7	25.0, CH ₂	1.20–1.23, ^g m	7	24.9, CH ₂	1.20–1.23, ^m m
8	29.0, ^a CH ₂	1.20–1.23, ^g m	8	29.0, ^h CH ₂	1.20–1.23, ^m m
9	28.9, ^a CH ₂	1.20–1.23, ^g m	9	28.9, ^h CH ₂	1.20–1.23, ^m m
10	28.7, ^a CH ₂	1.20–1.23, ^g m	10	28.7, ^h CH ₂	1.20–1.23, ^m m
11	28.7, ^a CH ₂	1.20–1.23, ^g m	11	28.7, ^h CH ₂	1.20–1.23, ^m m
12	31.3, CH ₂	1.20–1.23, ^g m	12	31.3, CH ₂	1.20–1.23, ^m m
13	22.1, CH ₂	1.20–1.23, ^g m	13	22.1, CH ₂	1.20–1.23, ^m m
14	14.0, CH ₃	0.83, t (6.9)	14	14.0, CH ₃	0.83, t (6.9)

^{a,h}Carbon chemical shifts may be interchanged.^{b,c,d,e,f,g,i,j,k,l,m}The proton resonances are overlapped.

5-hydroxytetradecanoic acid, Thr (1) or Ser (2), Ala, Pro, Gln, Tyr and Val, based on HMBC and ROESY (1) and NOESY (2) correlation. HMBC correlation of the oxymethine on 5-hydroxytetradecanoic acid with C-1 on Val demonstrated the formation of a cyclic structure (Figure 1). Thus, the compound 2 was proposed to be a new analog of verlamelin (verlamelin B) in which Ser was replaced with Thr.

To determine the absolute configurations in the peptide moiety of the verlamelins, Marfey's analysis was conducted.¹⁵ After degradation of verlamelin A and verlamelin B under acidic conditions, the

hydrolysate was labeled by Marfey's reagent (N α -(5-Fluoro-2,4-dinitrophenyl)-L-alaninamide). The retention times of the labeled products in C18 HPLC (29.3, 30.2, 34.5, 37.3, 40.4, 42.1, 62.2 min for verlamelin A; and 26.0, 30.2, 34.6, 37.3, 40.5, 42.2, 62.2 min for verlamelin B) indicated that the component amino acids were L-Gln, L-Pro, D-Ala, L-Val, D-Tyr and D-allo-Thr (verlamelin A) or D-Ser (verlamelin B) (see 'Experimental procedure' for standards).

As for the absolute configuration of C-5 in the 5-hydroxytetradecanoic acid moiety, a modified Mosher's method was used.^{16,17} After the reactive phenolic hydroxyl group was protected by methyl iodide






Figure 1 Key HMBC, ROESY (1) and NOESY (2) correlation.

(R) - (S) (ppm)



Derivative of (1)

(R) - (S) (ppm)



Derivative of (2)

Figure 2 Partial (R)–(S) values of derivatives of (1) and (2) synthesized by a modified Mosher's method.

and the macrolactone ring was cleaved by methanolysis, the reaction products (purified by C18 HPLC) showed m/z at 932.5 and 918.5, respectively, suggesting that methanolysis proceeded successfully. Reaction of the product from **1** with (R)- or (S)-M α NP acid (M.W.: 230.2) gave products (purified by C18 HPLC) of m/z at 1144.6, implying that the esterification with M α NP occurred at a single position. Similar esterification reaction of the product from **2** gave products of m/z 1342.7, suggesting that the two molecules of M α NP were incorporated. Subsequent partial hydrolysis with K₂CO₃ resulted in the loss of one M α NP at the primary alcohol of the Ser residue and yielded a product of m/z at 1112.6, corresponding to the monoester with an M α NP. Each monoester with an M α NP was analyzed by ¹H NMR and COSY to observe the anisotropic effect as the result of shielding by the M α NP group. The $\Delta\delta$ (δ_R – δ_S) values of the Mosher's esters indicated the 5-S configuration (Figure 2). Based on these results, the absolute structures of verlamin A and verlamin B were definitively determined for the first time, as shown in Figure 3.

Based on the previous results of *in vitro* antifungal activity of verlamin,^{11–13} we measured the antifungal activity of **1** and **2** by disc diffusion method (summary of results, Supplementary Figure S11).




Figure 3 Stereochemical structure of verlamin A (1) and B (2), refined in this study.

The potency of both compounds was equal when tested against *Cochliobolus miyabeanus* and *Alternaria solani*. However, the new derivative of verlamin, **2** was less active against *Fusarium oxysporum*, *Cladosporium cucumerinum* and *Ustilago maydis* (potency of **2**: 4, 0.5 and >64 μ g per disc, respectively), compared with **1** (potency of **1**: 2, 0.25 and 16 μ g per disc, respectively), suggesting that the methyl group on the first amino-acid residue connected to the fatty acid moiety should have an important role for antifungal activities against these fungi. As antifungal spectrum of **1** agreed well with that of the verlamin reported previously,¹³ **1** might possess identical absolute configuration to that of the original verlamin.

By detailed structure determination, the absolute structures of **1** and **2** were identified for the first time; they were found to consist of a cyclic lipodepsipeptide, cyclo(5S-hydroxytetradecanoic acid-D-alloThr/Ser-D-Ala-L-Pro-L-Gln-D-Tyr-L-Val) (Figure 3). The related compounds, W493 A/B, cyclo(3S-hydroxy-4R-methyltetradecanoic acid-D-alloThr-L-Ala-D-Ala-L-Gln-D-Tyr-L-Val/Ile), from *Fusarium* sp.,¹⁸ and cordycommunin, cyclo(5-hydroxytetradecanoic acid-D-alloThr-L-Ala-L-Ala-L-Gln-L-Tyr-L-Val), from *Ophiocordyceps communis*,¹⁴ have also similar partial structure to fengycin (pilastatin A1, inhibitor of phospholipase A₂), which was isolated from *Bacillus subtilis* as an antifungal lipopeptide compound containing a cyclic peptide moiety composed of L-Tyr-D-alloThr-L-Glu-D-Ala-L-Pro-L-Gln-D-Tyr-L-Ile.^{19,20} Judging from the correlation of the structure

and the antifungal activity among these cyclic peptide compounds, D-tyrosine after L-glutamine might be critical for antifungal activities because cordycommunin, which possesses L-Tyr instead of D-Tyr, did not show activity against *Candida albicans* and *Magnaporthe grisea*.¹⁴

EXPERIMENTAL PROCEDURE

General experimental procedure

NMR spectra were recorded on a JEOL JMN-ECS400 (JEOL, Tokyo, Japan) at 400 MHz for ¹H and 100 MHz for ¹³C. The ¹H and ¹³C chemical shifts were referenced to the solvent signal (δ H 2.49 and δ C 39.5 in dimethyl sulfoxide (DMSO)-d₆, δ H 3.31 in CD₃OD). HRFABMS were recorded on a JEOL JMS-700 spectrometer. Optical rotation was measured on a P-1020 polarimeter (Jasco, Tokyo, Japan). The UV spectrum was recorded on a Hitachi U-3200 spectrophotometer (Hitachi, Tokyo, Japan).

Fungal material

The entomopathogenic fungus *Lecanicillium* sp. HF627 was isolated from a chillie thrips cadaver. Fungal conidia that had developed on the surface of the cadaver were transferred onto solid Sabouraud maltose yeast-extract (SMY) medium (4% maltose, 1% yeast extract, 1% peptone) with 1.5% agar and incubated at 25 °C for several days. After several rounds of single-colony isolation, the isolated strain was identified as *Verticillium lecanii* (*Lecanicillium* sp.) according to its morphology. The fungus is deposited at the culture collection of Natural Institute of Fruit Tree Science.

Fermentation and isolation

The seed culture was prepared by inoculating conidia of strain HF627 into 50 ml of SMY medium in a 500-ml baffled flask, followed by cultivation on a rotary shaker at 28 °C and 160 r.p.m. for 3 days. For the main cultivation, the seed culture (2 ml) was inoculated into 100 ml of Medium #5 (8% sucrose, 4% yeast extract, 0.5% CaCO₃, 1% activated HP20 (Mitsubishi Chemical Co., Tokyo, Japan) by methanol) in 500-ml Erlenmeyer flasks, followed by static incubation for 21 days at 25 °C. The culture broth (1 liter) was mixed with *n*-butanol (500 ml) and stirred for 1 h, and the *n*-butanol layer was collected after centrifugation (3000 r.p.m., 10 min), dried over anhydrous sodium sulfate and concentrated *in vacuo* using a rotary evaporator, yielding 3 g of brown gum as the crude extract. The crude extract was fractionated on a silica gel 60 (70–230 mesh; Merck, Darmstadt, Germany) column (ϕ 30 mm) by stepwise elution with increasing ethyl acetate concentrations (hexane/ethyl acetate = 4:1, 2:1, 3:2, 1:1, 1:2, 1:4 and 0:10 v/v), followed by stepwise elution with increasing methanol concentrations (ethyl acetate/methanol = 40:1, 20:1, 10:1, 8:1, 4:1, 2:1, 1:1, and 0:10 v/v). Fractions containing verlamelins were detected by silica gel TLC (ethyl acetate/methanol = 2:1, R_f 0.21), combined and evaporated *in vacuo* until dryness. The residues were further fractionated with a Sep-Pak Silica 35cc Vac cartridge (10 g; Waters, Milford, MA, USA) by stepwise elution with increasing methanol concentrations (ethyl acetate/methanol = 2:1, 3:2, 1:1, 1:2, 1:4, and 0:10 v/v). Verlamelins in the Sep-Pak fractions were further separated by preparative reverse phase HPLC with a Shiseido Capcell-Pak C18 column (ϕ 10 × 250 mm) (Shiseido Co., LTD., Tokyo, Japan) with isocratic elution using 80% methanol at a flow rate of 4 ml min⁻¹. The peaks of 2 and 1, detected at 11 and 12 min by UV (210 nm), respectively, were collected individually and evaporated *in vacuo* to dryness, yielding 2.7 and 65.5 mg of colorless amorphous solid, respectively (chromatogram of HPLC, see Supplementary Figure S1).

Additional quadruplicate experiments without the Sep-Pak Silica fractionation were carried out further to obtain 2. As a result, a total of 97.7 mg of 2 was obtained.

Compound (1): a colorless solid; [α]_D²⁵ +2.5 (c 0.01, MeOH); UV (MeOH) λ_{max} (log ϵ) 224 (4.20), 278 (3.29) nm; HRFABMS m/z 886.5289 [M+H]⁺ (calcd for C₄₅H₇₂N₇O₁₁, 886.5290). For ¹H, ¹³C, see Table 1 (see also Supplementary Figures S2 and S3).

Compound (2): a colorless solid; [α]_D²⁵ -68.5 (c 0.01, MeOH); UV (MeOH) λ_{max} (log ϵ) 224 (4.12), 278 (3.26) nm; HRFABMS m/z 872.5140 [M+H]⁺ (calcd for C₄₄H₇₀N₇O₁₁, 872.5134). For ¹H, ¹³C, see Table 1 (see also Supplementary Figures S4 and S5).

Stereochemistry of amino acids constituting verlamelin A and B

One milligram of aliquots of 1 or 2 were hydrolyzed by heating at 115 °C for 8 h in 10 ml of 6 M HCl. After cooling to room temperature, they were completely dried *in vacuo* and dissolved in 150 μ l of water. Marfey's reagent (300 μ l of 10 mg ml⁻¹ solution in acetone) (N^α-(2,4-dinitro-5-fluorophenyl)-L-alaninamide) (Tokyo Chemical Industry Co., Ltd., Tokyo, Japan) was added, followed by the addition of 70 μ l of 1 M NaHCO₃. The reactions proceeded at 37 °C for 1 h and were quenched by addition of 70 μ l of 1 M HCl. The resulting mixture was dried *in vacuo* and then dissolved in 1 ml of DMSO to be analyzed by HPLC. Marfey's derivatives of amino acids as a standard were prepared by reacting 50 mM of amino acids in the same manner as described above. HPLC analysis was carried out on a Shiseido Capcell-Pak C18 column (ϕ 4.6 × 250 mm) with a linear gradient of CH₃CN from 10% to 50% in 0.05% aqueous TFA solution (60 min from 10% to 50%, 5 min at 50%, 3 min from 50% to 10%), at a flow rate of 1.0 ml min⁻¹, with detection at 340 nm. The glutamine (Gln) residues of 1 and 2 should be converted by acid hydrolysis to glutamic acid (Glu). Retention times (min) of Marfey's derivatives used as standards were as follows: L-Ser (25.6), D-Ser (26.0), L-Thr (27.0), D-Thr (31.8), L-alloThr (27.5), D-alloThr (29.3), L-Glu (30.1), D-Glu (31.9), L-Pro (34.3), D-Pro (36.0), L-Ala (33.2), D-Ala (37.2), L-Val (42.1), D-Val (48.1), L-Tyr (57.9) and D-Tyr (40.4, 62.2).

(R)- and (S)-M α NP acid ester derivatives of 1 and 2

A modified Mosher's method using (R)- and (S)-M α NP acid (Tokyo Chemical Industry Co., Ltd.) was used to clarify the stereogenic centers of the hydroxytetradecanoic acid moiety.¹⁶ To protect the hydroxyl group of the Tyr residue in advance, methylation was carried out as follows. To 1 and 2 dissolved in acetonitrile (10 mg per 500 μ l) were added 50 μ l of 1,8-diazabicyclo[5.4.0]undec-7-ene and 50 μ l of CH₃I. After incubation at 50 °C for 3 h and quenching by dilution with water, the reaction products were extracted three times with an equal volume of ethyl acetate. The ethyl acetate layer was washed with 1 M HCl and brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*, yielding crude methyl ethers of 1 and 2.

To cleave the ester linkage between hydroxytetradecanoic acid and Val residue, methanolysis was carried out as follows. Crude verlamelin methyl ethers were dissolved in 2 ml of 0.5 M methanolic sodium methoxide and stirred for 5 h at room temperature. After quenching with 1 M HCl, extraction with ethyl acetate (equal volume, three times) and evaporation *in vacuo*, the cleaved product and the uncleaved reactant were collected individually by reverse phase C18 HPLC using 85% methanol as eluent. The recovered reactant was reused for the same reaction, and the product was obtained as described above. The methyl ethers of 1 and 2 yielded 6.2 and 4.2 mg of methanolysis products, respectively.

(R)- and (S)-M α NP acid esters of 1 were prepared as follows. The methanolysis product of 1 methyl ether (in 200 μ l of CH₂Cl₂) was mixed with 2 mg of dimethyl amino pyridine, 3 mg of 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC), 2 μ l of triethyl amine and 2.1 mg of (R)-M α NP acid or 2.1 mg of (S)-M α NP acid. After standing at room temperature overnight and quenching by addition of water, the reaction mixture was extracted three times with an equal volume of ethyl acetate. After washing the ethyl acetate extract with 1 M HCl and brine, drying over anhydrous sodium sulfate and concentrating *in vacuo*, the (R)- and (S)-M α NP acid ester derivatives of 1 (3a and 3b, respectively) were purified by reverse phase HPLC on a Shiseido Capcell-Pak C18 column (ϕ 10 × 250 mm) with isocratic elution using 90% methanol as a mobile phase.

(R)- and (S)-M α NP acid ester derivatives of 2 (4a and 4b, respectively) were synthesized and purified in a similar manner as those of 1.

Compound 3a: HRFABMS m/z 1144.6558 [M+H]⁺ (calcd for C₆₁H₉₀N₇O₁₄, 1144.6546). For ¹H, see Supplementary Figures S6 and S7.

Compound 3b: HRFABMS m/z 1144.6543 [M+H]⁺ (calcd for C₆₁H₉₀N₇O₁₄, 1144.6546). For ¹H, see Supplementary Figures S6 and S8.

Compound 4a: HRFABMS m/z 1112.6274 [M+H]⁺ (calcd for C₆₀H₈₆N₇O₁₃, 1112.6283). For ¹H, see Supplementary Figures S6 and S9.

Compound 4b: HRFABMS m/z 1112.6284 [M+H]⁺ (calcd for C₆₀H₈₆N₇O₁₃, 1112.6283). For ¹H, see Supplementary Figures S6 and S10.

Antifungal assay

In vitro antifungal activity of **1** and **2** against six indicator strains (*Fusarium oxysporum* subsp. *cucumerinum* NBRC 31224, *Cladosporium cucumerinum* NBRC 6370, *Aspergillus niger* NBRC 105649, *Alternaria solani* NBRC 7516, *Cochliobolus miyabeanus* NBRC 9633 and *Ustilago maydis* NBRC 6907) was tested by disc diffusion method as follows. Paper discs loaded with 10 μ l of verlamin solution in DMSO were placed on the potato dextrose agar (Difco, Detroit, MI, USA) plate, which was homogeneously inoculated with conidia, mycelia or vegetative culture of the indicator strains in advance. Growth-inhibitory effect was observed after incubation at 28 °C for 1–3 days, depending on indicator strains. Nystatin (16 μ g per disc) and DMSO alone were used as the positive and negative controls, respectively.

- 1 Wang, B., Kang, Q., Lua, Y., Bai, L. & Wanga, C. Unveiling the biosynthetic puzzle of destruxins in *Metarhizium* species. *Proc. Natl Acad. Sci. USA* **109**, 1287–1292 (2012).
- 2 Xu, Y. *et al.* Biosynthesis of the cyclooligomer depsipeptide bassianolide, an insecticidal virulence factor of *Beauveria bassiana*. *Fungal Genet. Biol.* **46**, 353–364 (2009).
- 3 Xu, Y. *et al.* Biosynthesis of the cyclooligomer depsipeptide beauvericin, a virulence factor of the entomopathogenic fungus *Beauveria bassiana*. *Chem. Biol.* **15**, 898–907 (2008).
- 4 Azumi, M. *et al.* Aurovertins F–H from the entomopathogenic fungus *Metarhizium anisopliae*. *J. Nat. Prod.* **71**, 278–280 (2008).
- 5 Isaka, M., Palasarn, S., Tobwor, P., Boonruangprapa, T. & Tسانathai, K. Bioactive anthraquinone dimers from the leafhopper pathogenic fungus *Torribiella* sp. BCC 28517. *J. Antibiot.* **65**, 571–574 (2012).
- 6 Putri, S. P., Kinoshita, H., Ihara, F., Igarashi, Y. & Nihira, T. Farinomalein, a maleimide-bearing compound from the entomopathogenic fungus *Paecilomyces farinosus*. *J. Nat. Prod.* **72**, 1544–1546 (2009).
- 7 Roll, D. M. *et al.* The Lecanindoles, nonsteroidal progestins from the terrestrial fungus *Verticillium lecanii* 6144. *J. Nat. Prod.* **72**, 1944–1948 (2009).
- 8 Soman, A. G., Gloer, J. B., Angawi, R. F., Wicklow, D. T. & Dowd, P. F. Vertilecanins: new phenopicolinic acid analogues from *Verticillium lecanii*. *J. Nat. Prod.* **64**, 189–192 (2001).
- 9 Claydon, N., Grove, J. F., Pople, M. & Begley, M. J. New metabolic products of *Verticillium lecanii*. Part 1. 3 β -Hydroxy-4, 4, 14 α -trimethyl-5 α -pregna-7, 9 (11)-diene-20S-carboxylic acid and the isolation and characterisation of some minor metabolites. *J. Chem. Soc. Perkin Trans.* **1**, 497–502 (1984).
- 10 Grove, J. F. 23, 24, 25, 26, 27-Pentanorlanost-8-en-3 β , 22-diol from *Verticillium lecanii*. *Phytochemistry* **23**, 1721–1723 (1984).
- 11 Onishi, J. C., Rowin, G. L. & Miller, J. E. Antibiotic A43F. United States Patent 4201771. May 6 (1980).
- 12 Kim, J. C. *et al.* Verlamin, an antifungal compound produced by a mycoparasite *Acremonium strictum*. *Plant Pathol. J.* **18**, 102–105 (2002).
- 13 Rowin, G. L. *et al.* Verlamin, a new antifungal agent. *J. Antibiot.* **39**, 1772–1775 (1986).
- 14 Haritakun, R., Sappan, M., Suvannakad, R., Tسانathai, K. & Isaka, M. An antimycobacterial cyclodepsipeptide from the entomopathogenic fungus *Ophiocordyceps communis* BCC 16475. *J. Nat. Prod.* **73**, 75–78 (2010).
- 15 Bhushan, R. & Bruckner, H. Marfey's reagent for chiral amino acid analysis: a review. *Amino acids* **27**, 231–247 (2004).
- 16 Kasai, Y. *et al.* MxNP acid, a powerful chiral molecular tool for preparation of enantiopure alcohols by resolution and determination of their absolute configurations by the ^1H NMR anisotropy method. *Chirality* **16**, 569–585 (2004).
- 17 Kusumi, T., Yabuuchi, T., Takahashi, H. & Ooi, T. Chiral anisotropic reagents for determining the absolute configuration of secondary alcohols and carboxylic acids. *Yuki Gosei Kagaku Kyokaiishi* **63**, 1102–1114 (2005).
- 18 Nihei, K., Itoh, H., Hashimoto, K., Miyairi, K. & Okuno, T. Antifungal cyclodepsipeptides, W493 A and B, from *Fusarium* sp.: isolation and structural determination. *Biosci. Biotechnol. Biochem.* **62**, 858–863 (1998).
- 19 Honma, M. *et al.* Termination of the structural confusion between plipastatin A1 and fengycin IX. *Bioorg. Med. Chem.* **20**, 3793–3798 (2012).
- 20 Samel, S. A., Wagner, B., Marahiel, M. A. & Essen, L. O. The thioesterase domain of the fengycin biosynthesis cluster: a structural base for the macrocyclization of a non-ribosomal lipopeptide. *J. Mol. Biol.* **359**, 876–889 (2006).

Supplementary Information accompanies the paper on The Journal of Antibiotics website (<http://www.nature.com/ja>)