

Organic Synthesis

Enantioselective Organocatalytic Reduction of β -Trifluoromethyl Nitroalkenes: An Efficient Strategy for the Synthesis of Chiral β -Trifluoromethyl AminesElisabetta Massolo,^[a] Maurizio Benaglia,^{*[a]} Manuel Orlandi,^[a] Sergio Rossi,^[a] and Giuseppe Celentano^[b]

Abstract: An efficient organocatalytic stereoselective reduction of β -trifluoromethyl-substituted nitroalkenes, mediated by 3,5-dicarboxylic ester-dihydropyridines (Hantzsch ester type), has been successfully developed. A multifunctional thiourea-based (*S*)-valine derivative was found to be the catalyst of choice, promoting the reaction in up to 97% *ee*. The methodology has been applied to a wide variety of substrates, leading to the formation of differently substituted precursors of enantiomerically enriched β -trifluoromethyl amines. The mechanism of the reaction and the mode of action of the metal-free catalytic species were computationally investigated; on the basis of DFT transition-state (TS) analysis, a model of stereoselection was also proposed.

The introduction of fluorinated residues in a bioactive molecule may have a strong impact on its effectiveness, simultaneously modulating electronic, lipophilic and steric parameters, critically influencing both the pharmacodynamic and pharmacokinetic properties.^[1] Many drugs are characterized by the presence of a CF₃ group, as in Efavirenz, used against HIV infection, Telcagepant, employed in the treatment of the neurovascular disorder migraine and Prozac (Figure 1). Among several trifluoromethylated compounds known in medicinal chemistry, α - and β -trifluoromethylamine units are major components found in this class of bioactive species (Figure 1).^[2] Despite the great interest in the topic and progress in the field, highly ste-

reochemically efficient catalytic methods for the synthesis of chiral fluorinated amines are still deeply needed.^[3]

After focusing on the preparation of chiral α -trifluoromethylamine, by trichlorosilane-mediated enantioselective catalytic reduction of trifluoromethyl ketoimines,^[4] we turned our attention to chiral β -trifluoromethylamines. Recent works on innovative, nonstereoselective, methods of trifluoromethylation to generate β -trifluoromethyl-substituted aza-derivatives clearly testify for the great interest in the preparation of this class of compounds.^[5] However, really few strategies are available for the stereoselective synthesis of enantiopure β -trifluoromethylamines. In 2013, a chemo- and enantioselective CF₃SiMe₃ addition to α -imino ketones promoted by Cinchona-derived ammonium salts was reported (up to 71% *ee*).^[6] In the same year, based on the seminal contributions by the MacMillan group,^[7] the first visible light-driven intermolecular aminotrifluoromethylation of alkenes was developed, as efficient and regioselective one-step access to a variety of β -trifluoromethylamides in up to 82% yield and 78% d.r.^[8]

An alternative strategy to obtain chiral trifluoromethyl derivatives relies on the stereoselective transformation of trifluoromethylated prochiral, readily available substrates. Fluorinated nitroalkenes represent an obvious entry for the synthesis of β -trifluoromethylamino derivatives. They have been widely used in combination with a chiral reactant partner to diastereoisomerically produce β -nitro-trifluoromethylated products; typical examples include the synthesis of trifluoromethylated peptide analogues, using enantiopure α -amino acids as nucleophiles.^[9] Fluorinated nitroolefines have been also employed in organocatalytic Michael reactions with aldehydes,^[10] intermolecular oxa-Michael addition of oximes,^[11] and 3-alkylidene oxindoles,^[12] and in the addition of indole^[13] and β -dicarbonyl compounds.^[14]

However, as far as we know, no examples of enantioselective organocatalytic reduction of trifluoromethylated nitroalkenes are known. Here we wish to report our preliminary investigation on the synthesis of enantiomerically enriched- β -trifluoromethyl nitroolefines, by organocatalyzed addition of Hantzsch ester type reducing agents to fluorinated nitroalkenes. Considering that the starting materials are readily, and often commercially available, trifluoromethyl ketones the methodology offers an easy, catalytic, and practical entry to chiral β -trifluoromethylamines (Scheme 1).

In 2007, List and co-workers published the organocatalyzed stereoselective reduction of nitroalkenes affording the corre-

[a] E. Massolo, Prof. M. Benaglia, M. Orlandi, Dr. S. Rossi
Dipartimento di Chimica
Università degli Studi di Milano
via Golgi 19, 20133 Milano (Italy)
Fax: (+39)0250314159
E-mail: maurizio.benaglia@unimi.it

[b] G. Celentano
Dipartimento di Scienze Farmaceutiche
Università degli Studi di Milano
Via Mangiagalli, 25, Milano (Italy)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/chem.201405730>. It contains experimental details for catalysts synthesis, catalytic enantioselective reductions and synthetic modification for the absolute configuration determination. Analysis and characterization: NMR and HPLC traces of the reaction products. Cartesian coordinates, total electronic and Gibbs free energies of all reported structures.





Figure 1. Some biologically active trifluoromethylated compounds.

substituents clearly showed that an acceptable level of enantio-control could be achieved only with sterically hindered residues on the amino group (Table 1, entries 1–4 vs. 5–6). In particular, a pyrrole ring proved to be more effective than a phthalimide in shielding one of the two enantiotopic faces of the nitroalkene, probably due to the presence of the 2,5-dimethyl substituents, able to extend the influence of the heterocycle ring on the region of space just close to the site at which the substrate is supposed to be coordinated to the thiourea residue of the catalyst (Figure 3, see below, Figure 5, for further considerations based on computational studies). However, with catalyst




Scheme 1. Organocatalytic approach to the synthesis of chiral β -trifluoromethylamines.

sponding enantiomerically enriched nitroalkanes in high yields and up to 96% *ee*, the most efficient catalyst featuring a thiourea moiety linked to the diethylamide of (*S*)-*tert*-leucine and to the *N*-pyrrole derivative of (*R,R*)-diaminocyclohexane.^[15] Based on this previous work we have preliminarily explored a variety of bifunctional, thiourea-based molecules as catalyst for the reduction of trifluoromethylated nitroolefins (Figure 2).^[16] Catalysts 1–11 were easily synthesized in four steps through conversion of the *N-tert*-butoxycarbonyl (Boc) amino acid to the corresponding amide, followed by *N*-Boc deprotection, generation of the isothiocyanate group and reaction with the proper 1,2-diaminocyclohexane derivative.

The reduction of β -trifluoromethyl- β -nitrostyrene **12** was selected as model reaction and typically performed at 25 °C for 20 h in the presence of a stoichiometric amount of Hantzsch ester **A** and a catalytic amount (10 mol%) of the organocatalysts, to afford the expected product **13** (Scheme 2). The results of this preliminary screening are detailed in Table 1.

In looking for the best catalyst, we systematically investigated how the different parts constituting the skeleton of these thiourea-based structures influence their activity and stereoselectivity. Thus, the modification of the diaminocyclohexane *N*-



Scheme 2. Organocatalytic reduction of β -trifluoromethyl nitrostyrene.

Table 1. Preliminary studies on the catalytic reduction of **12** with organocatalysts 1–11.

Entry	Cat.	Conv. [%] ^[a]	<i>ee</i> [%] ^[b]
1	1	60	< 5
2	2	> 99	rac.
3	3	60	13
4	4	83	rac.
5	5	> 99	38
6	6	85	57
7	7	> 99 (99) ^[c]	59
8	8	> 99	43
9	9	95 (93) ^[c]	81
10	10	> 99 (98) ^[c]	77
11	11	80	55

[a] Reaction run in deuterated benzene at RT; conversion determined by ¹H and ¹⁹F NMR spectra. [b] Enantiomeric excess determined by HPLC analysis on a chiral stationary phase (for details see the Supporting Information). [c] In brackets isolated yields after chromatographic purification.

6 a modest enantioselectivity was obtained. Therefore, we decided to look at the amino acid portion of the catalyst; it was observed that, when replacing *tert*-leucine with valine, higher *ee* values were obtained. Table 1, entries 7–8, demonstrated the combination (*S*)-valine and (*R,R*)-diaminocyclohexane to be the match pair, which guarantees higher *ee*. The 2,5-dimethyl derivative **9** was able to promote the reduction of β -trifluoro-




Figure 2. A few selected chiral organocatalysts employed in the present study.




Figure 3. Proposed mode of action of the bifunctional thiourea-based catalyst.

methyl nitrostyrene at room temperature in 81% enantioselectivity, to afford (*R*)-2-trifluoromethyl-2-phenyl-nitroethane **13** in 95% yield; *N,N*-dibenzyl substituents on the amide function (catalyst **10**) afforded only slightly lower *ee* with respect to derivative **9**.

Once **9** and **10** had been identified as the most promising catalysts, a screening of reaction conditions was carried out. When the temperature was lowered to 0 °C, a diminished chemical activity was observed but no significant improvement in enantiocontrol (Table 2, entries 1, 2). To improve the reaction rate and to allow us to perform the reduction at lower temperature, *tert*-butyl ester derivative **B** (Figure 4), known to be a more active reducing agent,^[17] was employed. Higher *ee* values could not be achieved in dichloromethane (entries 3–6) at 0 or at –50 °C; however, by running the reaction in toluene, an increase of the enantioselectivity was observed for lower reaction temperatures. In the presence of the Hantzsch-type ester **B** at –50 °C, the stereoselectivity was improved to up to 87% *ee* with catalyst **9** and up to 93% *ee* with catalyst **10** (en-




Figure 4. Different reducing agents.

tries 9–10). Reductions in polar or protic solvents led to the products with low stereocontrol, while hexanes proved to be a suitable solvent for the reaction: with catalyst **9** at –50 °C, up to 83% *ee* was reached (entry 15).

The generality of the method was finally investigated (Scheme 3); differently 2-aryl- and 2-alkyl-substituted 3,3,3-trifluoro-1-nitropropene derivatives **14a–g** have been prepared and reacted with dihydropyridine **B** to afford the corresponding chiral trifluoromethyl nitroalkanes **15a–g**, in the presence of 10% mol equivalents of catalyst **10** in toluene, typically at –50 °C. Some selected results are reported in Table 3.

The reduction of differently substituted 2-aryl-2-trifluoromethyl nitroalkenes has been successfully accomplished, either

on substrates featuring electron-donating groups or electron-withdrawing residues; enantioselectivities usually higher than 80% and up to 90% were obtained (Table 3, entries 1–4).

Even better results were achieved with 2-alkyl-substituted fluorinated nitroalkenes **14e–g**; with benzyl derivative **14e** the reduction mediated by catalyst **10** afforded the expected nitroalkane **15e** in 90% *ee*. Excellent results were obtained in the reduction of alkyl-nitroalkenes **14f–g** that afforded products **15f** and **15g** in 97 and 93% *ee*, respectively.

Table 2. Screening of experimental conditions in the organocatalyzed reduction of **12**.

Entry	Cat.	Solvent	<i>T</i> [°C]	Hantzsch ester	Conv. [%] ^[a]	<i>ee</i> [%] ^[b]
1	7	CH ₂ Cl ₂	0	A (COOEt)	50	60
2	9	CH ₂ Cl ₂	0	A (COOEt)	60	80
3	9	CH ₂ Cl ₂	0	B (COOtBu)	>99	78
4	9	CH ₂ Cl ₂	–15	B (COOtBu)	>99	77
5	10	CH ₂ Cl ₂	–15	B (COOtBu)	>99	74
6	9	CH ₂ Cl ₂	–50	B (COOtBu)	>99	77
7	9	toluene	RT	B (COOtBu)	>99	71
8	10	toluene	RT	B (COOtBu)	>99	81
9	9	toluene	–50	B (COOtBu)	61	87
10	10	toluene	–50	B (COOtBu)	71	93
11	9	THF	0	B (COOtBu)	>99	9
12	9	EtOH	0	B (COOtBu)	>99	6
13	9	hexanes	0	B (COOtBu)	>99	80
14	10	hexanes	–50	B (COOtBu)	50	71
15	9	hexanes	–50	B (COOtBu)	70	83

[a] Conversion determined by ¹H and ¹⁹F NMR spectra. [b] Enantiomeric excess determined by HPLC on chiral stationary phase (for details see the Supporting Information).






Figure 5. B3LYP/6-31G(d) geometries of the TSs for catalyst **9**. In the “balls and springs” pictures, the COOMe groups of the Hantzsch ester are reported in “transparent” fashion; non-participating hydrogen atoms are omitted for clarity. Below a comparison between catalysts **6** and **9** is reported.



Scheme 3. Organocatalytic reduction of β -trifluoromethyl nitroalkenes **14a–g**.



Scheme 4. Reduction of α -substituted- β -trifluoromethyl nitroalkenes **16a–c**.

Possible extension of this reaction to α,β -disubstituted- β -trifluoromethylnitroalkenes has also been preliminary investigated (Scheme 4). It is worth mentioning that the stereoselective reduction of nitrostyrene derivatives featuring an α -substituent is still considered a challenging transformation that requires a controlled protonation step^[18] and leads to the formation of a rather labile stereocentre; indeed, to the best of our knowledge, no organocatalytic example has been reported so far.^[19]

Substrates were obtained as a differently enriched mixture of isomers that were shown to not be configurationally stable and involved an equilibrium.^[20] Due to this equilibration, H-H

NOESY experiments were not decisive for attributing the *E/Z* configuration; however, computational analysis showed the *E* isomer to be slightly more stable than the *Z* isomer (although by less than 2 kcal/mol, see the Supporting Information). However, it is important to note that almost identical results in terms of yield, d.r. and *ee* were obtained in the enantioselective reduction of **16**, independently from the diastereoisomeric ratio of the starting nitroalkene.

The β -trifluoromethyl- α -methyl-substituted nitroalkanes **17a–c** were obtained in modest yields and low diastereoisomeric ratio, but with an appreciable level of enantioselectivity, typically about 60–70% *ee* for both isomers, and up to 77% *ee* (Table 4). Further experiments are necessary to determine the configuration of these novel fluorinated nitroalkanes, featuring two stereocenters. Follow-up studies will be undertaken to design and develop a stereoselective catalytic system better suited for this transformation.

To establish the absolute configuration of the reduction products, **15e** was transformed by hydrogenation into the corresponding β -trifluoromethyl amine **18** that was converted in the known *N*-benzyl derivative **19**,^[21] thus determining the absolute configuration of the major isomer to be *R* (Scheme 5).


The different behaviour of *tert*-leucine derivative **6** and of the valine-based catalyst **9** was computationally investigated. The transition states (TSs) leading to the formation of both (*R*)- and (*S*)-**13**, for the reaction with both catalysts **6** and **9**, have been located at a B3LYP/6-31G(d) level of theory; finer electronic energies have successively been obtained increasing the basis set up to 6/311 + (2df,2pd) with two different functionals: B3LYP^[22] and M062X.^[23]

entry	Substrate	t [h]	Conv. [%] ^[b]	ee [%] ^[c]
1		36	81	83
2		36	75	73
3		20	85	81
4		20	81	90
5		20	89	90
6		20	80	97
7		36	75	93

[a] Reaction run in deuterated toluene, with 10 mol% of catalyst and 1.3 mol equivalents of reducing agent **B**. [b] Conversion determined by ¹H and ¹⁹F NMR spectra. [c] Enantiomeric excess determined by HPLC on chiral stationary phase (see the Supporting Information for details).

Preliminary conformational analysis on the catalysts underlined a very rigid structure: only three conformations for catalyst **9** and a unique conformation for catalyst **6** are energetically allowed. The selected TSs have been found, assuming the coordination of the nitro group to the thiourea moiety and of the Hantzsch ester NH group to the catalyst carboxamide group (Figure 5). The methyl-substituted Hantzsch ester **C** (see Figure 4) was used to reduce possible conformations of the system. In the tables and figures below, the notation **TS-S9** refers to the TS of catalyst **9** leading to the *S* isomer.

In the case of catalyst **9**, a first inspection of the two TSs clearly highlights the role of the pyrrole moiety, that exerts in **TS-S9** a repulsive interaction on the phenyl group of the nitroalkene, thus disfavoring the formation of the *S* enantiomer. Additionally, a deeper look at **TS-S9** and **TS-R9** shows that upon the coordination of the substrate to the thiourea, the ni-



Scheme 5. Determination of absolute configuration of product **15e**.

Entry	Substrate	E/Z ratio starting material	React. T [°C] (t [h])	conv. [%] ^[b]	d.r. ^[b]	ee [%] ^[c]
1		71:29	0 (48)	20	58:42	45–51
2		71:29	25	51	60:40	67–70
3		95:5	25	57	60:40	65–66
4		54:46	25	66	64:36	77–67
5		90:10	25	42	62:38	<5–<5

[a] Reaction run in deuterated toluene, with 10 mol% of catalyst and 1.3 mol equivalents of reducing agent **B**. [b] Conversion and diastereoisomeric ratio determined by ¹H and ¹⁹F NMR spectroscopy. [c] Enantiomeric excess determined by HPLC on chiral stationary phase (for details see the Supporting Information).

troolefin **C2** electrophilic carbon atom in **TS-S9** is closer to the bulky diaminocyclohexane moiety than in **TS-R9**. Therefore the reducing agent, in releasing the hydride, is forced in a hindered zone at which its ester group repulsively interacts with the 2,5-methyl pyrrole substituents, thus further favouring **TS-R9** (see Figure 5).

On the other hand, in the attempt to understand and compare the behaviour of catalysts **6** and **9**, it should be noted that in **TS-R** of both catalysts a further repulsive interaction may occur between the phenyl ring and *i*Pr or *t*Bu groups. The competition between this two negative interactions seems to be responsible for the worst behaviour of catalyst **6** with respect to catalyst **9**, in which the less hindered *i*Pr moiety may avoid a destabilizing interaction with the substrate (compare **TS-R6** and **TS-R9** structures in Figure 6).

In an attempt to rationalize the higher *ee* observed in the reduction of alkyl-substituted nitroalkenes it was noticed that in the disfavoured TSs the methyl group of the 2,5-dimethyl pyrrole moiety is forced toward the β -substituent of the substrate. Therefore, when such a substituent is an aryl group, a CH₃- π interaction might take place, reducing the energy difference between favoured and disfavoured TSs.^[24]

The presence of this kind of interaction was confirmed by performing a non-covalent interaction (NCI) analysis, as previously reported by Yang et al., using the NCIPLOT software;^[25] results are shown in Figure 7. The green surface between the phenyl ring and the catalyst's methyl substituent evidences a large region




Figure 6. B3LYP/6-31G(d) geometries of the disfavoured TSs for catalysts **6** and **9**. In the “balls and springs” pictures, the COOMe groups of the Hantzsch ester are reported in “transparent” fashion; non-participating hydrogen atoms are omitted for clarity.




Figure 7. NCIPLLOT analysis performed in a selected cube (Supporting Information) using the M06-2X/6-311 + G(2df,2pd) electron density calculated on the B3LYP/6-31G(d) geometry of the disfavoured TS for catalyst **9**. Non-participating hydrogen atoms are omitted for clarity.

of weak positive interaction due to the expected $\text{CH}_3\text{-}\pi$ stacking. Clearly, when the β -aryl group is replaced by an alkyl one, this interaction cannot occur. Hence, TSs leading to the *R* and *S* enantiomer should be energetically more differentiated, justifying the experimentally observed higher enantiomeric excesses with alkyl derivatives (Table 3, entries 6 and 7).

The calculated Gibbs free energies and the prediction of the enantioselectivities are reported in Table 5. Since the B3LYP functional is known to give a poor description for long-range dispersive interactions, as expected, a better prediction of the enantiomeric excesses was obtained with M06-2X. In particular, a remarkably good agreement between calculated and experimental *ee* obtained with catalyst **6** has been achieved with M06-2X (43 vs. 57% *ee*). Noteworthy, both the DFT functionals predict higher TSs energy differences for catalyst **9** compared to **6**, thus confirming the valine-derivative **9** to be the best-performing catalyst.^[26]

Table 5. Computationally determined energies and enantioselectivities.					
	B3LYP		M06-2X		Exptl. <i>ee</i> [%] ^[c]
	$\Delta\Delta G$ ^[a]	<i>ee</i> [%] ^[c]	$\Delta\Delta G$ ^[b]	<i>ee</i> [%] ^[c]	
TS-R6	0.0	97	0.0	43	57
TS-S6	2.4		0.5		
TS-R9	0.0	> 99	0.0	99	81
TS-S9	4.6		3.0		

[a] $\Delta\Delta G$ in kcal mol^{-1} at the B3LYP/6-311 + G(2df,2pd)//B3LYP/6-31G(d) level of theory. The thermal correction to the Gibbs free energy was calculated at the B3LYP/6-31G(d) level. [b] $\Delta\Delta G$ in kcal mol^{-1} at the M062X/6-311 + G(2df,2pd)//B3LYP/6-31G(d) level of theory. The thermal correction to the Gibbs free energy was calculated at the B3LYP/6-31G(d) level. [c] *ee* was calculated according to the formula: $ee = 100 \exp[-(\Delta\Delta G_s)/RT]$.

In conclusion, the enantioselective organocatalytic reduction of β -trifluoromethyl nitroalkenes was successfully accomplished, either on β -aryl- or β -alkyl-substituted substrates, in up to 97% *ee*. The stereochemical outcome of the reaction and the behaviour of the metal-free, thiourea-based catalyst were also rationalized and discussed, based on computational studies and DFT transition-state analysis. This easy to perform methodology offers a valuable entry for a straightforward synthesis of enantiomerically pure β -trifluoromethyl amines.^[27]

Experimental Section

General methods

All commercially available reagents including dry solvents were used as received. Organic extracts were dried over sodium sulfate, filtered and concentrated under vacuum using a rotatory evaporator. Non-volatile materials were dried under high vacuum. Reactions were monitored by TLC on precoated Merck silica gel 60 F254 plates and visualized either by UV or by staining with a solution of cerium sulfate (1 g) and ammonium heptamolybdate tetrahydrate (27 g) in a mixture of water (469 mL) and concentrated sulfuric acid (31 mL). Flash chromatography was performed on silica

gel 60. The NMR spectra were obtained at 300 MHz for ^1H -, 75 MHz for ^{13}C -, and 282 MHz for ^{19}F NMR spectra on a Bruker 300 spectrometer.

General procedure for organocatalytic reactions

The β -trifluoromethyl nitroalkene derivative (1 equiv) and the catalyst (0.1 equiv) were dissolved in the dry solvent (1 M solution) under a nitrogen atmosphere for 10 min at the desired temperature; Hantzsch ester (1 equiv) was added as a solid and the mixture stirred for 20 h at constant temperature. The solvent was then removed under reduced pressure and products were isolated by flash column chromatography on silica gel (hexane:ethyl acetate) for their NMR spectroscopic characterization. When low-boiling substrates were employed—yielding low boiling products—only the crude mixture was analyzed. Enantiomeric excess was determined by HPLC on chiral stationary phase.

Acknowledgements

M. B. thanks COST action CM9505 "ORCA" Organocatalysis and ZaCh Systems (Zambon Group) for financial support. E.M. and M.O. thank Università degli Studi di Milano for PhD fellowships, S.R. thanks Università degli Studi di Milano for post-doctoral fellowship.

Keywords: bifunctional catalyst · chiral fluorinated amines · nitroalkenes · reduction · trifluoromethyl derivatives

- [1] a) P. Kirsch, *Modern Fluoroorganic Chemistry: Synthesis Reactivity, Applications*, Wiley-VCH, Weinheim, **2013**; b) I. Ojima, *Fluorine in Medicinal Chemistry and Chemical Biology*, Wiley-Blackwell, Chichester, **2009**; c) J.-P. Beguè, D. Bonnet-Delpon, *Bioorganic and Medicinal Chemistry of Fluorine*, Wiley-VCH, Hoboken, **2008**; d) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320–330; e) K. Muller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881–1886.
- [2] H.-P. Chiu, B. Kokona, R. Fairman, R. P. Cheng, *J. Am. Chem. Soc.* **2009**, *131*, 13192–13193; d) B. C. Buer, J. Chugh, H. M. Al-Hashimi, E. N. G. Marsh, *Biochemistry* **2010**, *49*, 5760–5765; e) C. R. Hopkins, *ACS Chem. Neurosci.* **2012**, *3*, 3–4.
- [3] Review: J. Nie, H.-C. Guo, D. Cahard, J.-A. Ma, *Chem. Rev.* **2011**, *111*, 455–529. See also: D. Cahard, V. Bizet, *Chem. Soc. Rev.* **2014**, *43*, 135–147.
- [4] A. Genoni, M. Benaglia, E. Massolo, S. Rossi, *Chem. Commun.* **2013**, *49*, 8365–8367.
- [5] a) Y.-F. Wang, G. Hervé Lonca, S. Chiba, *Angew. Chem. Int. Ed.* **2014**, *53*, 1067–1071; *Angew. Chem.* **2014**, *126*, 1085–1089; b) H. Egami, S. Kawamura, A. Miyazaki, M. Sodeoka, *Angew. Chem. Int. Ed.* **2013**, *52*, 7841–7844; *Angew. Chem.* **2013**, *125*, 7995–7998; c) E. Kim, S. Choi, H. Kim, E. J. Cho, *Chem. Eur. J.* **2013**, *19*, 6209–6212.
- [6] E. Objalska, G. Mloston, A. Six, *Tetrahedron Lett.* **2013**, *54*, 2462–2465.
- [7] a) D. A. Nagib, M. E. Scott, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2009**, *131*, 10875–10877; b) A. E. Allen, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2010**, *132*, 4986–4987.
- [8] Y. Yasu, T. Koike, M. Akita, *Org. Lett.* **2013**, *15*, 2136–2139.
- [9] M. Molteni, M. C. Bellucci, S. Bigotti, S. Mazzini, A. Volonterio, M. Zanda, *Org. Biomol. Chem.* **2009**, *7*, 2286–2296; and references cited therein.
- [10] O. Andrey, A. Alexakis, A. Tomassini, G. Bernardinelli, *Adv. Synth. Catal.* **2004**, *346*, 1147–1168.
- [11] F.-L. Liu, J.-R. Chen, B. Feng, X.-Q. Hu, L.-H. Ye, L.-Q. Lu, W.-J. Xiao, *Org. Biomol. Chem.* **2014**, *12*, 1057–1060.
- [12] Q. Chen, G. Wang, X. Jiang, Z. Xu, L. Lin, R. Wang, *Org. Lett.* **2014**, *16*, 1394–1397.
- [13] J.-R. Gao, H. Wu, B. Xiang, W.-B. Yu, L. Han, Y.-X. Jia, *J. Am. Chem. Soc.* **2013**, *135*, 2983–2986.
- [14] Y. Zhao, X.-J. Wang, Y. Lin, C.-X. Cai, J.-T. Liu, *Tetrahedron* **2014**, *70*, 2523–2528.
- [15] N. J. A. Martin, L. Ozores, B. List, *J. Am. Chem. Soc.* **2007**, *129*, 8976–8977. For a recent organometallic-catalyzed nitroalkene reduction with Hantzsch ester see: L.-A. Chen, W. Xu, B. Huang, J. Ma, L. Wang, J. Xi, K. Harms, L. Gong, E. Meggers, *J. Am. Chem. Soc.* **2013**, *135*, 10598–10601. For other contributions in the field see: J. F. Schneider, F. C. Falk, R. Fröhlich, J. Paradies, *Eur. J. Org. Chem.* **2010**, 2265–2269 and J. F. Schneider, M. B. Lauber, V. Muhr, D. Kratzer, J. Paradies, *Org. Biomol. Chem.* **2011**, *9*, 4323–4327.
- [16] For experimental details on the synthesis and characterization of the catalysts, see the Supporting Information.
- [17] S. G. Ouellet, A. M. Walji, D. W. C. MacMillan, *Acc. Chem. Res.* **2007**, *40*, 1327–1339.
- [18] For a recent review on enantioselective protonation, see: S. Oudeyer, J.-F. Brière, V. Levacher, *Eur. J. Org. Chem.* **2014**, 6103–6119. For important contributions in the field, see also: D. Leow, S. Lin, S. K. Chittimalla, X. Fu, C.-H. Tan, *Angew. Chem. Int. Ed.* **2008**, *47*, 5641–5645; *Angew. Chem.* **2008**, *120*, 5723–5727 and S. Lin, D. Leow, K.-W. Huang, C.-H. Tan, *Chem. Asian J.* **2009**, *4*, 1741–1744.
- [19] For a recent example of an enzymatic stereoselective reduction of α -methylated *trans*-nitroalkenes, see: E. Burda, T. Reiß, T. Winkler, C. Giese, X. Kostrov, T. Huber, W. Hummel, H. Gröger, *Angew. Chem. Int. Ed.* **2013**, *52*, 9323–9326; *Angew. Chem.* **2013**, *125*, 9493–9496.
- [20] In fact, a change in the diastereoisomeric ratio of the two isomers, from roughly 50:50 to 90:10 was observed for all compounds **16a–c** to take place in few days, even when the isomeric mixture was kept neat at 8 °C.
- [21] See ref. [7a]. The reactions sequence did not affect the stereochemical integrity of the starting material, as confirmed by HPLC analysis on a chiral stationary phase (Supporting Information).
- [22] a) A. D. Becke, *J. Chem. Phys.* **1993**, *98*, 5648–5652; b) C. Lee, W. Yang, R. G. Parr, *Phys. Rev. B* **1988**, *37*, 785–789; c) P. J. Stephens, F. J. Devlin, C. F. Chabalowski, M. J. Frisch, *J. Phys. Chem.* **1994**, *98*, 11623–11627.
- [23] Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215–241.
- [24] We thank the referee for the nice suggestion to consider also this type of interaction.
- [25] a) E. R. Johnson, S. Keinan, P. Mori-Sánchez, J. Contreras-García, A. J. Cohen, W. Yang, *J. Am. Chem. Soc.* **2010**, *132*, 6498–6506; b) J. Contreras-García, E. R. Johnson, S. Keinan, R. Chaudret, J.-P. Piquemal, D. N. Beratan, W. Yang, *J. Chem. Theory Comput.* **2011**, *7*, 625–632.
- [26] For further details on the computational studies see the Supporting Information.
- [27] For a recent contribution on the synthesis of chiral trifluoromethyl amines, see: X. Dai, D. Cahard, *Adv. Synth. Catal.* **2014**, *356*, 1317–1328. See also: S. J. Shaw, D. A. Goff, L. A. Boralsky, M. Irving, R. Singh, *J. Org. Chem.* **2013**, *78*, 8892–8897. For a discussion on the unique features of β -fluoroamines see: L. E. Zimmer, C. Sparr, R. Gilmour, *Angew. Chem. Int. Ed.* **2011**, *50*, 11860–11871; *Angew. Chem.* **2011**, *123*, 12062–12074.

Received: October 20, 2014
Published online on ■■■■■, 0000


COMMUNICATION

Organic Synthesis

E. Massolo, M. Benaglia, M. Orlandi,
S. Rossi, G. Celentano*

■■ – ■■

 **Enantioselective Organocatalytic Reduction of β -Trifluoromethyl Nitroalkenes: An Efficient Strategy for the Synthesis of Chiral β -Trifluoromethyl Amines**



An efficient organocatalytic stereoselective reduction of β -trifluoromethyl-substituted nitroalkenes, mediated by 3,5-dicarboxylic ester-dihydropyridines (Hantzsch ester type), has been success-

fully developed. A thiourea-based (S)-valine derivative promoted the reaction in up to 97% *ee*, affording immediate precursors of enantiomerically enriched β -trifluoromethylamines (see scheme).