



Cite this: DOI: 10.1039/c7ob03109b

One-pot synthesis of diverse *N,N'*-disubstituted guanidines from *N*-chlorophthalimide, isocyanides and amines *via N*-phthaloyl-guanidines†

András Demjén,^{a,b} Anikó Angyal,^{a,b} János Wölfling,^b László G. Puskás^a and Iván Kanizsai^{✉*}

Received 15th December 2017,

Accepted 1st March 2018

DOI: 10.1039/c7ob03109b

rsc.li/obc

A sequential one-pot approach towards *N,N'*-disubstituted guanidines from *N*-chlorophthalimide, isocyanides and amines is reported. This strategy provides straightforward and efficient access to diverse guanidines in yields up to 81% through previously unprecedented *N*-phthaloylguanidines. This protocol also features wide substrate scope and mild conditions.


Introduction

The guanidine functionality is a privileged structure in many natural products, biochemical processes and pharmaceuticals, playing key roles in various biological functions.¹ Moreover, guanidines also serve as valuable scaffolds in organocatalysis² and precursors for the synthesis of heterocycles.³ The traditional synthesis of guanidines mainly relies on the addition of amines to carbodiimides,⁴ or utilizes thioureas (usually bearing electron-withdrawing substituents) in conjunction with thiophilic metal salts,⁵ Mukaiyama's reagent,⁶ coupling reagents,⁷ or other activating agents.⁸ *S*-Oxidized thiourea derivatives⁹ and guanylation agents¹⁰ (such as *S*-methylisothiureas, pyrazole-1-carboximidamide and its derivatives, or triflyl guanidines) are also commonly employed. Beyond the well-known¹¹ and recently¹² developed approaches, a few isocyanide-based procedures have also been established, albeit each method exclusively affords *N,N',N''*-substituted guanidines.¹³


Looking at the synthetic toolbox for the assembly of *N,N'*-disubstituted guanidines, *N*-protected *S*-methylisothiureas are often used as starting materials; however, the techniques available for the derivatization of isothiureas lack the achievable diversity (Scheme 1a).¹⁴ *N,N'*-Disubstituted guanidines can also be obtained from amines through cyanamides, but utilization of toxic cyanogen bromide and harsh conditions are required (Scheme 1b).¹⁵ The application of a commercially

available guanylation reagent di(imidazole-1-yl)methanimine offers a more convenient access to *N,N'*-disubstituted guanidines through the stepwise displacement of its imidazole groups by amines (Scheme 1c).¹⁶ Besides the necessary isolation of intermediates, the nucleophilicity of amines can strongly affect the sequence of substitution and the yield of products, or even limit the achievable substitution pattern. Therefore, the development of a facile and general one-pot procedure for the synthesis of diverse *N,N'*-substituted guanidines is still highly desired.

Classical approaches



Our approach



Scheme 1 Classical and new routes to *N,N'*-disubstituted guanidines.

^aAVIDIN Ltd, Alsó kikötő sor 11/D, Szeged, H-6726, Hungary.

E-mail: i.kanizsai@avidinbiotech.com; Tel: (+36)-62-202-107

^bDepartment of Organic Chemistry, University of Szeged, Dóm tér 8, Szeged, H-6720, Hungary

†Electronic supplementary information (ESI) available: Experimental procedures, mechanistic study, compound characterization data and copies of NMR spectra. See DOI: 10.1039/c7ob03109b

Herein, we report a new approach for the synthesis of *N,N'*-disubstituted guanidines employing *N*-chlorophthalimide, isocyanides and amines as substrates in a sequential one-pot protocol (Scheme 1d).


Results and discussion

At the outset of the study, the model reaction of *N*-chlorophthalimide (**1**) with *tert*-butyl isocyanide (**2a**) and *p*-anisidine (**4a**) was investigated (Table 1). On the basis of literature information on the addition of analogous *N*-chloroamines to isocyanides,^{13a} we presumed the need for the prior formation of the imidoyl chloride intermediate **3a**, which can react with *p*-anisidine to furnish *N*-phthaloylguanidine **5a**. Pleasingly, the addition of

N-chlorophthalimide to *tert*-butyl isocyanide took place rapidly in dichloromethane at 0 °C with full conversion. However, further reaction with *p*-anisidine led to the desired guanidine **5a** in only 12% HPLC yield, along with an unexpected side-product, which was identified as isoindolinone **6a** (Table 1, entry 1). In order to optimize the reaction conditions, various solvents were tested (entries 1–15). Acetonitrile was found to be the best medium affording **5a** in 75% HPLC yield, while non-polar solvents and ethers led to **6a** predominantly. In order to avoid the formation of the urea-type product, anhydrous solvents were used; however, “wet” acetonitrile provided **5a** in almost identical yield (entry 15). Decreasing the temperature gave **5a** in lower yields, while elevated temperatures had no significant effect on the outcome of the reaction (entries 16–20). It is noteworthy, that the replacement of **1** with *N*-bromo- or *N*-iodophthalimide resulted in complex reaction mixtures and no trace of **5a** or **6a**.¹⁷

Interestingly, performing the reaction with aromatic isocyanide **2b** under the optimized conditions failed to produce the corresponding guanidine **5b** (Table 2, entry 1). However, application of an equimolar amount of base (KO*t*Bu, DBU, Na₂CO₃ or tertiary aliphatic amine) in order to neutralize the liberated hydrogen chloride promoted the formation of **5b** (Table 2, entries 7–12). Other bases were ineffective and gave access only to isoindolinone **6a** (Table 2, entries 2–6). The best result

Table 1 Optimization of the model reaction^a




| Entry | Solvent | Temp. ^b | Yield of | |
|-------|---------------------------------|--------------------|----------------------------|----------------------------|
| | | | 5a ^c [%] | 6a ^c [%] |
| 1 | CH ₂ Cl ₂ | r.t. | 12 | 40 |
| 2 | DMSO | r.t. | 0 | 0 |
| 3 | MeOH | r.t. | 1 | 8 |
| 4 | 1,4-Dioxane | r.t. | 5 | 26 |
| 5 | THF | r.t. | 7 | 32 |
| 6 | Et ₂ O | r.t. | 11 | 24 |
| 7 | CHCl ₃ | r.t. | 16 | 30 |
| 8 | Toluene | r.t. | 4 | 46 |
| 9 | EtOAc | r.t. | 20 | 26 |
| 10 | DMF | r.t. | 26 | 3 |
| 11 | Acetone | r.t. | 45 | 18 |
| 12 | CH ₃ NO ₂ | r.t. | 48 | 8 |
| 13 | IPA | r.t. | 66 | 10 |
| 14 | MeCN | r.t. | 75 | 5 |
| 15 | MeCN ^d | r.t. | 72 | 7 |
| 16 | MeCN | −40 °C | 29 | 5 |
| 17 | MeCN | −20 °C | 30 | 13 |
| 18 | MeCN | 0 °C | 47 | 11 |
| 19 | MeCN | 40 °C | 73 | 2 |
| 20 | MeCN | 60 °C | 66 | 6 |

^a Reaction conditions: *N*-Chlorophthalimide (0.25 mmol), anhydrous solvent (0.50 ml), *t*-butyl isocyanide (1.1 equiv.), 15 min, 0 °C, then *p*-anisidine (1.2 equiv.), 2 h. ^b Temperature after the addition of *p*-anisidine. ^c Yield was determined by HPLC (each product was calibrated).

^d Non-dried solvent was used.

Table 2 The effect of base on the reaction performed with aromatic isocyanide **2b**^a



| Entry | Base | Yield of | |
|-------|---------------------------------|----------------------------|----------------------------|
| | | 5b ^b [%] | 6a ^b [%] |
| 1 | — | 0 | 32 |
| 2 | TMG | 0 | 13 |
| 3 | Proton Sponge | 0 | 23 |
| 4 | Pyridine | 0 | 45 |
| 5 | DMAP | 0 | 46 |
| 6 | <i>N</i> -Methylimidazole | 0 | 48 |
| 7 | KO <i>t</i> Bu | 1 | 16 |
| 8 | DBU | 2 | 13 |
| 9 | Na ₂ CO ₃ | 10 | 32 |
| 10 | DABCO | 39 | 28 |
| 11 | DIPEA | 47 | 41 |
| 12 | NEt ₃ | 48 | 31 |


^a Reaction conditions: *N*-Chlorophthalimide (0.25 mmol), anhydrous MeCN (0.50 ml), 4-methoxyphenyl isocyanide (1.1 equiv.), 15 min, 0 °C, then base (1.0 equiv.) and *p*-anisidine (1.2 equiv.), r.t., 2 h. ^b Yield was determined by HPLC (each product was calibrated).

was achieved when triethylamine was utilized, providing **5b** in an acceptable 48% HPLC yield (Table 2, entry 12).

To investigate the scope of the reaction and the cleavability of the phthaloyl moiety, six structurally different *N*-phthaloylguanidines **5a–f** were first synthesized by employing aliphatic (substrates **2a,c**), benzylic (substrate **2d**) or aromatic (substrates **2b,e**) isocyanides and anilines bearing both electron-donating (substrates **4a,c**) and electron-withdrawing groups (substrates **4b,d**) (Table 3, entries 1–6). The reactions proceeded smoothly in the presence of triethylamine under the optimized conditions providing **5a–f** in a non-protonated form in 28–68% isolated yields.¹⁸ To our delight, further treatment of **5a–f** with methylhydrazine completely removed the phthaloyl group in all cases, leading to the desired *N,N'*-disubstituted guanidines **7a–f** in excellent yields under mild conditions (Table 3, entries 1–6).^{19,20} Obtaining the products as hydrochloride salts facilitated the isolation procedure.


Although **5a–f** were readily formed, their isolation proved to be rather demanding and required individual chromatographic conditions (see the ESI† for details). Therefore, we decided to combine the steps of the *N,N'*-disubstituted guanidine synthesis into a sequential one-pot three-step protocol and omit the isolation of *N*-phthaloylguanidine intermediates. First, the scope of the combined method with respect to the isocyanide reagent was evaluated, using **4a** as an aniline input (Table 4). Gratifyingly, both aliphatic and benzylic, as well as

Table 4 Scope of the one-pot three-step synthesis with respect to isocyanide^a



^a Reaction conditions: *N*-Chlorophthalimide (1.0 mmol), anhydrous MeCN (2.0 ml), isocyanide (1.1 equiv.), 0 °C, 15 min, then Et₃N (1.0 equiv.) and aniline (1.2 equiv.), r.t., 2 h, then MeNHNH₂ (1.5 equiv.), 40 °C, 2 h. The products were isolated as hydrochloride salts. Isolated yield (NMR yield in parenthesis). NMR yield was determined by ¹H-NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.

Table 3 Synthesis and hydrazinolysis of *N*-phthaloyl-guanidines^{a,b}



| Entry | 2 | R ¹ | 4 | R ² | 5 | Yield ^c [%] | 7 | Yield ^c [%] |
|-------|-----------|------------------------------------|-----------|----------------|-----------|------------------------|-----------|------------------------|
| 1 | 2a | <i>t</i> -Bu | 4a | 4-MeO | 5a | 68 (73) | 7a | 98 (99) |
| 2 | 2b | 4-MeOC ₆ H ₄ | 4a | 4-MeO | 5b | 31 (49) | 7b | 94 (98) |
| 3 | 2b | 4-MeOC ₆ H ₄ | 4b | 4-Br | 5c | 28 (44) | 7c | 96 (99) |
| 4 | 2c | <i>c</i> -Hex | 4c | 3,5-Me | 5d | 29 (64) | 7d | 96 (98) |
| 5 | 2d | Bn | 4d | 4-F | 5e | 48 (54) | 7e | 97 (99) |
| 6 | 2e | 4-FC ₆ H ₄ | 4e | H | 5f | 30 (47) | 7f | 96 (99) |


^a Reaction conditions for the synthesis of **5a–f**: *N*-Chlorophthalimide (1.0 mmol), anhydrous MeCN (2.0 ml), isocyanide (1.1 equiv.), 0 °C, 15 min, then Et₃N (1.0 equiv.) and aniline (1.2 equiv.), r.t., 2 h.

^b Reaction conditions for the synthesis of **7a–f**: Guanidine **5a–f** (0.25 mmol), MeCN (0.5 ml), MeNHNH₂ (1.5 equiv.), 40 °C, 2 h, then HCl/EtOH (3 equiv.), r.t., 15 min. ^c Isolated yield (NMR yield in parenthesis). NMR yield was determined by ¹H-NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.

aromatic isocyanides could be subjected to the reaction; however, their electronic nature had a notable impact on the overall yield of the products. Benzylic and aliphatic isocyanides (including the sterically hindered 1,1,3,3-tetramethylbutyl isocyanide) provided the best results (**7a** and **7g–i**, 51–69% isolated yields), while aromatic isocyanides bearing electron-donating MeO or electron-withdrawing F substituents delivered the corresponding guanidine hydrochlorides **7b,j** and **7k** in moderate yields (33–48%).

Unfortunately, the strongly electron-deficient 4-nitrophenyl isocyanide was barely tolerated (**7l**, 22% isolated yield), while methyl isocyanoacetate, TosMIC and 2-isocyano- or 3-isocyano-pyridine did not afford the desired products.

To further explore the generality of our protocol, various anilines were subjected to the one-pot reaction applying the


Table 5 Scope of the one-pot three-step synthesis with respect to aniline^a


| Entry | 2 | R ¹ | 4 | R ² | R ³ | 7 | Yield ^b [%] |
|-------|----|---|----|----------------|------------------------|----|------------------------|
| 1 | 2a | <i>t</i> -Bu | 4f | H | 2,4-F | 7m | 73 (78) |
| 2 | 2a | <i>t</i> -Bu | 4g | Me | H | 7n | 66 (71) |
| 3 | 2f | <i>t</i> -Octyl | 4h | H | 4-CF ₃ | 7o | 55 (61) |
| 4 | 2f | <i>t</i> -Octyl | 4i | H | 3-I | 7p | 64 (68) |
| 5 | 2c | <i>c</i> -Hex | 4c | H | 3,5-Me | 7d | 56 (58) |
| 6 | 2c | <i>c</i> -Hex | 4j | H | 4-Me-3-NO ₂ | 7q | 35 (38) |
| 7 | 2d | Bn | 4d | H | 4-F | 7e | 52 (54) |
| 8 | 2d | Bn | 4k | H | 4-(NMe ₂) | 7r | 47 (51) |
| 9 | 2b | 4-MeOC ₆ H ₄ | 4e | H | H | 7s | 42 (43) |
| 10 | 2b | 4-MeOC ₆ H ₄ | 4b | H | 4-Br | 7c | 34 (42) |
| 11 | 2g | 3,4,5-MeOC ₆ H ₂ | 4e | H | H | 7t | 43 (45) |
| 12 | 2e | 4-FC ₆ H ₄ | 4e | H | H | 7f | 37 (39) |
| 13 | 2e | 4-FC ₆ H ₄ | 4l | H | 4-CN | 7u | 27 (33) |
| 14 | 2h | 4-NO ₂ C ₆ H ₄ | 4e | H | H | 7v | 35 (37) |

^a Reaction conditions: *N*-Chlorophthalimide (1.0 mmol), anhydrous MeCN (2.0 ml), isocyanide (1.1 equiv.), 0 °C, 15 min, then Et₃N (1.0 equiv.) and aniline (1.2 equiv.), r.t., 2 h, then MeNHNH₂ (1.5 equiv.), 40 °C, 2 h. The products were isolated as hydrochloride salts. ^b Isolated yield (NMR yield in parenthesis). NMR yield was determined by ¹H-NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.

previously utilized isocyanides (Table 5, entries 1–14). Interestingly, both electron-rich and electron-poor anilines were equally tolerated. The electronic effect of substituents was not significant, with the exception of the nitro group (substrate 4j, Table 5, entry 6). Even *N*-substituted anilines could be used, as exemplified by *N*-methylaniline (Table 5, entry 2). Guanidines derived from aliphatic and benzyl isocyanides were obtained in the highest yields (7d,e and 7m–r, up to 73% isolated yield), while aromatic isocyanides, especially with electron-withdrawing substituents, furnished the corresponding 7c,f and 7s–v products in lower yields ranging from 27 to 43%. These results suggest that the overall performance of our method principally depends on the reactivity of the isocyanide component.

As an extension of the one-pot protocol towards *N,N'*-di-alkylguanidines, we next surveyed the reaction of isobutylamine (8a) with 1 and 2a under standard conditions (Scheme 2). Surprisingly, no appreciable amount of guanidine 10a was produced and no trace of the expected intermediate 5g could be detected. Instead, compound 9a was isolated as the main product, presumably as a result of the subsequent reaction of 5g with an additional molecule of amine. Unfortunately, preventing the instantaneous ring-opening reaction by decreasing

**Scheme 2** Unexpected ring-opening with aliphatic amine 8a.

the temperature (–40 °C) was not successful. Nevertheless, we reasoned that product 9a might also be transformed to the desired *N,N'*-disubstituted guanidine 10a by a straightforward intramolecular cleavage. Our hypothesis was supported by the reaction mechanism of phthalimide deprotection with aliphatic amines.²¹ Indeed, simply heating 9a alone in refluxing acetonitrile for 10 h readily generated guanidine 10a in an almost quantitative yield (Scheme 2).


Afterwards, a series of primary aliphatic and aralkyl amines were tested in a combined one-pot three-step manner (Table 6, entries 1–9). Intermediates 9a–i were formed smoothly by reacting 1 with isocyanides and amines under slightly modified conditions (2.2 equivalents of primary amine were used). Subsequent heating of the reaction mixtures at reflux temperature gave complete conversion of 9a–i within 10 h furnishing, in all cases, guanidine hydrochlorides in moderate to good yields (44–81%). We were pleased to find that bifunctional aliphatic amines, such as aminoalcohol 8b and Boc-protected diaminobutane 8d, were compatible with the protocol and provided the corresponding guanidines 10b and 10d in 80% and 55% isolated yields, respectively (Table 6, entries 2 and 4). Moreover, propargylamine (8e) and the sterically demanding *tert*-butylamine (8f) were also well tolerated (Table 6, entries 5 and 6). Alternatively, *N*-alkyl-*N'*-aryl guanidines are readily accessible from aromatic isocyanides as well, as demonstrated by the synthesis of 7h (Table 6, entry 9). Although their synthetic routes are somewhat different, it is noteworthy that aliphatic and aralkyl amines provided the corresponding guanidines generally in better yields compared to anilines (see the two complementary synthesis of 7h). This, most probably, is due to their higher nucleophilicity.

Based on the above results and observations, a plausible mechanism is proposed (Scheme 3). In the first step, *N*-chlorophthalimide 1 undergoes α -addition to isocyanide to

Table 6 One-pot three-step synthesis of *N,N'*-disubstituted guanidines from aliphatic and aralkyl amines^a

| Entry | 2 | 8 | Product | | | Yield ^b [%] |
|-------|----|---|---------|--|--|------------------------|
| 1 | 2a | R ¹ = <i>t</i> -Bu | 8a | R ² = <i>i</i> -Bu | | 81 (84) |
| 2 | 2a | R ¹ = <i>t</i> -Bu | 8b | R ² = HO(CH ₂) ₅ | | 80 (85) |
| 3 | 2f | R ¹ = <i>t</i> -Octyl | 8c | R ² = C ₆ H ₅ (CH ₂) ₂ | | 72 (78) |
| 4 | 2f | R ¹ = <i>t</i> -Octyl | 8d | R ² = BocNH(CH ₂) ₄ | | 55 (62) |
| 5 | 2c | R ¹ = <i>c</i> -Hex | 8e | R ² = HCCCH ₂ | | 44 (47) |
| 6 | 2c | R ¹ = <i>c</i> -Hex | 8f | R ² = <i>t</i> -Bu | | 64 (68) |
| 7 | 2d | R ¹ = Bn | 8g | R ² = <i>c</i> -Hex | | 71 (75) |
| 8 | 2d | R ¹ = Bn | 8h | R ² = | | 53 (64) |
| 9 | 2b | R ¹ = 4-MeOC ₆ H ₄ | 8g | R ² = <i>c</i> -Hex | | 66 (70) |

^a Reaction conditions: *N*-Chlorophthalimide (1.0 mmol), anhydrous MeCN (2.0 ml), isocyanide (1.1 equiv.), 0 °C, 15 min, then Et₃N (1.0 equiv.) and amine (2.2 equiv.), r.t., 2 h, then reflux, 10 h. The products were isolated as hydrochloride salts. ^b Isolated yield (NMR yield in parenthesis). NMR yield was determined by ¹H-NMR spectroscopy with 1,3,5-trimethoxybenzene as an internal standard.

**Scheme 3** Proposed reaction mechanism.

form imidoyl chloride **B** through nitrilium species **A**. Then, nucleophilic attack of the amine takes place, which can occur either on the imidoyl carbon to provide guanidine products (route A), or on the carbonyl carbon to give intermediate **C** (route B). The subsequent rearrangement of **C** results in isoindolone **D** along with an isocyanate by-product. Finally, **D** undergoes tautomerization to afford the more stable²² isoindolinone **6**. To support the mechanism, the formation of isocyanate was confirmed by control experiments and a representative example of **B** was also isolated (see the ESI† for details).

Conclusions

In conclusion, we have developed a new and efficient synthesis of *N,N'*-disubstituted guanidines from readily available *N*-chlorophthalimide, isocyanides and amines in a sequential one-pot manner. The reactions proceed through the formation of *N*-phthaloylguanidines, which represent a novel class of guanidines. This operationally simple method tolerates both aromatic and aliphatic substrates in all possible combinations, providing general and diverse access to *N*-alkyl-*N'*-aryl, *N*-aryl-*N'*-aryl and *N*-alkyl-*N'*-alkyl guanidines with broad substrate scope.

Experimental section

General procedure for the one-pot synthesis of guanidines 7a–v

To a cooled suspension of *N*-chlorophthalimide (1.0 mmol, 182 mg) in anhydrous acetonitrile (2 mL) isocyanide (1.1 mmol) was added and stirred at 0 °C for 15 min. Then triethylamine (1.0 mmol, 140 µL) and subsequently the corresponding aniline (1.2 mmol) were added and the reaction mixture was allowed to warm to room temperature. After stirring for 2 h, methylhydrazine (1.5 mmol, 79 µL) was added, and the stirring was continued at 40 °C for 2 h. Then the reaction mixture was poured into aqueous NaOH solution (30 mL, 1 M) and extracted with chloroform (4 × 50 mL). The organic layers were combined, dried over anhydrous Na₂SO₄ and concentrated *in vacuo* until complete removal of the solvent and triethylamine. The residue was purified by flash column chromatography on neutral alumina (RediSep Rf; EtOAc:hexanes 0:100–100:0 gradient, then eluent switch to methanol:chloroform 0:100–1:10 gradient) to afford the pure guanidine base, which was then treated with HCl in ethanol (1 M, 2–3 equiv.) and stirred at room temperature for 15 min. Finally, evaporation to dryness followed by trituration with *n*-hexane or diisopropyl ether or diethyl ether (if necessary) gave pure guanidine hydrochlorides **7a–v**.

General procedure for the one-pot synthesis of guanidines 10a–h

To a cooled suspension of *N*-chlorophthalimide (1.0 mmol, 182 mg) in anhydrous acetonitrile (2 mL) isocyanide

(1.1 mmol) was added and stirred at 0 °C for 15 min. Then triethylamine (1.0 mmol, 140 µL) and subsequently primary amine (2.2 mmol) were added and the mixture was warmed to room temperature. After stirring for 2 h, the reaction mixture was warmed to reflux temperature and the stirring was continued for 10 h. Then the reaction mixture was poured into aqueous NaOH solution (30 mL, 1 M) and extracted with chloroform (4 × 50 mL). The organic layers were combined, dried over anhydrous Na₂SO₄ and concentrated *in vacuo* until the complete removal of the solvent and triethylamine. The residue was purified by flash column chromatography on neutral alumina (RediSep Rf; EtOAc:hexanes 0:100–100:0 gradient, then eluent switch to methanol:chloroform 0:100–1:10 gradient) to afford the pure guanidine base, which was then treated with HCl in ethanol (1 M, 2–3 equiv.) and stirred at room temperature for 15 min. Finally, evaporation to dryness followed by trituration with *n*-hexane or diisopropyl ether or diethyl ether (if necessary) gave pure guanidine hydrochlorides **10a–h**.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- For general reviews, see: (a) F. Saczewski and L. Balewski, *Expert Opin. Ther. Pat.*, 2009, **19**, 1417; (b) F. Saczewski and L. Balewski, *Expert Opin. Ther. Pat.*, 2013, **23**, 965; (c) L. Peterlin-Mašič and D. Kikelj, *Tetrahedron*, 2001, **57**, 7073; (d) D. Castagnolo, S. Schenone and M. Botta, *Chem. Rev.*, 2011, **111**, 5247; (e) R. G. S. Berlinck and S. Romminger, *Nat. Prod. Rep.*, 2016, **33**, 456.
- For general reviews, see: (a) T. Ishikawa and T. Kumamoto, *Synthesis*, 2006, 737; (b) J. E. Taylor, S. D. Bull and J. M. J. Williams, *Chem. Soc. Rev.*, 2012, **41**, 2109; (c) P. Selig, *Synthesis*, 2013, 703. For recent example, see: (d) X.-T. Gao, C.-C. Gan, S.-Y. Liu, F. Zhou, H.-H. Wu and J. Zhou, *ACS Catal.*, 2017, **7**, 8588.
- For selected examples, see: (a) W. Zeghida, J. Debray, S. Chierici, P. Dumy and M. Demeunynck, *J. Org. Chem.*, 2008, **73**, 2473; (b) X. Deng, H. McAllister and N. S. Mani, *J. Org. Chem.*, 2009, **74**, 5742.
- For general reviews, see: (a) C. Alonso-Moreno, A. Antiñolo, F. Carrillo-Hermosilla and A. Oterob, *Chem. Soc. Rev.*, 2014, **43**, 3406; (b) W.-X. Zhang, L. Xu and Z. Xi, *Chem. Commun.*, 2015, **51**, 254. For selected examples, see: (c) T.-G. Ong, G. P. A. Yap and D. S. Richeson, *J. Am. Chem. Soc.*, 2003, **125**, 8100; (d) F. Montilla, A. Pastor and A. Galindo, *J. Organomet. Chem.*, 2004, **689**, 993; (e) W.-X. Zhang, M. Nishiura and Z. Hou, *Synlett*, 2006, 1213; (f) H. Shen, H.-S. Chan and Z. Xie, *Organometallics*, 2006, **25**, 5515; (g) T.-G. Ong, J. S. O'Brien, I. Korobkov and D. S. Richeson, *Organometallics*, 2006, **25**, 4728; (h) Q. Li, S. Wang, S. Zhou,

- G. Yang, X. Zhu and Y. Liu, *J. Org. Chem.*, 2007, **72**, 6763; (i) W.-X. Zhang, M. Nishiura and Z. Hou, *Chem. – Eur. J.*, 2007, **13**, 4037; (j) W.-X. Zhang, D. Li, Z. Wang and Z. Xi, *Organometallics*, 2009, **28**, 882; (k) C. Alonso-Moreno, F. Carrillo-Hermosilla, A. Garcés, A. Otero, I. López-Solera, A. M. Rodríguez and A. Antiñolo, *Organometallics*, 2010, **29**, 2789; (l) D. Li, J. Guang, W.-X. Zhang, Y. Wang and Z. Xi, *Org. Biomol. Chem.*, 2010, **8**, 1816; (m) S. Pottabathula and B. Royo, *Tetrahedron Lett.*, 2012, **53**, 5156.
- 5 For selected examples, see: (a) C. Levallet, J. Lerpiniere and S. Y. Ko, *Tetrahedron*, 1997, **53**, 5291; (b) S. Cunha, B. R. de Lima and A. R. de Souza, *Tetrahedron Lett.*, 2002, **43**, 49; (c) D. H. O'Donovan and I. Rozas, *Tetrahedron Lett.*, 2011, **52**, 4117; (d) B. Kelly and I. Rozas, *Tetrahedron Lett.*, 2013, **54**, 3982.
- 6 Y. F. Yong, J. A. Kowalski and M. A. Lipton, *J. Org. Chem.*, 1997, **62**, 1540.
- 7 For selected examples, see: (a) B. R. Linton, A. J. Carr, B. P. Orner and A. D. Hamilton, *J. Org. Chem.*, 2000, **65**, 1566; (b) M. Li, L. J. Wilson and D. E. Portlock, *Tetrahedron Lett.*, 2001, **42**, 2273; (c) D. S. Ermolat'ev, J. B. Bariwal, H. P. L. Steenackers, S. C. J. De Keersmaecker and E. V. Van der Eycken, *Angew. Chem., Int. Ed.*, 2010, **49**, 9465.
- 8 For selected examples, see: (a) A. Porcheddu, L. D. Luca and G. Giacomelli, *Synlett*, 2009, 3368; (b) P. S. Dangate and K. G. Akamanchi, *Tetrahedron Lett.*, 2012, **53**, 6765; (c) S. Wangngae, M. Pattarawarapan and W. Phakhodee, *Synlett*, 2015, 1121.
- 9 For selected examples, see: (a) A. Miller and J. J. Bischoff, *Synthesis*, 1986, 777; (b) C. A. Maryanoff, R. C. Stanzione, J. N. Plampin and J. E. Mills, *J. Org. Chem.*, 1986, **51**, 1882; (c) N. Srinivasan and K. Ramadas, *Tetrahedron Lett.*, 2001, **42**, 343.
- 10 For a general review, see: (a) A. R. Katritzky and B. V. Rogovoy, *ARKIVOC*, 2005, 49. For selected examples, see: (b) C. R. Rasmussen, F. J. Villani Jr., B. E. Reynolds, J. N. Plampin, A. R. Hood, L. R. Hecker, S. O. Nortey, A. Hanslin, M. J. Costanzo, R. M. Howse Jr. and A. J. Molinari, *Synthesis*, 1988, 460; (c) K. Feichtinger, C. Zapf, H. L. Sings and M. Goodman, *J. Org. Chem.*, 1998, **63**, 3804; (d) H.-J. Musiol and L. Moroder, *Org. Lett.*, 2001, **3**, 3859.
- 11 For a general review, see: S. Tahir, A. Badshah and R. A. Hussain, *Bioorg. Chem.*, 2015, **59**, 39.
- 12 (a) C.-Y. Chen, H.-C. Lin, Y.-Y. Huang, K.-L. Chen, J.-J. Huang, M.-Y. Yeh and F. F. Wong, *Tetrahedron*, 2010, **66**, 1892; (b) R. E. Looper, T. J. Haussener and J. B. C. Mack, *J. Org. Chem.*, 2011, **76**, 6967; (c) J. Li and L. Neuville, *Org. Lett.*, 2013, **15**, 6124; (d) J. Li, H. Wang, Y. Hou, W. Yu, S. Xu and Y. Zhang, *Eur. J. Org. Chem.*, 2016, 2388; (e) M. Baeten and B. U. W. Maes, *Adv. Synth. Catal.*, 2016, **358**, 826.
- 13 (a) R. Abu-El-Halawa and J. C. Jochims, *Chem. Ber.*, 1983, **116**, 1834; (b) R. Bossio, S. Marcaccini and R. Pepino, *Tetrahedron Lett.*, 1995, **36**, 2325; (c) R. Bossio, S. Marcaccini and R. Pepino, *J. Org. Chem.*, 1996, **61**, 2202; (d) A. R. Katritzky, B. Rogovoy, C. Klein, H. Insuasty, V. Vvedensky and B. Insuasty, *J. Org. Chem.*, 2001, **66**, 2854; (e) A. Czarna, B. Beck, S. Srivastava, G. M. Popowicz, S. Wolf, Y. Huang, M. Bista, T. A. Holak and A. Dömling, *Angew. Chem., Int. Ed.*, 2010, **49**, 5352; (f) T.-H. Zhu, S.-Y. Wang, T.-Q. Wei and S.-J. Ji, *Adv. Synth. Catal.*, 2015, **357**, 823; (g) Z.-Y. Gu, Y. Liu, F. Wang, X. Bao, S.-Y. Wang and S.-J. Ji, *ACS Catal.*, 2017, **7**, 3893.
- 14 For selected examples, see: (a) H.-O. Kim, F. Mathew and C. Ogbu, *Synlett*, 1999, 193; (b) T. Suhs and B. König, *Chem. – Eur. J.*, 2006, **12**, 8150; (c) E. Tassoni, F. Giannessi, T. Brunetti, P. Pessotto, M. Renzulli, M. Travagli, S. Rajamäki, S. Prati, S. Dottori, F. Corelli, W. Cabri, P. Carminati and M. Botta, *J. Med. Chem.*, 2008, **51**, 3073.
- 15 For selected examples, see: (a) X. Bi, C. Lopez, C. J. Bacchi, D. Rattendi and P. M. Woster, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 3229; (b) L. Zhang, R. Sathunuru, T. Luong, V. Melendez, M. P. Kozar and A. J. Lin, *Bioorg. Med. Chem.*, 2011, **19**, 1541; (c) P. J. Klein, J. A. M. Christiaans, A. Metaxas, R. C. Schuit, A. A. Lammertsma, B. N. M. van Berckel and A. D. Windhorst, *Bioorg. Med. Chem.*, 2015, **23**, 1189.
- 16 (a) J. P. Ferris, C.-H. Huang and W. J. Hagan Jr., *Nucleosides Nucleotides*, 1989, **8**, 407; (b) Y.-Q. Wu, S. K. Hamilton, D. E. Wilkinson and G. S. Hamilton, *J. Org. Chem.*, 2002, **67**, 7553; (c) V. D. Jadhav and F. P. Schmidtchen, *J. Org. Chem.*, 2008, **73**, 1077; (d) A. Turočkin, R. Honeker, W. Raven and P. Selig, *J. Org. Chem.*, 2016, **81**, 4516.
- 17 Application of NCS led to the corresponding succinimidyl analogue of **5a**, however, in lower isolated yield (51%). On the other hand, no desired product was observed by means of NBS or NIS.
- 18 In order to evaluate the effectiveness of the isolation procedures, NMR yields obtained from crude reaction mixtures are also shown.
- 19 It should be noted that similar efficiencies were observed when hydrazine monohydrate was utilized, but the separation of the byproduct phthalhydrazide from *N,N'*-disubstituted guanidines was tedious.
- 20 The succinimidyl analogue of **5a** (see ref. 17) could not be transformed to the desired *N,N'*-disubstituted guanidine **7a** by hydrazinolysis (MeNHNH₂, MeCN) even at reflux temperature.
- 21 P. G. M. Wuts, *Greene's Protective Groups in Organic Synthesis*, Wiley, Hoboken, 5th edn, 2014.
- 22 S. Scherbakow, J. C. Namyslo, M. Gjikaj and A. Schmidt, *Synlett*, 2009, 1964.