Inorganic Chemistry

Dicoordinate Copper(I) Silanechalcogenolates

Iliana Medina, Heiko Jacobsen, Joel T. Mague, and Mark J. Fink*

Department of Chemistry, Tulane University, New Orleans, Louisiana 70118 Received August 21, 2006

The copper silanechalcogenolates 'Bu₃PCuESiPh₃ (1, E = O; 2, E = S; 3, E = Se) were prepared from the reaction of ['Bu₃PCu-(CH₃CN)₃]BF₄ with [Ph₃SiELi(THF)₂]₂ in acetonitrile. The compounds were obtained as colorless, crystalline, but thermally labile solids. X-ray crystallography shows that complexes 1–3 are monomeric in the solid state with no Cu···Cu interactions. The Cu atoms have either a linear or a near-linear coordination geometry in all three complexes. Interestingly, the O atom in complex 1 is also linear, which is in contrast to the highly bent S (2) and Se analogues (3). Density functional theory calculations suggest that both the linear geometry of 1 and an associated extremely short Cu–O distance [1.769(4) Å] are not the result of π delocalization but are the result of a fine balance of electrostatic interaction and Pauli repulsion.

Cu^I complexes with the group 16 elements are of great interest as models for metalloprotein active sites,¹ as catalysts,² and for use as photovoltaic materials.^{3,4} Because of the ability of the group 16 atom to adopt either terminal or bridge bonding modes, complexes of the type L_nCuX (L = monodentate phosphine; X = O, S, Se) display a rich coordination chemistry. Bridge bonding allows for the formation of Cu-based oligomers, cluster compounds, and polymers.⁵ However, monomeric Cu^I complexes are relatively rare and generally exist either as loose bimolecular aggregates with short Cu···Cu or Cu···X contacts or as structures with a cyclic Cu₂X₂ core.⁶ Nonetheless, discrete monomeric copper(I) chalcogenides may be isolated in

Scheme 1

$^{\prime}\mathrm{Bu}_{3}\mathrm{P}$ + [Cu(CH ₃ CN) ₄]BF ₄	CH ₃ CN	->	[^t Bu ₃ PCu(CH ₃ CN) ₃]BF ₄
$\frac{1}{2}$ [Ph ₃ SiELi(thf) ₂] ₂ + ['Bu ₃ PCu(CH ₃ CN) ₃]BF ₄	CH3CN 0°C	->	 'Bu₃PCuESiPh₃ + LiBF₄ (1) E = O; yield: 75% (2) E = S; yield: 94% (3) E = Se; yield 60%

systems in which the Cu atom is coordinatively saturated by phosphine ligands and the chalcogen is sterically protected. For instance, the solid-state structures of the copper chalcogenide complexes $(R_3P)_3CuESiMe_3$ (R = Et, "Pr; E = S, Se, Te) show discrete monomeric units with a coordination number of 4 about the Cu atom.⁷ These thermally labile complexes are prone to the removal of the silvl group via acetoxydesilvlation or thermolysis, resulting in the formation of clusters with extensive chalcogen bridge bonding.⁸

We now report the *first* series of monomeric *dicoordinate* copper(I) silanolates and silanechalcogenolate complexes, ^{*t*}-Bu₃PCuESiPh₃ (E = O, S, Se). The sterically demanding 'Bu₃P ligand in conjunction with the bulky silyl chalcogenide ligand both promotes coordinative unsaturation at the Cu center and also prevents association with other molecules. The empty orbitals on the Cu atom are, therefore, potentially available for novel π -bonding interactions with the lone pairs of the chalcogen atoms.

The compounds 'Bu₃PCuESiPh₃ (1, E = O; 2, E = S; 3, E = Se) have been synthesized in moderate to high yields from the reaction of ['Bu₃PCu(CH₃CN)₃]BF₄ with [Ph₃SiELi-(THF)₂]₂ (E = O, S, Se) in acetonitrile (Scheme 1).

Compounds 1-3 were isolated as colorless crystalline, airsensitive solids that decompose before melting (~100 °C), generating black powders. The thermal stability of 1 and 2 is significantly greater than that observed for 3. Whereas compounds 1 and 2 can be handled briefly at room temperature, compound 3 slowly decomposes in solution even at 0 °C.⁹ All compounds gave satisfactory analytical

^{*} To whom correspondence should be addressed. E-mail: fink@mailhost.tcs.tulane.edu.

Winge, D. R.; Dameron, C. T.; George, G. N.; Pickering, I. J.; Dance, I. G. Bioinorg. Chem. Copper 1993, 110.

^{(2) (}a) Grigg, R.; Cooper, D. M.; Holloway, S.; McDonald, S.; Millington, E.; Sarker, M. A. B. *Tetrahedron* 2005, *81*, 8677. (b) Percec, V.; Grigoras, C. J. Polym. Sci., Part A 2005, *43*, 3920. (c) Corma, A.; Palomares, A. E.; Rey, F.; Márquez, F. J. Catal. 1997, 170, 140. (d) Linder, J.; Garcia, M. A. V.; Achdev, A.; Schwank, J. J. Chem. Soc., Chem. Commun. 1989, 1833.

⁽³⁾ Komuro, T.; Kawaguchi, H.; Tatsumi, K. Inorg. Chem. 2002, 41, 5083.

 ^{(4) (}a) Hirpo, W.; Dhingra, S.; Sutorik, A. C.; Kanatzidas, M. G. J. Am. Chem. Soc. 1993, 115, 1597. (b) Banger, K. K.; Hollingsworth, J. A.; Harris, J. D.; Cowan, J.; Buhro, W. E.; Hepp, A. F. Appl. Organomet. Chem. 2002, 16, 617. (c) Raffaelle, R. P.; Castro, S. L.; Hepp, A. F.; Bailey, S. G. Prog. Photovolt. Res. Appl. 2002, 10, 433. (d) Stanbery, B. J. Crit. Rev. Solid State Mater. Sci. 2002, 27, 73.

⁽⁵⁾ Dance, I. G.; Scudder, M. L.; Fitzpatrick, L. J. Inorg. Chem. 1985, 24, 2547.

^{(6) (}a) Liu, X.-Y.; Mota, F.; Alemany, P.; Novoa, J. J.; Alvarez, S. Chem. Commun. 1998, 1149. (b) Carvajal, M. A.; Novoa, J. J.; Alverez, S. J. Am. Chem. Soc. 2004, 126, 1465.

⁽⁷⁾ Tran, D. T. T.; Corrigan, J. F. Organometallics 2000, 19, 5202.

⁽⁸⁾ Tran, D. T. T.; Taylor, N. J.; Corrigan, J. F. Angew. Chem., Int. Ed. 2000, 39, 935.

⁽⁹⁾ Because of their thermal instability, all compounds should be stored at -25 °C.



Figure 1. Molecular structure of **1** with atomic numbering (ORTEP, 50% probability ellipsoids; H atoms are omitted for clarity). Selected bond lengths (Å) and angles (deg) for **1**: O-Cu, 1.769(4); Cu-P, 2.1496(17); O-Si, 1.578(4); Cu-O-Si, 180.00(1); O-Cu-P, 180.00(1).

and spectroscopic data.¹⁰ Interestingly, the ³¹P{¹H} NMR spectra of all three complexes show a single broadened resonance due to quadrupolar interactions with the multiple I > 1/2 Cu isotopes in a low-coordinate environment.¹¹

X-ray-quality crystals of complexes 1-3 were grown from benzene/pentane (1:3) mixtures at low temperatures. The X-ray structures of 1-3 reveal that these compounds are monomeric in the solid state and possess a linear coordination environment about the Cu atom. The isolation of coordinatively unsaturated Cu^I complexes is a likely result of the combined steric demand of both the phosphine and the chalcogenide moieties.

The copper siloxide (1) crystallizes in the $P\bar{3}$ point group; the linear P-Cu-O-Si skeleton lies exactly on a crystallographic 3-fold rotation axis (Figure 1). Two monomeric units of 1 pack in a head-to-tail fashion in the unit cell. Although there are some nonbonding interactions between the peripheral phenyl and isopropyl groups, there are no apparent d¹⁰-d¹⁰ Cu···Cu interactions (shortest contact, 8.077 Å).¹² This contrasts with an analogous linear gold siloxide system, Ph₃PAuOSiPh₃, which exists as head-to-tail monomers in the unit cell but with a Au···Au interaction distance of 3.376(2) Å.¹³ The observed Cu-O bond distance of 1.769-(4) Å in **1** is significantly shorter than the Cu–O distances reported for related compounds (hfac)Cu(PMe₃) [1.990(8) and 2.034(7) Å], (dmp)Cu(PMe₃) [1.941(5) and 2.030(5) Å], and (dmb)Cu(PMe₃) [1.985(2) Å] (where hfac = hexafluoroacetylacetonate, dmp = dipivaloylmethanate, and dmb = dibenzoylmethanate).14

(13) Bauer, A.; Schneider, W.; Angermeier, K.; Schier, A.; Schmidbauer, H. Inorg. Chem. Acta 1996, 251, 249.



Figure 2. Molecular structure of **2** with atomic numbering (ORTEP, 50% probability ellipsoids; H atoms are omitted for clarity). Selected bond lengths (Å) and angles (deg) for **2**: S–Cu, 2.1578(11); Cu–P, 2.2016(10); S–Si, 2.0916(14); Cu–S–Si, 97.63(5); P–Cu–S, 174.12(4).



Figure 3. Molecular structure of **3** with atomic numbering (ORTEP, 50% probability ellipsoids; H atoms are omitted for clarity). Selected bond lengths (Å) and angles (deg) for **3**: Se–Cu, 2.2650(6); Se–Si, 2.2326(10); Cu–P, 2.2014(10); Cu–Se–Si, 95.27(3); Se–Cu–P, 173.28(3).

The molecular geometries of the S (2) and Se (3) analogues are shown in Figures 2 and 3, respectively. These isomorphic structures of 2 and 3 show a linear geometry at the Cu atom similar to the siloxide 1; however, they strikingly differ from the linear siloxide structure by the nearly orthogonal angle at the chalcogen atom [97.63(5)° for 2 and 95.27(3)° for 3]. Complex 3 is one of the few examples of a terminally bonded copper(I) selenolate complex.

The length of the Cu–S bond in **2** [2.1578(11) Å] is in the normal range for homoleptic copper thiolates: [Cu-(SSiMe₂Bu^{*i*})]₄¹⁵ [2.1666(4) Å], [Cu(SSiPh'Bu₂)]₄ [2.1597-(7) Å],¹⁶ and [Cu(SSiPh₃)]₄¹⁷ [2.1633(5) Å]. However, the Cu–S bond of **2** is significantly shorter than that reported for (Et₃P)₃CuSSiMe₃ [2.4022(5) Å] and ("Pr₃P)₃CuSSiMe₃ [2.3970(10) Å]. Complex **3** also shows a shorter Cu–Se bond length when compared with the Cu–Se bond lengths in the analogous complexes (Et₃P)₃CuSeSiMe₃ [2.5124(6) Å] and ("Pr₃P)₃CuSeSiMe₃ [2.5160(4) Å].⁷ The shorter Cu–chalcogen bond lengths in **2** and **3** may be attributed to a combination of diminished sterics at the Cu atom (dicoor-

- (16) Kückmann, T. I.; Hermsen, M.; Bolte, M.; Wagner, M.; Lerner, H.-W. *Inorg. Chem.* **2005**, *44*, 3449.
- (17) Komuro, T.; Kawaguchi, H.; Tatsumi, K. Inorg. Chem. 2002, 41, 5083.

⁽¹⁰⁾ **Compound Characterization. 1.** ¹H NMR (δ , C₆D₆): 0.98 (d, 27H, ³J_{H-P} = 13 Hz), 7.32 (m, 9H), 8.16 (m, 6H). ¹³C NMR (δ , C₆D₆): 142.5 (o-Ph), 135.6 (m-Ph), 128.4 (p-Ph), 36.2 (d, C(CH₃)₃), 31.6 (d, C(CH₃)₃). ³¹P NMR (δ , C₆D₆): 72.17. ²⁹Si NMR (δ , C₆D₆): -24.29. Elem anal. Calcd: C, 66.57; H, 7.82. Found: C, 66.65; H, 7.25. Mp: 126 °C (dec). **2.** ¹H NMR (δ , C₆D₆): 0.90 (d, 27H, ³J_{H-P} = 13 Hz), 7.20 (m, 9H), 8.12 (m, 6H). ¹³C NMR (δ , C₆D₆): 141.6 (o-Ph), 135.9 (m-Ph), 128.4 (p-Ph), 36.3 (d, C(CH₃)₃), 31.7 (d, C(CH₃)₃). ³¹P NMR (δ , C₆D₆): 62.4. Elem anal. Calcd: C, 64.65; H, 7.59. Found: C, 64.53; H, 7.07. Mp: >100 °C (dec). **3.** ¹H NMR (δ , C₆D₆): 0.90 (d, 27H, ³J_{H-P} = 13 Hz), 7.25 (m, 9H), 7.75 (m, 6H). ¹³C NMR (δ , C₆D₆): 140.9 (o-Ph), 136.2 (m-Ph), 128.6 (p-Ph), 36.5 (d, C(CH₃)₃), 31.8 (d, C(CH₃)₃). ³¹P NMR (δ , C₆D₆): 61.2. Mp: >100 °C (dec).

⁽¹¹⁾ Verkade, J. G.; Mosbo, J. E. In *Phosphorus-13 NMR Spectroscopy in Stereochemical Analysis*; Verkade, J. G., Quinn, L., Eds.; VCH Publishers: Deefield, FL, 1987; p 441.

⁽¹²⁾ Pyykö, P. Chem. Rev. 1997, 97, 597.

⁽¹⁴⁾ Shin, H. K.; Chi, K. M.; Farkas, J.; Hampden-Smith, M. J.; Kodas, T. T.; Duesler, E. N. *Inorg. Chem.* **1992**, *31*, 424.

⁽¹⁵⁾ Komuro, T.; Matsuo, T.; Kawaguchi, H.; Tatsumi, K. *Dalton Trans.* **2004**, 1618.

COMMUNICATION

dinate vs tetracoordinate) and an increase in the nominal "s" character at the Cu ("sp" vs "sp³").

Trends in the Cu–E–Si bond angles for complexes **1–3** closely parallel those for the Si–E–Si bond angles of (Ph₃-Si)₂E. Thus, for (Ph₃Si)₂E, θ (Si–E–Si) varies in the series $E = O (180^{\circ})$,¹⁸ S [111.94(8)°],¹⁹ Se [110.51(2)°].²⁰ In addition, the unique bonding in the Cu–O–Si fragment of **1** is manifested not only by its linear geometry but also in some *extremely* short Cu–O and Si–O bond distances. To our knowledge, the Cu–O bond distance of 1.769(4) Å in **1** is the shortest known for a Cu^I–O bond.²¹ The very short Si–O bond distance of 1.578(4) Å is also notable. In contrast, Cu–E and Si–E (E = S and Se) bond distances found for **2** and **3**, respectively, are more in the expected range.

Traditionally, the siloxane linkage is believed to have considerable multiple-bond character because of dative π bonding from the O lone pairs to low-lying acceptor orbitals on the Si, thus giving rise to shorter than expected Si–O bonds and near-linear geometries about the O atom. However, density functional theory (DFT) calculations on H₃-PCuOSiH₃ suggest that almost no π bonding occurs between the O and Cu atoms and that the primary interaction is electrostatic in nature.²² The combined effects of ionic bonding and Pauli repulsion result in a very flexible bending

- (19) Wojnowski, W.; Peters, K.; Peters, E. M.; Von Schnering, H. G. Z. Anorg. Allg. Chem. 1985, 525, 121.
- (20) Medina, I.; Mague, J. T.; Donahue, J. P.; Fink, M. J. Acta Crystallogr., Sect. E 2005, E61, 02687.
- (21) A very short Cu–O bond is also found for the dicoordinate cuprate (o-xylyl–O)₂Cu⁻, 1.798(8) Å. Fiaschi, P.; Floriani, C.; Pasquali, M.; Chiesi-Villa, A.; Guastini, C. *Inorg. Chem.* **1986**, 25, 462.

potential at a significantly bent Cu–O–Si bond. Thus, the equilibrium bond angle θ (Cu–O–Si) found for H₃PCuOSiH₃ is 135° but requires only 3 kJ/mol to deform to a linear geometry. In the case of **1**, it is likely that crystal packing forces would be sufficient to stabilize the observed linear geometry. In contrast, DFT calculations indicate predominately covalent interactions for H₃PCuSSiH₃ and H₃-PCuSeSiH₃ with nearly orthogonal bond angles (97.1° and 93.1°, respectively) and a large difference in energy between the bent ground state and the linear transition state (67 and 85 kJ/mol, respectively).²³

The chemistry of low-coordinate copper(I) silanechalcogenolates is currently being investigated. These coordinatively unsaturated complexes may serve as building blocks to heterobimetallic complexes or as chemical vapor deposition precursors to CuE (E = O, S, Se).

Acknowledgment. Financial support from NASA (Grant NCC3-946) through the Tulane Institute for Macromolecular Engineering and Science, NASA (Grant NCC3-97) through the Ohio Aerospace Institute, and the Louisiana Board of Regents (LaSPACE Grant C164092) is gratefully acknowledged. We thank Prof. L. Cavallo for providing access to the MoLNaC Computing Facilities at Dipartimento di Chimica, Università di Salerno, Salerno, Italy.

Supporting Information Available: Synthetic procedures and crystallographic data (CIF) for compounds 1-3. This material is available free of charge via the Internet at http://pubs.acs.org.

IC061579Q

⁽¹⁸⁾ Glidewell, C.; Liles, D. C. Acta Crystallogr., Sect. B 1978, B34 (1), 124.

⁽²²⁾ Ionic bonding has also been proposed as an alternative model for siloxane bonds. Apeloig, Y. *Theoretic Aspects of Organosilicon Compounds*. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappaport, Z., Eds.; Wiley: Chichester, U.K., 1989; Vol. 1, Chapter 2, p 57.

⁽²³⁾ Details of the calculations will be published in a full paper.