This article was downloaded by: [University of Auckland Library] On: 06 November 2014, At: 12:25 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Effect of lodine on the Antimicrobial Activity of New Spiro and Isolated β-Lactam Thiazolidinone Derivatives

N. A. A. El-Kanzi^a, A. K. Khalafallah^a & M. Younis^b ^a Chemistry Department, Faculty of Science, Aswan, Egypt

^b Botany Department, Faculty of Science, Aswan, Egypt Published online: 19 Apr 2007.

To cite this article: N. A. A. El-Kanzi , A. K. Khalafallah & M. Younis (2007) Effect of lodine on the Antimicrobial Activity of New Spiro and Isolated β -Lactam Thiazolidinone Derivatives, Phosphorus, Sulfur, and Silicon and the Related Elements, 182:5, 1163-1181, DOI: <u>10.1080/10426500601149929</u>

To link to this article: http://dx.doi.org/10.1080/10426500601149929

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages,

and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Effect of lodine on the Antimicrobial Activity of New Spiro and Isolated β -Lactam Thiazolidinone Derivatives

N. A. A. El-Kanzi A. K. Khalafallah

Chemistry Department, Faculty of Science, Aswan, Egypt

M. Younis

Botany Department, Faculty of Science, Aswan, Egypt

The reaction of 1,4-naphthoquinone with a mixture of urea and ethylcyanoacetate in ethanol containing a piperidine catalyst afforded compound 1, which reacted with acetonitrile and steam of hydrogen chloride to yield compound 2, which readily condensed with different aromatic amine to give 3a-c. Compound 3a-c reacted with chloroacetyl chloride and/or mercaptoacetic acid to give spiro β -Lactam 4a-c, and spiro thiazolidinone 5a-c also isolated β -Lactam and thiazolidinone 7a-d and 8a-d prepared from a reaction of isolated schiff base 6a-d with chloroacetyle chloride to give to give spiro β -Lactam 4a-c, and spiro thiazolidinote 5a-c also isolated β -Lactam and thiazolidinote 7a-d and 8a-d prepared from a reaction of isolated schiff base 6a-d with chloroacetyle chloride to give compound 9, which used to synthesis of new shiff bases 10a-c and 13a-d. The new schiff bases 10a-c and 13a-d reacted with chloroacetyl chloride or mercaptoacetic acid to give B-Lactams 11a-c and 14a-d and thiazolidinone derivatives 12a-c and 15a-d.

Keywords Spiro B lactam, spiro thiazolidinone, shief base

INTRODUCTION

The explosive interest in β -Lactams and related derivatives stems from the fact that some monocyclic β -Lactams exhibit antibacterial activities. Some examples comprise of the naturally occuring monobactams and nocardicins.¹ Contrary to penicillins, cephalosporins, or nocardicins, monobactams were not produced by fungi or actinomycetes, but from bacteria, for example, *Bacillus pseudomonas*.² Nocardicins proved the relationship with cephalosporins and penicillins via the corresponding β -configuration at C₍₃₎, but they have no therapeutic significance. Synthetic oxamazins,^{3–5} thiamazins,⁶ and monosulfactams⁷ showed some antibacterial activity, which raised again a major interest

Received December 4, 2004; accepted October 19, 2006.

Address correspondence to N. A. A. El-Kanzi, Chemistry Department, Faculty of Science, South Valley University, Aswan, Egypt. E-mail: kanzi2nour@yahoo.com

in the area of 2-azetidinone chemistry. Also thiazole derivatives such as pencillins, which have fused thiazolidine and β -Lactam rings, were known and used as potent antibiotics.⁸

RESULTS AND DISCUSSION

Our initial strategy in this research project for the synthesis of different structural formulas corresponding to the isolated, spiro, and fused systems depended upon the synthesis of the new fused 2,4diamino-5, 10-dioxo-1,5,10,11-tetrahydrobenz[g]quinoline-3-ethyl carboxylate 1, which was synthesized by a cycloaddition reaction of equimolecular ratios of the appropriate urea and ethylcyano acetate with 2 mole of 1,4-naphthoquinone (one mole used as an oxidizing agent), in ethanol containing piperidine as catalyst. At first our theortical conception of the obvious cycloaddition reaction led to the formation of compound 1, but the experimental evidence that depends on the different types of analysis to the reaction product proved that the cycloaddition reaction leads to the formation of compound 1 through the electronic cyclization according to the suggested mechanism (Scheme 1 illustrates the formation of compound 1).

The structure of the new synthesized compound 1 was confirmed by its elemental analysis and IR which revealed a carbonyl group of ester at 1745 cm⁻¹ and ¹H NMR spectra, which revealed the presence of NH group at δ 11.02, NH₂ group at δ 6.82, NH₂ group at δ 4.53, a triplet at δ 1.23 assigned for methyl group, and quartet at δ 3.55 for CH₂ group. The mass spectrum showed the molecular ion peak (M⁺, C₁₆H₁₅O₄N₃) at mz 313. The inteligible bases of ¹H NMR led us to descide that the isolable structural formula produced from the reaction of the appropriate urea and malononitrile with 1,4-naphthoquinone is the structural formula of compound 1 because of the appearance of the signal at δ 3.21 as a result of the fussion of 1,4-naphthoquinone with the unsaturated substituted piperidine ring.

Compound **2** was prepared by the reaction of compound **1** and acetonitrile with a stream of dry hydrogen chloride gas in the presence of dioxane for about 12 h, and the reaction mixture was heated on a water bath for 2 h after passing hydrogen chloride gas. The structure of the new synthesized compound **2** was confirmed by its elemental analysis and IR spectra, which revealed a signal of the carbonyl group at 1680 cm⁻¹ and ¹H NMR spectra, which revealed the presence of NH group at δ 10.56, NH₂ group at δ 6.81, singlet signal at 1.25 for the methyl group. The absence of the signals assigned for the ethyl group indicate the structure of compound **2**; the mass spectrum showed the molecular ion peak (M⁺, C₁₆H₁₂O₃N₄) at m/z 308.





Compound **2** underwent a fussion reaction with ethyl iodide to give compound **9**. The structure of compound **9** was confirmed by their elemental analysis, IR, ¹H NMR spectra, and mass spectra (cf. Table I, II, and III).

It is very important to know that the formation of Schiff bases corresponding to the newly heterocyclic compounds is the cornerstone in the synthesis of the corresponding spiro and isolated β -Lactams and thiazolidinone compounds. The activity of the carbonyl group in position 4 and also the activity of the methyl group at C_2 render compound 2 to react with different aromatic amine and different nitroso compound to give new Schiff bases **3a-c** and **6a-d**. The structures of these newly synthesized Schiff bases **3a-c** and **6a-d** were confirmed by their elemental analysis, IR, ¹H NMR, and mass spectra (cf. Tables I, II, and III.) The activity of the azamethine center in compound **3a-c** is more avaliable than the activity of the NH group toward the addition process of chloroacetyl chloride, and this mentioned phenomena is due to the presence of the π electron, which makes the foundation of the δ positive and δ negative charge on the carbon and nitrogen atom, respectively. more easy than the presence of this phenomena on the NH group in which the bonding between nitrogen and hydrogen wheather strong according to the nature of this bonding which leads to decreasing of the mobility desire of the hydrogen atom of this NH group. Thus compound **3a-d** reacted with chloroacetyl chloride or mercaptoacetic acid to give spiro β -Lactams and spiro thiazolidinone compounds⁹ 4a–c and 5a–c. The structures of these spiro compounds 4a-c and 5a-c were confirmed by their elemental analysis, IR, ¹H NMR and mass spectra (cf. Tables I, II, III). The formation of spiro azitidine derivatives **4a–c** was suggested to proceed according to the following mechanisms (Scheme 2).

It is reported that β -Lactams play a vital role in medicinal chemistry as a key intermediate for the synthesis of penicillin, cephalosporins, and their analogs. On the other hand, thiazolidinones derivatives are used in biological activities such as bactericidal, pesticidal, and fungicidals, insecticidal activities.^{10,11} Thus the activity of azamethine center in compound **6a–d** renders it avaliable to react with chloroacetyl chloride and/or mercaptoacetic acid to give new isolated β -Lactams and thiazolidinone compounds **7a–d** and 8a–d. The structures of compounds **7a–d** and **8a–d** were confirmed by their elemental analysis, IR, ¹H NMR and mass spectra [cf. Tables 1,2,3].

The application of heterocyclic compounds in various aspects is beyond estimation, and there is always continuous need for the discovery of new heterocyclic compounds, which are used in industrial and various biological fields. Thus we used the new compound **9** in the synthesis of spiro and isolated β -lactams and thiazolidinone compounds. Compound

| Compound | | | Crystal | | | |
|----------|---------|------|---------|--|-----------|-----|
| No. | Yield % | M.P. | solvent | Mol. formula | (Mol. Wt) | MS |
| 1 | 75 | >300 | DMF | C ₁₆ H ₁₅ O ₄ N ₃ | (313.31) | 313 |
| 2 | 65 | >300 | DMF | $C_{16}H_{12}O_{3}N_{4}$ | (308.30) | 308 |
| 3a | 63 | >300 | DMF | $C_{22}H_{17}O_{2}N_{5}$ | (383.41) | 383 |
| 3b | 60 | >300 | DMF | $C_{22} = 17 O_2 N_5$ | (399.41) | 399 |
| 3c | 66 | >300 | DMF | $C_{22}H_{16}O_4N_5$ | (414.40) | 414 |
| 4a | 70 | >300 | DMF | $C_{24}H_{18}O_{2}N_{5}Cl$ | (459.90) | 459 |
| 4b | 69 | >300 | DMF | $C_{24}H_{10}O_4N_5Cl$ | (475.90) | 475 |
| 4c | 73 | >300 | DMF | $C_{24}H_{17}O_5N_5Cl$ | (490.89) | 490 |
| 5a | 75 | >300 | DMF | $C_{24}H_{10}O_{2}N_{5}S$ | (457.51) | 457 |
| 5b | 72 | >300 | DMF | $C_{24}H_{19}O_4N_5S$ | (473.51) | 473 |
| 5c | 74 | >300 | DMF | C24H18O5N5S | (488.50) | 488 |
| 6a | 67 | >300 | DMF | $C_{24} = 13 \circ 3 \cdot 35 \circ 35 \circ 35 \circ 35 \circ 35 \circ 35 \circ 35$ | (463.37) | 463 |
| 6b | 68 | >300 | DMF | $C_{26}H_{17}O_4N_5$ | (463.37) | 463 |
| 6c | 70 | >300 | DMF | $C_{20} H_{15} O_4 N_5$ | (413.39) | 413 |
| 6d | 66 | >300 | DMF | $C_{24}H_{20}O_{2}N_{6}$ | (440.46) | 440 |
| 7a | 69 | >300 | DMF | C241205N5Cl | (539.94) | 539 |
| 7b | 68 | >300 | DMF | $C_{24}H_{18}O_5N_5Cl$ | (539.94) | 539 |
| 7c | 65 | >300 | DMF | $C_{24}H_{16}O_5N_5Cl$ | (489.88) | 489 |
| 7d | 77 | >300 | DMF | C26H21O4N6Cl | (516.87) | 516 |
| 8a | 76 | >300 | DMF | $C_{28}H_{10}O_5N_5S$ | (537.55) | 537 |
| 8b | 74 | >300 | DMF | $C_{28}H_{19}O_5N_5S$ | (537.55) | 537 |
| 8c | 73 | >300 | DMF | $C_{24}H_{17}O_5N_5S$ | (487.49) | 487 |
| 8d | 72 | >300 | DMF | $C_{26}H_{22}O_4N_6S$ | (514.56) | 514 |
| 9 | 75 | >300 | DMF | $C_{18}H_{17}O_3N_4I$ | (464.26) | 464 |
| 10a | 72 | >300 | DMF | C ₂₄ H ₃₂ O ₂ N ₅ I | (549.45) | 549 |
| 10b | 73 | >300 | DMF | C24 32 22 35 | (565.45) | 565 |
| 10c | 70 | >300 | DMF | $C_{24}H_{31}O_4N_5I$ | (580.44) | 580 |
| 11a | 70 | >300 | DMF | C ₂₆ H ₂₃ O ₃ N ₅ ClI | (615.86) | 615 |
| 11b | 71 | >300 | DMF | $C_{26}H_{23}O_4N_5CII$ | (631.86) | 631 |
| 11c | 73 | >300 | DMF | C ₂₆ H ₂₂ O ₅ N ₅ ClI | (646.85) | 646 |
| 12a | 69 | >300 | DMF | $C_{26}H_{24}O_3N_5SI$ | (613.47) | 613 |
| 12b | 70 | >300 | DMF | $C_{26}H_{24}O_4N_5SI$ | (629.47) | 629 |
| 12c | 72 | >300 | DMF | $C_{26}H_{23}O_5N_5SI$ | (644.46) | 644 |
| 13a | 73 | >300 | DMF | $C_{28}H_{22}O_4N_5I$ | (619.41) | 619 |
| 13b | 71 | >300 | DMF | $C_{28}H_{22}O_4N_5I$ | (619.41) | 619 |
| 13c | 69 | >300 | DMF | $C_{24}H_{20}O_4N_5I$ | (569.35) | 569 |
| 13d | 68 | >300 | DMF | $C_{26}H_{25}O_3N_6I$ | (596.42) | 596 |
| 14a | 70 | >300 | DMF | $C_{30}H_{23}O_5N_5ClI$ | (695.90) | 695 |
| 14b | 72 | >300 | DMF | C ₃₀ H ₂₃ O ₅ N ₅ ClI | (695.90) | 695 |
| 14c | 69 | >300 | DMF | $C_{26}H_{21}O_5N_5ClI$ | (645.84) | 645 |
| 14d | 71 | >300 | DMF | C ₂₈ H ₂₆ O ₄ N ₆ ClI | (672.91) | 672 |
| 15a | 69 | >300 | DMF | $C_{30}H_{24}O_5N_5SI$ | (693.51) | 693 |
| 15b | 67 | >300 | DMF | $C_{30}H_{24}O_5N_5SI$ | (693.51) | 693 |
| 15c | 70 | >300 | DMF | $C_{26}H_{22}O_5N_5SI$ | (743.45) | 743 |
| 15d | 68 | >300 | DMF | $\mathrm{C}_{28}\mathrm{H}_{27}\mathrm{O}_{4}\mathrm{N}_{6}\mathrm{SI}$ | (670.52) | 670 |
| | | | | | | |

TABLE I

| Compound No. | ¹ H NMR (DMSO) ppm | | | | | | | |
|--------------|--|--|--|--|--|--|--|--|
| 1 | $\delta1.23(t, CH_3), \delta$ 3.55(q, CH ₂), δ 4.53(s, NH ₂), 6.82(s, NH ₂), δ 11.02(s, NH), 8.1–7.02(m, 5H aromatic hydrogen) | | | | | | | |
| 2 | δ1.25(s, CH ₃), δ 6.81(s, NH ₂), δ 10.56(brs, NH), 8.01–7.1(m,7H aromatic protons) | | | | | | | |
| 3a | δ1.24(s, CH ₃), δ 6.79(s, NH ₂), δ 10.55(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |
| 3b | δ1.259(s, CH ₃), δ 6.77(s, NH ₂), δ 10.56(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |
| 3c | δ1.26(s, CH ₃), δ 6.78(s, NH ₂), δ 10.57(brs, NH), 8.1–7.01(m, 10H aromatic protons) | | | | | | | |
| 4a | δ1.24(s, CH ₃), δ 6.75(s, NH ₂), δ 10.43(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 4b | δ1.23(s, CH ₃), δ 6.73(s, NH ₂), δ 10.44(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 4c | δ 1.24(s, CH ₃), δ 6.74(s, NH ₂), δ 10.45(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |
| 5a | δ1.24(s, CH ₃), 2.5(s, CH ₂ protons), δ 6.76(s, NH ₂), δ 10.55(brs, NH), 8.1-7.01(m, 10H aromatic protons) | | | | | | | |
| 5b | δ1.23(s, CH ₃), 2.5(s, CH ₂ protons), δ 6.75(s, NH ₂), 10.54(brs, NH), 8.1-7.01(m, 10H aromatic protons) | | | | | | | |
| 5c | δ1.24(s, CH ₃), 2.5(s, CH ₂ protons), δ 6.75(s, NH), 10.54(brs, NH), 8.1-7.01(m, 9H aromatic protons) | | | | | | | |
| 6a | $\delta 6.75(s, NH_2), \delta 10.25(brs, NH), \delta 8.1-7.01(m, 14H aromatic protons)$ | | | | | | | |
| 6b | $\delta 6.73(s, NH_2), \delta 10.26(brs, NH), 8.1-7.01(m, 14H aromatic protons)$ | | | | | | | |
| 6c | $\delta 6.69(s, NH_2), \delta 10.27(brs, NH), 8.1-7.01(m, 12H aromatic protons)$ | | | | | | | |
| 6d | δ1.25(ds, 2CH ₃), δ 6.66(s, NH ₂), δ 10.26(brs, NH), 8.1–7.01(m, 11H aromatic protons). | | | | | | | |
| 7a | $\delta 6.65(s, NH_2), \delta 10.23(brs, NH), 8.1-7.01(m, 15H aromatic protons)$ | | | | | | | |
| 7b | δ6.67(s, NH ₂), δ 10.24(brs, NH), 8.1–7.01(m, 15H aromatic protons) | | | | | | | |
| 7c | δ6.66(s, NH ₂), 10.25(brs, NH), 8.1–7.01(m, 13H aromatic protons) | | | | | | | |
| 7d | δ1.23(ds, 2CH ₃), δ 6.65(s, NH ₂), 10.25(brs, NH), 8.1–7.01(m, 15H aromatic protons) | | | | | | | |
| 8a | $\delta 2.5$ (s, CH ₂ of thiazolidinone), $\delta 6.6$ (s, NH ₂), $\delta 10.26$ (brs, NH), $8.1-7.01$ (m, 14H aromatic protons) | | | | | | | |
| 8b | $\delta 2.5$ (s, CH ₂ of thiazolidinone), $\delta 6.65$ (s, NH ₂), $\delta 10.25$ (brs, NH), 8.1–7.01(m, 14H aromatic protons) | | | | | | | |
| 8c | $\delta 2.5(s, CH_2 \text{ of thiazolidinone}), \delta 6.64(s, NH_2), \delta 10.23(brs, NH), 8.1-7.01(m, 12H aromatic protons)$ | | | | | | | |
| 8d | δ1.23(ds, 2CH ₃), δ 2.5(s, CH ₂ of thiazolidinone), δ 6.66(s, NH2), 10.25(brs, NH), 8.1–7.01(m, 11 aromatic protons) | | | | | | | |
| 9 | δ1.1(t, CH ₃), δ 1.25(s, CH ₃), δ 3.35(q, CH ₂), δ 6.67(s, NH ₂), δ 10.35(brs, NH), 8.1–7.01(m, 6H aromatic protons) | | | | | | | |
| 10a | δ1.12(t, CH ₃), δ 1.24(s, CH ₃), δ 3.36(q, CH ₂), δ 6.65(s, NH ₂), δ 10.36(brs, NH), 8.1–7.01(m, 21 aromatic protons) | | | | | | | |
| 10b | δ1.11(t, CH ₃), δ 1.23(s, CH ₃), δ 3.35(q, CH ₂), δ 6.66(s, NH ₂), δ 10.35(brs, NH), 8.1–7.01(m, 21H aromatic protons) | | | | | | | |
| 10c | δ1.12(t, CH ₃), δ 1.24(s, CH ₃), δ 3.37(q, CH ₂), δ 6.67(s, NH ₂), 10.34(brs, NH), 8.1–7.01(m, 20H aromatic protons) | | | | | | | |

TABLE II

(Continued on next page)

| Compound No. | ¹ H NMR (DMSO) ppm | | | | | | | |
|--------------|---|--|--|--|--|--|--|--|
| 11a | δ 1.11(t, CH ₃), δ 1.23(s, CH ₃), δ 3.36(q, CH ₂), δ 6.68(s, NH ₂), 10.36(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 11b | δ1.12(t, CH ₃), δ 1.24(s, CH ₃), 3.35(q, CH ₂), δ 6.67(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 11c | δ1.11(t, CH ₃), δ 1.23(s, CH ₃), 3.35(q, CH ₂), δ 6.68(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 13H aromatic protons) | | | | | | | |
| 12a | δ1.12(t, CH ₃), δ 1.23(s, CH ₃), δ 2.5(s, CH ₂ of thiazolidinone), δ 3.35(q, CH ₂), δ 6.65(s, NH ₂), 10.33(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |
| 12b | δ1.1(t, CH₂), δ 1.23(s, CH₃), δ 2.5(s, CH₂ of thiazolidinone), δ 3.36(q, CH₂), δ 6.66(s, NH₂), 10.34(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |
| 12c | δ1.12(t, CH ₃), δ 1.22(s, CH ₃), δ 2.5(s, CH ₂ of thiazolidinone), δ 3.33(q, CH ₂), δ 6.65(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 10H aromatic protons) | | | | | | | |
| 13a | δ1.12(t, CH ₃), δ 3.35(q, CH ₂), δ 6.66(s, NH ₂), 10.36(brs, NH), 8.1–7.01(m, 14H aromatic protons) | | | | | | | |
| 13b | δ1.11(t, CH ₃), δ 3.34(q, CH ₂), δ 6.65(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 14H aromatic protons) | | | | | | | |
| 13c | δ1.12(t, CH ₃), δ 3.35(q, CH ₂), δ 6.67(s, NH ₂), 10.36(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 13d | δ1.12(t, CH ₃), δ 1.25(s, 2CH ₃), δ 3.36(q, CH ₂), δ 6.66(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 17H aromatic protons) | | | | | | | |
| 14a | δ1.12(t, CH ₃), δ 3.35(q, CH ₂), δ 6.65(s, NH ₂), 10.33(brs, NH), 8.1–7.01(m, 15H aromatic protons) | | | | | | | |
| 14b | δ 1.11(t, CH ₃), δ 3.36(q, CH ₂), δ 6.66(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 15H aromatic protons) | | | | | | | |
| 14c | δ1.1(t, CH ₃), δ 3.35(q, CH ₂), δ 6.64(s, NH ₂), 10.34(brs, NH), 8.1–7.01(m, 13H aromatic protons) | | | | | | | |
| 14d | δ1.12(t, CH ₃), δ 1.23(s, 2CH ₃), δ 3.36(q, CH ₂), δ 6.65(s, NH ₂), 10.35(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 15a | δ1.12(t, CH ₃), δ 2.5(s, CH ₂ of thiazolidinone), δ 3.34(q, CH ₂), δ 6.67(s, NH ₂), 10.36(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 15b | δ1.11(t, CH ₃), δ 2.5(s, CH ₂ of thiazolidinone), δ 3.35(q, CH ₂), δ 6.66(s, NH ₂), 10.33(brs, NH), 8.1–7.01(m, 12H aromatic protons) | | | | | | | |
| 15c | δ1.05(t, CH ₃), δ (s, CH ₂ of thiazolidinone), δ 3.36(q, CH ₂), δ 6.64(s, NH ₂), 10.34(brs, NH), 8.1–7.01(m, 14H aromatic protons) | | | | | | | |
| 15d | $1.12(t, CH_3), 1.24(s, 2CH_3), \delta$ 2.5(s, CH ₂ of thiazolidinone), δ 3.35(q, CH ₂), δ 6.66(s, NH ₂), 10.36(brs, NH), 8.1–7.01(m, 11H aromatic protons) | | | | | | | |

TABLE II (Continued)

9 was prepared by the reaction of compound **2** with ethyl iodide, which was heated together under fussion for 1 h. The structure of compound **9** was confirmed by elemental analysis, IR spectra, and ¹H NMR which reaveled the presense of a singlet signal for the methyl group at δ 1.25, a quarted signal at 3.45 for CH₂, and a triplet signal at 1.02 for the

| Compound | | (Calcd.) Found | | | | | |
|----------|---|----------------|--------|---------|--------------|---------|--|
| No. | IR $v_{\rm max}/{\rm cm}^{-1}$ | С | Н | Ν | \mathbf{S} | Cl | |
| 1 | 3400–3100(NH, NH ₂), 1745 (C=O | (61.34) | (4.83) | (13.41) | _ | _ | |
| | ester), 1645 (C=O) | 61.30 | 4.78 | 13.39 | _ | _ | |
| 2 | 3400–3150(NH, NH ₂), 1685(C=O), | (62.34) | (3.92) | (18.17) | _ | — | |
| | 1660(C=O) | 62.29 | 3.89 | 18.13 | _ | — | |
| 3a | 3400–3100(NH, NH ₂), 1665(C=O), | (68.92) | (4.47) | (18.27) | _ | _ | |
| | 1585(C=N) | 68.88) | 4.42 | 18.23 | _ | _ | |
| 3b | 3400–3100(NH, NH ₂ , OH), 1660(C=O), | (66.16) | (4.29) | (17.53) | — | — | |
| | 1590(C=N) | 66.13 | 4.25 | 17.49 | — | — | |
| 3c | 3400–3100(NH, NH ₂), 1650(C=O), | (63.77) | (3.89) | (16.90) | - | - | |
| | 1580(C=N) | 63.73 | 3.86 | 16.86 | — | — | |
| 4a | 3400–3100(NH, NH ₂), 1655(C=O) | (62.68) | (3.95) | (15.23) | _ | (7.71) | |
| | | 62.65 | 3.91 | 15.19 | — | 7.58 | |
| 4b | 3450–3100(NH, NH ₂ , OH), 1665(C=O) | (60.57) | (3.81) | (14.72) | _ | (7.45) | |
| | | 60.52 | 3.78 | 14.68 | — | 7.41 | |
| 4c | 3400–3100(NH, NH ₂), 1660(C=O) | (58.72) | (3.49) | (14.27) | — | (7.22) | |
| | | 58.69 | 3.45 | 14.23 | — | 7.19 | |
| 5a | $3400-3150(NH, NH_2), 1650(C=O)$ | (63.01) | (4.19) | (15.31) | (7.01) | — | |
| | | 62.99 | 4.18 | 15.29 | 6.98 | _ | |
| 5b | 3450–3100(NH, NH ₂ , OH), 1655(C=O) | (60.88) | (4.04) | (14.79) | (6.77) | — | |
| | | 60.83 | 4.01 | 14.75 | 6.74 | _ | |
| 5c | 3400–3150(NH, NH ₂), 1650(C=O) | (59.01) | (3.71) | (14.34) | (6.56) | — | |
| | | 58.98 | 3.68 | 13.99 | 6.52 | — | |
| 6a | 3450–3100(NH, NH ₂ , OH), 1670(C=O), | (67.38) | (3.70) | (15.11) | — | — | |
| | 1595(C=N) | 67.34 | 3.67 | 15.07 | — | — | |
| 6b | 3450–3100(NH, NH ₂ , OH), 1675(C=O), | (67.38) | (3.70) | (15.11) | — | — | |
| | 1585(C=N) | 67.35 | 3.65 | 15.05 | — | — | |
| 6c | 3450–3100(NH, NH ₂ , OH), 1660(C=O), | (63.92) | (3.66) | (16.94) | — | — | |
| | 1590(C=N) | 63.88 | 3.62 | 16.90 | _ | _ | |
| 6d | 3450–3100(NH, NH ₂), 1655(C=O), | (65.45) | (4.58) | (19.08) | _ | _ | |
| | 1580(C=N) | 65.41 | 4.53 | 19.04 | _ | — | |
| 7a | 3450–3100(NH, NH ₂ , OH), 1660(C=O) | (62.29) | (3.36) | (12.97) | _ | (6.57) | |
| | | 62.26 | 3.32 | 12.93 | _ | 6.54 | |
| 7b | 3400–3100(NH2NH ₂ , OH), | (62.29) | (3.36) | (12.97) | _ | (6.57) | |
| | 1690–1660(C=O) | 62.25 | 3.31 | 12.92 | _ | 6.52 | |
| 7c | 3400–3100(NH, NH ₂ , OH), | (58.84) | (3.29) | (14.30) | _ | (7.24) | |
| | 1717–1665(C=O) | 58.81 | 3.24 | 14.26 | _ | 7.19 | |
| 7d | 3400–3100(NH, NH ₂), | (60.40) | (4.10) | (16.27) | _ | 6.86 | |
| | 1718—1660(C=O) | 60.35 | 3.97 | 3.97 | _ | 6.82 | |
| 8a | 3400–3100(NH, NH ₂ , OH), | (62.56) | (3.56) | (13.03) | (5.94) | — | |
| | 1717-1655(C=O) | 62.51 | 3.53 | 12.99 | 5.90 | — | |
| 8b | 3450–3100(NH, NH ₂ , OH), | (62.56) | (3.56) | (13.03) | (5.94) | — | |
| | 1720–1660(C=O) | 62.53 | 3.52 | 12.97 | 5.89 | — | |
| | | | (Cor | ntinued | on next | t page) | |

TABLE III

| ~ . | | (Calcd.) Found | | | | |
|-----------------|--|----------------|--------|---------|--------|--------|
| Compound No. | IR $v_{\rm max}/{\rm cm}^{-1}$ | С | Н | Ν | S | Cl |
| 8c | 3400–3100(NH, NH ₂ , OH), | (59.13) | (3.52) | (14.37) | (6.58) | |
| | 1717–1665(C = O) | 59.09 | 3.49 | 14.33 | 6.54 | _ |
| 8d | 3400–3100(NH, NH ₂), | (60.69) | (4.31) | (16.33) | (6.23) | _ |
| | 1720–1655(C=O) | 60.64 | 4.28 | 16.29 | 6.19 | _ |
| 9 | 3400–3100(NH, NH ₂), 1680(C=O), | (46.57) | (3.69) | (12.07) | _ | _ |
| | 1655(C=O) | 46.53 | 3.66 | 12.03 | _ | _ |
| 10a | 3450–3100(NH, NH ₂), 1686(C=O), | (52.46) | (5.87) | (12.75) | _ | _ |
| | 1590(C=N) | 52.43 | 5.84 | 12.70 | _ | _ |
| 10b | 3450–3100(NH, NH ₂ , OH), 1680(C=O), | (50.98) | (5.70) | (12.39) | — | — |
| | 1580(C=N) | 50.93 | 5.67 | 12.35 | — | — |
| 10c | $3400-3100(NH, NH_2), 1685(C=O),$ | (49.66) | (5.38) | (12.07) | | _ |
| | 1590(C=N) | 49.62 | 5.35 | 12.03 | — | |
| 11a | $3450-3100(NH, NH_2),$ | (50.71) | (3.76) | (11.37) | | (5.76) |
| | 1725 - 1655(C=O) | 50.68 | 3.72 | 11.34 | | 5.73 |
| 11b | 3450–3100(NH, NH ₂ , OH), | (49.42) | (3.67) | (11.08) | _ | (5.61) |
| | 1720–1660(C=O) | 49.38 | 3.64 | 11.03 | — | 5.57 |
| 11c | $3400-3100(NH, NH_2),$ | (48.28) | (3.43) | (10.83) | _ | (5.48) |
| | 1725 - 1650(C=O) | 48.23 | 3.38 | 10.79 | | 5.45 |
| 12a | $3400-3100(NH, NH_2),$ | (50.91) | (3.94) | (11.42) | (5.23) | — |
| | 1730-1660(C=O) | 50.87 | 3.90 | 11.38 | 5.19 | — |
| 12b | 3450–3100(NH, NH ₂ , OH), | (49.61) | (3.84) | (11.13) | (5.09) | _ |
| | 1725 - 1655(C=O) | 49.58 | 3.81 | 11.90 | 5.04 | — |
| 12c | 3400–3100(NH, NH ₂), | (48.46) | (3.60) | (10.87) | (4.98) | — |
| | 1720-1650(C=O) | 48.42 | 3.55 | 10.83 | 4.92 | — |
| 13a | 3450–3100(NH, NH ₂ , OH), 1675(C=O), | (54.30) | (3.58) | (11.31) | _ | _ |
| | 1587(C=N) | 54.27 | 3.54 | 11.28 | _ | _ |
| 13b | 3450–3100(NH, NH ₂ , OH), 1665(C=O), | (54.30) | (3.58) | (11.31) | _ | _ |
| | 1589(C=N) | 54.26 | 3.52 | 11.26 | | _ |
| 13c | 3450–3100(NH, NH ₂ , OH), 1670(C=O), | (50.63) | (3.54) | (12.30) | _ | _ |
| _ | 1585(C=N) | 50.60 | 3.49 | 12.27 | — | — |
| 13d | $3400-3100(\text{NH}, \text{NH}_2), 1668(\text{C=O}),$ | (52.36) | (4.23) | (14.09) | _ | — |
| | 1590(C=N) | 52.32 | 4.18 | 14.06 | — | . — . |
| 14a | 3450–3100(NH, NH ₂ , OH), | (51.78) | (3.33) | (10.06) | _ | (5.10) |
| | 1717–1650(C=O) | 51.73 | 3.29 | 10.03 | _ | 4.98 |
| 14b | 3450–3100(NH, NH ₂ , OH), | (51.78) | (3.33) | (10.06) | _ | (5.10) |
| | 1720-1655(C=O) | 51.75 | 3.27 | 10.01 | — | 4.96 |
| 14c | 34540–3100(NH, NH ₂ , OH), | (48.39) | (3.28) | (10.84) | _ | (5.49) |
| | 1725 - 1660(C=O) | 48.35 | 3.23 | 10.80 | — | 5.46 |
| 14d | $3400-3100(NH, NH_2),$ | (49.98) | (3.89) | (12.49) | _ | (5.27) |
| | 1720-1655(C=O) | 49.92 | 3.84 | 12.45 | | 5.23 |
| 15a | 3450–3100(NH, NH ₂ , OH), | (51.96) | (3.49) | (10.10) | (4.62) | — |
| | 1720-1666(C=O) | 51.93 | 3.45 | 9.98 | 4.58) | — |
| 15b | 3450–3100(NH, NH ₂ , OH), | (51.96) | (3.49) | (10.10) | (4.62) | — |
| | 1717 - 1655(C=O) | 51.91 | 3.43 | 9.96 | 4.55 | — |
| 15c | 3450–3100(NH, NH ₂ , OH), | (42.01) | (2.98) | (9.42) | (4.31) | — |
| | 1717–1650(C=O) | 41.99 | 2.94 | 9.38 | 4.27 | — |
| 15d | $3400-3100(NH, NH_2), 1717-1660(CO)$ | (50.16) | (4.06) | (12.53) | (4.78) | — |
| | | 50.12 | 4.03 | 12.48 | 4.72 | _ |



SCHEME 2

 CH_3 group, and NH_2 group at δ 6.75. The mass spectrum showed the molecular ion peak $(M^+,\,C_{18}H_{17}O_3N_4I)$ at m/z 464.

The activity of the carbonyl group in position 4 and the activity of the methyl group at C_2 rendered compound 9, which contained iodine to react with different aromatic amine and different nitroso compounds to give new Schiff bases **10a–c** and **13a–d**. The structures of these newly synthesized Schiff bases compounds **10a–c** and **13a–d** were confirmed by their elemental analysis, IR, ¹H NMR, and mass spectra (cf. Tables I,



Downloaded by [University of Auckland Library] at 12:25 06 November 2014

SCHEME 3



SCHEME 4

II, III). The activity of the azamethine center in compounds **10a–c** and **13a–d** is more avaliable than the activity of the NH group toward the addition process of chloroacetyl chloride, and this mentioned phenomena is due to the presence of the π electron, which makes the foundation of the δ positive and δ negative charge on the carbon and nitrogen atom, respectively, more easy than the presence of this phenomena on the NH group in which the bonding between nitrogen and hydrogen whether strong according the nature of this bonding, which leads to decreased mobility devise of the hydrogen atom of this NH group. Thus compound 10a-c reacted with chloroacetylchloride or mercaptoacetic acid⁹ to give spiro β -Lactams and spiro thiazolidinone compounds **11a–c** and **12a– c**. The structure of compounds **11a–c** and **12a–c** were confirmed by their elemental analysis, IR, ¹H NMR, and mass spectra (cf. Tables I, II, III). Also we synthesized isolated β -lactams and thiazolidinone compounds by the reaction of new Schiff bases 13a-d with chloroacetyl chloride and/or mercaptoacetic acid to give 14a-d and 15a-d. The structures of these compounds were confirmed by their elemental analysis, IR, ¹H NMR, and mass spectra (cf. Tables I, II, III). We studied the effect of iodine on the antimicrobial activity of the synthesized compound. Thus the activity of eight synthesized compounds was examined in vito against three groups of microorganisms, including six strains of Gram positive bacteria, five strains of Gram negative bacteria, and three strains of fungi. The compounds were divided into two groups, four compounds with iodine in their structures and the other four compounds without iodine.

Several organo compounds are well authenticated to have antimicrobial activity versus many species of bacteria and fungi.^{12–17} These authors pointed out that some of these compounds are useful in the treatment and/or control of human, animal, and plant diseases. The present investigation was carried out to investigate the antimicrobial activity of the eight synthesized compounds and shed new light on the effect of iodine on the antimicrobial activity of the eight synthesized compounds.

MATERIAL AND METHODS

Organisms

The microorganisms (No S. 1, 4, 6, 8, 9, 13, and 14) were obtained from the culture collection of the United State Department of Agriculture, Northern Regional Research Laboratory (Peoria, Illinois. USA), while others were obtained from the Botany Department microorganism collection and identified according to a key given in Bergy's Manual of

| | Compound | | | | | | | | | |
|--|----------|----|----|----|-----|-----|-----|-----|------------|----------|
| Name | 2 | 3a | 4a | 5a | 9 | 10a | 11a | 12a | Ampicillin | Nystatin |
| 1-Bacillus subtilis NRS-744 | + | + | + | + | ++ | ++ | +++ | +++ | + | _ |
| 2-Micrococcus luteus SW 712 | ++ | ++ | + | ++ | ++ | +++ | +++ | +++ | + | - |
| 3-Bacillus megaterium SW 354 | ++ | ++ | + | + | +++ | ++ | +++ | +++ | + | _ |
| 4-Staphylococcus aureus B-767 | ++ | + | + | + | +++ | ++ | +++ | +++ | + | _ |
| 5-Streptomyces sp. SW 123 | ++ | - | + | + | ++ | ++ | ++ | +++ | + | - |
| 6-Bacillus cereus ATCC-9634 | ++ | + | + | ++ | ++ | ++ | +++ | +++ | + | _ |
| 7-Serratia Mar SW 98 | + | + | - | + | ++ | + | ++ | +++ | + | _ |
| 8-Pseudomonas aeruginosa ATCC- 6NA 10245 | - | + | _ | + | + | ++ | ++ | +++ | + | _ |
| 9-Escherichia coli B-3704 | — | + | - | + | + | ++ | ++ | +++ | + | - |
| 10-Salmonella sp. SW 476 | + | + | + | + | + | ++ | +++ | +++ | + | _ |
| 11-Pseudomonas sp. SW 653 | - | + | + | + | + | ++ | +++ | +++ | + | _ |
| 12-Sacharomyces cerevisiae SW 201 | + | + | - | + | ++ | +++ | ++ | ++ | _ | + |
| 13-Condida albicans IMRU3669 | + | + | + | - | ++ | +++ | +++ | ++ | _ | + |
| 14-Aspergillus flouus S-C 43313 | + | - | + | + | ++ | ++ | +++ | +++ | _ | + |

 TABLE IV Antimicrobial Potentialities of the Tested Compounds

 Expressed as Size (mm) of the Inhibition Zone

Determinative Bacteriology.¹⁸ Sacharomyces cerevisiae were identified as ascosporogenous.

Media

For the disc diffusion method, nutrient (NB) was used for cultivating the bacteria. It contained (g/L) the following: beef extract, 3; peptone, 5; and pH, 7.0. Czapek-Dox's agar medium was used for cultivating the fungi. It contained the following: (g/L) sucrose, 30.0; NaNO₃, 2.0; KCl, 0.5;

 K_2 HPO₄, 1.0; MgSO₄, 7H₂O, 0.5; FeSO₄, 0.0004; agar, 20; and pH, 7.0–7.2. malt extractagar was used for cultivating the yeast. It contained the following: (g/L). malt extract, 20; peptone, 1; dextrose, 20; agar, 20; and pH, 7.0–7.2.

Preparation of Bacterial Suspensions

Suspensions of the microorganisms were prepared by suspending each bacteria in 5 mL sterile nutrient broth media, using a standard loop and then incubating the inoculated nutrient broth at 37°C for 2 h. One mL of each suspension was added to the center of the sensitivity testing plate. A sterile, dry cotton-wool swap was used to spread the inoculum on the media. The inocula were allowed to dry for a few minutes.

Preparation of Discs

Five compounds were tested as 200 μ g/mL (W/V) solutions in sterile DMSO. Discs of 6-mm diameter of filter paper were placed in a petri dish (each one contained 10 discs) and then sterilized in a hot air oven at 180°C for 1 h. After cooling, 11 mL of the chemical solution was added onto each of the 10 discs to make a 20- μ g concentration perone discs. The discs were dried in the incubator at 35–37°C, for 1 h, or dried over phosphorous pentaoxide (P₂O₅) in a dissector under vacuum, and then distributed on the inocula by sterile forceps.

Each disc should be pressed down on the medium and should not be moved once in place. The plates were incubated at 37° C overnight. The diameters of the clear zones of inhibition were measured to the nearest 0.5 mm, compared to DMSO, under the same standardized conditions. The data obtained are expressed as the size (mm) of inhibition zone. The diameter of the inhibition zones were high: (+ + +) (20–16 mm), moderate (++): (15–12 mm), slight (+): (11–1 mm), and no zone of inhibition (negative) (-).

Disc Diffusion Method for Three Tested Fungi

For the disc diffusion,¹⁹ the discs of standard concentration (100 μ g/disc) of each of the five tested compounds were appropriately placed on the surface of an agar plate freshly seeded with standard inoculum of young culture (3 days old). The plates were kept at 5°C for 1 h to allow diffusion of the compounds through the agar media. The plates of the fungal test organisms were maintained at 30°C for 4 days. At the end of the incubation period, the inhibition zones were measured.

Standard Antimicrobial and Antifungal

The preliminary antimicrobial²⁰ was recorded in comparison to standard antibacterial ampicillin (10 mg/ml) and antifungal nystatin (10 mg/ml) in distiled water.

RESULTS AND DISCUSSION

Many antimicrobial agents have been introduced into therapy;^{21,22} however, the field still needs extensive efforts for the development of new antimicrobial agents of superior activity and less toxic side effects as well as to overcome the highly resistant strains of microorganisms. The data of the disc susceptibility tests for the compounds containing iodine, i.e., **9**, **10a**, **11a**,**12a**, clearly showed significant and potent antimicrobial activity (bactericidal and fungicidal) against the all-tested Gram positive bacteria and Gram negative bacteria and fungi. The compounds did not contain iodine, i.e., **2**, **3a**, **4a**, **5a** revealed weak susceptibility for the tested organisms compared with those of iodine, indicating that the presence of iodine in the compounds' structures enhanced the antimicrobial activity of the eight synthesized compounds. Compounds **11a** and **12a** exhibited the highest antimicrobial activity and were capable of inhibiting the growth of the all examined organisms.

EXPERIMENTAL SECTION

Melting points are uncorrected. IR spectra were measured as KBr pellets on a pye-unicam sp 1,000 spectrophotometer. ¹H NMR spectra were recorded in (²H₆) dimethylsulfoxide at 200 MHz on a Varian Gemini NMR spectrometer using Mesi as an internal reference. Mass spectra were obtained on a Shimadzu GCMS-QP 1000 EX mass spectrometer at 70 eV. Elemental analysis were carried out at the Microanalytical Center of Cairo University.

Synthesis of 2,4-Diamino 5,10-dioxo-1,5,10,11tetrahydrobenzo[g]quino-line-3-ethyl Carboxylate 1

A solution of the appropriate ethylcyano acetate and urea (0.01 mole), which was prepared in situ in ethanol (30 mL) containing (0.5 mL) piperidine, was treated with 1,4-naphthoquinone (0.02 mole, 1.58 g). The reaction mixture was heated under reflux for 10-12 h. The solvent was then evaporated under reduced pressure. Poured onto ice water acidified by HCl, the solid product that formed was collected by filtration and crystallized from DMF.

Synthesis of Compound 2

A stream of dry hydrogen chloride gas was passed through a mixture of compound 1 (3.13 gm, 10 mmol) and an appropriate acetonitrile (0.41 gm, 10 mmol) in dioxane (30 mL) for about 12 h. The reaction mixture was heated on a water bath for 2 h after passing hydrogen chloride gas. The reaction mixture was diluted with ice water and basified with 10% aq. ammonium hydroxide. The solid obtained as washed with water, dried, and crystallized from dimethyl formamide.

Synthesis of New Schiff Bases 3a-c and 6a-d

A solution of 2(0.01 mole, 3.08 gm) in DMF (30 mL) was treated with different aromatic amines (0.01 mole) and/or different nitroso compounds (0.01 mole) in the presence of a catalytic amount of piperidine (0.5 mL). The reaction mixture was heated under reflux for 8–9 h (monitored by TLC). The solvent was then evaporated under reduced pressure, and the residue was treated with ice water acidified by HCl. The solid product was collected and crystallized from the proper solvent (cf. Table I).

Synthesis of Spiro β -Lactam 4a–c and Thiazolidinone 5a–c

A solution of **3a–c** (0.01 mole) in DMF (20 mL) was treated with chloroacetyl chloride or mercaptoacetic acid (0.01 mole) in the presence of (0.5 ml) triethylamine catalyst. The reaction mixture was heated under reflux for 15–17 h (monitored by TLC). The solvent was then evaporated under reduced pressure, and the residue was treated with ice water. The solid product was collected and crystallized from the proper solvent (cf. Table I).

Synthesis of Isolated β -Lactam 7a–d and Thiazolidinone 8a–d

A solution of **6a–d** (0.01 mole) in DMF (30 mL) was treated with chloroacetyl chloride or mercaptoacetic acid (0.01 mole), which was added drop by drop and stirred for 1 h in the presence of triethyl amine catalyst. The reaction mixture was heated under reflux for 18–20 h (monitored by TLC). The solvent was then evaporated under reduced pressure, and the residue was treated with ice water. The solid product was collected by filtration and crystallized from the proper solvent (cf. Table I).

Synthesis of Compound 9

A solution of ethyl Iodide (0.01 mole, 1.56 gm) was treated to compound 2 (0.01 mole, 3.08 gm) drop by drop and heated together by fussion for

1 h; the solid product was collected by ethanol and crystallized from the proper solvent (cf. Table I).

Synthesis of New Schiff Bases 10a-c and 13a-d

A solution of 9 (0.01 mole, 4.64 gm) in DMF (10 mL) and ethanol (20 mL) was treated with different aromatic amine (0.01 mole) and/or different nitroso compounds (0.01 mole) in the presence of a catalytic amount of piperidine (0.5 ml). The reaction mixture was heated under reflux for 7–8 h (monitored by TLC). The solvent was then evaporated under reduced pressure, and the residue was treated with ice water acidified by HCl. The solid product was collected by filtration and crystallized from the proper solvent (cf. Table I).

Synthesis of Spiro β -Lactam 11a–c and Thiazolidinone 12a–c

A solution of **10a–c** (0.01 mole) in DMF (30 mL) was treated with chloroacetyl chloride or mercaptoacetic acid (0.01 mole) in the presence of (0.5 ml) triethylamine catalyst. The reaction mixture was heated under reflux for 16–18 h (monitored by TLC). The solvent was then evaporated under reduced pressure, and the residue treated with ice water. The solid product was collected by filtration and crystallized from the proper solvent (cf. Table I).

Synthesis of Isolated β -Lactam 14a–d and Thiazolidinone 15a–d

A solution of **13a-d** (0.01 mole) in DMF (30 mL) was treated with chloroacetyl chloride or mercaptoacetic acid (0.01 mole), which was added drop by drop and stirred for 1 h in the presence of triethylamine catalyst. The reaction mixture was heated under reflux for 19–20 h (monitroed by TLC). The solvent was then evaporated under reduced pressure, and the residue was treated with ice water. The solid product was collected by filtration and crystallized from the proper solvent (cf. Table I).

REFERENCES

- W. Durckheimer, J. Blumbach, R. Lattrell, and K. H. Scheunemann, Angew. Chem., 97, 183 (1985).
- [2] C. Cimarusti and R. Sykes, Chem. Br., 19, 302 (1983).
- [3] S. R. Woulfe and M. Miller, Tetrahedron Lett., 25, 3293 (1984).
- [4] S. R. Woulfe and M. Miller, J. Med. Chem., 28, 1447 (1985).

- [5] H. Breure, H. Straub, U. Treuner, J. M. Drossard, H. Hoehn, and K. Lindler, J. Antibiot., 38, 813 (1985).
- [6] S. R. Woulfe and M. J. Miller, J. Org. Chem., 51, 3133 (1986).
- [7] M. J. Miller, A. Biswas, and M. A. Krook, Tetrahedron, 39, 2571 (1983).
- [8] W. O. Foye, *Principles of Medicinal Chemistry*. 2nd ed., (Lea and Febiger, Philadelphia, 1981).
- [9] A. K. Khalafallah, R. M. Abd El Aal, and N. A. A. El-Kanzi, J. Chin. Chem. Soc., 49, 2002.
- [10] A. K. Khalafallah, A. I. M. Koraiem, M. A. Elmaghraby, and H. A. Shindy, *Egypt J. Chem.*, **36**, 159 1 (1993).
- [11] A. K. Khalafallah, F. M. Abd Latif, M. A. Salim, and M. A. El Maghraby, Asian, Journal of Mem. 5, (1993).
- [12] R. M. Abd El-Aal and M. Younis, Dyes and Pigments, 60, 205 (2004).
- [13] Gamal A. Elmegeed, W. W. Wagnat, M. Younis, and L. Nadia, Archive der Pharmazie, 337, 140 (2004).
- [14] N. R. Mohamed, G. A. Elmegeed, and M. Yonis, *Phosphorus, Sulfur, and Silicon*, 178, 2003 (2003).
- [15] R. K. Pettit, G. D. Cage, G. R. Pettit, and J. A. Liebman, Int. J. Antimicrob Agent, 15, 299 (2000).
- [16] A. D. Russel, I. Ahonkhai, and D. T. Rogers, J. Appl. Bact., 46, 207 (1979).
- [17] M. Younis, G. A. Elmegeed, and M. I. Nassar, Egypt J. Microbial., 38, (2003).
- [18] Bergy's Manual of Determinative Bacteriology, 8th ed., R. E. Buchanan and N. E. Gibbons, eds., (Williams and Wilkins, Baltimore, 1974).
- [19] T. D. Wilkins, L. V. Holdman, I. J. Abramson, and W. E. C. Moore, Antimicrobial Agents and Chemotherapy, 1, 451, 9 (1972).
- [20] D. S. Reeves and L. O. White, Principles of Methods of Assaying Antibiotics in Pharmaceutical Microbiology, 3rd ed., pp. 140–162, (Blackwell Scientific Publication, 1983).
- [21] L. Z. Yan, A. C. Gibbs, M. E. Stiles, D. S. Wishart, and J. C. Vederas, J. Med. Chem., 43, 4579 (2000).
- [22] K. C. Fang, Y. L. Chen, J. Y. Sheu, T. C. Wang, and C. C. Tzeng, J. Med. Chem., 43, 3809.