

† *Selected physical data for cycloadducts:* M.p.s, b.p.s (Kugelrohr distillation), and diagnostic  $^1\text{H}$  n.m.r. ( $\text{CDCl}_3$ ) signals were recorded as follows: **(4a)** and **(5a)**, b.p.  $95^\circ\text{C}$  (0.02 mbar); **(4a)**,  $\delta$  4.42 (d,  $J$  4 Hz, 3-H); **(5a)**,  $\delta$  3.30 (br. s, 3-H); **(6a)**, b.p.  $110\text{--}120^\circ\text{C}$  (0.03 mbar),  $\delta$  3.61 (t,  $J$  6 Hz, 2-H); **(4b)** + **(5b)**, m.p.  $81\text{--}82^\circ\text{C}$ ; **(4b)**,  $\delta$  5.10 (d,  $J$  4 Hz, 3-H); **(5b)**,  $\delta$  4.04 (br. s, 3-H); **(6b)**, m.p.  $110\text{--}112^\circ\text{C}$ ,  $\delta$  3.65 (t,  $J$  5 Hz, 2-H); **(4c)** and **(5c)**, m.p.  $88\text{--}89^\circ\text{C}$ ; **(4c)**,  $\delta$  5.12 (d,  $J$  4 Hz, 3-H); **(5c)**,  $\delta$  4.07 (br. s, 3-H); **(6c)**,  $\delta$  4.50 (t,  $J$  6 Hz, 2-H); **(4d)** + **(5d)**, b.p.  $90^\circ\text{C}$  (0.02 mbar); **(4d)**,  $\delta$  4.27 (d,  $J$  4 Hz, 3-H); **(5d)**,  $\delta$  3.28 (br. s, 3-H); **(6d)**, b.p.  $120^\circ\text{C}$  (0.02 mbar),  $\delta$  3.76 (t,  $J$  5 Hz, 2-H); **(4e)** + **(5e)**, m.p.  $76\text{--}78^\circ\text{C}$ ; **(4e)**,  $\delta$  4.96 (d,  $J$  4.5 Hz, 3-H); **(5e)**,  $\delta$  4.04 (br. s, 3-H); **(6e)**, m.p.  $64\text{--}65^\circ\text{C}$ ,  $\delta$  4.06 (t,  $J$  7 Hz, 2-H).

**Table 1.** Cycloadducts of thioaldehydes, ZCHS (**2**), with conjugated dienes.

Bunte salt ( <b>1</b> ; X = SO <sub>3</sub> Na) Z	Cyclopentadiene		2,3-Dimethylbuta-1,3-diene Yield of ( <b>6</b> ), %
	Yield of ( <b>4</b> ) + ( <b>5</b> ), %	( <b>4</b> ) : ( <b>5</b> ) ratio <sup>d</sup>	
<b>a</b> EtO <sub>2</sub> C	67 <sup>a,b</sup>	7 : 3 <sup>c</sup>	68
<b>b</b> PhNHCO	66 <sup>c</sup>	6 : 1	63
<b>c</b> PhCO	65 <sup>b</sup>	7 : 3	65
<b>d</b> NC	77 <sup>c</sup>	2 : 1 <sup>c</sup>	51
<b>e</b> 4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	88 <sup>c</sup>	7 : 1	55

<sup>a</sup> EtOH was used as solvent. <sup>b</sup> Separable by t.l.c. <sup>c</sup> Characterised as a mixture. <sup>d</sup> Determined by <sup>1</sup>H n.m.r. spectroscopy. <sup>e</sup> Unchanged by Kugelrohr distillation.

yield of adducts (**4e**) and (**5e**) whereas the unsubstituted benzyl derivative (**1**; Z = Ph, X = SO<sub>3</sub>Na) gave no detectable amounts of the corresponding adducts. This illustrates the importance of methylene hydrogen acidity; conjugation of the thioformyl group with a benzene ring is not, in itself, adequate to facilitate elimination using triethylamine.

When cyclopentadiene was replaced as trapping agent by the less reactive 2,3-dimethylbuta-1,3-diene, the yields of cycloadducts (**6**), formed under the foregoing conditions, were consistently lower. However, acceptable yields were obtained using a less polar solvent system. Thus, triethylamine in benzene (5 ml) was added to the Bunte salt (typically 4 mmol) in ethanol (5 ml) and benzene (10 ml) containing calcium chloride dihydrate and 2,3-dimethylbuta-1,3-diene, with heating under reflux. Heating was continued for 4 h and the reaction mixtures were worked-up as before. The tabulated yields of the adducts (**6**)<sup>†</sup> are for purified materials. Apparently, competitive attack by nucleophiles (*e.g.* SO<sub>3</sub><sup>2-</sup>) on the thioaldehydes is less important in less polar solvents, but the elimination reactions are generally slower.

We earlier reported<sup>2</sup> that the kinetically determined mixture of adducts (**4a**) and (**5a**) (1.4 mmol), when heated with 2,3-dimethylbuta-1,3-diene (1.54 mmol) in toluene under nitrogen at 120 °C (sealed tube) for 24 h, gave the adduct (**6a**) in 82% yield (after purification). Similarly, the adducts (**4b**)

and (**5b**) gave (**6b**) (84%) and (**4c**) and (**5c**) gave (**6c**) (84%). This thermal transfer of thioaldehydes was slower with the cyano-adducts (**4d**) and (**5d**) but was complete in 48 h to give (**6d**) (79%). In contrast, transfer of 4-nitrothiobenzaldehyde from (**4e**) and (**5e**) to dimethylbutadiene was accomplished in benzene under reflux for 48 h to give (**6e**) (78%). It appears therefore that cyclopentadiene adducts are generally synthetically valuable, auxiliary sources of reactive thioaldehydes.

We thank the S.E.R.C. for financial support.

Received, 25th April 1984; Com. 567

## References

- 1 C. M. Bladon, I. E. G. Ferguson, G. W. Kirby, A. W. Lohead, and D. C. McDougall, *J. Chem. Soc., Chem. Commun.*, 1983, 423.
- 2 G. W. Kirby and A. W. Lohead, *J. Chem. Soc., Chem. Commun.*, 1983, 1325.
- 3 H. Bunte, *Chem. Ber.*, 1874, **7**, 646.
- 4 B. Milligan and J. M. Swan, *J. Chem. Soc.*, 1959, 2969.
- 5 B. Saville and M. Steer, *J. Chem. Soc., Chem. Commun.*, 1972, 616.
- 6 K. Thimm and J. Voss, *Tetrahedron Lett.*, 1975, 537; K. Thimm, W. Schmöser, and J. Voss, *J. Chem. Res.*, 1977, (*S*), 244; (*M*), 2837.