# Characterization and biological effects of di-hydroxylated compounds deriving from the lipoxygenation of ALA

Miao Liu,\* Ping Chen,<sup>†</sup> Evelyne Véricel,\* Moreno Lelli,<sup>§</sup> Laetitia Béguin,<sup>§</sup> Michel Lagarde,\* and Michel Guichardant<sup>1,\*</sup>

Université de Lyon,\* UMR 1060 Inserm (CarMeN), IMBL/INSA-Lyon, 69621 Villeurbanne, France; Tea Science Department,<sup>†</sup> College of Agriculture and Biotechnology, Zhejiang University, 310029 Hangzhou, China; and Centre de RMN à Très Hauts Champs,<sup>§</sup> Université de Lyon (CNRS/ENS Lyon/UCB Lyon 1), 69100 Villeurbanne, France

Abstract We have recently described a di-hydroxylated compound called protectin DX (PDX) which derives from docosahexaenoic acid (DHA) by double lipoxygenation. PDX exhibits anti-aggregatory and anti-inflammatory properties, that are also exhibited by similar molecules, called poxytrins, which possess the same E,Z,E conjugated triene geometry, and are synthesized from other polyunsaturated fatty acids with 22 or 20 carbons. Here we present new biological activities of di-hydroxylated metabolites deriving from  $\alpha$ -linolenic acid (18:3n-3) treated by soybean 15-lipoxygenase (sLOX). We show that 18:3n-3 is converted by sLOX into mainly 13(S)-OH-18:3 after reduction of the hydroperoxide product. But surprisingly, and in contrast to DHA which is metabolized into only one di-hydroxylated compound, 18:3n-3 leads to four di-hydroxylated fatty acid isomers. We report here the complete characterization of these compounds using high field NMR and GC-MS techniques, and some of their biological activities. These compounds are: 9(R),16(S)-dihydroxy-10E,12E,14E-octadecatrienoic acid, 9(S),16(S)-dihydroxy-10E,12E,14E-octadecatrienoic acid, 9(S),16(S)-dihydroxy-10E,12Z,14E-octadecatrienoic acid, and 9(R),16(S)-dihydroxy-10E,12Z,14E-octadecatrienoic acid. They can also be synthesized by the human recombinant 15-lipoxygenase (type 2). Their inhibitory effect on blood platelet and anti-inflammatory properties were compared with those already reported for PDX.-Liu, M., P. Chen, E. Véricel, M. Lelli, L. Béguin, M. Lagarde, and M. Guichardant. Characterization and biological effects of di-hydroxylated compounds deriving from the lipoxygenation of ALA. J. Lipid Res. 2013. 54: 2083-2094.

**Supplementary key words** octadecatrienoic acid • 15-lipoxygenase • platelet aggregation • anti-inflammatory activity •  $\alpha$ -linolenic acid

N-3 polyunsaturated fatty acids (PUFAs) have been well studied during the last decades because they regulate in

Manuscript received 18 December 2012 and in revised form 29 May 2013. Published, JLR Papers in Press, June 5, 2013 DOI 10.1194/jtr.M035139

Copyright © 2013 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at http://www.jlr.org

vivo many physiological functions. It is assumed that longchain n-3 PUFAs have beneficial effects on cardiovascular and coronary artery diseases (1, 2), rheumatoid arthritis (3–5), blood pressure control and heart hypertension (6, 7), and they might prevent multiple sclerosis and Alzheimer's disease (8-11). More recently, new lipid mediators deriving from eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have been described. Among them, resolvins are oxygenated molecules that derive from EPA and DHA (12, 13) and are separated into two series: Eseries (from EPA) and D-series (from DHA) (14). In addition to the D-series resolvins, protectin D1 (PD1) is produced from DHA via a mechanism involving an epoxide intermediate (15), as well as maresin which can be produced by macrophages (16). PD1 has been described as a potent anti-inflammatory agent without an anti-aggregatory effect (15, 17). In contrast, protectin DX (PDX), an isomer of PD1, which derives from DHA via a double lipoxygenation, reveals inhibitory effects on blood platelet aggregation. PDX inhibits platelet cyclooxygenase (COX)-1 and antagonizes the thromboxane A<sub>2</sub>-induced aggregation (18). We found that di-hydroxylated compounds called poxytrins, synthesized from C20 and C22 PUFAs via the soybean 15-lipoxygenase (sLOX) and having the same E,Z,E conjugated triene motif, including PDX, are all anti-aggregatory agents (19). On the other hand, very few data are available concerning the oxygenated metabolism of  $\alpha$ -linolenic acid (ALA) (18:3n-3). Investigations relating to ALA are particularly relevant because

e-mail: michel.guichardant@insa-lyon.fr

This study was supported by Inserm and the French Ministry of Education and Research. M.L. is a Ph.D. student granted from the China Scholarship Council.

Abbreviations: ALA,  $\alpha$ -linolenic acid (9Z,12Z,15Z-octadecatrienic acid); COX, cyclooxygenase; DHA, docosahexaenoic acid; EPA, eicosapentaenoic acid; 9(S)-HOTE, 9(S)-hydroxy-10*E*,12Z,15Z-octadecatrienoic acid; 13(S)-HOTE, 13(S)-hydroxy-9Z,11*E*,15Z-octadecatrienoic acid; 9(S)-HpOTE, 9(S)-hydroperoxy-octadeca-10*E*,12Z,15Z-trienoic acid; 13(S)-HpOTE, 13(S)-hydroperoxy-9Z,11*E*,15Z-octadecatrienoic acid; 13(S)-HpOTE, 13(S)-hydroperoxy-9Z,11*E*,15Z-octadecatrienoic acid; 13(S)-HpOTE, 13(S)-hydroperoxy-9Z,11*E*,15Z-octadecatrienoic acid; 13(S)-HpOTE, 13(S)-hydroperoxy-9Z,11*E*,15Z-octadecatrienoic acid; LOX, lipoxygenase; LTB<sub>4</sub>, leukotriene B<sub>4</sub>; PD1, protectin D1; PDX, protectin DX; PGD<sub>2</sub>, prostaglandin D<sub>2</sub>; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; RP-HPLC, reverse-phase high-performance liquid chromatography; sLOX, soybean 15-lipoxygenase; 9,16-di-HOTE, 9,16-dihydroxyoctadecatrienoic acid; UV, ultraviolet.

To whom correspondence should be addressed.



this fatty acid is taken in large amounts (1-2 g/day in)human adults) as a nutrient and hardly accumulates in plasma, suggesting an active metabolism. Moreover, its conversion into EPA and DHA is relatively limited. A conversion of ALA into its long-chain product DHA was found to be around 1% in men (20) and 2% in young women (21). Formation of 9,16-dihydroxyoctadecatrienoic acid (9,16-di-HOTE) isomers from ALA by sLOX-2 have already been reported first in 1984 by Feiters et al. (22) without establishing the double bond configuration or the stereochemistry of the asymmetric carbons. Further studies presented by Sok and Kim (23, 24) reported four isomers, but again their characterization was not fully established, especially that of the all trans isomers. In 1991 Grechkin et al. (25) showed that potato tuber lipoxygenase (LOX) generated mainly 9(S)-hydroperoxy-octadeca-10E,12Z,15Z-trienoic acid [9(S)-HpOTE] which is different from the sLOX which produces mainly 13(S)-hydroperoxy-9Z,11E,15Zoctadecatrienoic acid [13(S)-HpOTE]. Also, two compounds after reduction by NaBH<sub>4</sub> were characterized as 9,16-di-HOTE (10E,12Z,14E) except for the stereochemistry of the carbon 16 which remains undetermined and trans isomers were not detected. Moreover, the biological properties of the 9,16-di-HOTE isomers have not been investigated yet. Because ALA is the most abundant n-3 PUFA in human diet, it is relevant to investigate more thoroughly its oxygenated metabolism, especially by 15-LOX that is expressed in blood leukocytes and most endothelial cells, and compare the biological activity of the metabolites to that of PDX, the main 15-LOX product of DHA.

#### MATERIALS AND METHODS

#### Materials

2084

ALA, sLOX (E.C. 1.13.11.12, Type 1-B, 131,000 units/mg), platinum oxide (PtO<sub>2</sub>), *N*,*O*-bis(trimethylsilyl)-trifluoroacetamide, COX-1, and COX-2 (E.C. 1.14.99.1) were from Sigma-Aldrich. Human recombinant 15-LOX-2 (0.23 units/mg), prostaglandin D<sub>2</sub> (PGD<sub>2</sub>)-d<sub>4</sub>, and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)-d<sub>4</sub> were from Cayman. 9(S)-hydroxy-10*E*,12*Z*,15*Z*-octadecatrienoic acid [9(S)-HOTE], 9(S)-HPOTE, and 13(S)-hydroxy-9*Z*,11*E*,15*Z*-octadecatrienoic acid [13(S)-HOTE] were from Interchim. Organic solvents were from Carlo-Erba. All chemicals used were reagent grade or the highest quality available.

#### Biosynthesis of 9,16-dihydroxy-ALA derivatives

ALA was incubated with sLOX (type 1-B) in sodium-borate buffer. Synthesized hydroperoxides were reduced by sodium borohydride (NaBH<sub>4</sub>) as previously described (18). After acidification, mono- and di-hydroxylated fatty acids were extracted on a C18 solid-phase cartridge as previously described (18). Commercial 9(S)-HOTE, 9(S)-HpOTE, and the homemade racemic 9( $\pm$ )-HOTE chemically produced by oxygen treatment of ALA and isolated by reverse-phase high-performance liquid chromatography (RP-HPLC) were further treated by sLOX in order to generate 9,16-di-HOTEs with defined stereochemistry.

#### Purification of mono- and di-hydroxylated fatty acids

Mono- and di-hydroxylated fatty acids were analyzed by RP-HPLC on a Waters XBridge C18 column  $(4.6 \times 250 \text{ mm}, 3.5 \text{ }\mu\text{m})$ 

#### GC-MS analysis of mono- and di-hydroxylated fatty acids

Mono- and di-hydroxylated fatty acids isolated by RP-HPLC were hydrogenated using  $PtO_2$  as a catalyst, and then derivatized into methyl esters and TMS ethers. Samples were then analyzed by GC-MS using the electron impact ionization (EI) mode in order to localize the hydroxyl groups on the fatty chain (18).

#### NMR of 9,16-di-HOTEs

The NMR experiments were performed on a Bruker-Biospin NMR spectrometer operating at 23.4 Tesla (1,000 MHz of Proton Larmor frequency) and equipped with a Cryoprobe. All the spectra were acquired in a CDCl3 solution, working at 298 K and referencing the chemical shifts to tetramethylsilane as an internal standard. Standard sequences were used for 1D, DQF COSY (26–28), and J-resolved (29–31) spectra. The <sup>1</sup>H pulse lengths of 9.0 µs, 1 s, and 5 s were used as recycle delay for J-resolved and COSY spectra. A weak selective decoupling of 270 Hz of power during the acquisition time (900 ms) was used in 1D selective homo-decoupling experiments to simplify some coupling patterns. 2D DQF COSY was acquired with 8 scans and  $2,048 \times 1,024$ real points (F2  $\times$  F1) with acquisition times of 114 ms and 57 ms for the direct and indirect dimensions. J-resolved 2D was acquired with 16 scans and 8,192 × 256 real points, acquiring 458 ms in the direct dimension and 1,280 ms in the indirect J-coupling dimension. COSY spectra were processed with a  $4,096 \times 4,096$  point matrix using square cosine and square sine window function for the direct and indirect dimensions. J-resolved 2D spectra were processed with an  $8,192 \times 2,048$  point matrix with square cosine window function for both the dimensions.

## Incubation of ALA, 9(S)-HOTE, and $9(\pm)$ -HOTE with human recombinant 15-LOX-2

ALA, 9(S)-HOTE, and 9(±)-HOTE were incubated separately with human recombinant 15-LOX-2 in Tris-HCl buffer as previously described (32, 33). Resulting hydroperoxides were reduced by NaBH<sub>4</sub>. Lipids were extracted after acidification to pH 3 with acetic acid and di-hydroxylated metabolites were purified as described above. Downloaded from www.jlr.org at Carleton Univ - OCUL, on June 19, 2014

#### Preparation of cell suspensions

*Platelet suspensions.* Venous blood from healthy volunteers who had not taken any medication for at least one week was collected at the local blood bank (Etablissement Français du Sang). Blood was drawn onto anticoagulant and centrifuged at 200 g for 15 min at 20°C to obtain the platelet-rich plasma upper phase, and platelets were isolated as described previously (34). Briefly, platelet-rich plasma was acidified to pH 6.4 with 0.15 M citric acid and immediately centrifuged at 900 g for 10 min, and platelet pellets were suspended into Tyrode-HEPES buffer (140 mM NaCl, 2.7 mM KCl, 1.0 mM MgCl<sub>2</sub>, 12 mM NaHcO<sub>3</sub>, 0.41 mM NaH<sub>2</sub>PO<sub>4</sub>, 5 mM HEPES, and 5.5 mM glucose, pH 7.35).

*Leukocyte suspensions.* Leukocytes were isolated from blood obtained from Etablissement Français du Sang, as previously described (35). Dextran (2% final concentration) was added to the lower phase obtained after the first blood centrifugation at 200 g. Erythrocytes were then sedimented under gravity for 45 min at 15°C. The erythrocyte-depleted supernatant containing leukocytes

was centrifuged at 400 g for 10 min. The resulting leukocyte-rich pellets were resuspended into saline buffer, and layered over a Ficoll-Paque Plus (density 1.077). After centrifugation, the fraction above the erythrocyte layer was removed and centrifuged, and resuspended in water for 20 s to lyse the contaminant erythrocytes. Osmotic pressure was immediately restored with NaCl, followed by centrifugation, which provided a white pellet consisting of whole leukocytes that was resuspended into Tyrode-HEPES buffer. The morphological examination and trypan blue exclusion tests were performed to determine the cell count and purity, and evaluate the viability of neutrophils.

### Measurement of platelet aggregation and incubation with human leukocytes

Blood platelet aggregation was determined by the turbidimetric method of Born (36) using a dual aggregometer (ChronoLog, Havertown, PA). To measure changes in the light transmission rate, 400  $\mu$ l of platelet suspension was heated at 37°C for 1 min with stirring, incubated in the presence or absence of di-hydroxylated compounds (added in 1  $\mu$ l of ethanol) for one more minute, and then with collagen as a platelet aggregatory agent for 4 min at 37°C. Concentrations of collagen were adjusted to obtain nearly 75% aggregations for control platelets.

Isolated human leukocytes, suspended in a Tyrode-HEPES buffer containing 2 mM Ca<sup>2+</sup> were preincubated in the presence or absence of 1  $\mu$ M of different 9,16-di-HOTE isomers for 3 min at 37°C, and triggered with 1  $\mu$ M ionophore A23187 plus 10  $\mu$ M arachidonic acid for 10 min. The reaction was stopped by acidifying the media

to pH 3 with acetic acid. The oxygenated metabolites were then extracted as previously described (18).

### GC-MS measurement of purified COX-1 and COX-2 products

Commercial COX-1 (40 units) and COX-2 (40 units) were preincubated for 15 min in a 0.1 M Tris-HCl buffer pH 8.1 with or without 1 µM of different di-hydroxylated metabolites and incubated for 10 min with 10 µM arachidonic acid used as a substrate. Albumin (45 g/l) was added at the end of incubation to facilitate the formation of PGD<sub>9</sub> and PGE<sub>9</sub> from PGH<sub>9</sub> (37). After acidification to pH 3 with acetic acid, PGE<sub>9</sub> and PGD<sub>9</sub> were extracted by 10 vol of diethyl ether stabilized by 10% ethanol. Deuterated PGD<sub>2</sub> and PGE<sub>2</sub> used as internal standards were added for the quantification. Prostaglandins were derivatized into methoxime, pentafluorobenzyl esters, and TMS and further analyzed by negative-ion chemical ionization GC-MS. PGD<sub>2</sub> and PGE<sub>2</sub> were measured using the SIM mode. Selected ions corresponding to  $[M-181]^{-}$  (loss of the pentafluorobenzyl group): m/z 524 for both derivatized PGD<sub>2</sub> and PGE<sub>2</sub> and m/z 528 for their corresponding deuterated internal standards (38, 39) were measured.

#### Statistical analysis

All results are expressed as means  $\pm$  standard error of the mean (SEM) or means  $\pm$  standard deviation (SD). P < 0.05 was set as the level of significance as determined by two-tailed Student's *t*-tests. One-way ANOVA analysis was used to assess differences between groups.



Fig. 1. Typical RP-HPLC chromatogram of ALA metabolites. ALA was incubated with sLOX, and further treated by NaBH<sub>4</sub> to reduce hydroperoxides into hydroxylated derivatives. The lipid extract was then analyzed by RP-HPLC. Mono-hydroxylated fatty acids (A) and di-hydroxylated fatty acids (B) were detected at  $\lambda$ = 235 nm and 270 nm, respectively. On the right side of the chromatograms, the UV spectra acquired on the top of chromatographic peaks are reported. mAU, milli absorbance units.

#### RESULTS

### Characterization of metabolites obtained from ALA incubated with sLOX

From ALA incubated with sLOX as described in Materials and Methods, hydroxylated derivatives analyzed by RP-HPLC provided one main compound detected at 50 min with a  $\lambda_{max}$  of 235 nm (**Fig. 1A**). Its retention time and its ultraviolet (UV) spectrum were similar to that of commercial 13(S)-HOTE (result not shown).

Regarding di-hydroxylated products (Fig. 1B), and in contrast to DHA which is only converted into one major dihydroxylated metabolite called PDX, four di-hydroxylated isomers were detected from ALA. They all exhibited characteristic UV spectra with a maximum absorption  $(\lambda_{max})$  at 270 nm with two shoulder peaks at 260 nm and 280 nm, indicating the presence of a conjugated triene. Moreover, the UV spectra of the first two isomers eluted at 32.1 and 32.6 min are superimposable to that of commercial 12-epiall trans-leukotriene  $B_4$  (LTB<sub>4</sub>) with an *E,E,E* conjugated triene geometry (result not shown), whereas the UV spectra of the two other isomers eluted at 33.2 and 33.5 min are superimposable to that of PDX, suggesting an *E*,*Z*,*E* geometry of these conjugated trienes (Fig. 1B). The spectra of these four isomers differ from those with a Z,E,E configuration such as LTB<sub>4</sub> and 12-epi-LTB<sub>4</sub> (result not shown).

### Localization by GC-MS of hydroxyl groups on the fatty chain

Mono- and di-hydroxylated compounds from ALA mentioned above were hydrogenated and derivatized as methyl esters and TMS ethers as described in Materials and Methods and analyzed by GC-MS using the EI mode.

The EI mass spectrum of the hydrogenated monohydroxylated ALA derivative (**Fig. 2A**) shows characteristic ions at: m/z 371 (M-15) loss of CH<sub>3</sub>, m/z 315 (M-71) which corresponds to the loss of CH<sub>2</sub>-(CH<sub>2</sub>)<sub>3</sub>-CH<sub>3</sub>, and m/z 173 [TMS-O-CH-(CH<sub>2</sub>)<sub>4</sub>-CH<sub>3</sub>]. Such a fragmentation pattern clearly indicates the presence of the hydroxyl group on carbon 13 of the fatty chain. So ALA was mainly converted into 13(S)-HpOTE, further reduced by NaBH<sub>4</sub> into hydroxylated product for analytical purposes.

All four hydrogenated di-hydroxylated compounds from ALA exhibited the same mass spectra (Fig. 2B). Their fragmentation pattern shows the following characteristic ions: at m/z 459 (M-15, loss of CH<sub>3</sub>); at m/z 445 (M-29, loss of CH<sub>2</sub>-CH<sub>3</sub>); at m/z 317 [M-157, loss of CH<sub>2</sub>-(CH<sub>2</sub>)<sub>6</sub>-COOCH<sub>3</sub>]; at m/z 259 [TMS-O-CH-(CH<sub>2</sub>)<sub>7</sub>-COOCH<sub>3</sub>]; at m/z 131 (TMS-O-CH-CH<sub>2</sub>-CH<sub>3</sub>); and at m/z 73 (TMS) which indicates an oxygenation at carbons 9 and 16.



Fig. 2. Electron impact mass spectrum of the Me-TMS derivative of the hydrogenated HOTE (A) and hydrogenated di-HOTE (B) synthesized from ALA by sLOX followed by reduction of hydroperoxides into hydroxylated derivatives by NaBH<sub>4</sub> treatment.

### Determination of carbon stereochemistry of di-hydroxylated metabolites synthesized from ALA

Commercial 9(S)-HOTE and racemic  $9(\pm)$ -HOTE, prepared as described in Materials and Methods, were incubated separately with sLOX as described with ALA, and further reduction by NaBH<sub>4</sub>. 9(S)-HOTE was metabolized into a main compound eluted at 33.4 min (Fig. 3A). This peak was attributed to 9(S), 16(S)-di-HOTE because sLOX produces almost exclusively the S enantiomer via a double dioxygenation mechanism (40), although soybean LOX generates almost exclusively 13(S)-HpOTE from ALA (40) as it was already shown (Fig. 1A). On the other hand, the 9(R),16(S)-di-HOTE peak, eluting at 33.7 min, was also formed (Fig. 3B) when racemic  $9(\pm)$ -HOTE was incubated with sLOX, compared with 9(S)-HOTE. The two diastereoisomers 9(S),16(S)-di-HOTE and 9(R),16(S)-di-HOTE eluted at 33.4 min and 33.7 min (Fig. 3B), respectively, exhibited the same intensity in agreement with the equivalent amounts of 9(S)- and 9(R)-HOTE present in the racemic mixture. The latter chromatogram (Fig. 3B) is similar to that presented in Fig. 1B, suggesting that the two peaks (retention time 33.2 and 33.5 min, Fig. 1B) observed with ALA as a substrate correspond to the two diastereoisomers 9(S), 16(S)di-HOTE and 9(R),16(S)-di-HOTE, respectively.

This assignment was also confirmed by the fact that chromatograms of di-hydroxylated isomers from 9(S)-HpOTE and 9(S)-HOTE were almost identical, but different However, we may assume that the two all trans diastereoisomers synthesized from ALA, as this will be confirmed by NMR below (retention time 32.1 and 32.6 min) (Fig. 1B), were likely to be all trans 9(R),16(S)-di-HOTE and 9(S),16(S)-di-HOTE, respectively, according to the separation of 8(R),15(S)- and 8(S),15(S)-all trans-diHETEs reported in (41, 42). Such an elution order has also been established for the separation of 5(S),12(R)- and 5(S),12(S)all trans LTB<sub>4</sub> and 5(S),12(R)- and 5(S),12(S)-all trans LTB<sub>5</sub> diastereoisomers (43–45).

# NMR determination of the conjugated triene double bound geometry

A NMR investigation was performed to determine the E/Z configuration in the triene moiety. A change in the E/Z configuration significantly modifies the <sup>1</sup>H-<sup>1</sup>H J-couplings among the protons bound to the  $sp^2$  carbons in C=C double bonds, assuming values around ~17 Hz for *E* configuration and around ~10 Hz for *Z* configuration (46). Furthermore, molecules like 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid (33.2 min, Fig. 1B) and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid



**Fig. 3.** RP-HPLC profile of 9(S), 16(S)-di-HOTE synthesized from 9(S)-HOTE by sLOX (A); RP-HPLC profile of 9(S), 16(S)-di-HOTE and 9(R), 16(S)-di-HOTE synthesized from  $9(\pm)$ -HOTE by sLOX (B). The intermediate hydroperoxide products were reduced by NaBH<sub>4</sub> prior to analysis. mAU, milli absorbance units.



**Fig. 4.** RP-HPLC profile of 9,16-di-HOTEs synthesized by sLOX from ALA (A); from 9(S)-HPOTE (B); and from 9(S)-HOTE (C). The intermediate hydroperoxide products were reduced by NaBH<sub>4</sub> prior to analysis. mAU, milli absorbance units.

(33.5 min, Fig. 1B), which only differ for the inversion of the stereochemical configuration at the chiral center of position 9, should show very similar spectra with minor differences in chemical shift and J-couplings, especially in the triene region. On this basis, the inspection of the 1D NMR spectra, and in particular of the triene resonance region (4-7 ppm), allows us to rapidly distinguish among the ensemble of molecules having the same triene configuration [e.g., the E,E,E compounds (32.1 min and 32.6 min, Fig. 1B) from the E,Z,E compounds (33.2 min and 33.5 min, Fig.1B)], as it can be seen in Figs. 5 and 6. The resonance assignment has been determined by using DQF COSY, while a large part of the J-couplings were measured through the analysis of 2D J-resolved spectra (data not shown). In such a way, it was possible to determine that the <sup>1</sup>H-<sup>1</sup>H <sup>3</sup>J-couplings between protons in positions 10-11 and 14-15 is around 15 Hz for all the examined cases, evidencing that the protons are in trans position with respect to the double bond. The determination of the <sup>1</sup>H-<sup>1</sup>H <sup>3</sup>J-coupling among protons 12 and 13 is complicated by the quasi-equivalence of their chemical shifts (strong-coupling condition) and by the presence of additional couplings with the closer protons 11 and 14. To simplify the problem, we observed the resonances of protons 12 and 13 in the 1D spectra of 9(R), 16(S)-dihydroxy-10E, 12Z, 14Eoctadecatrienoic acid (33.5 min, Fig. 1B) upon selective decoupling of the protons 11 and 14. This made a significant simplification of the multiplet pattern making evident a  ${}^{3}J_{12-13}$  coupling of  $\sim 10$  Hz typical of Z configuration. The same approach could not be done in the case of the *E*,*E*,*E* molecules because of the strong overlap among the protons 11, 12, 13, and 14 that have a very similar chemical

shift. Nevertheless, because the determination of the configuration in the E,Z,E molecules is demonstrated above, we can safely assign by exclusion the configuration of the E,E,E molecules on the basis of the marked differences among the spectra of E,E,E and E,Z,E.

#### Metabolism of ALA by human recombinant 15-LOX-2

This experiment was done to check whether the human recombinant 15-LOX type 2 is also able to synthesize such metabolites. For this purpose ALA, 9(S)-HOTE, and  $9(\pm)$ -HOTE were separately incubated with this enzyme. We found that human recombinant 15-LOX-2 is able to convert ALA into 13-HOTE as previously described with the sLOX (result not shown). However, ALA was less markedly metabolized into di-hydroxylated fatty acids, the main ones being all trans 9,16-di-HOTE isomers. Interestingly enough, the mono-hydroxylated derivatives 9(S)-HOTE and 9(R)-HOTE were well converted into 9(S),16(S)-*E*,*Z*,*E*-di-HOTE, respectively (result not shown) by the human recombinant LOX.

#### Inhibitory effect of 9,16-di-HOTEs on COX-1 and antiaggregatory effects

The inhibition of sheep COX-1 by different 9,16-di-HOTE isomers was tested at 1  $\mu$ M by measuring both PGD<sub>2</sub> and PGE<sub>2</sub> as described in Materials and Methods. Results reported in **Table 1** show that 1  $\mu$ M of 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acids decreased significantly the synthesis of both PGD<sub>2</sub> and PGE<sub>2</sub> by around 28% and 38%, respectively. Such results are in agreement with previous data showing that poxytrins, characterized by a *E*,*Z*,*E* 



**Fig. 5.** 1D spectra and J-couplings for 9(S),16(S)-dihydroxy-10*E*,12*E*,14*E*-octadecatrienoic acid (A) and 9(R),16(S)-dihydroxy-10*E*,12*E*, 14*E*-octadecatrienoic acid (B). Resonance assignment and J-couplings were determined from the analysis of a DQF COSY, and 2D J-resolved spectra (data not shown). Spectra were acquired with eight scans and 2 s of recycle delay, the acquisition time was 1.8 s. Spectra were processed with exponential line broadening of 0.3 Hz as window function.

ASBMB



**Fig. 6.** 1D spectra and J-couplings for 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid (A) and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid (B). Resonance assignments and J-couplings were determined from the analysis of a DQF COSY, and 2D J-resolved spectra (data not shown). Spectra were acquired with eight scans and 2 s of recycle delay, the acquisition time was 1.8 s. Spectra were processed with exponential line broadening of 0.3 Hz as window function.

H

ASBMB

	Control	9(S),16(S)-di-HOTE E,Z,E	9(R),16(S)-di-HOTE <i>E</i> , <i>Z</i> , <i>E</i>
COX-1			$a a a a a^{h}$
$(PGD_2 + PGE_2)$ COX-2	$134.9 \pm 15.8$	$96.8 \pm 14.0^{\circ}$	$83.2 \pm 9.0^{\circ}$
$(PGD_2 + PGE_2)$	$136.1 \pm 11.0$	$124.5\pm8.8^a$	$107.4\pm7.2^b$

Purified COX-1 and COX-2 were incubated with or without 1  $\mu$ M of different 9,16-di-HOTE isomers and with 10  $\mu$ M arachidonic acid. PGD<sub>2</sub> and PGE<sub>2</sub> were measured by GC-MS. Results expressed in ng of PGD<sub>2</sub> and PGE<sub>2</sub> represent the mean ± SEM of four determinations. Only the inhibitory effects of *E*, *Z*,*E* isomers are shown. <sup>*a*</sup>*P* < 0.05.

 ${}^{b}P < 0.01$  versus control.

I < 0.01 versus control.

triene geometry, are inhibitors of platelet COX-1 (19), whereas the all trans isomers are inactive.

Such results are in agreement with data relating to platelet aggregation induced by collagen. Platelet aggregation was significantly decreased by 64% and 65% in presence of 1  $\mu$ M 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid, respectively (**Fig. 7**). In addition, the aggregation started to be reversed from 2 min following the agonist addition. As previously shown (19), all trans isomers had no effect on platelet aggregation.

#### Anti-inflammatory effect of 9,16-di-HOTEs

After showing that 9,16-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid isomers possess anti-aggregatory properties as those previously described for poxytrins (19), it was important to know whether such di-hydroxylated compounds issued from ALA may exhibit anti-inflammatory properties. We then investigated the effects of 9,16-di-HOTEs on COX-2 as well as the 5-LOX pathway.

Effect of 9,16-di-HOTEs on human recombinant COX-2. The inhibition of human recombinant COX-2 by 1  $\mu$ M of 9,16-di-HOTE isomers was assessed by measuring PGD<sub>2</sub> and PGE<sub>2</sub> as described above. Results reported in Table 1 show that 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acids significantly inhibited COX-2 activity by 9% and 21%, respectively.

Effect of 9,16-di-HOTEs on the metabolism of arachidonic acid by the human polymorphonuclear leukocyte 5-LOX. The inhibition of the human 5-LOX pathway by 1  $\mu$ M of 9,16-di-HOTE isomers was assessed by measuring the following metabolites: 5(S)-HETE, LTB<sub>4</sub> and its two all trans isomers. As reported in **Table 2**, only the 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid isomer significantly decreased the formation of leukotrienes and 5(S)-HETE by around 10% and 20%, respectively. All other isomers did not inhibit such a production (results not shown). Such data suggest that the inhibition concerns only 5-LOX but not leukotriene A<sub>4</sub> hydroxylase. Taken together, these data suggest that the 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid isomer may have a potential anti-inflammatory effect.

#### DISCUSSION

ALA is an essential fatty acid present in vegetable oils, and as such available in human diet. It is the precursor of EPA and DHA, but its conversion into EPA and DHA is rather limiting with around 10% to the former and a further 10% to the latter, which means only 1% conversion of ALA to DHA (47, 48). ALA is also rather well  $\beta$ -oxidized (49, 50), that could partially explain its low accumulation in plasma and tissues. In addition, we describe here a pathway which could explain some beneficial effects of ALA in the vascular system at large by reducing both the



 $\rightarrow$ 

**Fig. 7.** Effect of *E*,*Z*,*E*9,16-di-HOTEs on platelet aggregation triggered by collagen. Platelet suspension (400  $\mu$ l) was preincubated for 1 min at 37°C and triggered by ~0.05 ng/ $\mu$ l collagen in the presence or absence of 1  $\mu$ M 9,16-di-HOTEs. Platelet aggregation was monitored for 4 min. The aggregation profiles shown are representative of three independent experiments. T, transmission.

TABLE 2. Effect of 9,16-di-HOTEs on human leukocyte 5-LOX

	Control	9(R),16(S)-di-HOTE <i>E,Z,I</i>
LTB <sub>4</sub> s	$9.57 \pm 0.66$	$8.78\pm0.78^a$
5-HETE	$0.89 \pm 0.03$	$0.71 \pm 0.05^{a}$
5-LOX products	$10.46\pm0.68$	$9.49 \pm 0.81^{a}$

Isolated human leukocytes were incubated in the presence or absence of 1  $\mu$ M of different 9,16-di-HOTE isomers at 37°C and triggered with 1  $\mu$ M of ionophore A23187 and 10  $\mu$ M of arachidonic acid. Oxygenated metabolites were then extracted and quantified by RP-HPLC. Results expressed in nmol of total LTB<sub>4</sub>s (LTB<sub>4</sub> + LTB<sub>4</sub> isomers) and 5-HETE represent the mean ± SEM of five determinations. Only the inhibitory effect of 9(R),16(S)-di-HOTE is shown.

 $^{a}P < 0.05$ , versus control.



production of pro-inflammatory eicosanoids  $PGD_2/PGE_2$ and  $LTB_4$  (51–53). We recently showed that a new di-hydroxylated metabolite from DHA called PDX, which is an isomer of the anti-inflammatory PD1, exhibits anti-aggregatory properties, both by inhibiting collagen-induced prostanoid production, including thromboxane A<sub>2</sub>, and thromboxane A<sub>2</sub>-induced aggregation (19). In the present paper we report the full characterization of the lipoxygenation of ALA, and the anti-inflammatory and anti-aggregatory effects of two LOX end-products. The structure of the four LOX metabolites that we herein describe have already been partially reported (23, 24), but the proposed stereochemistry structures of the 9(S),16(R)-di-HOTE do not fit with our experimental data. On the other hand, their biological properties have not been investigated.

Our data show that ALA is mainly converted into its 13oxygenated derivative 13-HOTE by 15/ω6-LOX. However, formation of a small amount of the 9-oxygenated product 9-HOTE could also be observed. In contrast to DHA, which is converted into only one main di-hydroxylated compound (PDX) via a double lipoxygenation, four di-hydroxylated compounds were observed from ALA following incubation with 15-LOX, characterized as 9,16-di-HOTEs. The stereochemistry of these di-hydroxylated compounds was assessed by incubating 9(S)-HOTE and racemic  $9(\pm)$ -HOTE separately with sLOX (see the above results). The UV spectra of the two main di-hydroxylated products are superimposed to that of PDX, which possesses a *E*,*Z*,*E* conjugated triene geometry. The diastereoisomers are likely produced via a dioxygenation mechanism as previously described for PDX (18) and these two compounds are likely to have an E,Z,Econjugated triene geometry. Such a geometry was further confirmed by high field NMR which allowed us to determine precisely the <sup>1</sup>H-<sup>1</sup>H J-couplings among the protons bound to the  $sp^2$  carbons in C=C double bonds. The geometry of the conjugated trienes of two additional but less abundant (around half the amount of the E,Z,E-9(S),16(S)di-HOTE and 9(R),16(S)-di-HOTE) di-hydroxylated products was supposed to be *E,E,E* because their UV spectra were superimposed to that of all trans LTB<sub>4</sub>, which has such an E,E,E conjugated triene geometry. This all trans geometry was also confirmed by NMR to determine the <sup>1</sup>H-<sup>1</sup>H J-couplings among the protons bound to the  $sp^2$  carbons in C=C double bonds. Based on data from the literature, we may suggest that the stereochemistry of these two minor products (Fig. 1B) is 9(R),16(S)- and 9(S),16(S)-all trans-di-HOTE,

in order of increasing retention time, respectively. Such a configuration is also in agreement with the 5(S), 12(R)- and 5(S), 12(S)-all trans LTB<sub>4</sub> and the 5(S), 12(R)- and 5(S), 12(S)all trans LTB<sub>5</sub> diastereoisomers already described (43–45). We conclude that the four di-hydroxylated compounds issued from ALA treated by sLOX are 9(R),16(S)-dihydroxy-10E,12E, 14E-octadecatrienoicacid,9(S),16(S)-dihydroxy-10E,12E,14Eoctadecatrienoic acid, 9(S),16(S)-dihydroxy-10E,12Z,14Eoctadecatrienoic acid, and 9(R),16(S)-dihydroxy-10E,12Z, 14E-octadecatrienoic acid, in order of increasing retention time, respectively. Such data suggest that there are two different mechanisms explaining the formation of these 9,16-di-HOTE isomers. The *E*, *Z*,*E* compounds 9(S),16(S)-di-HOTE and 9(R), 16(S)-di-HOTE would result from a dioxygenation mechanism as previously shown for PDX (18), whereas the all trans 9,16-di-HOTE isomers might derive from unknown intermediates.

We also investigated the metabolism of ALA incubated with human recombinant 15-LOX-2. Interestingly, 9(S)-HOTE or 9(R)-HOTE were both well converted into the corresponding 9,16-di-HOTEs whereas ALA was less efficiently metabolized into those di-hydroxylated fatty acids than by sLOX. Such a result might be biologically relevant because 9(S)-HOTE can be produced through cyclooxygenases, and 9(R)-HOTE can be produced via aspirinated COX-2, both in an aborted cyclooxygenation process (54).

The present study also shows some biological effects of the E,Z,E di-hydroxylated 9,16-di-HOTE isomers. We observed that 9(S),16(S)-dihydroxy-10E,12Z,14E-octadecatrienoic and 9(R),16(S)-dihydroxy-10E,12Z,14E-octadecatrienoic acids decreased significantly prostaglandin synthesized by recombinant COX-1. Moreover, these E,Z,E isomers inhibit platelet aggregation triggered by collagen both by decreasing the extent of aggregation and making it partially reversible (Fig. 7). However, these E,Z,E di-hydroxylated ALA metabolites appear to be less potent inhibitors than PDX under the same conditions, although the anti-aggregatory effects observed are in agreement with our previous data showing that poxytrins, characterized by an E,Z,E conjugated triene, are inhibitors of COX-1 (19).

Downloaded from www.jlr.org at Carleton Univ - OCUL, on June 19, 2014

COX-2 is an inducible enzyme becoming abundant at sites of inflammation, and is upregulated in various carcinomas, having a central role in tumorigenesis (55). The inhibition of COX-2 is a major mode of action to reduce inflammation (56). In our study, we found that 9(S),16(S)dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic and 9(R),16(S)dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acids exhibit an inhibitory effect on the human recombinant COX-2.

Moreover, 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid also decreased significantly the formation of LTB<sub>4</sub> and 5(S)-HETE from arachidonic acid incubated with human polymorphonuclear leukocytes, indicating the inhibition of 5-LOX.

It is interesting to note that in all the tests evaluated, the  $E,Z,E9(\mathbf{R}),16(\mathbf{S})$  isomer was a more potent inhibitor than the  $E,Z,E9(\mathbf{S}),16(\mathbf{S})$  isomer. This reinforces the interest of the 15-lipoxygenation of  $9(\mathbf{R})$ -HOTE produced by aspirinated COX-2 (54) to synergize with aspirin in atherothrombogenesis. In conclusion, we observe from the present study that four di-hydroxylated compounds are produced from ALA treated by sLOX. They are 9(R),16(S)-dihydroxy-10*E*,12*E*,14*E*-octadecatrienoic acid, 9(S),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid, and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadeca trienoic acid, and 9(R),16(S)-dihydroxy-10*E*,12*Z*,14*E*-octadeca trienoic acid. Interestingly, 9,16-dihydroxy-10*E*,12*Z*, 14*E*-octadecatrienoic acid isomers exhibit anti-aggregatory properties as other poxytrins (19). In addition, 9(R),16(S)dihydroxy-10*E*,12*Z*,14*E*-octadecatrienoic acid, which can be produced from 9(R)-HOTE by aspirinated COX-2, inhibits both cyclooxygenases and the 5-LOX pathway and appears to be a potential anti-inflammatory compound in vitro.

The authors acknowledge the use of the 1 GHz spectrometer installed at the French high field NMR Center in Lyon (TGE RMN THC Fr3050).

#### REFERENCES

- Mozaffarian, D., and J. H. Wu. 2011. Omega-3 fatty acids and cardiovascular disease: effects on risk factors, molecular pathways and clinical events. J. Am. Coll. Cardiol. 58: 2047–2067.
- Kris-Etherton, P. M., W. S. Harris, and L. J. Appel. 2002. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation.* 106: 2747–2757.
- Rennie, K. L., J. Hughes, R. Lang, and S. A. Jebb. 2003. Nutritional management of rheumatoid arthritis: a review of the evidence. *J. Hum. Nutr. Diet.* 16: 97–109.
- McCann, K. 2007. Nutrition and rheumatoid arthritis. *Explore (NY)*. 3: 616–618.
- Venkatraman, J. T., and W. C. Chu. 1999. Effects of dietary omega-3 and omega-6 lipids and vitamin E on serum cytokines, lipid mediators and anti-DNA antibodies in a mouse model for rheumatoid arthritis. J. Am. Coll. Nutr. 18: 602–613.
- Holm, T., A. K. Andreassen, P. Aukrust, K. Andersen, O. R. Geiran, J. Kjekshus, S. Simonsen, and L. Gullestad. 2001. Omega-3 fatty acids improve blood pressure control and preserve renal function in hypertensive heart transplant recipients. *Eur. Heart J.* 22: 428–436.
- Weisinger, H. S., J. A. Armitage, A. J. Sinclair, A. J. Vingrys, P. L. Burns, and R. S. Weisinger. 2001. Perinatal omega-3 fatty acid deficiency affects blood pressure later in life. *Nat. Med.* 7: 258–259.
- James, M. J., R. A. Gibson, and L. G. Cleland. 2000. Dietary polyunsaturated fatty acids and inflammatory mediator production. *Am. J. Clin. Nutr.* 71(1 Suppl.): 343S–348S.
- Furse, R. K., R. G. Rossetti, and R. B. Zurier. 2001. Gammalinolenic acid, an unsaturated fatty acid with anti-inflammatory properties, blocks amplification of IL-1 beta production by human monocytes. *J. Immunol.* 167: 490–496.
- Song, C., X. Li, B. E. Leonard, and D. F. Horrobin. 2003. Effects of dietary n-3 or n-6 fatty acids on interleukin-1beta-induced anxiety, stress, and inflammatory responses in rats. J. Lipid Res. 44: 1984–1991.
- Conquer, J. A., M. C. Tierney, J. Zecevic, W. J. Bettger, and R. H. Fisher. 2000. Fatty acid analysis of blood plasma of patients with Alzheimer's disease, other types of dementia, and cognitive impairment. *Lipids.* 35: 1305–1312.
- Serhan, C. N., C. B. Clish, J. Brannon, S. P. Colgan, N. Chiang, and K. Gronert. 2000. Novel functional sets of lipid-derived mediators with antiinflammatory actions generated from omega-3 fatty acids via cyclooxygenase 2-nonsteroidal antiinflammatory drugs and transcellular processing. J. Exp. Med. 192: 1197–1204.
- Serhan, C. N., N. Chiang, and T. E. Van Dyke. 2008. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat. Rev. Immunol.* 8: 349–361.
- Serhan, C. N. 2007. Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators end pathways. *Annu. Rev. Immunol.* 25: 101–137.
- Serhan, C. N., K. Gotlinger, S. Hong, Y. Lu, J. Siegelman, T. Baer, R. Yang, S. P. Colgan, and N. A. Petasis. 2006. Anti-inflammatory

actions of neuroprotectin D1/protectin D1 and its natural stereoisomers: assignments of dihydroxy-containing docosatrienes. *J. Immunol.* **176:** 1848–1859. [Erratum. 2006. *J. Immunol.* **176:** 3843.]

- Serhan, C. N., R. Yang, K. Martinod, K. Kasuga, P. S. Pillai, T. F. Porter, S. F. Oh, and M. Spite. 2009. Maresins: novel macrophage mediators with potent antiinflammatory and proresolving actions. *J. Exp. Med.* **206**: 15–23.
- Dona, M., G. Fredman, J. M. Schwab, N. Chiang, M. Arita, A. Goodarzi, G. Cheng, U. H. von Andrian, and C. N. Serhan. 2008. Resolvin E1, an EPA-derived mediator in whole blood, selectively counterregulates leukocytes and platelets. *Blood.* **112**: 848–855.
- Chen, P., B. Fenet, S. Michaud, N. Tomczyk, E. Véricel, M. Lagarde, and M. Guichardant. 2009. Full characterization of PDX, a neuroprotectin/protectin D1 isomer, which inhibits blood platelet aggregation. *FEBS Lett.* 583: 3478–3484.
- Chen, P., E. Véricel, M. Lagarde, and M. Guichardant. 2011. Poxytrins, a class of oxygenated products from polyunsaturated fatty acids, potently inhibit blood platelet aggregation. *FASEB J.* 25: 382–388.
- Hussein, N., E. Ah-Sing, P. Wilkinson, C. Leach, B. A. Griffin, and D. J. Millward. 2005. Long-chain conversion of [13C]linoleic acid and alpha-linolenic acid in response to marked changes in their dietary intake in men. J. Lipid Res. 46: 269–280.
- Burdge, G. C., and S. A. Wootton. 2002. Conversion of alpha-linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women. *Br. J. Nutr.* 88: 411–420.
- 22. Feiters, M. C., G. A. Veldink, and J. F. G. Vliegenthart. 1984. Conjugated dihydroperoxyoctadecatrienoic fatty acids formed upon double dioxygenation of α-linolenic acid by lipoxygenase-2 from soybeans. *In* Oxidative Damage and Related Enzymes, G. Rotilio and J. V. Bannister, editors. Harwood Academic Publishers, Chur, Switzerland; 132–138.
- Sok, D. E., and M. R. Kim. 1990. Enzymatic formation of 9,16dihydro(pero)xyoctadecatrienoic acid isomers from alpha-linolenic acid. Arch. Biochem. Biophys. 277: 86–93.
- Sok, D. E., and M. R. Kim. 1994. Conversion of alpha-linolenic acid to dihydro(pero)xyoctadecatrienoic acid isomers by soybean and potato lipoxygenases. J. Agric. Food Chem. 42: 2703–2708.
- Grechkin, A. N., R. A. Kuramshin, E. Y. Safonova, Y. J. Yefremov, S. K. Latypov, A. V. Ilyasov, and I. A. Tarchevsky. 1991. Double hydroperoxidation of alpha-linolenic acid by potato tuber lipoxygenase. *Biochim. Biophys. Acta.* 1081: 79–84.
- Piantini, U., O. W. Sorensen, and R. R. Ernst. 1982. Multiple quantum filters for elucidating NMR coupling networks. J. Am. Chem. Soc. 104: 6800–6801.
- Shaka, A. J., and R. Freeman. 1983. Simplification of NMR spectra by filtration through multiple-quantum coherence. *J. Magn. Reson.* 51: 169–173.
- Derome, A. E., and M. P. Williamson. 1990. Rapid-pulsing artifacts in double-quantum-filtered COSY. J. Magn. Reson. 88: 177–185.
- Aue, W., J. Karhan, and R. Ernst. 1976. Homonuclear broad-band decoupling and 2-dimensional J-resolved NMR spectroscopy. J. Chem. Phys. 64: 4226–4227.
- Nagayama, K., P. Bachmann, K. Wüthrich, and R. R. Ernst. 1978. The use of cross-sections and of projections in two-dimensional NMR spectroscopy. J. Magn. Reson. 31: 133–148.
- Wider, G., R. Baumann, K. Nagayama, R. R. Ernst, and K. Wüthrich. 1981. Strong spin-spin coupling in the two-dimensional *J*-resolved 360-MHz <sup>1</sup>H NMR spectra of the common amino acids. *J. Magn. Reson.* 42: 73–87.
- 32. Kühn, H., J. Barnett, D. Grunberger, P. Baecker, J. Chow, B. Nguyen, H. Bursztyn-Pettegrew, H. Chan, and E. Sigal. 1993. Over-expression, purification and characterization of human recombinant 15-lipoxygenase. *Biochim. Biophys. Acta.* **1169**: 80–89.
- 33. Shappell, S. B., R. A. Gupta, S. Manning, R. Whitehead, W. E. Boeglin, C. Schneider, T. Case, J. Price, G. S. Jack, T. M. Wheeler, et al. 2001. 15S-hydroxyeicosatetraenoic acid activates peroxisome proliferator-activated receptor γ and inhibits proliferation in PC3 prostate carcinoma cells. *Cancer Res.* **61**: 497–503.
- Lagarde, M., P. A. Bryon, M. Guichardant, and M. Dechavanne. 1980. A simple and efficient method for platelet isolation from their plasma. *Thromb. Res.* 17: 581–588.
- Maclouf, J., B. F. de Laclos, and P. Borgeat. 1982. Stimulation of leukotriene biosynthesis in human blood leukocytes by plateletderived 12-hydroperoxy-icosatetraenoic acid. *Proc. Natl. Acad. Sci.* USA. 79: 6042–6046.

BMB

- Born, G. V. R. 1962. Aggregation of blood platelets by adenosine diphosphate and its reversal. *Nature*. 194: 927–929.
- Watanabe, T., S. Narumiya, T. Shimizu, and O. Hayaishi. 1982. Characterization of the biosynthetic pathway of prostaglandin D2 in human platelet-rich plasma. *J. Biol. Chem.* 257: 14847–14853.
- Turk, J., J. R. Colca, N. Kotagal, and M. L. McDaniel. 1984. Arachidonic acid metabolism in isolated pancreatic islets: I. Identification and quantitation of lipoxygenase and cyclooxygenase products. *Biochim. Biophys. Acta.* **794**: 110–124.
- Schweer, H., J. Kammer, and H. W. Seyberth. 1985. Simultaneous determination of prostanoids in plasma by gas chromatographynegative-ion chemical-ionization mass spectrometry. *J. Chromatogr.* 338: 273–280.
- 40. Van Os, C. P. A., G. P. M. Rijke-Schilder, H. V. Halbeek, J. Verhagen, and J. F. G. Vliegenthart. 1981. Double dioxygenation of arachidonic acid by soybean lipoxygenase-1. Kinetics and regiostereo specificities of the reaction steps. *Biochim. Biophys. Acta.* 663: 177–193.
- 41. Maas, R. L., and A. R. Brash. 1983. Evidence for a lipoxygenase mechanism in the biosynthesis of epoxide and dihydroxy leukotrienes from 15(S)-hydroperoxyicosatetraenoic acid by human platelets and porcine leukocytes. *Proc. Natl. Acad. Sci. USA.* 80: 2884–2888.

SBMB

JOURNAL OF LIPID RESEARCH

- Maas, R. L., A. R. Brash, and J. A. Oates. 1981. A second pathway of leukotriene biosynthesis in porcine leukocytes. *Proc. Natl. Acad. Sci.* USA. 78: 5523–5527.
- Lee, C. W., R. A. Lewis, E. J. Corey, A. Barton, H. Oh, A. I. Tauber, and K. F. Austen. 1982. Oxidative inactivation of leukotriene C<sub>4</sub> by stimulated human polymorphonuclear leukocytes. *Proc. Natl. Acad. Sci. USA*. **79**: 4166–4170.
- Mencia-Huerta, J. M., E. Razin, E. W. Ringel, E. J. Corey, D. Hoover, K. F. Austen, and R. A. Lewis. 1983. Immunologic and ionophoreinduced generation of leukotriene B<sub>4</sub> from mouse bone marrowderived mast cells. *J. Immunol.* 130: 1885–1890.
- 45. Lee, T. H., J. M. Menica-Huerta, C. Shih, E. J. Corey, R. A. Lewis, and K. F. Austen. 1984. Characterization and biologic properties of

5,12-dihydroxy derivatives of eicosapentaenoic acid, includingleukotriene  $B_5$  and the double lipoxygenase product. *J. Biol. Chem.* **259**: 2383–2389.

- Silverstein, R. M., and F. X. Webster. 1998. Spectrometric Identification of Organic Compounds. 6th edition. Wiley, New York. 212–213.
- Brenna, J. T. 2002. Efficiency of conversion of alpha-linolenic acid to long chain n-3 fatty acids in man. *Curr. Opin. Clin. Nutr. Metab. Care.* 5: 127–132.
- Sinclair, A. J., N. M. Attar-Bashi, and D. Li. 2002. What is the role of alpha-linolenic acid for mammals? *Lipids*. 37: 1113–1123.
- Burdge, G. C. 2006. Metabolism of α-linolenic acid in humans. Prostaglandins Leukot. Essent. Fatty Acids. 75: 161–168.
- Leyton, J., P. J. Drury, and M. A. Crawford. 1987. Differential oxidation of saturated and unsaturated fatty acids in vivo in rat. *Br. J. Nutr.* 57: 383–393.
- 51. Ren, J., E. J. Han, and S. H. Chung. 2007. In vivo and in vitro anti-inflammatory activities of alpha-linolenic acid isolated from Actinidia polygama fruits. *Arch. Pharm. Res.* **30**: 708–714.
- Ishihara, K., W. Komatsu, H. Saito, and K. Shinohara. 2002. Comparison of the effects of dietary alpha-linolenic, stearidonic, and eicosapentaenoic acids on production of inflammatory mediators in mice. *Lipids.* 37: 481–486.
- Cherian, G. 2007. Metabolic and cardiovascular disease in poultry: role of dietary lipids. *Poult. Sci.* 86: 1012–1016.
- 54. Schneider, C., and A. R. Brash. 2000. Stereospecificity of hydrogen abstraction in the conversion of arachidonic acid to 15R-HETE by aspirin-treated cyclooxygenase-2. Implications for the alignment of substrate in the active site. *J. Biol. Chem.* 275: 4743–4746.
- Aggarwal, B. B., S. Shishodia, S. K. Sandur, M. K. Pandey, and G. Sethi. 2006. Inflammation and cancer: how hot is the link? *Biochem. Pharmacol.* 72: 1605–1621.
- Masferrer, J. L., B. S. Zweifel, P. T. Manning, S. D. Hauser, K. M. Leahy, W. G. Smith, P. C. Isakson, and K. Seibert. 1994. Selective inhibition of inducible cyclooxygenase 2 in vivo is antiinflammatory and nonulcerogenic. *Proc. Natl. Acad. Sci. USA*. 91: 3228–3232.