

Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. M. Nielsen, P. Maa, E. Þ. Baldursson and C. M. Pedersen, *Chem. Sci.*, 2019, DOI: 10.1039/C9SC00857H.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Self-promoted and stereospecific formation of *N*-glycosidesMichael Martin Nielsen,^a Patrycja Mała,^{a,b} Eiríkur Þórir Baldursson,^a and Christian Marcus Pedersen^{a*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


A stereoselective and self-promoted glycosylation for the synthesis of various *N*-glycosides and glycosyl sulfonamides from trichloroacetimidates is presented. No additional catalysts or promoters are needed in what is essentially a two-component reaction. When α -glucosyl trichloroacetimidates are employed, the reaction resulted in the stereospecific formation of the corresponding β -*N*-glucosides in high yields at ambient conditions. On the other hand when equatorial glucosyl donors were used, the stereospecificity decreased and resulted in a mixture of anomers. By NMR-studies, it was concluded that this decrease in stereospecificity was due to an, until now, unrepresented anomerization of the trichloroacetimidate under the very mildly acidic conditions. The mechanism and kinetics of the glycosylations have been studied by NMR-experiments, which gave an insight into the activation of trichloroacetimidates, suggesting an S_N1 -like mechanism involving ion pairs. The scope of glycosyl donors and sulfonamides was found to be very broad including popular *N*-protective groups and common glycosyl donors of various reactivity. Peracetylated GlcNAc trichloroacetimidate could be used without the need for any promoters or additives and a tyrosine side chain was glycosylated as an *N*-glycosyl carbamate. The *N*-carbamates and the *N*-sulfonyl groups functioned as orthogonal protective groups of the *N*-glycoside and hence allowed further *N*-functionalization without risking mutarotation of the *N*-glycoside. The *N*-glycosylation was also performed on a gram scale, without a drop in stereoselectivity nor yield.

Introduction

Trichloroacetimidates (TCAs) remain one of the most popular glycosyl donors in catalytic glycosylations.¹ Since the introduction of TCA donors by Schmidt,² numerous Lewis- and Brønsted acids have been shown to catalyze their activation.^{2–4} In common for many of the methods is the formation of an oxocarbenium ion intermediate or similar, i.e. with no memory of the stereochemistry of the donor.⁵ The TCAs are synthesized from reacting the hemiacetal with trichloroacetonitrile under basic conditions and depending on the conditions, mainly the α - or the β -anomer can be synthesized.^{6–8} Since it is possible to access the TCAs stereoselectively, several researchers have been interested in utilizing this particular glycosyl donor in stereospecific glycosylations.⁹

Stereospecific glycosylations have previously been performed via pre-complexation in *O*-glycoside synthesis by using additives such as chloral,³ boron fluorides¹⁰ or silanes¹¹ to complex the acceptor alcohol making it more acidic and hence able to activate the trichloroacetimidate. This method was however initially limited to a few glycosyl donor types and simple glycosyl acceptors. Still, the early results sparked a wide

interest in this concept that has since led to substantial advances, which has recently been reviewed in detail.^{9,12}

Scheme 1 Self-promoted *O*- and *N*-glycosylations.

In the early eighties, the Schmidt group reported that certain carboxylic acids and phosphorous acid derivatives could activate the α -trichloroacetimidates and substitute these with inversion of stereochemistry (Scheme 1).¹³ The inversion of stereochemistry was explained by pre-complexation of the TCA donor and acid, leading to a pseudo six-membered transition state³ and a

^a Department of Chemistry, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen O, Denmark

^b Faculty of Chemistry, Adam Mickiewicz University, Umultowska 89b, 61614 Poznań, Poland

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



concerted mechanism, without the formation of an oxocarbenium intermediate.

Likewise, the Miller group have shown more recently¹⁴ that a wide range of carboxylic acids can act as catalytic activators of TCA donors, concluding that the catalytic efficiency of these catalysts was inversely proportional to their pK_a -value; carboxylic acids with higher pK_a -values simply resulted in self-condensation of the catalyst and TCA donor, thus yielding undesired glycosyl esters. It is important to note that these glycosyl ester by-products were produced with inverted stereochemistry compared to the TCA donors (Scheme 1). There have been numerous reports of the *in situ* formation of several different intermediates,¹⁵ but the concept of self-condensation of the glycosyl donor and an acidic reactant in the context of glycoside synthesis has only received little attention. However, it seems quite clear that this reaction could have interesting applications considering the apparent stereospecific formation of the undesired by-products.

As part of our ongoing interest in stereospecific catalytic glycosylations, we have studied acidic N-H groups as potential self-promoters in glycosylation (Scheme 1).

Glycosyl sulfonamides are an important group of compounds that have been shown to be carbonic anhydrase inhibitors^{16–20} and more recently, the antitumor activity of this class of compounds has been recognized.^{17,21,22} Their synthesis has mainly been from reduction of the corresponding β -glycosyl azide,^{23–27} reductive amination of the lactol,²⁸ from 1,2 diols using Burgess reagent,²⁹ from glycals^{16,19,21,30–36} or via anomeric substitution under strongly acidic conditions.^{17,37,38} Thus, it would be of interest to get access to this class of compounds via a milder, higher-yielding route.

The Miller group have reported that sulfonamides are sufficiently acidic to act as catalytic activators of TCA glycosyl donors, enabling the synthesis of *O*-glycosides in good yields, albeit with low stereoselectivity.³⁹ This inspired us to develop a glycosylation procedure in which a glycosyl TCA donor is condensed with a sulfonamide, leading to the stereoselective formation of the desired *N*-glycosides. This reaction takes place in common organic solvents without the addition of any additives and is, to the best of our knowledge, the first example of a self-promoted *N*-glycosylation. This has resulted in the stereoselective formation of various glycosyl sulfonamides in high yields on a gram scale, including the synthesis of an amino acid-functionalized *N*-glycoside. We believe that this method serves as the simplest possible method for selective *N*-glycosylations and seek to further broaden its scope and possibilities in future studies.

Synthesis of *N*-glycosides

Initial glycosylation reactions

Glycosyl donors **1–6** (Chart 1) were investigated in a series of self-promoted *N*-glycosylations to investigate the effect of various protecting group patterns, donor reactivities and configurations; **1–3** represents the three most common glycosyl donors employed in carbohydrate synthesis, deoxysugar **4** represents a highly reactive glycosyl donor whereas donors **5** and

6 exemplifies unreactive, but synthetically interesting, glycosyl donors.

DOI: 10.1039/C9SC00857H




Chart 1 Glycosyl donors used in this study.

The scope of glycosyl acceptors consisted of various functional groups (Chart 2), namely tosylcarbamates **7–11**, **16** and **17** and nosylcarbamates **12–15**. These carbamates are selected for various reasons; **7–15** represent a wide pK_a -range, which could give an indication of the limits of the self-promoted glycosylation procedure, but also vary significantly in size, allowing an investigation of how steric bulk influences this particular glycosylation reaction. Also, it was found that both the tosyl- and nosyl groups can be selectively removed after the glycosylation using two different procedures, further widening the scope of post-glycosylational *N*-functionalization (*vide infra*).

An amino acid-derived glycosyl acceptor was also investigated in this study, namely tyrosine-derivative, **16**. Also carbamate **17** was found as an interesting *O*-functionalized carbamate for the self-promoted glycosylations. Compounds **7–11**, **13–15**, **16** and **17** were all obtained in a single step from commercially available starting materials (Scheme 2).





Chart 2 Glycosyl acceptors employed for self-promoted *N*-glycosylations





Scheme 2 General reaction scheme for the preparation of tosylcarbamates and nosylcarbamates.

Certain trends were clear from the initial series of glycosylations carried out from the α -glycosyl donors (Table 1). Glycosylations with **1a** (Entries 1-10) were essentially β -stereospecific, resulting in yields of 78-95% at various concentrations. Even notoriously unreactive glycosyl donors such as **5** and **6** (Entries 25-27) served as suitable glycosyl donors in this glycosylation procedure, both resulting in formation of the corresponding β -glycosides in good yields, albeit at elevated temperatures. No formation of the common oxazoline by-product^{40,41} was observed during the glycosylation with donor **6**.

Boc-derivatives **8** and **13** (Table 1, entries 11-13) were unsuitable for this type of reaction and led to the formation of by-products which made both purification and determination of stereoselectivity troublesome. This indicated a certain limit for the size of the *N*-substituents for this particular reaction to proceed successfully. Despite this, the desired *N*-glycosides were obtained, although in relatively lowered yields.

Glycosylations involving donor **2a** (Entries 14-18) proceeded in high yields of 61-90%, with a clear preference for the formation of the desired β -glycoside. Glycosylations with mannopyranosyl TCA donor **3a** were α -selective, indicating that the formation of 1,2-*cis* glycosidic linkages were likely disfavored by the steric factors due to the bulkiness of the acceptors. This was also the case when the same glycosylation was carried out at lowered temperature (Entry 24). The steric bulk of the acceptors was also apparent when analyzing the ¹H-NMR spectra of the α -glycosides since these were found to adopt a skew boat conformation, whereas all the β -glycosides all adopted a ⁴C₁-like conformation (Figure 1).

It was investigated whether altering the reaction concentration would have an effect on the yield and selectivity (Table 1). This gave rise to alternating variations in yield depending on which acceptor was used, but the stereoselectivity seemingly remained unaffected by the change in concentration. In practice, this glycosylation method was incredibly easy as it only required mixing two compounds in CH₂Cl₂. Furthermore, the excess of acceptor could be removed by an extraction with 1M aq. NaOH.⁴² The excess of acceptor could alternatively be recovered by flash column chromatography.

Table 1 Overview of initial glycosylations employing α -TCA donors **1-6**

DOI: 10.1039/C9SC00857H

Entry	Donor	Acceptor	t (h)	c (M) ^a	Yield (%) ^b	α/β ^c
1	1 α	7	o.n.	0.15	92	0:100
2	1 α	9	16	0.1	80	0:100
3	1 α	9	16	0.2	90	0:100
4	1 α	10	o.n.	0.2	95	0:100
5	1 α	11	o.n.	0.2	93	0:100
6	1 α	12	o.n.	0.08	67	1:99
7	1 α	14	4.5	0.2	62	6:94
8	1 α	15	1	0.04	85	0:100
9	1 α	15	1	0.08	95	6:94
10	1 α	15	0.75	0.2	78	6:94
11	1 α	8	24	0.2	35 ^d	0:100 ^d
12	1 α	13	4	0.2	33 ^d	n.d. ^d
13	1 α	13	o.n.	0.05	79 ^d	0:100 ^d
14	2 α	9	18	0.2	63	29:71
15	2 α	11	16	0.2	61	34:66
16	2 α	14	2.5	0.2	79	25:75
17	2 α	15	3	0.04	90	26:74
18	2 α	15	5	0.2	90	33:67
19	3 α	7	24	0.2	78	100:0
20	3 α	9	20	0.2	64	100:0
21	3 α	10	o.n.	0.2	74	100:0
22	3 α	11	24	0.2	72	100:0
23	3 α	14	6.5	0.2	79	100:0
24 ^e	3 α	14	o.n.	0.2	66	100:0
25 ^f	5	15	3.25	0.2	70	0:100
26 ^f	6	14	1.5	0.2	44	19:81
27 ^f	6	15	1.5	0.2	45	5:95

^aConcentration of glycosyl donor. ^bIsolated yield. ^cDetermined from crude ¹H-NMR.

^dThe Boc-derivatives **8** and **13** were found to give rise to various by-products, making it challenging to precisely determine yield and selectivity. ^eReaction was stirred at -78 °C for 3 h and then allowed to reach 0 °C. ^fReaction carried out in 1,2-dichloroethane at 80 °C.




Figure 1 Observed conformations of *N*-glycosides determined from ¹H-NMR coupling constants.

Next, it was investigated whether a glycosylation with β -TCA glycosyl donors would produce the corresponding α -glycosides with a similar stereospecific inversion of stereochemistry as was found in the initial glycosylations with α -TCA donors. These results are summarized in Table 2.



Table 2 Overview of glycosylations employing β -TCA donors **1** and **2**.

Entry	Donor	Acceptor	t (h)	c (M) ^a	Yield (%) ^b	α/β ^c
1	1β	7	4 d	0.15	96	66:34
2	1β	9	16	0.2	73	43:57
3	1β	10	o.n.	0.2	80	50:50
4	1β	11	o.n.	0.2	82	45:55
5	1β	12	24	0.04	45	76:24
6	1β	14	4.5	0.2	78	52:48
7	2β	7	o.n.	0.2	71	77:23
8	2β	14	2.5	0.2	67	80:20



^aConcentration of glycosyl donor. ^bIsolated yield. ^cDetermined from crude ¹H-NMR.

Generally, the glycosylations proceeded in high yields of 67–82% and with a higher degree of α -selectivity. It was clear that although a significantly higher selectivity for the desired α -glycosides was achieved, the stereospecificity was lower than in the previous glycosylations summarized in Table 1, indicative of unfavorable configuration of 1,2-*cis*-glycosides with bulky glycosyl acceptors. It is also noteworthy that the galactosylations (Entries 7 and 8) proceeded with a higher degree of stereospecificity than the corresponding glucosylations (Entries 1–6).

In an attempt to rule out whether it was a sterically repulsive effect from the C2-substituents that was inhibiting the formation of 1,2-*cis* glycosidic linkages, 2-deoxyglucosyl TCA donors **4 α** and **4 β** were synthesized (Scheme 3). It was however found that both glycosylations gave rise to essentially identical stereoselectivity independent on the configuration of the glycosyl donor, which strongly suggests the formation a common intermediate, which reacts with a reaction rate near the diffusion limit, seemingly negating the stereochemical information from the glycosyl donors.


Glycosylations with functionalized carbamates

With a well-established glycosylation procedure in hand, it was investigated whether this glycosylation strategy could be employed to make biologically relevant model-compounds derived from carbohydrates. It was found that the general

**Scheme 3** Glycosylations employing 2-deoxy-glycosyl TCA donors **4 α** and **4 β** .**Scheme 4** Synthesis of carbamates **16** and **17**.

conditions for synthesizing the carbamate acceptors (Scheme 2) were also suitable for the synthesis of two amino acid- or carbohydrate-derived carbamates, **16** and **17** from cheap, commercially available starting materials (Scheme 4).

Glycosyl acceptors **16** and **17** were found to perform convincingly in glycosylations, following similar conditions as previously described for the glycosylations in Table 1 and Table 2, underlining that even in these more complex systems, there was still no need for further additives. The glycosylations with TCA donor **1 α** (Scheme 5) proceeded with complete β -stereoselectivity and in high yields, whereas the glycosylation starting from **1 β** was reacting slower and resulted a mixture of anomers.

**Scheme 5** Glycosylations of carbamates **18** and **19** with donors **1 α** and **1 β** .


Orthogonal deprotections of *N*-glycosides

With access to the fully protected or selectively functionalized *N*-glycosides, the orthogonality of the protective groups was evaluated. Initially, the deprotection of the tosyl derivatives was investigated (Scheme 6). Using Zemplén conditions on **20** (K₂CO₃ in MeOH) under reflux resulted in the deprotected *N*-glycoside as a 1:1 mixture of anomers. The same results and ratio was achieved when using an anomeric mixture of **20**. The *in situ* anomerization was confirmed by applying the same conditions on the pure β -anomer of the desired glycosyl sulfonamide, which again gave the same anomeric ratio. The anomeric ratio was eventually 1:0.6 after five days of heating (see SI).




The orthogonal *N*-deprotections were then studied using different carbamates, which are more commonly used as protective groups. The Troc protective group in **21** was deprotected using Zn in AcOH at 70°C giving the glycosyl sulfonamide in 82% yield albeit still with some anomerization (α/β 28:72). The Boc protective group in **22** was deprotected under acidic conditions, TFA in CH₂Cl₂, and gave the glycosyl sulfonamide in 84% as a mixture of anomers. The problems with anomerization under acidic or basic conditions could be overcome by using the Alloc protective group (**23**), as it was deprotected under milder conditions using palladium catalysis in combination with a mild acid or nucleophile.²⁹ The removal of the Alloc protective group was performed on a variety of substrates, with different stereochemistry, and both the Ts or Ns group. No anomerization was observed under these conditions and the glycosyl sulfonamides could be isolated in high yields (89 and 87% respectively, Schemes 6 and 7). With the protocols for carbamate deprotections in place, the attention was turned to the desulfonation to give the corresponding *N*-glycosyl carbamates.

The Ts-group has a somewhat bad reputation as an *N*-protective group and the first attempts to remove it were indeed not very successful. As an example, deprotection using lithium naphthalide only gave 22% of the desired product. Instead, employing the protocol by Nyasse et al.⁴³ turned out to be the solution (Scheme 6). Magnesium turnings were used as the reductant in anhydrous MeOH at rt and under these mild conditions, the desired carbamate-protected *N*-glycosides could be obtained in high yields. The major byproduct was found to be the glycosyl sulfonamides, i.e. carbamate deprotection, which increased if water was present. The reductive removal of the Ts-group worked well on the various *N*-glycosides with different carbamate protective groups. Removal of the Ns-group (Scheme 7) was more straightforward using thiophenolate in DMF, employing either K₂CO₃ or KOH as the base. This reaction was generally fast and clean giving isolated yields in the range of 58–96% depending on the conditions. Caution should however be taken not to leave the reaction at prolonged reaction times, as anomerization eventually takes place.



Scheme 6 Orthogonal deprotection of tosyl-protected *N*-glycosides. Reaction conditions: 1) K₂CO₃ (2 equiv.), 0.02m in MeOH, reflux, 4 h. 2) Zn (8 equiv.), 0.12m in AcOH, 70 °C, 2 h. 3) TFA (20 equiv.), 0.03m in CH₂Cl₂, rt, 8 h. 4) PhSiH₃ (4 equiv.), Pd(PPh₃)₄ (0.05 equiv.), 0.75m in CH₂Cl₂, rt, overnight.




Scheme 7 Orthogonal Deprotection of Nosyl-Protected *N*-Glycosides. Reaction conditions 1) Sodium *p*-toluenesulfonate (1.1 equiv.), Pd(PPh₃)₄ (0.1 equiv.), 0.012M in MeOH/THF 1:2, rt, 1.5 h. 2) Zn (20 equiv.), 0.025M in AcOH, rt, 0.5 h. 3) TFA (20 equiv.), 0.4m in CH₂Cl₂, rt, o.n. ^aResulted in reduction of the Ns nitro group to the corresponding amine. ^bGlycosylation and deprotection carried out in a single step, 70% overall yield. ^cK₂CO₃ used as base. ^dKOH used as base.

The carbamate-deprotection of nosyl-protected *N*-glycosides **25–28** (Scheme 7) proved to be somewhat more challenging compared to the tosyl-derivatives, since the Ns-group is generally more sensitive towards various reaction conditions. Thus it was reduced to the amine with Zn in AcOH when trying to remove the Troc group in **26**. The deprotection of the Alloc- and Boc-groups (**27** and **28** respectively) was however efficient, resulting in yields of 70–77%. Thus it seemed that for both tosyl- and nosyl-protected *N*-glycosides, the ideal combination with both was the Alloc protective group.

Scale-up and *N*-Functionalization


With optimized deprotection-procedures in hand, a significant upscaling of the reaction was carried out (Scheme 8). It was found that *N*-deprotected glycosides **29** and **30** could be obtained in 82% and 76% yields accordingly in a glycosylation/deprotection procedure following optimized reaction conditions from Scheme 6. Furthermore, the two products, **29** and **30**, are crystalline which simplifies purification and isolation on a large scale, yielding two ideal starting materials for further *N*-functionalization.

Worth mentioning is the formation of a minor byproduct during the Alloc-deprotection to give **29**, namely allyl *p*-tolyl sulfone,⁴⁴ via a sulfonate-sulfone rearrangement that has previously been shown to take place under similar conditions.^{45,46}



Scheme 8 Gram-scale synthesis of 1-amino-1-deoxy-glycosides.



Scheme 9 Synthesis of glycopeptide model **32**.

With the deprotected *N*-glycoside **29** in hand, it was investigated whether this would be a suitable starting material for synthesis of glycopeptides. Several methods were investigated, but it was found that the NH of **29** could undergo a DIC coupling with a commercially available aspartic acid building block **31**, facilitating the synthesis of **32** in an 82% yield with no anomerization taking place. This underlines the efficiency of the self-promoted glycosylations, since relatively complex compounds such as **32** can be achieved in very few simple and high-yielding steps.

Mechanistic studies

Reaction kinetics

Having established a solid understanding of the scope of the reaction, the reaction mechanism was scrutinized via NMR-experiments and kinetic studies. Since the glycosyl acceptors all have different pK_a -values ranging from 3.5–6.0, a series of glycosylations were followed by $^1\text{H-NMR}$ to clarify whether the pK_a -values corresponded to the observed rate of reaction. All reactions follow a first-order kinetic profile in the glycosyl donors (See SI) and the relative rate constants are summarized in Table 3.

From Table 3 it is clear that there is no clear correlation between the acidity of the glycosyl acceptors and the relative rate of glycosylation, which underlines that steric parameters may be similarly important. It is however noteworthy that all the nosyl-derivatives **14–15** react considerably faster than the tosyl-derivatives **7–11**.

Table 3 Comparison of relative rate constants and pK_a -values of glycosyl acceptors

Acceptor	pK_a	k_{rel}^a	R^{2b}
7	3.9	0.019	0.99
9	5.0	0.014	0.99
10	3.8	0.53	0.99
11	4.8	0.15	0.95
13	4.2	0.0053	1.00
14	3.7	1.00	1.00
15	3.5	$\sim 25^c$	$-^c$

^aRelative rate. k_{rel} for consumption of **1a** under reaction with acceptor **14** set to 1.00. Determined from $^1\text{H-NMR}$ in CD_2Cl_2 at 300 K using mesitylene as internal standard. Donor concentration was 0.032 mM and 1.5 equiv. of acceptor was used in all cases. ^bLinear fit to $\ln[\mathbf{1a}]$ vs. time. ^cStarting material was already consumed when two $^1\text{H-NMR}$ spectrums had been acquired.

To determine the overall reaction order, a series of glycosylation reactions with varying concentrations were carried out and followed by $^1\text{H-NMR}$ -spectroscopy. Acceptor **10** was chosen as the model compound for this experiment as it expressed mid-range reactivity according to Table 3. When doubling the concentration of either **1a** or **10**, the estimated initial rate of product formation was approximately doubled, hence suggesting a second order reaction mechanism. Graph S1 and S2 (see SI) clearly demonstrates the dependence of initial rates on changing concentrations of **1a** and **10**.

Table 4 Overview of estimated initial rates at various concentrations.

$c_{(1a)}(\text{mM})$	$c_{(10)}(\text{mM})$	$v_{(initial)}(\text{mM hr}^{-1})^a$
29	58	15
58	58	25
117	58	58
234	58	157
58	29	15
58	117	55
58	234	162


^aDetermined from $^1\text{H-NMR}$ in CD_2Cl_2 at 300 K using mesitylene as internal standard.

NMR-studies

Next, the reaction mechanism was investigated by $^1\text{H-NMR}$. It was decided to use carbamate **7** as a model acceptor since this particular acceptor reacted somewhat slowly, thus being suitable for the $^1\text{H-NMR}$ time scale. Furthermore, the progress of the reaction could also easily be determined due to the $-\text{OCH}_3$ group, which is distinguished from other signals in the crude spectra. The glycosylation of glycosyl donors **1a** and **1b** were followed by $^1\text{H-NMR}$ at room temperature to identify possible intermediates or by-products formed during the reaction. The glycosylation of **1a** proceeded as expected (Scheme 10), yielding only the desired product **20b**. However, during the glycosylation of **1b**, it was found that the β -TCA donor partly anomerized *in situ* to the corresponding α -TCA donor as a competing reaction, which was then glycosylated, yielding the β -*N*-glycoside (Figure



2A and 2B). This *in situ* anomerization is unprecedented in organic solvents and has only previously been reported by Poletti when performing glycosylations in ionic liquids.^{47,48} Thus, it seems that the β -TCA donor reacts via two competing reaction pathways: Either it reacts in a glycosylation, apparently yielding primarily the α -N-glycoside, but the competing reaction yields the α -TCA donor that is then converted exclusively into the β -N-glycoside, thus giving rise to anomeric mixtures of **20 α** and **20 β** .



Scheme 10 Observed difference in the glycosylation behavior of glycosyl donors **1 α** and **1 β** . Conditions: T=300K 0.075m in dry, neutralized CDCl₃ (passed through basic Al₂O₃) using 3 equiv. of acceptor **7**.





Figure 2 A) Graphical illustration of the anomerization of **1 β** to **1 α** under the reaction conditions shown in scheme 10. B) Graphical illustration of the formation of either **20 α** or **20 β** during the anomerization of **1 β** . Mesitylene was used as an internal standard for concentration determination.

The mechanistic details of the glycosylation reaction have been discussed thoroughly for more than half a century.^{49,50,51} Recently, the interest in mechanistic studies in glycosylation chemistry received renewed interest with the appearance of new methods for selective glycosylations, suggesting S_N2-like mechanisms.^{5,52} Also, the conformation of the oxocarbenium ion has been found to influence the stereochemical outcome of glycosylations.⁵³


From the results of the mechanistic investigation, it is clear that the glycosylation of the α - and the β -TCA donors do not necessarily pass through the same intermediate and hence can follow different reaction paths. In many cases, the reaction gave inversion of stereochemistry, which was most pronounced using the axial TCA in 1,2-*cis* glycosyl donors. On the other hand, the

equatorial TCA donors gave lower selectivity, which could be because of certain steric requirements for a donor-acceptor complex. Therefore, it was investigated whether a complex between donor **1 α** and acceptor **7** was initially formed during the glycosylations (Scheme 11). However, no change in the ¹H-NMR spectrum of either **1 α** or **7** was observed when titrating the donor with acceptor (0.2-1.0 equiv., See SI), ruling out the formation of a stable donor-acceptor complex.

The kinetics are of first order for both glycosyl donor and acceptor. The selectivity observed, with inversion of stereochemistry, suggests an S_N2-like reaction, which could be on the protonated acetimidate, i.e. involving associated ion-pair and therefore an S_Ni-type of mechanism as recently suggested by Tanaka et al. for a glycosylation reaction.⁵⁴ When the solvent polarity is increased, e.g. by using acetonitrile (Scheme 12), **33** is formed. This indicates the formation of a glycosyl cation, and competition from the solvent when the ion-pair is more dissociated. It should be emphasized that the reaction mechanism can change and these conditions cannot be directly transferred to reactions in CH₂Cl₂.



Scheme 11 Proposed complex favoring an S_N2-like reaction mechanism.



Scheme 12 Trapping of glycosyl cation by using acetonitrile as the solvent.

Two simplified mechanistic scenarios for both the α - and β -anomer are shown in Scheme 13. Two different conformers (**A** and **D**) have been proposed for the α -TCA donors of which **A** has previously been found to be energetically favorable.³ Upon activation of **A**, the associated ion pair, **B**, will be formed. In this ion pair, the amide ion is in close proximity to the glycosyl cation intermediate, and should thus react to form **C** much more readily than its counterpart **E** in which the nucleophilic attack is sterically hindered by the C-2 substituent, blocking the required trajectory leading to **F**.


Also, the back reaction to form the TCA donor **D** could be a competing reaction (orange dotted arrow). This can explain why such a high preference for the formation of 1,2-*trans*-glycosides was found for both the *gluco* and *galacto* α -TCA donors. The *manno* donors (not shown) also yielded 1,2-*trans*-glycosides, as the associated ion pair **B** would be obscured by the axial C-2



substituent, instead making the dissociated ion pair **E** the more favorable intermediate leading to **F**.

For the β -TCA donors (Scheme 13, right side), two different conformers (**G** and **J**) are also presented. Here, however, the associated ion pair **H** will not be as favorable as its counterpart **B** since the C-2 substituent of **H** will block the trajectory of the incoming nucleophile, impeding the formation of **I**. On the other hand, the dissociated ion pair, **K**, should have less hindered trajectory for the nucleophilic attack, leading to **L**, but since the nucleophile will be further away from the glycosyl cation, and possibly hydrogen-bonded to the leaving group, the anomerization to the corresponding α -TCA donor (as observed

by $^1\text{H-NMR}$) will be a competing reaction. Thus, it seems that both reaction pathways leading to products **I** and **L** will have certain drawbacks, possibly explaining the lowered stereoselectivity observed during glycosylations with β -TCA donors. The lowered stereoselectivity is also a natural consequence of the observed formation of the α -TCA as this will result in a loss of stereochemical information from the glycosyl donor.



Scheme 13 Proposed intermediates during an S_Ni -like glycosylation with both α - and β -TCA glycosyl donors.

Conclusions

In conclusion, we have developed a self-promoted N -glycosylation using TCA donors and easily obtainable N -sulfonyl carbamates. It has been shown that tosylcarbamates, easily obtained from amino acids or carbohydrates, are capable of reacting in this self-promoted glycosylation, which could be of great interest in the glycosylation of e.g. peptide side chains or other biomolecules. The obtained N -glycosides are stable and hence could be a useful linker for biomolecules in general as they can be used for classic coupling reactions with carboxylic acids, giving glycosyl amides in high yields.


Besides the very mild reaction conditions, it was found that the reaction often was stereospecific, especially when substituting an axial TCA, even in the absence of neighboring group participation. In addition, this glycosylation method allows for the orthogonal deprotection of the N -glycosides,

enabling easily accessible 1-amino-1-deoxy glycosides for further functionalization.

The selectivity and reaction mechanism has been investigated on basis and NMR-experiments and an S_Ni -like mechanism has been proposed.

The self-promoted glycosylation was shown to allow acceptors with a wide range of pK_a -values (3.5-6.0) and very different steric bulk. It was also shown that the procedure was easily scaled up to a column-free procedure on multiple grams without observing a drop in selectivity and yield, which underlines that this glycosylation protocol represents a way of simplifying N -glycosylations, whilst still achieving high yield and selectivity.

It is our belief that this method could prove very valuable in the synthesis of various biomolecules and a further investigation into the scope of this method for the functionalization of proteins and other relevant biomolecules is currently on-going in our laboratory.



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Christian Tortzen for his help with NMR-experiments. Professor Richard R. Schmidt is acknowledged for fruitful discussions. PM acknowledge grant no. POWR.03.02.00-00-1026/16 co-financed by the European Union through the European Social Fund under the Operational Program Knowledge Education Development, for support. We also thank the anonymous reviewers for their comments and valuable input for the discussion of the mechanistic studies.

Notes and references

- Nielsen, M. M.; Pedersen, C. M. *Chem. Rev.* **2018**, *118*, 8285–8358.
- Schmidt, R. R.; Michel, J. *Angew. Chemie Int. Ed. English* **1980**, *19*, 731–732.
- Schmidt, R. R.; Gaden, H.; Jatzke, H. *Tetrahedron Lett.* **1990**, *31*, 327–329.
- Urban, F. J.; Moore, B. S.; Breitenbach, R. *Tetrahedron Lett.* **1990**, *31*, 4421–4424.
- Adero, P. O.; Amarasekara, H.; Wen, P.; Bohé, L.; Crich, D. *Chem. Rev.* **2018**, *118*, 8242–8284.
- Schmidt, R. R.; Michel, J. *Tetrahedron Lett.* **1984**, *25*, 821–824.
- Schmidt, R. R. *Angew. Chemie Int. Ed. English* **1986**, *25*, 212–235.
- Schmidt, R. R.; Michel, J.; Roos, M. *Liebigs Ann. der Chemie* **1984**, *1984*, 1343–1357.
- Peng, P.; Schmidt, R. R. *Acc. Chem. Res.* **2017**, *50*, 1171–1183.
- Kumar, A.; Kumar, V.; Dere, R. T.; Schmidt, R. R. *Org. Lett.* **2011**, *13*, 3612–3615.
- Kumar, A.; Geng, Y.; Schmidt, R. R. *Adv. Synth. Catal.* **2012**, *354*, 1489–1499.
- Nielsen, M. M.; Pedersen, C. M. *Chem. Rev.* **2018**, *118*, 8285–8358.
- Schmidt, R. R.; Stumpp, M.; Michel, J. *Tetrahedron Lett.* **1982**, *23*, 405–408.
- Gould, N. D.; Liana Allen, C.; Nam, B. C.; Schepartz, A.; Miller, S. J. *Carbohydr. Res.* **2013**, *382*, 36–42.
- Frihed, T. G.; Bols, M.; Pedersen, C. M. *Chem. Rev.* **2015**, *115*, 4963–5013.
- Colinas, P. A.; Bravo, R. D. *Carbohydr. Res.* **2007**, *342*, 2297–2302.
- Rodríguez, O. M.; Maresca, A.; Témpora, C. A.; Bravo, R. D.; Colinas, P. A.; Supuran, C. T. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 4447–4450.
- Lavecchia, M. J.; Diez, R. P.; Colinas, P. A. *Carbohydr. Res.* **2011**, *346*, 442–448.
- Ombouma, J.; Vullo, D.; Supuran, C. T.; Winum, J.-Y. *Bioorg. Med. Chem.* **2014**, *22*, 6353–6359.
- Winum, J.-Y.; Colinas, P. A.; Supuran, C. T. *Bioorg. Med. Chem.* **2013**, *21*, 1419–1426. DOI: 10.1039/C9SC00857H
- Crespo, R.; de Bravo, M. G.; Colinas, P. A.; Bravo, R. D. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6469–6471.
- Xavier, N. M.; Porcheron, A.; Batista, D.; Jorda, R.; Řezníčková, E.; Kryštof, V.; Oliveira, M. C. *Org. Biomol. Chem.* **2017**, *15*, 4667–4680.
- Marks, G. S.; Neuberger, A. J. *Chem. Soc.* **1961**, 4872–4879.
- Garg, H. G.; Jeanloz, R. W. *Adv. Carbohydr. Chem. Biochem.* **1985**, *43*, 135–201.
- Garg, H. G.; Von Dem Bruch, K.; Kunz, H. *Adv. Carbohydr. Chem. Biochem.* **1994**, *50*, 277–310.
- Srivastava, A.; Varghese, B.; Loganathan, D. *Chem. – Eur. J.* **2013**, *19*, 17720–17732.
- Franco, L. L.; Brandão, M. C.; Filho, J. D. S.; Alves, R. J. *Química Nova*. scielo 2015, pp 1044–1052.
- Likhoshesterov, L. M.; Novikova, O. S.; Derevitskaja, V. A.; Kochetkov, N. K. *Carbohydr. Res.* **1986**, *146*, C1–C5.
- Nicolaou, K. C.; Snyder, S. A.; Nalbandian, A. Z.; Longbottom, D. A. *J. Am. Chem. Soc.* **2004**, *126*, 6234–6235.
- Colinas, P. A.; Bravo, R. D. *Org. Lett.* **2003**, *5*, 4509–4511.
- Nardi, M.; Cano, N. H.; De Nino, A.; Di Gioia, M. L.; Maiuolo, L.; Oliverio, M.; Santiago, A.; Sorrentino, D.; Procopio, A. *Tetrahedron Lett.* **2017**, *58*, 1721–1726.
- Ding, F.; William, R.; Liu, X.-W. *J. Org. Chem.* **2013**, *78*, 1293–1299.
- Ombouma, J.; Vullo, D.; Dumy, P.; Supuran, C. T.; Winum, J.-Y. *ACS Med. Chem. Lett.* **2015**, *6*, 819–821.
- Dai, Y.; Zheng, J.; Zhang, Q. *Org. Lett.* **2018**, *20*, 3923–3927.
- Battina, S. K.; Kashyap, S. J. *Carbohydr. Chem.* **2015**, *34*, 133–144.
- Bradshaw, G. A.; Colgan, A. C.; Allen, N. P.; Pongener, I.; Boland, M. B.; Ortin, Y.; McGarrigle, E. M. *Chem. Sci.* **2019**, *10*, 508–514.
- Suthagar, K.; Polson, M. I. J.; Fairbanks, A. J. *Org. Biomol. Chem.* **2015**, *13*, 6573–6579.
- Colinas, P. A.; Núñez, N. A.; Bravo, R. D. *J. Carbohydr. Chem.* **2008**, *27*, 141–147.
- Griswold, K. S.; Horstmann, T. E.; Miller, S. J. *Synlett* **2003**, *2003*, 1923–1926.
- Krag, J.; Christiansen, M. S.; Petersen, J. G.; Jensen, H. H. *Carbohydr. Res.* **2010**, *345*, 872–879.
- Marqvorsen, M. H. S.; Pedersen, M. J.; Rasmussen, M. R.; Kristensen, S. K.; Dahl-Lassen, R.; Jensen, H. H. *J. Org. Chem.* **2017**, *82*, 143–156.
- Heuckendorff, M.; Jensen, H. H. *Carbohydr. Res.* **2017**, *439*, 50–56.
- Nyasse, B.; Grehn, L.; Ragnarsson, U. *Chem. Commun.* **1997**, No. 11, 1017–1018.
- Chang, M.-Y.; Wu, M.-H.; Chen, Y.-L. *Org. Lett.* **2013**, *15*, 2822–2825.
- Hiroi, K.; Kitayama, R.; Sato, S. *J. Chem. Soc. Chem. Commun.* **1984**, No. 5, 303–305.
- Hiroi, K.; Makino, K. *Chem. Lett.* **1986**, *15*, 617–620.




ARTICLE

Journal Name

- (47) Rencurosi, A.; Lay, L.; Russo, G.; Caneva, E.; Poletti, L. *J. Org. Chem.* **2005**, *70*, 7765–7768.
- (48) Rencurosi, A.; Lay, L.; Russo, G.; Caneva, E.; Poletti, L. *Carbohydr. Res.* **2006**, *341*, 903–908.
- (49) BOCHKOV, A. F.; ZAIKOV, G. E. BOCHKOV, A. F., ZAIKOV, G. E. B. T.-C. of the O. B., Eds.; Pergamon, 1979; pp 1–4.
- (50) BOCHKOV, A. F.; ZAIKOV, G. E. BOCHKOV, A. F., ZAIKOV, G. E. B. T.-C. of the O. B., Eds.; Pergamon, 1979; pp 5–79.
- (51) Capon, B. *Chem. Rev.* **1969**, *69*, 407–498.
- (52) Huang, M.; Garrett, G. E.; Birlirakis, N.; Bohé, L.; Pratt, D. A.; Crich, D. *Nat. Chem.* **2012**, *4*, 663.
- (53) Walvoort, M. T. C.; Dinkelaar, J.; van den Bos, L. J.; Lodder, G.; Overkleef, H. S.; Codée, J. D. C.; van der Marel, G. A. *Carbohydr. Res.* **2010**, *345*, 1252–1263.
- (54) Tanaka, M.; Nakagawa, A.; Nishi, N.; Iijima, K.; Sawa, R.; Takahashi, D.; Toshima, K. *J. Am. Chem. Soc.* **2018**, *140*, 3644–3651.

View Article Online
DOI: 10.1039/C9SC00857H





164x65mm (600 x 600 DPI)

